

QUARTERLY INTERNATIONAL JOURNAL

ECOLOGY

ENVIRONMENT & CONSERVATION

ISSN 0971-765 X

Vol 26
No. (3)
2020



EM INTERNATIONAL

ECOLOGY, ENVIRONMENT AND CONSERVATION

VOL. 26 (3) : 2020

CONTENTS

- 957–962 Short-term experimenton effects of sulfuric acid on atranorin concentration and upper surface structure in *Pyxine cocoes* (Sw.) Nyl.
—*Chitsupang Kheawsalab, Wanaruk Saipunkaew, Somporn Chantara, Pitchaya Mungkornasawakul and Nattakarn Sransupphasirigul*
- 963–968 Effect of skip irrigation and nano potassium treatments on maize yield
—*Ali H. Jasim, Sura R. Husain and Hanaa H, Mohammed*
- 969–973 Computation model of electrostatic spraying in Agriculture Industry
—*Murtadha Al-Mamury, Balachandran, Wamadeva, Al-Raweshidy, Hamed and Manivannan, Nadarajah*
- 974–979 Study of the dynamics of natural stands of *Pinus halepensis* in the Beni Oudjana forest (Khenchela, Algeria)
—*Insaf Hani, Malika Rached-Kanouni, Hichem Khamar and Menasri Ammar*
- 980–986 Assessment of vulnerability of climate change on crop and livestock production, adaption in Vietnam
—*Pham Thi To Oanh and Tran Ngoc Thin*
- 987–994 The determination of radon gas concenentration distributed North of Baghdad Governorate: spatial analysis by applying remote Sensing (RS) and geographic information system (GIS) techniques
—*Fouad K. Mashee Al Ramahi, Muaid Jassim Rasheed and Mazin Shakir Jasim*
- 995–999 Single modified fractionation technique of lead in sediments of Shatt Al-Hilla-Iraq
—*Hazim Aziz AL-Robai and Russel Jabbar Al-Jbouri*
- 1000–1004 Contribution to the study of the adaptation of *Casuarina equisetifolia* to heat stress using biochemical markers
—*Malika Rached-Kanouni, Zahia Kadi, Hichem Khammar, Rima Bouzobra and Fadhila Belghit*
- 1005–1011 Effect of Nano fertilizer and Gibberellic acid application in vegetative and floral growth of *Gazania rigens* L.
—*Raghad Adnan AL-Qadi, Mateen Yilmaz I. AL-Bayati, Ala, aHussien AL-Bakkar and Ali Mohammad Ahmed*
- 1012–1017 Developing collective awareness in providing local-based preservation of coastal area
—*Fenty Puluwulawa and Amanda Adelina Harun*
- 1018–1026 Effect of spraying organic fertilizer, untrigrin and alkaren algae extract on vegetative floral and chemical growth of two types of tulip plants
—*Alaa Hashim Younis Altaee and Kefaia Ghazi saeed*
- 1027–1035 Community perception and participation in mangrove ecotourism development in Lembar area west Lombok regency
—*Sukuryadi, Nuddin Harahab, Mimit Primyastanto and Bambang Semedi*
- 1036–1042 The resistance of tomato plants from seed treated with a magnetic field of 0.2 m T against *Fusarium* sp.
—*Rochmah Agustrina, Endang Nurcahyani, Bambang Irawan, Eko Pramono, Ika Listiani, EkoNastiti and Sutopo Hadi*
- 1043–1048 The assessment of the efficiency of environmental activities in Moscow
—*Konstantin Zakharov*
- 1049–1054 Characteristics of growing sites *Anisoptera thurifera* in South Manokwari West Papua
—*Rima H.S. Siburian, Anna Tampang, Agustunus Murdjoko and Doni Djitmau*

-
- 1055–1058 First record of threatened Asian Catfish, *Clarias batrachus* (Linnaeus, 1758) (Siluriformes, Clariidae) from Kangean Island, Indonesia
—*Aditya Mirzapahlevi Saptadjaja, Veryl Hasan, Muhammad Arief, Wahyu Isoni and Rozi*
- 1059–1067 Study the effect of Carvacrol, Eugenol and Thymol on *Fusariums* sp responsible for *Loliumperenne* fusariosis
—*Hamza Saghrouchni, Azeddin El Barnossi, Hanane Chefchaou, Aouatef Mzabi, Mariam Tanghort, Adnane Remmaland and Chami Fouzia*
- 1068–1077 Flood risk management of Urban Territories
—*Elena Sierikova, Elena Strelnikova, Leonid Pishnia and Elena Pozdnyakova*
- 1078–1084 Understanding identity of a tourist region through Travelsketch event
—*Augustinus Madyana Putra, Gagoek Hardiman and Agung Budi Sardjono*
- 1085–1089 Environmental management: A study on the precautionary principle in siak regency of Indonesia towards sustainable development
—*Sri Maulidiah, Monalisa, Zaini Ali, Sharifah Zarina Syed Zakaria, Nuriah Abd. Majid, Kadir Arifin, Zuliskandar Ramli, Emrizal and Muhammad Rizal Razman*
- 1090–1097 Value chain analysis of Bee Jay mangrove ecotourism in Probolinggo
—*Candra Adi Intyas, Mochammad Fattah and Tiwi Nurjannati Utami*
- 1098–1099 Effect of the Bacteria *Bacillus thuringiensis* var kurstaky on second larval instar of Angoumois moth *Sitotroga cerealella*
—*Ahmed M. Tarek*
- 1100–1109 The implementation of conservation village model program in mount Rinjani National Park (A Regulation Perspective)
—*Andi Chairil Ichsan, Irwan Mahakam Lesmono Aji, Hairil Anwar, Tenri Waru and Indra Gumay Febryano*
- 1110–1114 Prospects for the use of fallow lands in the TVER region for sowing long-stalked flax
—*Alexey Alekseevich Akimov, Natalya Nikolaevna Ivanyutina, Alexander Sergeevich Vasiliev, Ilya Alexandrovich Drozdov and Yuri Teodorovich Farinyuk*
- 1115–1118 Presence of the invasive Nile Tilapia *Oreochromis niloticus* Linnaeus, 1758 (Perciformes, Cichlidae) in the Yamdena Island, Indonesia
—*Liga Insani, Veryl Hasan, Fitri Sil Valen, Fajar Surya Pratama, Maheno Sri Widodo, Abdul Rahem Faqih, R. Adharyan Islamy, Akhmad Taufiq Mukti and Wahyu Isoni*
- 1119–1123 Heavy metal (Mercury and Plumbum) accumulation of two fish species in Sipin and Teluk Lake, Jambi Province
—*Siswanta Kaban, M. Edi Armanto, M. Rasyid Ridho and Poedji L. Hariani*
- 1124–1127 Realization of the biological potential of Potato varieties in the agricultural conditions of the upper Volga region
—*Z.I. Usanova, M.N. Pavlov, N.S. Chernikova and S.E. Pryadein*
- 1128–1138 Analysis of Islamic performance index on Sharia business unit in Indonesia towards sustainable development
—*Hamdi Agustin, Sri Indrastuti, Amris Rusli Tanjung, Syahdanur, Pipin Kurnia, Sharifah Zarina Syed Zakaria, Nuriah Abd. Majid, Kadir Arifin, Zuliskandar Ramli, Emrizal and Muhammad Rizal Razman*
- 1139–1144 Improvement of regional cooperation in overcoming the problem of water resources in the decentralization Era in East Java Province, Indonesia
—*Lutfi Agus Salim*
- 1145–1157 Actinobacteria isolated from Algerian hot spring waters: A potential source of important enzymes
—*Meissa Medjemadj, Juan-José Escuder-Rodríguez, Allaoueddine Boudemagh and María-Isabel González-Siso*

- 1158–1161 A review on aquatic macrophytes as bio indicators of Water Quality of Lakes
—*Kalpita Bhatta and Hemant Kumar Patra*
- 1162–1165 Study and first report on the occurrence of six species of coccoidal cyanobacteria - *Merismopedia meyen* 1839 from Bardhaman East and Bardhaman West Districts of West Bengal, India
—*Sudip Chatterjee*
- 1166–1169 The number of main species of bark beetles (Scolytinae) and protective measures in the mountain forests of the Zaili Alatau
—*Nurzhan Mukhamadiyev, J. Bolat, A. Koigeldina and Gulnaz Mengdibayeva*
- 1170–1174 Macroinvertebrate diversity of effluent affected Dhansiri and Kaliani Rivers of Assam, India
—*Janmoni Moran*
- 1175–1183 Understanding climate change in terms of rainfall fluctuations and status of agricultural productivity in Northeastern States of India
—*Niranjan Roy, Avijit Debnath and Sunil Nautiyal*
- 1184–1188 Effect of plastic mulches and irrigation levels on yield parameters of tomato (*Solanum lycopersicum*) in Madurai District of Tamil Nadu
—*M. Jayalalitha, M. Rajeswari, P. Saravanapandian and R. Lalitha*
- 1189–1192 The population of Bark Beetles (Scolytinae) and their entomophages on endemic spruce in the Zaili Alatau
—*Nurzhan Mukhamadiyev, J. Bolat, A. Koigeldina and Gulnaz Mengdibayeva*
- 1193–1199 Exogeneous application of *Moringa oleifera* leaf extract and Salicylic acid on modulation of Salinity Stress effects in Pea (*Pisum sativum*) plants
—*Vishal Sharma, Sarita and Alka*
- 1200–1203 Water potability assessment of Boranakanive Reservoir in Chikkanayakanahalli Taluk of Tumkur District, Karnataka, India
—*R. Shruthi and S. Ramakrishna*
- 1204–1207 Integrated system of protection of soybeans from insect pests in the fields of “Ontustik Agropark” LLP
—*A.O. Sagitov, N.S. Mukhamadiyev, N.Zh. Sultanova, A.S. Dinassilov, M.À. Uzakbayeva and G.Zh. Mengdibayeva*
- 1208–1211 Multivariate analysis for genetic diversity estimation among tomato (*Solanum lycopersicum* L.) Genotypes
—*Ankita Debnath, Rajesh Kumar, Sita Kumari Prasad, Nisha Sharma and Jitendra Kumar Kushwah*
- 1212–1217 Effect of Arsenic contaminated ground water on the developmental phenomena of primary-school children in Assam
—*Ananta Kumar Jena, Munmi Barman, Somnath Gupta, Jaishree Devi, Satarupa Bhattacharjee and Sabina Yeasmin Barbhuiya*
- 1218–1221 Biological activity and emission of carbon dioxide from dark chestnut soil of Western Kazakhstan
—*Nurlan Sergaliyev, Aliya Nagiyeva and Anvar Tlepov*
- 1222–1227 Density and population growth of the Black sea cucumber, *Holothuria (Halodeima) atra* (Jaeger, 1833) in two different microhabitats of Jepara Waters, Indonesia
—*Retno Hartati, Muhammad Zainuri, Ambariyanto Ambariyanto, Ita Riniatsih, Hadi Endrawati, Sri Redjeki, Ria AT Nuraini, Agoes Soegianto and Sekar Widyaningsih*
- 1228–1234 Floristic diversity of lacustrine ecotone of Lakes in Valsad District, Gujarat, India
—*Ayantika Das and Jigna Desai*

- 1235–1238 The effect of spring wheat cultivation methods on some fertility indicators of Southern carbonate chernozem in Northern Kazakhstan
—*Nadezhda Filippova, Galina Churkina, Maral Utebayev, Irina Rukavitsina and Kairat Kunanbayev*
- 1239–1243 Noise pollution and human health: A case study in Bangkok City, Thailand
—*N. Vattanaprateep*
- 1244–1248 Surabaya strategy to Answer air pollution improvement
—*Intan Ayu Pratiwi and Rusdiana Puspa Ayu*
- 1249–1258 Local ecological knowledge of invasive Alien species *Opuntia ficus-indica* in three local municipalities in the Eastern Cape Province, South Africa
—*Luleka Mdweshu and Alfred Maroyi*
- 1259–1270 Plant bacterial endophytes as a potent source of plant growth promoters and other bioactive compounds: A Review
—*K.K. Dayamrita and Nivya Mariam Paul*
- 1271–1275 Monitoring of population density of *Vibrio* sp. and health condition of hepatopancreas pacific white shrimp (*Litopenaeus vannamei*) cultivated with intensive systems in Bulukumba regency, South Sulawesi, Indonesia
—*Mayadita Dwi Sani, Agnes Yuantin Maharani, Muhammad Inas Riandy, Raden Joko Kuncoroningrat Susilo, Putu Angga Wiradana and Agoes Soegianto*
- 1276–1281 People within Park: forest cover dynamics and management strategies in Dibru-Saikhowa National Park, India
—*Rekib Ahmed*
- 1282–1289 Spatial relationship modeling for urban environmental factors analysis to population density change
—*Nutchanat Buasri, Patiwat Littidej, Naruset Prasertsri and Benjamabhorn Pumhirunroj*
- 1290–1299 Adoption of united nations sustainable development goals 2030 in environmental management system by earth moving and construction equipment industries in India
—*Dheeraj Verma, Vartika Singh, Prodyut Bhattacharya and Jagdish Kishwan*
- 1300–1304 Development of hydroxyapatite from corals obtained from contamination waters of Northern Java by precipitation method
—*Siswanto, Siti Nurmala, Dyah Hikmawati, Aminatun and Retna Apsari*
- 1305–1308 Physico-chemical characteristics of Groundwater in Byramangala and Harobele command areas of Arkavathi sub-basin of River Cauvery, Karnataka
—*Suma Rani S.N. and B.C. Nagaraja*
- 1309–1312 Studying the biological features of development and the technology of breeding the *Orius laevigatus* Fieb Predatory Bug
—*Aizhan Mukashevna Chadinova, Abay Orazuly Sagitov, Karlygash Azirbekovna Alpysbayeva and Balzhan Zhasulanovna Naimanova*
- 1313–1318 Larvicidal and Pupicidal activities from *Citrus hystrix* against *Aedes aegypti* Mosquitoes
—*Niken Subekti, Asni Puraedah, Dyah Rini Indriyanti and Agoes Soegianto*
- 1319–1323 Comparison of remediation performance of *Chaetoceros calcitrans* on heavy metals and diesel fuel exposure
—*Diana Arfiati, Nunik Cokrowati, Dwi Candra Pratiwi, Niken Pratiwi, Qurrota A'yunin, Trisnadi W. C. Putranto and Thin Soedarti*
- 1324–1326 Effect of microbial inoculants on soil quality, growth and yield of pea plant
—*Simmi Goel and Yasmeen Bano*
- 1327–1330 Biosurfactants production using bacterial isolate from Cangar hot spring, East Java, Indonesia
—*Almando Geraldi, Aulia Azzahra, Lillah Asritafriha, Rizki Amaliah Zain, Us Watun Nurul Khasanah, Nastiti Trikurniadewi and Ni'matuzahroh*

-
- 1331–1341 Wetland degradation and its impact on life and livelihood of people in the Majuli River Island, Assam
—*Ranjan Das and Jhimli Bhattacharjee*
- 1342–1347 Greenhouse Gas (GHG) Emission Reduction Model in Surabaya
—*Intan Ayu Pratiwi, Eka Prasetyo and Helmi Dadang Ardiansyah*
- 1348–1351 Difference of Hg (II) removal efficiency in leachate by Immobilized *Skeletonema* sp.
—*Isna Arifatus Shalihah, Eko Prasetyo Kuncoro, Tini Surtiningsih, Nur Indradewi Oktavitri and Thin Soedarti*
- 1352–1361 Classification of air pollutants caused by e-waste and health risk evaluation
—*Sajeev Ram, C.S. Shylaja and R. Anandan*
- 1362–1367 Application of probiotics and microalgae (*Chaetoceros calcitrans*) to stimulate non-specific Immune responses in white Shrimp (*Litopenaeus vannamei*) infected with *Vibrio harveyi*
—*Mila Ayu Ambarsari and Woro Hastuti Satyantini*
- 1368–1371 *Pseudomonas* community in white syndrome diseases of *Echinopora lamellosa* coral at nature reserve Pulau Sempu, Indonesia
—*Oktiyas Muzaky Luthfi, Muliawati Handayani, Andik Isdianto, Muhammad Arif Asadi, Tajuddin Atho'illah and Moch Affandi*
- 1372–1382 Effect of locally made subsoiler tines on some energy requirements
—*Ghazwan Ahmed Dahham, Othman Muayad Mohammed Taofeeq and Layth Mahmood Yahya*
- 1383–1388 Copepod growth populations (*Acartia* sp.) in outdoor mass culture tanks: Exploring natural feed potentials for sustainable aquaculture
—*Putu Angga Wiradana, Sephia Anjani, Deny Suhernawan Yusup, Joko Wiryatno, Regina Melianawati, Sin War Naw, Aondohemba Samuel Nege and Agoes Soegianto*
- 1389–1395 Modelling risks using Quantum physics principles for health projects: Mangaung Metropolitan Municipality, South Africa
—*Bernard Moeketsi Hlalele*
- 1396–1400 Productivity of corn hybrids in the North-eastern upper volga region, Russia
—*Z.I. Usanova, P.I. Migulev, Yu.T. Farinyuk and M.N. Pavlov*
- 1401–1404 Realization of the biological potential of potato varieties in the agricultural conditions of the upper Volga Region
—*Z.I. Usanova, M.N. Pavlov, N.S. Chernikova and S.E. Pryadein*
- 1405–1419 Economic and social analysis of the proposed programme of measures in order to achieve good environmental status in Turkish Seas: The Case of the EU Marine strategy framework directive
—*Ömer Eyuboglu*
- 1420–1423 Development and evaluation of bacterial consortia for biodegradation of polycyclic aromatic hydrocarbons
—*E.R. Faizulina, S.A. Aitkeldiyeva, L.G. Tatarkina, K. Ashimuly, A.V. Alimbetova and A.K. Sadanov*
- 1424–1431 Oil palm empty fruit bunch as the selected organic matter in developing the Swampy forest system for passive treatment of acid mine drainage
—*Ihsan Noor, Yudi Firmanul Arifin, Bambang Joko Priatmadi and Akhmad Rizalli Saidy*

Application of probiotics and microalgae (*Chaetoceros calcitrans*) to stimulate non-specific Immune responses in white Shrimp (*Litopenaeus vannamei*) infected with *Vibrio harveyi*

Mila Ayu Ambarsari¹ and Woro Hastuti Satyantini^{2*}

¹Post Graduate Program of Marine Biotechnology, Faculty of Fisheries and Marine, Universitas Airlangga, Surabaya 60115, Indonesia

²Department of Fish Health Management and Aquaculture Faculty of Fisheries and Marine, Universitas Airlangga, Surabaya 60115, Indonesia

(Received 10 January, 2019; Accepted 28 February, 2020)

ABSTRACT

This study aims to determine whether the administration of probiotics and microalgae *Chaetoceros calcitrans* can improve the non-specific immune response of white shrimp after *Vibrio harveyi* bacterial infection. This research is experimental with a completely randomized design (CRD) consisting of 5 treatments and 4 replications. -C: Rearing shrimp without the administration of probiotics or microalgae *Chaetoceros calcitrans*, the 15th day was injected with PBS (*Phosphate Buffer Saline*). +C: Rearing shrimp without the provision of probiotics or microalgae *Chaetoceros calcitrans*, and on the 15th day infected with *Vibrio harveyi*. PA: Shrimp maintenance by providing Probiotics (Pro). PB: Shrimp maintenance by giving Microalgae (Mic). PC: Shrimp maintenance by providing Probiotics and Microalgae (Promic) for 14 days, and on 15th day infected with *V. harveyi*. The parameters observed included *Total Haemocyte Count* (THC), *Phagocytic activity*. Data analysis used Analysis of Variance (ANOVA) followed by Duncan's multiple range test (DMRT). The results of this study indicate that the administration of probiotics and microalgae gave significantly different results ($p < 0.05$) on non-specific immune responses in white shrimp. The best research results on the treatment of probiotics and microalgae *Chaetoceros calcitrans* are able to increase the non-specific immune response in white shrimp after *Vibrio harveyi* bacterial infection THC value of 42.86×10^6 cells/mL, phagocytic activity (PA) activity by 95.50%, SR 94.75%. Probiotics (Pro), Microalgae (Mic) *Chaetoceros calcitrans* and Combination Probiotics and Microalgae (Promic) on the maintenance media can increase non-specific immune response and survival rate in white shrimp after *V. harveyi* bacterial infection.

Key words: Probiotics, *Bacillus subtilis*, *Bacillus mycoides*, *Pseudomonas diminuta*, *Litopenaeus vannamei*

Introduction

White Shrimp (*Litopenaeus vannamei*) was firstly imported into Indonesia in 2000 to replace the Tiger shrimp (*Penaeus monodon*) which is widely affected by diseases, especially bacteria and viruses

(Marwiyah *et al.*, 2019; Kharisma *et al.*, 2020). But in the course of this effort, losses from disease attacks continued. This Vibriosis disease can cause death in shrimp 90 – 100% of the total population, one of which is by *Vibrio harveyi* bacteria. A number of methods have been applied in the effort to control

diseases such as control of the immune system through the administration of probiotics (Jefri *et al.*, 2020; Rangka and Gunarto, 2012; Sriwulan *et al.*, 2019; Latifah *et al.*, 2019) and microalgae *Chaetoceros calcitrans* that can be a substitute antibiotic to reduce pathogenic bacteria. Probiotics have a cell wall structure of lipopolysaccharides (LPS) and have antioxidant activity (Trisnawati *et al.*, 2018; Rusmarilin *et al.*, 2018; Silalahi *et al.*, 2018), whereas microalgae *Chaetoceros calcitrans* contain a content of β -1-3 glucan. Both materials can be used to enhance non-specific immune responses. β -1-3 glucan and LPS may increase PO activity after β -1-3 glucan and LPS react with β -glucan binding protein (BGBP) or LPS binding proteins. After the bind, proPO will be re-activated into a PO enzyme that further performs its function in the process of melanization, while in microalgae that have a content of β -1-3 glucan can increase cell activating factors in hemocytes, so that it can increase the activity of PO and phagocytosis in shrimp (Smith *et al.*, 2014). The objectives of this study were to investigate the effects of probiotics and microalgae (*Chaetoceros calcitrans*) on the stimulation of non-specific immune responses in white shrimp (*Litopenaeus vannamei*) infected with *Vibrio harveyi*.

Materials and Methods

Probiotics used (*Bacillus subtilis*, *Bacillus mycoides*, *Pseudomonas diminuta*), Microalgae *Chaetoceros calcitrans*, *Vibrio harveyi* bacterial isolate. The size of aquarium used in this experimentation was $60 \times 30 \times 35 \text{ cm}^3$. Whiteshrimp (*Litopenaeus vannamei*) were in healthy condition and free from *Vibrio harveyi* infection with body length 6-8cm, and weights 6-6.5g.

Research Design

This research was conducted using a Completely Randomized Design consisting of 5 treatments and 4 replications. The dosage of *Vibrio harveyi* used in this experiment was 10^6 CFU/mL Cahayati (2012), and the dosage of *Chaetoceros calcitrans* microalgae was as much as 67.50×10^5 cells/mL (Yeh *et al.*, 2015).

The treatments used in this study were

Negative control (-C): Shrimp rearing without the provision of probiotics or microalgae *Chaetoceros calcitrans*, then on the day 15 was injected with PBS (Phosphate Buffer Saline).

- Positive control (+C): Maintenance of shrimp without the provision of probiotics and microalgae *C. calcitrans*, and on the day 15 infected with *Vibrio harveyi*.
- PA: Shrimp maintenance by providing Probiotics (Pro), for 14 days, and on the 15th day infected with *V. harveyi*.
- PB: Shrimp maintenance by giving Microalgae (Mic), for 14 days, and on the 15th day infected with *V. harveyi*.
- PC: Shrimp maintenance by providing Probiotics and Microalgae (Promic), for 14 days, and on the 15th day infected with *V. harveyi*.

Observation of Research Parameters

Observation parameters of white shrimp immune response were consisted of Total Hemocyte Count (THC), Phagocytic activity (PA), Survival Rate (SR).

Hemolymph collection

Hemolymph for measurement of immune parameters was taken from all shrimps of each treatment unit. About 0.1 mL of hemolymph was taken from the ventral sinus at the base of the 5th leg using a 1mL syringe after 0.1 mL of anticoagulant was added (30 mM trisodium citrate, 0.34 M sodium chloride, 10 mM EDTA, pH 7.5) (Mannopo, 2014).

Total Haemocyte Count (THC)

After hemolymph collection, approximately of 100 μ L of hemolymph was taken and dropped into the hemocytometer to calculate the Total Haemocyte Count (THC) under a light microscope with 400x magnification. Measurement of total haemocyte was carried out four times: at day 0 (at the beginning of the experiment, H0), at day 14 (H-14), after bacterial infection of *Vibrio harveyi* (at day 16, H-16) and at day 25 (H-25). THCs were expressed as number of cells/ml of hemolymph.

Phagocytic activity (PA)

Phagocytic activity was determined as described below. Retrieval of fresh shrimp hemolymph (20 μ L), inserted into the microtube and added with 20 μ L suspension of *Staphylococcus aureus* with density of 10^8 cells/mL, incubated at room temperature for 30 minutes. Furthermore, 5 μ L was taken to make a smear on the preparation object glass and let it dry. The preparations were then soaked with methanol 95% for 5 minutes and rinsed with 0.85% NaCl and

dried again. Then painted using 10% Giemsa for 15 minutes and air dried. The curing preparation was then watered for about 5 minutes to remove the remaining color of Giemsa. The preparations were observed under a microscope at 400x magnification. PA values was calculated using the following formula:

$$AF = a/b \times 100\%$$

Where:

PA = Phagocytic Activity (%)

a = Number of cells undergoing phagocytosis (cell/mL)

b = Total number of cells observed (cell/mL)

Data analysis

Measurement data were expressed as mean \pm SD. THC dan PA data then analyzed using One-way ANOVA by using statistical software SPSS-21. If significant difference were observed, the Duncan test then used to analyze which treatment caused significantly different to immune responses of white shrimp $p < 0.05$.

Water quality

Water quality parameters (temperature, dissolved oxygen, pH, Salinity, Ammonium, Nitrite, Ammonia, and Nitrate) were measured during the study are presented in. Table 3. The water quality was within the normal range based on standards for white shrimp culture.

Results and Discussion

Survival Rate

The survival rates of infected shrimp by *Vibrio harveyi* after provision of probiotic (Pro), microalgae (Mic) and probiotic+microalgae (Promic) are presented in Figure 1. Both at day 16 and day 25, the provision of probiotic+microalgae (Promic) treatments had a significantly different on survival rate of shrimp com-

pared to negative control (-C), positive control(+C) and microalgae (Mic) treatment. There were not significantly different between Pro and Promic treatment. Both at H-16 and H-25, the lowest survival rates were noted at +C.

Total Haemocyte Count (THC)

The total number of hemocyte count of *Litopenaeus vannamei* at H0, H-14, H-16 and H-25 are presented in Table 1. Haemocyte play an important role in the immune system of crustacea. At H0 the THC values ranged from 4.21×10^6 cell/ml to 4.47×10^6 cell/ml. All treatments were not significantly different at H0. Furthermore, the THC values of Pro and Mic treatments at H-14 were higher than those of -C, +C and Promic treatments.




Fig. 1. Application of probiotics and microalgae on survival rate of *Litopenaeus vannamei* infected by *Vibrio harveyi*

One day after *Vibrio harveyi* infection (at H-16) the THC values of Pro and Promic were not significantly different. The lowest value of THC was noted at -C. Furthermore, 10 days after *Vibrio harveyi* infection (at H-25) the THC values of Pro, Mic and Promic were not significantly different, and their levels were higher than -C and +C.

Phagocytic activity

Phagocytosis is the process of digesting particulate matter, especially bacteria, into the cytoplasm of white blood cells (Gullian *et al.*, 2014). The phago-

Table 1. Average Total Haemocyte Count of White Shrimp during the study

Treatment	Average value of haemocytes on day to- ($\times 10^6$) cell/mL			
	H-0	H-14	H-16	H-25
-C	4.21 ^a \pm 0.492	4.71 ^c \pm 0.512	9.16 ^d \pm 3.568	21.40 ^b \pm 3.290
+C	4.28 ^a \pm 0.543	6.16 ^b \pm 0.191	11.63 ^{cd} \pm 2.391	13.75 ^b \pm 1.946
Probiotics	4.30 ^a \pm 0.486	7.96 ^a \pm 1.346	18.64 ^a \pm 2.122	34.41 ^a \pm 8.372
Microalgae	4.27 ^a \pm 0.505	7.50 ^a \pm 0.493	13.81 ^{bc} \pm 2.121	34.50 ^a \pm 4.328
Probiotics and Microalgae	4.47 ^a \pm 0.408	4.54 ^c \pm 0.327	17.66 ^{ab} \pm 2.673	42.86 ^a \pm 9.677

Table 2. Average values of *phagocytic activity* (PA) during the study

Treatment	The Average value of PA (%) on observation			
	H-0	H-14	H-16	H-25
-C	61.25 ^a ± 4.788	67.25 ^c ± 3.594	77.50 ^{ab} ± 2.082	84.50 ^c ± 1.000
+C	60.75 ^a ± 5.620	68.25 ^c ± 4.646	74.75 ^c ± 2.500	81.00 ^d ± 1.155
Probiotics	62.50 ^a ± 2.082	78.50 ^{ab} ± 2.380	83.75 ^a ± 1.500	90.25 ^b ± 1.258
Microalgae	62.25 ^a ± 1.708	73.75 ^b ± 0.500	79.75 ^b ± 0.500	88.00 ^b ± 2.160
Probiotics and Microalgae	63.00 ^a ± 2.449	80.50 ^a ± 4.203	85.00 ^a ± 3.559	95.50 ^a ± 1.732

Table 3. The Range of Water Quality Value of Media Maintenance of Shrimp

Variable	Treatment					The Range of Normal values
	K ⁻	K ⁺	Probiotics	Microalgae	Probiotics and Microalgae	
pH	7.7	7.6	7.7	7.8	7.7	7.5 – 8.5
Salinity (ppt)	22	21	21.5	20	21	16 – 30
DO (mg/L)	6.21	5.48	6.01	6.11	6.03	>3 – 7
Temperature (°C)	28	27	29.1	26.5	28.7	27 – 32
Ammonium	0.25	0.3	0.18	0.17	0.17	0.1 – 5
Nitrit	0.17	0.2	0.15	0.15	0.1	0.1 – 1
Nitrat	0.4	0.5	0.5	0.6	0.6	>0.2 – 0.8
Ammonia	0.018	0.021	0.013	0.012	0.012	< 1

cytic activity values of all treatments starting from H-0, H-14, H-16 and H-25 are presented in Table 2. At H-0, all phagocytic activity of all treatments were not significantly different. At H-14, the highest phagocytic activity was recorded at Promic, and the lowest were observed at -C and +C. At H-16 there were not significantly different between -C, Pro and Promic, the lowest phagocytic activity was note at +C. At H-25 the highest phagocytic activity was noted at Promic and the lowest was observed at +C.

Discussion

The level of survival is an opportunity for an individual’s life in a certain time. At the end of the study (H-25) the Promic treatment received ansurvival rate (SR) value of around 94.75%, the Pro treatment 94.25%, the Mic treatment 90.75%, the -C treatment 92.37%, and lowest was noted in +C treatments around 82.25%. The high observations of SR values in the Promictreatments compared to other treatments especially to Pro and Mic allegedly due to a balanced work between probiotics and microalgae.

Maximum probiotic work can function as a biological control agent that produces antibacterial molecules such as bacteriocin directly capable of inhibiting other bacteria such as *Vibrio harveyi* and actively participating in fighting infections.

Probiotics are also able to inhibit the movement of other bacteria in the intestinal wall (translocation), so they can improve the function of the mucosal barrier by increasing the production of non-specific immune responses or modulating inflammation (Cerezuela *et al.*, 2011). In addition, the microalgae *Chaetoceros calcitrans* has the ability to inhibit the growth of gram-negative bacteria, by increasing the permeability of bacterial membranes (Zheng *et al.*, 2015).

Shrimp blood does not contain haemoglobin, so the blood is not red (Person *et al.*, 2014). In shrimp the function is replaced by haemocyanin, a protein that contains Cu and can bind with oxygen. Haemocyanin functions as oxygen transport, as a buffer in shrimp blood and plays an important role in blood osmotic (Maynard, 2014). White shrimp has nonspecific immunity (innate) which can recognize and destroy foreign objects that enter the body, so that the nonspecific immune system plays an important role in the shrimp immune system. Haemocytes play an important role in the crustacean immune system. Haemocytes play a role in phagocytosis, encapsulation, degranulation and aggregation of pathogens or foreign particles (Sahoo *et al.*, 2014).

This haemocyte value can be measured and can be used as a health assessment of shrimp through the activity of the defense system against infectious

agents. This total haemocyte is very important in resistance to pathogens. From the results of data during the research on the observation of THC H0 ranged from 4.21×10^6 cells/mL to 4.47×10^6 cells/mL. The H-14 before *Vibrio harveyi* infection, in Promic treatment at 4.54×10^6 sel/mL showed the lowest results compared to Pro treatment 7.96×10^6 sel/mL, Mic treatment at 7.50×10^6 cells/mL and positive control 6.16×10^6 cells/mL. This is suspected that the treatment of Promic given during maintenance influence the work of competitors in the utilization of nitrogen nutrients by probiotic bacteria and microalgae, so that the number of probiotics entering the digestive tract becomes less than in the Pro treatment. At H-16 (1 day after *Vibrio harveyi* infection) where the THC value of Promic increased compared to H-14. This is because as a form of shrimp immune system reaction in response to foreign objects that enter is the pathogenic bacterium *Vibrio harveyi*. However, after 10 days after the occurrence of *Vibrio harveyi* infection (H-25) THC value of Promic, Pro and Mic treatments were not significantly different, but they were significantly different compared to positive control treatments. This is because the mechanism of action of probiotics can also provide resistance to the body against *Vibrio harveyi* attacks. Where the *Pseudomonas diminuta* bacterium contains *Lipopolysaccharide*, LGBP (*Lipopolysaccharide β -glucan Binding Protein*) will induce haemocyte degranulocytes and stimulate the activation of the ProPO system into PO, hence the opsonin factor protein will produce an increase in phagocytosis. Whereas *Chaetoceros calcitrans* Microalgae also contains β -1,3 *Glucan* from carbohydrate group which through the introduction of proteins activated by PPA, and PPA itself can be activated by *lipopolysaccharides* from microorganisms through recognition will form β -1,3 *Glucan Binding Protein* found in granulocytes so as to increase haemocytes in white shrimp. The results of the study reported by Sahoo *et al.* (2014) also stated that white shrimp treated with Probiotics and Microalgae (Promic) had higher THC values compared to other treatments. The high THC value in white shrimp is due to the high mobilization of hemolymph in the body of the shrimp so that it can increase immunity and the introduction of foreign objects that enter the shrimp body. While the low THC value greatly affects the susceptibility of shrimp to pathogens (Le Moullac *et al.* 2014).

This study showed that increased THC values in the treatment of Promic also followed by an increase

in the value of phagocytic activity (PA). Determination of the value of PA is to determine the increase in shrimp endurance because in the phagocytosis process is a non-specific defense mechanism that is generally able to protect against disease (Fontaine and Lightner, 2013). The PA values of H-25, the Promic treatments showed the highest PA values of 95.50% compared to the Pro treatment 90.25% and the Mic treatment at 88%. In the Promic treatment, the effect was significantly different compared to controls. So that the treatment of Promic can improve shrimp immune response by increasing phagocytic cells to carry out the process of phagocytosis, namely the ability to phagocyte in destroying the *Vibrio harveyi* attack. The higher of phagocytic activity value, the immune system of white shrimp is suspected to be getting better, this is because in probiotics there is an antibacterial and antimicrobial content that can inhibit the formation of nucleic acids which causes the target cell membrane to be disrupted so that *Vibrio harveyi* lysis. Whereas the *Pseudomonas* bacteria are able to produce extracellular enzymes in the form of bacteriocin, siderophore and antibiotics which can inhibit pathogenic bacteria. Furthermore Itami and Takeuchi, (2013) support the theory which states that the administration of immunostimulants can prevent disease infections in the host body and cause increased phagocyte activity of haemocytes and proPO enzymes. From the data obtained during the H-25 maintenance period that white shrimp treated with Promic can provide significant survival rate values on the immune response that can be seen from the data there is a linear increase in THC, and PA values and quality water is one of the factors that greatly influences the life of shrimp, so in this study water quality parameters become one of the considerations of the results obtained. The water quality of the maintenance media during the study was in the ideal range for the maintenance of white shrimp. So that with good water quality parameters, and an increased immune response can provide shrimp growth and good survival value, resulting in an increase in production of white shrimp culture.

Conclusion

Based on the results of the research that has been done, the following conclusions can be drawn: In the combination treatment gives a very real effect on the non-specific immune response in white shrimp. Which results in an increase in the number of

haemocytes by 42.86×10^6 cells/mL, *phagocytic activity* (PA). To obtain an effective protocol for administering *Chaetoceros calcitrans* microalgae, further research is needed to observe.

Acknowledgments

Special thank due to professor Agoes Soegianto for constructive comments to improve this manuscript.

References

- Cerezuela. 2011. Immune Response of *Litopenaeus vannamei* after Infection with *Vibrio harveyi*. *Aquaculture*. 406-407: 115-120.
- Fontaine and Lightner, 2013. A Review of Some Major Disease Significant on Penaeid Prawn Shrimp of The American and Indopacific. Diseases in Asia Aquaculture. Fish and Health Section Asian Fisheries Society, Manila, Philippines. 57-62 p.
- Fontaine and Lightner, 2014. *In vivo* antiviral activity of Potential implication in shrimp disease management. *World Journal of Fish and Marine Sciences*. 1 (4): 278-282.
- Itami and Takeuchi, T. 2013. Body defence system of penaeid. *Seminar Avertebrata Physiology and Prevention partemen of Aquaculture and Biology*. Shimonoseki University of Fisheries, Japan. 7 : 59-65.
- Jefri, M., Satyantini, W. H., Sahidu, A. M., Nindarwi, D. D. and Rozi, 2020. Application of Probiotics for Organic Matter and Enhancement of Growth Performance in White Shrimp (*Litopenaeus vannamei*). *Jurnal Ilmiah Perikanan and Kelautan*. 12 (1): 97-104. doi.org/10.20473/jipk.v12i1.16618.
- Kharisma, A., Tjahjaningsih, W. W. and Sigit, S. 2020. Determination of minimum inhibitory and minimum bactericidal concentration of ketapang (*Terminatiacatappa*) leaves extract against *Vibrio harveyi*. *IOP Conf. Series: Earth and Environmental Science*. 441 (2020) 012012. doi:10.1088/1755-1315/441/1/012012
- Latifah, L.A., Soekamto, N.H. and Tahir, A. 2019. Preliminary study: *Padina australis* Hauck's antibacterial activity and phytochemical test against pathogenic shrimp bacteria. *Journal of Physics: Conference Series*. 1341 (2019) 022005. doi:10.1088/1742-6596/1341/2/022005.
- Le Moullac, G., Soyez, C., Saulnier, D., Ansquer, D., Avarre J.C. and Levy, P. 2014. Effect of hypoxic stress on the immune response and the resistance to vibriosis of the shrimp *Penaeus stylirostris*. *Fish Shellfish and Immunology*. 8 : 621-629.
- Manoppo, Hengky. 2011. Perannukleotidase sebagai immunostimulan terhadap respon imun nonspesifik dan resistensi udang white (*Litopenaeus vannamei*). Disertasi Pascasarjana Institut Pertanian Bogor.
- Marwiyah, U.C., Mahasri, G., Ratnasari, R.E. and Wiradana, P.A. 2019. Total plate count and identification of *vibrio* in pacific white shrimp (*Litopenaeus vannamei*) from ponds and in those exposed to immunogenic protein membrane *Zoothamnium penaei*. *IOP Conf. Series: Earth and Environmental Science*. 236 (2019) 012087. doi:10.1088/1755-1315/236/1/012087.
- Maynard, 2014. *Immunostimulants*. Japan Sci Soc. Press. Tokyo, p 41-56.
- Person, R. G., Gaxiola, G. and Taboada, M. Pascual. 2014. Effect of immunostimulants on hemolymph of *Litopenaeus vannamei*. *Aquaculture Research*. 38 : 1339-1345.
- Rangka, N.A. and Gunarto, 2012. The effect of bio floc growing on vannamei shrimp culture at intensive system pond. *Jurnal Ilmiah Perikanan and Kelautan*. 4 (2): 141-149.
- Rusmarilin, H., Nurhasanah and Andayani, R. Y. 2018. Soy-yangurt probiotic drink as a natural potential of antioxidant. *IOP Conf. Series: Earth and Environmental Science*. 122 (2018) 012087 doi :10.1088/1755-1315/122/1/012087.
- Sahoo, B., Sethi, S., Mishra, B.K. and Das, B.K. 2014. Effects elicitors on Prophenoloxidase and Superoxidase anion activities of *Litopenaeus vannamei*. *Asian Fish. Sci*. 18 : 345-353.
- Silalahi, J., Nadarason, D. and Silalahi, Y.C.E. 2018. The effect of storage condition on antioxidant activity of probiotics in yogurt drinks. *Asian J Pharm Clin Res*. 11 (12) : 280-283.
- Smith, V.J., Brown, J.H., Scoderhall, K.A. and Hauton, C. 2014. Physiological, Nutritional, and Immunological Role of Dietary β -glucan and Ascorbic Acid Monophosphate in *Litopenaeus vannamei* juveniles. *Fish and Shellfish Immunology*. 15(1): 223-243.
- Sriwulan, Azwar, A., Rantetondok, A. and Anshary, H. 2019. Screening and application of lactic acid bacteria isolated from vanamei shrimp (*Litopenaeus vannamei*) intestine as a probiotic potential for tiger shrimp (*Penaeus monodon*). *AACL Bioflux*. 12 (5): 1866-1881.
- Trisnawita, Y., Silalahi, J. and Sinaga, S.M. 2018. The effect of storage condition on viability of lactic acid bacteria in probiotic product. *Asian J Pharm Clin Res*. 11 (Special issue 1): 84-86.
- Yeh, S.P., Hsieh, S.L., Liu, C.H. and Cheng W. 2015. Immune response of shrimp, *Litopenaeus vannamei* after a concurrent infection with white spot syndrome virus and hepatopietic necrosis virus. *Fish and Shellfish Immunology*. 26 : 582-588.
- Zheng, Y. H., Yoshida, T., Isobe, K., Rahman, M. J., Nagase, F., Ding, L. and Nakashima, I. 2015. Modulation by Glycyrrhizin of the cell-surface expression of H-2 Class I Antigens on Marine Tumor Cell Lines and Normal Cell Populations. *Immunology*. 70 : 405-410.