

4

by Mila Ayu Ambarsari

Submission date: 29-Dec-2020 04:27PM (UTC+0800)

Submission ID: 1481838471

File name: 4._Application_of_probiotics_and_microalgae.pdf (187.99K)

Word count: 3920

Character count: 20605

Application of probiotics and microalgae (*Chaetoceros calcitrans*) to stimulate non-specific Immune responses in white Shrimp (*Litopenaeus vannamei*) infected with *Vibrio harveyi*

Mila Ayu Ambarsari¹ and Woro Hastuti Satyantini^{2*}

¹Post Graduate Program of Marine Biotechnology, Faculty of Fisheries and Marine, Universitas Airlangga, Surabaya 60115, Indonesia

²Department of Fish Health Management and Aquaculture Faculty of Fisheries and Marine, Universitas Airlangga, Surabaya 60115, Indonesia

(Received 10 January, 2019; Accepted 28 February, 2020)

ABSTRACT

This study aims to determine whether the administration of probiotics and microalgae *Chaetoceros calcitrans* can improve the non-specific immune response of white shrimp after *Vibrio harveyi* bacterial infection. This research is experimental with a completely randomized design (CRD) consisting of 5 treatments and 4 replications. -C: Rearing shrimp without the administration of probiotics or microalgae *Chaetoceros calcitrans*, the 15th day was injected with PBS (Phosphate Buffer Saline). +C: Rearing shrimp without the provision of probiotics or microalgae *Chaetoceros calcitrans*, and on the 15th day infected with *Vibrio harveyi*. PA: Shrimp maintenance by providing Probiotics (Pro). PB: Shrimp maintenance by giving Microalgae (Mic). PC: Shrimp maintenance by providing Probiotics and Microalgae (Promic) for 14 days, and on 15th day infected with *V. harveyi*. The parameters observed included Total Haemocyte Count (THC), Phagocytic activity. Data analysis used Analysis of Variance (ANOVA) followed by Duncan's multiple range test (DMRT). The results of this study indicate that the administration of probiotics and microalgae gave significantly different results ($p < 0.05$) on non-specific immune responses in white shrimp. The best research results on the treatment of probiotics and microalgae *Chaetoceros calcitrans* are able to increase the non-specific immune response in white shrimp after *Vibrio harveyi* bacterial infection THC value of 42.86×10^6 cells/mL, phagocytic activity (PA) activity by 95.50%, SR 94.75%. Probiotics (Pro), Microalgae (Mic) *Chaetoceros calcitrans* and Combination Probiotics and Microalgae (Promic) on the maintenance media can increase non-specific immune response and survival rate in white shrimp after *V. harveyi* bacterial infection.

Key words: Probiotics, *Bacillus subtilis*, *Bacillus mycoides*, *Pseudomonas diminuta*, *Litopenaeus vannamei*

Introduction

White Shrimp (*Litopenaeus vannamei*) was firstly imported into Indonesia in 2000 to replace the Tiger shrimp (*Penaeus monodon*) which is widely affected by diseases, especially bacteria and viruses

(Marwiyah *et al.*, 2019; Kharisma *et al.*, 2020). But in the course of this effort, losses from disease attacks continued. This Vibriosis disease can cause death in shrimp 90 – 100% of the total population, one of which is by *Vibrio harveyi* bacteria. A number of methods have been applied in the effort to control

diseases such as control of the immune system through the administration of probiotics (Jefri *et al.*, 2020; Rangka and Gunarto, 2012; Sriwulan *et al.*, 2019; Latifah *et al.*, 2019) and microalgae *Chaetoceros calcitrans* that can be a substitute antibiotic to reduce pathogenic bacteria. Probiotics have a cell wall structure of lipopolysaccharides (LPS) and have antioxidant activity (Trisnawati *et al.*, 2018; Rusmarilin *et al.*, 2018; Silalahi *et al.*, 2018), whereas microalgae *Chaetoceros calcitrans* contain a content of β -1-3 glucan. Both materials can be used to enhance non-specific immune responses. β -1-3 glucan and LPS may increase PO activity after β -1-3 glucan and LPS react with β -glucan binding protein (BGBP) or LPS binding proteins. After the bind, proPO will be re-activated into a PO enzyme that further performs its function in the process of melanization, while in microalgae that have a content of β -1-3 glucan can increase cell activating factors in hemocytes, so that it can increase the activity of PO and phagocytosis in shrimp (Smith *et al.*, 2014). The objectives of this study were to investigate the effects of probiotics and microalgae (*Chaetoceros calcitrans*) on the stimulation of non-specific immune responses in white shrimp (*Litopenaeus vannamei*) infected with *Vibrio harveyi*.

Materials and Methods

Probiotics used (*Bacillus subtilis*, *Bacillus mycoides*, *Pseudomonas diminuta*), Microalgae *Chaetoceros calcitrans*, *Vibrio harveyi* bacterial isolate. The size of aquarium used in this experimentation was $60 \times 30 \times 35 \text{ cm}^3$. Whiteshrimp (*Litopenaeus vannamei*) were in healthy condition and free from *Vibrio harveyi* infection with body length 6-8cm, and weights 6-6.5g.

Research Design

This research was conducted using a Completely Randomized Design consisting of 5 treatments and 4 replications. The dosage of *Vibrio harveyi* used in this experiment was 10^6 CFU/mL Cahayati (2012), and the dosage of *Chaetoceros calcitrans* microalgae was as much as 67.50×10^5 cells/mL (Yeh *et al.*, 2015).

The treatments used in this study were

Negative control (-C): Shrimp rearing without the provision of probiotics or microalgae *Chaetoceros calcitrans*, then on the day 15 was injected with PBS (Phosphate Buffer Saline).

- Positive control (+C): Maintenance of shrimp without the provision of probiotics and microalgae *C. calcitrans*, and on the day 15 infected with *Vibrio harveyi*.
- PA: Shrimp maintenance by providing Probiotics (Pro), for 14 days, and on the 15th day infected with *V. harveyi*.
- PB: Shrimp maintenance by giving Microalgae (Mic), for 14 days, and on the 15th day infected with *V. harveyi*.
- PC: Shrimp maintenance by providing Probiotics and Microalgae (Promic), for 14 days, and on the 15th day infected with *V. harveyi*.

Observation of Research Parameters

Observation parameters of white shrimp immune response were consisted of Total Hemocyte Count (THC), Phagocytic activity (PA), Survival Rate (SR).

Hemolymph collection

Hemolymph for measurement of immune parameters was taken from all shrimps of each treatment unit. About 0.1 mL of hemolymph was taken from the ventral sinus at the base of the 5th leg using a 1mL syringe after 0.1 mL of anticoagulant was added (30 mM trisodium citrate, 0.34 M sodium chloride, 10 mM EDTA, pH 7.5) (Mannopo, 2014).

Total Haemocyte Count (THC)

After hemolymph collection, approximately of 100 μL of hemolymph was taken and dropped into the hemocytometer to calculate the Total Haemocyte Count (THC) under a light microscope with 400x magnification. Measurement of total haemocyte was carried out four times: at day 0 (at the beginning of the experiment, H0), at day 14 (H-14), after bacterial infection of *Vibrio harveyi* (at day 16, H-16) and at day 25 (H-25). THCs were expressed as number of cells/mL of hemolymph.

Phagocytic activity (PA)

Phagocytic activity was determined as described below. Retrieval of fresh shrimp hemolymph (20 μL), inserted into the microtube and added with 20 μL suspension of *Staphylococcus aureus* with density of 10^8 cells/mL, incubated at room temperature for 30 minutes. Furthermore, 5 μL was taken to make a smear on the preparation object glass and let it dry. The preparations were then soaked with methanol 95% for 5 minutes and rinsed with 0.85% NaCl and

dried again. Then painted using 10% Giemsa for 15 minutes and dried. The curing preparation was then watered for about 5 minutes to remove the remaining color of Giemsa. The preparations were observed under a microscope at 400x magnification. PA values was calculated using the following formula:

$$AF = a/b \times 100\%$$

Where:

PA = Phagocytic Activity (%)

a = Number of cells undergoing phagocytosis (cell/mL)

b = Total number of cells observed (cell/mL)

Data analysis

Measurement data were expressed as mean \pm SD. THC dan PA data then analyzed using One-way ANOVA by using statistical software SPSS-21. If significant difference were observed, the Duncan test then used to analyze which treatment caused significantly different to immune responses of white shrimp $p < 0.05$.

Water quality

Water quality parameters (temperature, dissolved oxygen, pH, Salinity, Ammonium, Nitrite, Ammonia, and Nitrate) were measured during the study are presented in. Table 3. The water quality was within the normal range based on standards for white shrimp culture.

Results and Discussion

Survival Rate

The survival rates of infected shrimp by *Vibrio harveyi* after provision of probiotic (Pro), microalgae (Mic) and probiotic+microalgae are presented in Figure 1. Both at day 16 and day 25, the provision of probiotic+microalgae (Promic) treatments had a significantly different on survival rate of shrimp com-

pared to negative control (-C), positive control(+C) and microalgae (Mic) treatment. There were not significantly different between Pro and Promic treatment. Both at H-16 and H-25, the lowest survival rates were noted at +C.

Total Haemocyte Count (THC)

The total number of hemocyte count of *Litopenaeus vannamei* at H0, H-14, H-16 and H-25 are presented in Table 1. Haemocyte play an important role in the immune system of crustacea. At H0 the THC values ranged from 4.21×10^6 cell/ml to 4.47×10^6 cell/mL. All treatments were not significantly different at H0. Furthermore, the THC values of Pro and Mic treatments at H-14 were higher than those of -C, +C and Promic treatments.




Fig. 1. Application of probiotics and microalgae on survival rate of *Litopenaeus vannamei* infected by *Vibrio harveyi*

One day after *Vibrio harveyi* infection (at H-16) the THC values of Pro and Promic were not significantly different. The lowest value of THC was noted at -C. Furthermore, 10 days after *Vibrio harveyi* infection (at H-25) the THC values of Pro, Mic and Promic were not significantly different, and their levels were higher than -C and +C.

Phagocytic activity

Phagocytosis is the process of digesting particulate matter, especially bacteria, into the cytoplasm of white blood cells (Gullian *et al.*, 2014). The phago-

Table 1. Average Total Haemocyte Count of White Shrimp during the study

Treatment	Average value of haemocytes on day to- ($\times 10^6$) cell/mL			
	H-0	H-14	H-16	H-25
-C	$4.21^a \pm 0.492$	$4.71^c \pm 0.512$	$9.16^d \pm 3.568$	$21.40^b \pm 3.290$
+C	$4.28^a \pm 0.543$	$6.16^b \pm 0.191$	$11.63^{cd} \pm 2.391$	$13.75^b \pm 1.946$
Probiotics	$4.30^a \pm 0.486$	$7.96^c \pm 1.346$	$18.64^a \pm 2.122$	$34.41^a \pm 8.372$
Microalgae	$4.27^a \pm 0.505$	$7.50^c \pm 0.493$	$13.81^{bc} \pm 2.121$	$34.50^a \pm 4.328$
Probiotics and Microalgae	$4.47^a \pm 0.408$	$4.54^c \pm 0.327$	$17.66^{ab} \pm 2.673$	$42.86^a \pm 9.677$

Table 2. Average values of phagocytic activity (PA) during the study

Treatment	The Average value of PA (%) on observation			
	H-0	H-14	H-16	H-25
-C	61.25 ^a ± 4.788	67.25 ^c ± 3.594	77.50 ^{ab} ± 2.082	84.50 ^c ± 1.000
+C	60.75 ^a ± 5.620	68.25 ^c ± 4.646	74.75 ^c ± 2.500	81.00 ^d ± 1.155
Probiotics	62.50 ^a ± 2.082	78.50 ^{ab} ± 2.380	83.75 ^a ± 1.500	90.25 ^b ± 1.258
Microalgae	62.25 ^a ± 1.708	73.75 ^b ± 0.500	79.75 ^b ± 0.500	88.00 ^b ± 2.160
Probiotics and Microalgae	63.00 ^a ± 2.449	80.50 ^a ± 4.203	85.00 ^a ± 3.559	95.50 ^a ± 1.732

Table 3. The Range of Water Quality Value of Media Maintenance of Shrimp

Variable	Treatment					The Range of Normal values
	K ⁻	K ⁺	Probiotics	Microalgae	Probiotics and Microalgae	
pH	7.7	7.6	7.7	7.8	7.7	7.5 – 8.5
Salinity (ppt)	22	21	21.5	20	21	16 – 30
DO (mg/L)	6.21	5.48	6.01	6.11	6.03	>3 – 7
Temperature (°C)	28	27	29.1	26.5	28.7	27 – 32
Ammonium	0.25	0.3	0.18	0.17	0.17	0.1 – 5
Nitrit	0.17	0.2	0.15	0.15	0.1	0.1 – 1
Nitrat	0.4	0.5	0.5	0.6	0.6	>0.2 – 0.8
Ammonia	0.018	0.021	0.013	0.012	0.012	< 1

cytic activity values of all treatments starting from H-0, H-14, H-16 and H-25 are presented in Table 2. At H-0, all phagocytic activity of all treatments were not significantly different. At H-14, the highest phagocytic activity was recorded at Promic, and the lowest were observed at -C and +C. At H-16 there were not significantly different between -C, Pro and Promic, the lowest phagocytic activity was note at +C. At H-25 the highest phagocytic activity was noted at Promic and the lowest was observed at +C.

Discussion

The level of survival is an opportunity for an individual’s life in a certain time. At the end of the study (H-25) the Promic treatment received ansurvival rate (SR) value of around 94.75%, the Pro treatment 94.25%, the Mic treatment 90.75%, the -C treatment 92.37%, and lowest was noted in +C treatments around 82.25%. The high observations of SR values in the Promictreatments compared to other treatments especially to Pro and Mic allegedly due to a balanced work between probiotics and microalgae.

Maximum probiotic work can function as a biological control agent that produces antibacterial molecules such as bacteriocin directly capable of inhibiting other bacteria such as *Vibrio harveyi* and actively participating in fighting infections.

Probiotics are also able to inhibit the movement of other bacteria in the intestinal wall (translocation), so they can improve the function of the mucosal barrier by increasing the production of non-specific immune responses or modulating inflammation (Cerezuela *et al.*, 2011). In addition, the microalgae *Chaetoceros calcitrans* has the ability to inhibit the growth of gram-negative bacteria, by increasing the permeability of bacterial membranes (Zheng *et al.*, 2015).

Shrimp blood does not contain haemoglobin, so the blood is not red (Person *et al.*, 2014). In shrimp the function is replaced by haemocyanin, a protein that contains Cu and can bind with oxygen. Haemocyanin functions as oxygen transport, as a buffer in shrimp blood and plays an important role in blood osmotic (Maynard, 2014). White shrimp has nonspecific immunity (innate) which can recognize and destroy foreign objects that enter the body, so that the nonspecific immune system plays an important role in the shrimp immune system. Haemocytes play an important role in the crustacean immune system. Haemocytes play a role in phagocytosis, encapsulation, degranulation and aggregation of pathogens or foreign particles (Sahoo *et al.*, 2014).

This haemocyte value can be measured and can be used as a health assessment of shrimp through the activity of the defense system against infectious

agents. This total haemocyte is very important in resistance to pathogens. From the results of data during the research on the observation of THC H0 ranged from 4.21×10^6 cells/mL to 4.47×10^6 cells/mL. The H-14 before *Vibrio harveyi* infection, in Promic treatment at 4.54×10^6 sel/mL showed the lowest results compared to Pro treatment 7.96×10^6 sel/mL, Mic treatment at 7.50×10^6 cells/mL and positive control 6.16×10^6 cells/mL. This is suspected that the treatment of Promic given during maintenance influence the work of competitors in the utilization of nitrogen nutrients by probiotic bacteria and microalgae, so that the number of probiotics entering the digestive tract becomes less than in the Pro treatment. At H-16 (1 day after *Vibrio harveyi* infection) where the THC value of Promic increased compared to H-14. This is because as a form of shrimp immune system reaction in response to foreign objects that enter is the pathogenic bacterium *Vibrio harveyi*. However, after 10 days after the occurrence of *Vibrio harveyi* infection (H-25) THC value of Promic, Pro and Mic treatments were not significantly different, but they were significantly different compared to positive control treatments. This is because the mechanism of action of probiotics can also provide resistance to the body against *Vibrio harveyi* attacks. Where the *Pseudomonas diminuta* bacterium contains Lipopolysaccharide, LGBP (Lipopolysaccharide β -glucan Binding Protein) will **15**uce haemocyte degranulocytes and stimulate **the activation of the ProPO** system into **PO**, hence **the** opsonin factor protein will produce an increase in phagocytosis. Whereas *Chaetoceros calcitrans* Microalgae also contains β -1,3 Glucan from carbohydrate group which through the introduction of proteins activated by PPA, and PPA itself can be activated by lipopolysaccharides from microorganisms through recognition will form β -1,3 Glucan Binding Protein found in granulocytes so as to increase haemocytes in white shrimp. The results of the study reported by Sahoo *et al.* (2014) also stated that white shrimp treated with Probiotics and Microalgae (Promic) had higher THC values compared to other treatments. The high THC value in white shrimp **3** due to the high mobilization of hemolymph in **the body of the shrimp** so that **it** can **increase** immunity and **the** introduction **of** foreign objects that enter the shrimp body. While the low THC value greatly affects the susceptibility of shrimp to pathogens (Le Moullac *et al.* 2014).

This study showed that increased THC values in the treatment of Promic also followed by an increase

in the value of phagocytic activity (PA). Determination of the value of PA is to determine the increase in shrimp **5** endurance because in the phagocytosis process **is a non-specific defense mechanism that is generally** able to protect **against disease** (Fontaine and Lightner, 2013). The PA values of H-25, the Promic treatments showed the highest PA values of 95.50% compared to the Pro treatment 90.25% and the Mic treatment at 88%. In the Promic treatment, the effect was significantly different compared to controls. So that the treatment of Promic can improve shrimp immune response by increasing phagocytic cells to carry out the process of phagocytosis, namely the ability to phagocyte in destroying the *Vibrio harveyi* attack. The higher of phagocytic activity value, the immune system of white shrimp is suspected to be getting better, this is because in probiotics there is an antibacterial and antimicrobial content that can inhibit the formation of nucleic acids which causes the target cell membrane to be disrupted so that *Vibrio harveyi* lysis. Whereas the *Pseudomonas* bacteria are able to produce extracellular enzymes in the form of bacteriocin, siderophore and antibiotics which can inhibit pathogenic bacteria. Furthermore Itami and Takeuchi, (2013) support the theory which states that the administration **5** immunostimulants can prevent disease infections **in the host body and** cause **increased phagocyte** activity **of haemocytes and proPO enzymes**. From the data obtained during the H-25 maintenance period that white shrimp treated with Promic can provide significant survival rate values on the immune response that can be seen from the data there is a linear increase in THC, and PA values and quality water is one of the factors that greatly influences the life of shrimp, so in this study water quality parameters become one of the considerations of the results obtained. The water quality of the maintenance media during the study was in the ideal range for the maintenance of white shrimp. So that with good water quality parameters, and an increased immune response can provide shrimp growth and good survival value, resulting in an increase in production of white shrimp culture.

7 Conclusion

Based on the results of the research that has been done, the following conclusions can be drawn: **13** **the** combination treatment gives a very real effect **on the non-specific immune response in white shrimp**. Which results in an increase in the number of

haemocytes by 42.86×10^6 cells/mL, *phagocytic activity* (PA). To obtain an effective protocol for administering *Chaetoceros calcitrans* microalgae, further research is needed to observe.

Acknowledgments

Special thank due to professor Agoes Soegianto for constructive comments to improve this manuscript.

References

- Cerezuela. 2011. Immune Response of *Litopenaeus vannamei* after Infection with *Vibrio harveyi*. *Aquaculture*. 406-407: 115-120.
- Fontaine and Lightner, 2013. A Review of Some Major Disease Significant on Penaeid Prawn Shrimp of The American and Indopacific. Diseases in Asia Aquaculture. Fish and Health Section Asian Fisheries Society, Manila, Philippines. 57-62 p.
- Fontaine and Lightner, 2014. *In vivo* antiviral activity of Potential implication in shrimp disease management. *World Journal of Fish and Marine Sciences*. 1 (4): 278-282.
- Itami and Takeuchi, T. 2013. Body defence system of penaeid. *Seminar Avvertebrata Physiology and Prevention partemen of Aquaculture and Biology*. Shimonoseki University of Fisheries, Japan. 7 : 59-65.
- Jefri, M., Satyantini, W. H., Sahidu, A. M., Nindarwi, D. D. and Rozi, 2020. Application of Probiotics for Organic Matter and Enhancement of Growth Performance in White Shrimp (*Litopenaeus vannamei*). *Jurnal Ilmiah Perikanan dan Kelautan*. 12 (1): 97-104. doi.org/10.20473/jipk.v12i1.16618.
- Kharisma, A., Tjahjaningsih, W. W. and Sigit, S. 2020. Determination of minimum inhibitory and minimum bactericidal concentration of ketapang (*Terminatiacatappa*) leaves extract against *Vibrio harveyi*. *IOP Conf. Series: Earth and Environmental Science*. 441 (2020) 012012. doi:10.1088/1755-1315/441/1/012012
- Latifah, L.A., Soekamto, N.H. and Tahir, A. 2019. Preliminary study: *Padina australis* Hauck's antibacterial activity and phytochemical test against pathogenic shrimp bacteria. *Journal of Physics: Conference Series*. 1341 (2019) 022005. doi:10.1088/1742-6596/1341/2/022005.
- Le Moullac, G., Soyez, C., Saulnier, D., Ansquer, D., Avarre J.C. and Levy, P. 2014. Effect of hypoxic stress on the immune response and the resistance to vibriosis of the shrimp *Penaeus stylirostris*. *Fish Shellfish and Immunology*. 8 : 621-629.
- Manoppo, Hengky. 2011. Perannukleotidasebagai immunostimulanterhadapresponimunnonspesifik dan resistensidang white (*Litopenaeusvannamei*). Disertasi Pascasarjana Institut Pertanian Bogor.
- Marwiyah, U.C., Mahasri, G., Ratnasari, R.E. and Wiradana, P.A. 2019. Total plate count and identification of *vibrio* in pacific white shrimp (*Litopenaeus vannamei*) from ponds and in those exposed to immunogenic protein membrane *Zoothamniumpenaei*. *IOP Conf. Series: Earth and Environmental Science*. 236 (2019) 012087. doi:10.1088/1755-1315/236/1/012087.
- Maynard, 2014. *Immunostimulants*. Japan Sci Soc. Press. Tokyo, p 41-56.
- Person, R. G., Gaxiola, G. and Taboada, M. Pascual. 2014. Effect of immunostimulants on hemolymph of *Litopenaeus vannamei*. *Aquaculture Research*. 38 : 1339-1345.
- Rangka, N.A. and Gunarto, 2012. The effect of bio floc growing on vannamei shrimp culture at intensive system pond. *Jurnal Ilmiah Perikanan dan Kelautan*. 4 (2): 141-149.
- Rusmarilin, H., Nurhasanah and Andayani, R. Y. 2018. Soy-yangurt probiotic drink as a natural potential of antioxidant. *IOP Conf. Series: Earth and Environmental Science*. 122 (2018) 012087 doi :10.1088/1755-1315/122/1/012087.
- Sahoo, B., Sethi, S., Mishra, B.K. and Das, B.K. 2014. Effects elicitors on Prophenoloxidase and Superoxidase anion activities of *Litopenaeus vannamei*. *Asian Fish. Sci*. 18 : 345-353.
- Silalahi, J., Nadarason, D. and Silalahi, Y.C.E. 2018. The effect of storage condition on antioxidant activity of probiotics in yogurt drinks. *Asian J Pharm Clin Res*. 11 (12) : 280-283.
- Smith, V.J., Brown, J.H., Scoderhall, K.A. and Hauton, C. 2014. Physiological, Nutritional, and Immunological Role of Dietary β -glucan and Ascorbic Acid Monophosphate in *Litopenaeus vannamei* juveniles. *Fish and Shellfish Immunology*. 15(1): 223-243.
- Sriwulan, Azwar, A., Rantetondok, A. and Anshary, H. 2019. Screening and application of lactic acid bacteria isolated from vanamei shrimp (*Litopenaeus vannamei*) intestine as a probiotic potential for tiger shrimp (*Penaeus monodon*). *AAFL Bioflux*. 12 (5): 1866-1881.
- Trisnawita, Y., Silalahi, J. and Sinaga, S.M. 2018. The effect of storage condition on viability of lactic acid bacteria in probiotic product. *Asian J Pharm Clin Res*. 11 (Special issue 1): 84-86.
- Yeh, S.P., Hsieh, S.L., Liu, C.H. and Cheng W. 2015. Immune response of shrimp, *Litopenaeus vannamei* after a concurrent infection with white spot syndrome virus and hepatopoeitic necrosis virus. *Fish and Shellfish Immunology*. 26 : 582-588.
- Zheng, Y. H., Yoshida, T., Isobe, K., Rahman, M. J., Nagase, F., Ding, L. and Nakashima, I. 2015. Modulation by Glycyrrhizin of the cell-surface expression of H-2 Class I Antigens on Marine Tumor Cell Lines and Normal Cell Populations. *Immunology*. 70 : 405-410.

ORIGINALITY REPORT

17 %	15 %	7 %	4 %
SIMILARITY INDEX	INTERNET SOURCES	PUBLICATIONS	STUDENT PAPERS

PRIMARY SOURCES

1	repository.unair.ac.id Internet Source	9 %
2	mafiadoc.com Internet Source	2 %
3	G Mahasri, P D W Sari, Prayogo. " Immune response and parasitic infestation on Pacific white shrimp () in immuno-probio circulation system (SI-PBR) in ponds ", IOP Conference Series: Earth and Environmental Science, 2018 Publication	1 %
4	www.envirobiotechjournals.com Internet Source	1 %
5	G Mahasri, R Kusdarwati, Kismiyati, Rozi, H Gustrifandi. " Effectivity of immunostimulant from protein membrane for decreasing the mortality rate of white shrimp () in traditional plus pond ", IOP Conference Series: Earth and Environmental Science, 2018 Publication	1 %

Renjie Qiu, Yongxu Cheng, Xuxiong Huang,

6	Xugan Wu, Xiaozhen Yang, Rui Tong. "Effect of hypoxia on immunological, physiological response, and hepatopancreatic metabolism of juvenile Chinese mitten crab <i>Eriocheir sinensis</i> ", <i>Aquaculture International</i> , 2010 Publication	1%
7	www.journalajfar.com Internet Source	<1%
8	Submitted to Universitas Brawijaya Student Paper	<1%
9	Submitted to Universiti Malaysia Terengganu UMT Student Paper	<1%
10	Submitted to University of Mosul Student Paper	<1%
11	worldwidescience.org Internet Source	<1%
12	repository.ipb.ac.id Internet Source	<1%
13	www.science.gov Internet Source	<1%
14	link.springer.com Internet Source	<1%
15	Zhang, Xing, You-Ting Zhu, Xue-Jie Li, Shi-Chuang Wang, Dan Li, Wei-Wei Li, and Qun	<1%

Wang. "Lipopolysaccharide and beta-1, 3-glucan binding protein (LGBP) stimulates prophenoloxidase activating system in Chinese mitten crab (*Eriocheir sinensis*)", *Developmental & Comparative Immunology*, 2016.

Publication

16

Subuntith Nimrat, Weerasith Khaopong, Jumlong Sangsong, Traimat Boonthai, Verapong Vuthiphandchai. " Improvement of growth performance, water quality and disease resistance against of postlarval whiteleg shrimp () by administration of mixed microencapsulated probiotics ", *Aquaculture Nutrition*, 2019

<1%

Publication

Exclude quotes Off

Exclude matches Off

Exclude bibliography On