

QUARTERLY INTERNATIONAL JOURNAL

ECOLOGY

ENVIRONMENT & CONSERVATION

ISSN 0971-765 X

Vol 26
No. (3)
2020



EM INTERNATIONAL

Shopping Bag (Items)

[Home](#)[International Journals](#)[For Authors](#)[Books](#)[About Us](#)[Contact](#)

Ecology, Environment and Conservation Editorial Advisory Board

Chief Editor

Prof.(Dr.) R.K.Trivedy, Pune, India**Email id- rktrivedy@gmail.com****Tel: 91-20-46745119****Mobile No.-9975703363**

EDITORIAL ADVISORY BOARD	
1. Dr. Prof. Moses Inbaraj, Professor, Department of Botany (Retired), Madras Christian College, Chennai, India	Indiainbarajmoses2004@yahoo.com
2. Dr. Prof. D.J. Lee, Professor, Department of Chemical Engineering, National Taiwan University, Taiwan	djlee@ntu.edu.tw
3. Dr. T. Bahorun, Professor, Department of Biosciences, Mauritius University, Mauritius	tbahorun@uom.ac.mu
4. Dr. Prof. A.R. Ghosh, Professor, Department of Environmental Sciences, University of Burdwan, Burdwan, India	apurbaghosh2010@gmail.com
5. Dr. Prof. M. Zaman, Professor Department of Botany, Dhaka University, Dhaka, Bangladesh	zashim@du.ac.bd
6. Dr. Marcantonio Bragadin, Professor Department of Chemistry, University of Venice, Venice, Italy	bragadin@unive.it
7. Dr. Z. Fuat Topark, Professor Dicle University, Diyarbakir, Turkey	toprakzey@itu.edu.tr
8. Dr. Wilson S. Tisera, Associate Professor Christian University, Kupang, Indonesia	wilson_tisera@yahoo.com
9. Dr. Prof. Philip C. Reid, Scientist Marine Biological Association, U.K	pcre@mba.ac.uk
10. Dr. Prof. Mohd. Yusuf, Professor Deptt. of Environmental Engineering, University Sans Malaysia, Malaysia	danielmy@iium.edu.my
11. Dr. Prof. Azni Idris, Deputy Vice-Chancellor, Research and Innov., Univ. of Putra Malaysia, Malaysia	azni@upm.edu.my
12. Dr. Vikas Sharma, Associate Professor, RARS of SKUAST, Jammu, J&K, India	vsagron@gmail.com
13. Dr. A.C. Pandey, Senior Scientist, Birsa Agricultural University, Ranchi, India	azni@upm.edu.my
14. Dr. Prof. Agoes Soegianto, Professor , Dept. Biology, Airlangga Univ. Kampus C, Surabaya, Indonesia	agoes_soegianto@fst.unair.ac.id
15. Dr. Ahmed L. Mahmoudi, Professor, Water Studies Centre, King Faisal University, Saudi Arabia	mahmoudi@kfu.edu.sa
16. Dr. Siroop Chaudhuri, Associate Professor Center for Environment, Sustainability and Human Development (CESH) O.P. Jindal Global University, Sonipat, India	schaudhuri@jgu.edu.in
17. Dr. Amin L. Setyo, Professor, Graduate School of International Crop, Hiroshima University, Hiroshima, Japan	amin28@ub.ac.id
18. Dr. Francis Gbogbo, Associate Professor, Department of Animal Biology, University of Ghana, Ghana	fgbogbo@ug.edu.gh
19. Dr. Prof. Deenbandhu Sahoo, Professor Department of Botany, university of Delhi, Delhi, India	dbsahoo@yahoo.com
20. Dr. Prof. J.K. Ladha, Professor Emeritus University of California, Davis, California, U.S.A.	jkladha@ucdavis.edu
21. Dr. Prof. M.Z.M. Nomani, Professor Faculty of Law, Aligarh Muslim University, Aligarh, India	zafarnomani@rediffmail.com
22. Dr. G.R. Pathade, Principal (Retired) H.V. Desai College, Pune, India	girishpathade@yahoo.co.in
23. Dr. Alfred Maroyi, Associate Professor Department of Botany, University of Fort Hare, Alice 5700, South Africa	amaroyi@ufh.ac.za
24. Dr. Prof. A.V.L.N.S.H. Hariharan, Professor Dept. of Chemistry, College of Science, GITAM University, Visakhapatnam, India	ahharan@gmail.com

25. Dr. Zeinab El-Sayed Zayed, Associate Professor Department of Biotechnology, The Central Lab of Date Palm Researches and Development Agriculture Research Center, Cairo, Egypt	zemz2005@yahoo.com
26. Dr. Raj Singh, Professor Department of Botany, MMD University, Mullana, Ambala, Haryana, India	dr.rajsingh09@gmail.com
27. Dr. Prof. Mohammed Latif Khan, Professor, Department of Botany Dr. Hari Singh Gour Vishwavidyalaya Sagar, India	khanml61@gmail.com
28. Dr. R. Dinesh Principal Scientist, ICAR-Indian Institute of Spices Research, Marikunnu, P.O. Kozhikode, India	rdinesh2005@gmail.com
29. Dr. Suresha Gududappa, Senior Environmental Scientist, Saudi Arabia	suresh@ebe.com.sa
30. Dr. Azma Hanim Ismail, School of Biological Sciences, Universiti Sains Malaysia, 11800 USM, Pulau Pinang, Malaysia.	
Back to EEC Journal Details	

[Home](#) | [International Journals](#) | [Books](#) | [About Us](#) | [Contact Us](#) | [Submit Paper](#) | [Search Journal Article](#) |

[Become a fan](#) on Facebook

[Follow us](#) on Twitter



© EM International 2012-2019 | **Developed by Eneblur Consulting**

Shopping Bag (Items)

[Home](#)[International Journals](#)[For Authors](#)[Books](#)[About Us](#)[Contact](#)

Ecology, Environment and Conservation Journal Papers

Issue: Vol 26, Aug Suppl. Issue, 2020**Archived Issues**

[THE EFFECTS OF FARMING BEEF CATTLE ON BALURAN NATIONAL PARK CONSERVATION \(CASE STUDY IN KARANG TEKOK, BANYUPUTIH DISTRICT, SITUBONDO REGENCY\)](#)

Siti Azizah and Yahya Bagus Kawedar

[Get Abstract](#)[Get Paper](#)

[THE RELATIONSHIP BETWEEN GENE EXPRESSION AND BIODEGRADATION OF CLORSPAN AND GROUND UP PESTICIDES BY PSEUDOMONAS AERUGINOSA BACTERIA](#)

Beadaa Abdalqader Mahdii and Ahmed Mohammed Turke

[Get Abstract](#)[Get Paper](#)

[INFLUENCE OF SODIUM AZIDE ON MORPHOLOGICAL TRAITS OF PROSO MILLET \(PANICUM MILIACEUM L.\) GENOTYPES](#)

Aiman Rysbekova, Elmira Dyussibayeva Abilbashar Seitkhodzheyev, Irina Zhirnova, Aiym Zhakenova, Gulzat Yessenbekova, Sholpan Bekenova and Damira Yussayeva

[Get Abstract](#)[Get Paper](#)

[PREPARATION AND APPLICATION OF TIO₂/AL ELECTRODE FOR THE REMOVAL OF DISPERSE DYE IN EC PROCESS](#)

Parameswari Kalivel, Jegathambal Palanichamy and Vijila Moses

[Get Abstract](#)[Get Paper](#)

[THE INFLUENCE OF INNOVATION ADOPTION LEVEL OF INTEGRATED PLANT MANAGEMENT ON RICE PRODUCTIVITY OF NATIONAL STRATEGIC FOOD CENTER](#)

Tri Prajawahyudo, Kliwon Hidayat, Yayuk Yuliati and Edi Dwi Cahyono

[Get Abstract](#)[Get Paper](#)

[HIGHLIGHTING THE POSSIBLE SIDE EFFECTS OF CONSUMING GREEN TEA: A HERBAL DRINK](#)

Jigisha Anand, Somya Sinha and Nishant Rai

[Get Abstract](#)[Get Paper](#)

[DETECTION OF PATHOGENIC LEPTOSPIRA BACTERIA IN THE AIRPORT ENVIRONMENT USING POLYMERASE CHAIN REACTION](#)

Syamsuar Manyullei, Hasnawati Amqam, Alwiyah and andi Imam Arundhana Thahir

[Get Abstract](#)[Get Paper](#)

[STUDIES ON MORPHOLOGICAL AND GENETIC DIVERSITY OF JASMINE ECOTYPES OF COASTAL KARNATAKA](#)

Search Articles

Journal Issues[Vol. 29, May, Suppl. Issue 2023](#)[Vol. 29, Apr. Suppl. Issue 2023](#)[Vol 29, Issue 1, 2023](#)[Vol. 29, Jan. Suppl. Issue 2023](#)[Vol. 28, Dec. Suppl. Issue, 2022](#)[Vol 28, Issue 4, 2022](#)[Vol. 28, Nov Suppl. Issue 2022](#)[Vol. 28, oct Suppl. Issue 2022](#)[Vol. 28, Sep Suppl. Issue 2022](#)[Vol 28, Issue 3, 2022](#)[Vol. 28, Aug Suppl. Issue 2022](#)[Vol 28, Issue 2, 2022](#)[Vol. 28, May Suppl. Issue 2022](#)[Vol 28, Issue 1, 2022](#)[Vol 28, Feb Suppl. Issue, 2022](#)[Vol 28, Jan Suppl. Issue, 2022](#)[Vol 27, Issue 4, 2021](#)[Vol 27, Dec Special Issue, 2021](#)[Vol 27, Nov Suppl. Issue, 2021](#)[Vol 27, Oct Suppl. Issue, 2021](#)[Vol 27, Issue 3, 2021](#)[Vol 27, Aug Suppl. Issue, 2021](#)[Vol 27, Issue 2, 2021](#)[Vol 27, May Suppl. Issue, 2021](#)[Vol 27, Issue 1, 2021](#)[Vol 27, Feb Suppl. Issue, 2021](#)[Vol 26, Issue 4, 2020](#)[Vol 26, Nov Suppl. Issue, 2020](#)[Vol 26, Oct Suppl. Issue, 2020](#)[Vol 26, Issue 3, 2020](#)[Vol 26, Aug Suppl. Issue, 2020](#)[Vol 26, Issue 2, 2020](#)[Vol 26, June Suppl. Issue, 2020](#)[Vol 26, April Suppl. Issue, 2020](#)[Vol 26, Issue 1, 2020](#)[Vol 26, Feb Suppl. Issue, 2020](#)[Vol 25, Issue 4 2019](#)[Vol 25, Issue 3 2019](#)[Vol 25, Nov Suppl. Issue, 2019](#)[Vol 25, Sept Suppl. Issue, 2019](#)[Vol 25, Aug Suppl. Issue, 2019](#)[Vol 25, July Suppl. Issue, 2019](#)[Vol 25, Issue 2 2019](#)

H.S. Chaitanya, Nataraja S. and M. Krishnappa

[Get Abstract](#)

[Get Paper](#)

BACTERIAL COMMUNITIES AS INDUCED BY VOLATILE GASES RELEASING FROM ORGANIC COMPOUNDS IN FOOD WASTES AND BIO-WASTES IN BANGKOK, THAILAND

Apichaya Phadpi¹, Thanit Pattamapitoon and Kasem Chunkao

[Get Abstract](#)

[Get Paper](#)

IMPACT OF MINING ON TREE DIVERSITY OF THE COAL MINING FOREST AREA AT RANIGANJ COAL FIELD AREA OF WEST BENGAL, INDIA

Saikat Mondal, Debnath Palit and Pinaki Chattopadhyay

[Get Abstract](#)

[Get Paper](#)

PRESENCE OF STRIPED FLYING BARB ESOMUS METALLICUS (TELEOSTEI, CYPRINIDAE) FROM WEST SUMATRA, INDONESIA

Veryl Hasan, Maheno S. Widodo, Abdul R. Faqih, Gunanti Mahasri, Muhammad Arief, Fitri S. Valen, Muhammad B. Tamam, Danang Yonarta, Fajar S. Pratama and Ren Fitriadi

[Get Abstract](#)

[Get Paper](#)

POLLINATION ECOLOGY AND BREEDING SYSTEM IN DALBERGIA LATIFOLIA AND D. SISSOIDES

K.R. Sasidharan, D. Thangamani, S. Prakash and K. Muraleekrishnan

[Get Abstract](#)

[Get Paper](#)

THE INVESTIGATION OF THE PROSPECT OF USING POTENTIAL ABANDONED WELLS IN SOUTH SUMATERA

Eddy Ibrahim, Maulana Yusup and R.R. Harminuke E.H.

[Get Abstract](#)

[Get Paper](#)

HEXAVALENT CHROMIUM CONTAMINATION OF GROUNDWATER IN BHILAI CITY: POTENTIAL PUBLIC HEALTH CRISIS IN CENTRALEAST INDIA

Alka Banchhor, Madhurima Pandey and Piyush Kant Pandey

[Get Abstract](#)

[Get Paper](#)

SPATIAL DISTRIBUTION AND VULNERABILITY OF SEA WATER INTRUSION IN MAKASSAR CITY

Meyke¹, Soemarno, Harsuko Riniwati and Andi Tamsil

[Get Abstract](#)

[Get Paper](#)

IMPACT OF GREEN TEA COMPOST ON SOIL QUALITY AND GROWTH OF PLANTS

Jigisha Anand, Sumedha Rawat, Sujata Teshwar, Soumya Gupta, Mohammad Sahil and Nishant Rai

[Get Abstract](#)

[Get Paper](#)

ISOLATION AND OPTIMIZATION OF CELLULASE PRODUCTION BY ASPERGILLUS PENICILLIODES 12 ASZ USING EXPERIMENTAL DESIGN

Aman A. Sorour, Safaa M. Ali, Moustafa Y. El-Naggar and Zakia A. Olama

[Get Abstract](#)

[Get Paper](#)

ASSESSING THE YIELD AND JUICE QUALITY OF SUGARCANE UNDER AGRO-CYCLING OF PAPER MILL SOLID WASTES AND TREATED EFFLUENT IRRIGATION

M. Suguna Devakumari

[Vol 25, May Suppl. Issue, 2019](#)

[Vol 25, April Suppl. Issue, 2019](#)

[Vol 25, Issue 1 2019](#)

[Vol 24, Issue 4 2018](#)

[Vol 24, Issue 3 2018](#)

[Vol 24, Issue 2 2018](#)

[Vol 24, Issue 1 2018](#)

[Vol 24, March Suppl. Issue 2018](#)

[Vol 24, Feb. Suppl. Issue 2018](#)

[Vol 23, Issue 4, 2017](#)

[Vol 23, Nov. Suppl. Issue 2017](#)

[Vol 23, Sept. Suppl. Issue 2017](#)

[Vol 23, Issue 2, 2017](#)

[Vol 23, Issue 3, 2017](#)

[Vol 23, Issue 1, 2017](#)

[Vol 23, Feb 2017 Suppl. Issue](#)

[Vol 22, Dec 2016 Suppl. Issue](#)

[Vol 22, Issue 4, 2016](#)

[Vol 22, Sept. Suppl. Issue, 2016](#)

[Vol 22, Issue 3, 2016](#)

[Vol 22, June Suppl. Issue 2016](#)

[Vol 22, Issue 2, 2016](#)

[Vol 22, April Suppl. Issue 2016](#)

[Vol 22, Issue 1, 2016](#)

[Vol 21, Issue 4, 2015](#)

[Vol 21 Dec. 2015 Suppl. Issue](#)

[Vol 21 Nov. 2015 Suppl. Issue](#)

[Vol 21, Issue 3, 2015](#)

[Vol 21, Issue 2, 2015](#)

[Vol 21 Suppl. Issue August 2015](#)

[Vol 21. Suppl. Issue June 2015](#)

[Vol 21, Issue 1, 2015](#)

[Supplement Issue, Dec. 2014](#)

[Special Issue-2014](#)

[Vol 20, Issue 4, 2014](#)

[Vol 20, Issue 3, 2014](#)

[Vol 20, Issue 2, 2014](#)

[Vol. 20 Issue 01, 2014](#)

[Vol. 19 Issue 04, 2013](#)

[Vol. 19 Issue 03, 2013](#)

[Vol. 19, Issue 02, 2013](#)

[Vol. 19, Issue 01, 2013](#)

[Vol. 18, Issue 04, 2012](#)

[Vol. 18, Issue 3, 2012](#)

[Vol. 18, Issue 2, 2012](#)

[Vol. 18, Issue 1, 2012](#)

[Vol. 17, Issue 4, 2011](#)

[Vol. 17, Issue 3, 2011](#)

[Vol. 17, Issue 2, 2011](#)

[Vol. 17, Issue 1, 2011](#)

[Vol. 16, Issue 4, 2010](#)

[Vol. 16, Issue 3, 2010](#)

[Vol. 16, Issue 2, 2010](#)

[Vol. 16, Issue 1, 2010](#)

[Vol. 15, Issue 04, 2009](#)

[Vol. 15, Issue 03, 2009](#)

[Vol. 15, Issue 02, 2009](#)

[Vol. 15, Issue 1, 2009](#)

[Vol. 14, Issue 04, 2008](#)

[Vol. 14, Issue 2-3, 2008](#)

[Vol. 14, Issue 2-3, 2008](#)

[Vol. 14, Issue 1, 2008](#)

[Vol. 14, Issue 2-3, 2008](#)

[Vol. 13, Issue 04, 2007](#)

[Vol. 13, Issue 2, 2007](#)

[Vol. 13, Issue 1, 2007](#)

[Get Abstract](#)[Get Paper](#)**DEVELOPMENT OF ENVIRONMENTAL PROFESSIONAL BEHAVIOR BASED ON NATURAL SCHOOL STUDY CASE NATURAL SCHOOL OF UNIVERSE (SOU), INDONESIA**

Rukiah, Zulfan Saam, 3Sukendi, Nurfid Junus and Sawaluddin

[Get Abstract](#)[Get Paper](#)**EFFECT OF CARTAP HYDROCHLORIDE (50% SP) INSECTICIDE ON GILL HISTOLOGY OF THE FISH, CIRRHINUS MRIGALA (HAMILTON)**

G. Vani, K. Veeraiah, M. Vijaya Kumar and S.K. Parveen

[Get Abstract](#)[Get Paper](#)**FAT, POLYPHENOLS, FERMENTED AND NON-FERMENTED CACAO BEANS (THEOBROMA CACAO L.) GROWN UNDER THREE TYPES OF STAND STRUCTURE WITH THE ENGINEERING OF LIGHT DISTRIBUTION ENHANCEMENT THROUGH PRUNING MANAGEMENT IN SOPPENG REGENCY OF SOUTH SULAWESI, INDONESIA**

Fatmawati, Ariffin, Setyono Yudho Tyasmoro and Roedy Soelistyono

[Get Abstract](#)[Get Paper](#)**PITCHER IRRIGATION IN SALINITY MANAGEMENT FOR PRODUCTION OF BRINJAL CROP IN COASTAL SOIL OF WEST BENGAL, INDIA**

Rahul Adhikary, Monisankar Bera, Susanta Kumar De and Arunabha Pal

[Get Abstract](#)[Get Paper](#)**IMPACT OF CLIMATE CHANGE ON POTENTIAL DISPERSAL OF PAEONIAOBOVATA (PAEONIACEAE), A CRITICALLY ENDANGERED MEDICINAL PLANT OF SOUTH KOREA**

Ja-Young Jeon, Pradeep Adhikari and Changwon Seo

[Get Abstract](#)[Get Paper](#)**CONSERVATION AND MANAGEMENT OF URBAN LAKES; A CASE STUDY OF UDAIPUR CITY**

Shagufta Jabin, Sonal Gupta, Anupama Rajput and Mukta Sharma

[Get Abstract](#)[Get Paper](#)**DETERMINATION OF CROP COEFFICIENT (KC) OF SAGO PALM (METROXYLON SAGU) ON SEVERAL GROWTH STAGES**

Yumna, Sugeng Prijono, Zaenal Kusuma and Soemarno

[Get Abstract](#)[Get Paper](#)**PHYTOCHEMICAL PROFILING AND BIOLOGICAL ACTIVITY OF LEUCAS LANATA BENTH. AN IMPORTANT ETHNOMEDICINAL PLANT OF WESTERN HIMALAYA**

Rachna Vermaa, Ashwani Tapwalb, Dinesh Kumara and Sunil Puria

[Get Abstract](#)[Get Paper](#)**THE SOIL MACROFAUNA DIVERSITY IN 'NEGERI LAMA' OIL PALM PLANTATION AREA, INDONESIA**

Arlen Hanel John, Abdul Rauf, T. Sabrina and Erwin Nyak Akoeb

[Get Abstract](#)[Get Paper](#)**DISTRIBUTION OF AVIAN SPECIES IN MANDAKINI RIVER VALLEY IN RUDRAPRAYAG DISTRICT, UTTARAKHAND (WESTERN HIMALAYA), INDIA**

Tribhuvan Chandra and Dinesh Bhatt

[Vol.12, Issue 4, 2006](#)[Vol.12, Issue 3, 2006](#)[Vol.12, Issue 2, 2006](#)[Vol.12, Issue 1, 2006](#)[Vol.12, Issue 01, 2006](#)[Vol.11, Issue 3,4, 2005](#)[Vol.11, Issue 2, 2005](#)[Vol.11, Issue 1, 2005](#)[Vol.10, Issue 04, 2004](#)[Vol.10, Issue 03, 2004](#)[Vol.10, Issue 02, 2004](#)[Vol.10, Issue 01, 2004](#)[Vol.09, Issue 04, 2003](#)[Vol.09, Issue 03, 2003](#)[Vol.09, Issue 02, 2003](#)[Vol.08, Issue 04, 2002](#)[Vol.08, Issue 03, 2002](#)[Vol.08, Issue 01, 2002](#)[Vol.07, Issue 04, 2001](#)[Vol.07, Issue 03, 2001](#)[Vol.07, Issue 02, 2001](#)[Vol.07, Issue 01, 2001](#)[Vol.06, Issue 04, 2000](#)[Vol.06, Issue 03, 2000](#)[Vol.06, Issue 02, 2000](#)[Vol.06, Issue 01, 2000](#)[Vol.05, Issue 04, 1999](#)[Vol.05, Issue 03, 1999](#)[Vol.05, Issue 02, 1999](#)[Vol.05, Issue 01, 1999](#)[Vol.04, Issue 1,2, 1998](#)[Vol.03, Issue 3,4, 1997](#)[Vol.03, Issue 01, 1997](#)[Vol.02, Issue 1,2, 1996](#)[Vol.01, Issue 14, 1995](#)

Looking for Past Issues?

[Click here to get them!!](#)

[Get Abstract](#)[Get Paper](#)

EVALUATION OF EFFICACY OF ENTOMOPATHOGENS AGAINST SOLENOPSIS GEMINATA (FABRICIUS).(HYMENOPTERA: FORMICIDAE).

T. Nalini and S. Sasinathan

[Get Abstract](#)[Get Paper](#)

WATER VAPOR CONDUCTANCE OF DRY SOIL – ANALYZING TRANSPORT PROCESSES BY ANALOGY WITH OHM'S LAW

Mohammad Nazari-Sharabian, Moses Karakouzian and Parastoo Pashmchi

[Get Abstract](#)[Get Paper](#)

HETEROGENEITY OF ZOOPLANKTONS IN BORANAKANIVE RESERVOIR OF TUMAKURU DISTRICT, KARNATAKA, INDIA

R. Shruthi and S. Ramakrishna

[Get Abstract](#)[Get Paper](#)

EVALUATION OF TOXICITY OF BOTANICALS AND ESSENTIAL OILS AGAINST SOLENOPSIS GEMINATA (FABRICIUS).(HYMENOPTERA: FORMICIDAE)

T. Nalini and S. Sasinathan

[Get Abstract](#)[Get Paper](#)

EARTHWORM POPULATION DENSITY AND DIVERSITY WITH RESPECT TO SOIL PHYSICO CHEMICAL PROPERTIES, MICROBIAL POPULATION AND EXOENZYME DYNAMICS IN TWO AGROCLIMATIC ZONES OF ODISHA, INDIA

Pratik Acharya and C.S.K. Mishra

[Get Abstract](#)[Get Paper](#)

MEASUREMENT OF LONGSHORE SEDIMENT TRANSPORT IN THE COASTAL REGIONS USING WIRELESS SENSOR NETWORK

Pearlin Sam Jinoj T., Mahija Hann J. and Ramesh R.

[Get Abstract](#)[Get Paper](#)

SEASONAL VARIATIONS IN DIVERSITY OF AQUATIC MACROPHYTES OF UPPER LAKE, BHOPAL

Purna Gupta, Sadhna Tamot, Vinoy K Shrivastava and Rajendra Chakarde

[Get Abstract](#)[Get Paper](#)

COMPARATIVE STUDY ON SPECIES COMPOSITION AND DIVERSITY OF INSECTS IN THE SOUTH PARK STREET CEMETERY AND BANABITAN PARK, WEST BENGAL, INDIA

Hiroj Kumar Saha and Labani Islam

[Get Abstract](#)[Get Paper](#)

IMPROVING THE METHODS OF BREEDING, STORING, AND USING BEAKED MITES FOR BIOLOGICAL PEST CONTROL

Irina Sergeevna Agasieva, Vladimir Yaëovlevih Ismailov, Elena Valentinovna Fedorenco, Abay Orazuly Sagitov, Aida Askarkyzy Mukhtarkhanova and Aizhan Mukashevna Chadinova

[Get Abstract](#)[Get Paper](#)

BIO REMEDIATION OF TEXTILE EFFLUENT USING BACTERIAL CONSORTIUM OBTAINED FROM INDUSTRIALLY POLLUTED SITE

Poonam Ranga, Deepansh Sharma and Baljeet Singh Saharan

[Get Abstract](#)

[Get Paper](#)

ASSESSMENT OF WATER QUALITY STATUS OF A TROPICAL FRESH WATER MANU SWAMP-RAMBHA RIVER, RISHIKESH, DEHRADUN, INDIA

Shail Kulshrestha

[Get Abstract](#)

[Get Paper](#)

CAUSES OF MAN-ELEPHANT CONFLICTS AND ITS IMPACT ON LIVELIHOOD IN FRINGE VILLAGES OF MOMAN RESERVE FOREST, KAMRUP DISTRICT, ASSAM, INDIA

Meghna Das and Gogoi Barnali

[Get Abstract](#)

[Get Paper](#)

STUDY OF GEOPOLYMER CONCRETE OF RICE HUSK ASH AND FULLER'S EARTH

Lisha Kurup, Manas Nair, Ayush Vyas, Ayush Narnaware, Ayan Agrawal and Harshwardhan Gupta

[Get Abstract](#)

[Get Paper](#)

SOIL CHEMICAL PROPERTIES OF OPENCAST COAL MINING SITE IN INDONESIA AND ITS EFFECT ON PLANT GROWTH

Dewi Wulandari, Saridi, Doddy Herika, Cahyono Agus, Weiguo Cheng and Keitaro Tawaraya

[Get Abstract](#)

[Get Paper](#)

FROM SUGAR PRODUCTION TO SUSTAINABLE ENERGY PRODUCTION: EXPLORING SCENARIOS AND POLICY IMPLICATIONS FOR BIOENERGY IN THE SUGAR BOWL OF INDIA

Dhanraj A. Patil

[Get Abstract](#)

[Get Paper](#)

DIET AND FEEDING HABIT OF ASIATIC BLACK BEAR (URSUS THIBETANUS) IN NANDA DEVI BIOSPHERE RESERVE, UTTARAKHAND, INDIA

Vijay Kumar Yadav, D.S. Chauhan and P.C. Lakhera

[Get Abstract](#)

[Get Paper](#)

EXPLAINING THE RELATIONSHIP BETWEEN THE CONCEPTS OF SMART CITY AND CITIZENS' MENTAL HEALTH

Atiyeh Asgari, Mostafa Behzadfar and Asadollah Naghdi

[Get Abstract](#)

[Get Paper](#)

ENRICHMENT POTENTIAL OF BRACHIONUS PLICATILIS WITH FISH OIL AND SQUID OIL FOR THE ACCELERATION OF THE EARLY GIANT GROUPER DEVELOPMENT (EPINEPHELUS FUSCOGUTTATUS × EPINEPHELUS LANCEOLATUS)

Dwi Mirnasari, Woro Hastuti Satyantini and Endang Dwi Masithah

[Get Abstract](#)

[Get Paper](#)

GENETIC VARIABILITY AND AROMA STUDIES IN LANDRACES AND CULTIVARS OF ORYZA SATIVA

Akhtar Sabina

[Get Abstract](#)

[Get Paper](#)

GROWTH PROMOTION EFFECTS OF HUMIC ACIDS ISOLATED FROM CO-COMPOST PREPARED USING FISH PROCESSING WASTE AND SUGARCANE BAGASSE ON SEEDS OF FABACEAE FAMILY

Enrichment potential of *Brachionus plicatilis* with fish Oil and squid oil for the acceleration of the early giant Grouper development (*Epinephelus fuscoguttatus* × *Epinephelus lanceolatus*)

Dwi Mirnasari¹, Woro Hastuti Satyantini^{1*} and Endang Dwi Masithah²

¹Department of Marine, Faculty of Fisheries and Marine, Universitas Airlangga, Jl. Mulyorejo, Sutabaya, East Java, 60115, Indonesia

²Department of Fish Health Management and Aquaculture, Faculty of Fisheries and Marine, Universitas Airlangga

(Received 20 January, 2020; accepted 6 March, 2020)

ABSTRACT

The development of early stadia giant grouper larvae is a problem that often occurs in grouper hatcheries. The highly DHA and EPA content in feed is very necessary to the grouper larvae. However, the natural feed that is given, namely *Brachionus plicatilis*, has low EPA and DHA content. This study aims to determine the development of early stage grouper abstinence by giving *Brachionus plicatilis* enriched with a combination of fish oil and squid oil. This study used an experimental method and the test animals were divided into two groups, namely control (feed *B. plicatilis* without enrichment plus *Nannochloropsis sp.*) and treatment (feed *B. plicatilis* by enriching the combination of fish oil and squid oil plus *Nannochloropsis sp.*). Statistical test used Independent-samples, T-Test. The acceleration of the grouper larvae development in the treatment group, especially in the dorsalis spina and ventral spines formed earlier on the seventh day (D7) compared to the control group larvae of the newly formed control group on the eighth day (D8). In addition, it was found that enrichment of *B. plicatilis* with fish oil and squid oil could accelerate the growth of absolute length (L) of the flak grouper larvae by 3.27 ± 0.17 mm. The administration of *B. plicatilis* which was given squid oil and fish oil as larvae feed that made the early stages growth of the grouper larvae faster in several stages. In addition, the absolute lengths of larvae that are fed with enrichment are longer than the larvae that are fed without enrichment.

Key words : Giant grouper, *B. plicatilis* enrichment, Squid oil, Fish oil, Early larval stage

Introduction

Grouper is a leading Indonesian fishery commodity and has high demand reaching 35,000 tons per year with a price range of US \$ 25 - US \$ 125. BPS also noted that the contribution of live grouper exports in 2016 reached 1.11% of the total export value of fisheries commodities. Total value of live grouper

exports in 2016 reached 32.18 million US \$ (Kementrian Kelautan and Perikanan Republik Indonesia, 2013). However, there are still many problems in breeding process. The most common problem encountered in hatchling grouper fish is the slow development of the initial larval stage. This is caused by the low content of EPA and DHA in the natural food that was given, namely *Brachionus*

plicatilis, whereas EPA and DHA are important components for the growth of giant grouper larvae.

The development of grouper larvae can be divided into 4 phases, such as; i) yolk sac phase, which starts from hatching until the egg yolk runs out; ii) the pre-flexion phase, which starts from the yolk absorbed completely until the spine is formed; iii) the flexion phase, which starts from the formation of the dorsal spine and ventral spine to reduce the spine; iv) the post-flexion phase, which starts from the loss or reduction of the spine until it becomes juvenile.

One of the nutrients that are highly needed by marine fish larvae is fat (Halver and Hardy, 2002). The main constituent of fat is fatty acids (Purwaningsih *et al.*, 2014). Seawater fish larvae require long unsaturated fatty acids or Highly Unsaturated Fatty Acids (HUFA) of more than 4% (Léger *et al.*, 1986). HUFA that is needed by sea water fish larvae are Eicosanoic Acid (EPA) and Docosanoic Acid (DHA) (Tocher, 2003). EPA and DHA are very instrumental in the development of grouper larvae.

EPA and DHA is long chain polyunsaturated fatty acids that play an important role in the largest structural component of the phospholipid membrane that regulates membrane fluidity and ion transport (Chapkin *et al.*, 2008). The need of HUFA increases in the grouper larval development stage because it is widely used in the formation of cell membranes and tissues (Leaver *et al.*, 2008). *B. plicatilis* as a natural feed that is often given to grouper larvae has relatively small amount of HUFA, especially EPA and DHA, hence, the nutritional quality of fatty acids is very low (Watanabe, 1993). *B. plicatilis* that is cultured with bread yeast or *Nannochloropsis* has a low nutritional content. The content of *B. plicatilis* essential fatty acids fed with bread yeast gives a low nutritional value of only 1% EPA and 0.1% DHA (Watanabe *et al.*, 1983), while enrichment with *Nannochloropsis* only has an EPA value of 0.94%-1.46% and DHA in very small amounts. The low content of essential fatty acids especially EPA and DHA can be increased by utilizing *B. plicatilis*. *B. plicatilis* is a non-selective filter feeder, which take all the food around it without selection. Thus, the type of feed given during *B. plicatilis* culture greatly influences its nutritional value. By utilizing these properties, the nutritional value of *B. plicatilis* can be improved by enrichment. One of the enrichment in *B. plicatilis* can be done by

giving oil emulsions that are rich in EPA and DHA content. Fish oil and squid oil are rich of essential fatty acids such as EPA and DHA. *B. plicatilis* enriched by lemuru fish oil has an EPA content of 8.8% and DHA 0.1%. The content is higher than *B. plicatilis* enriched by *Nannochloropsis salina*, *Isochrysis galbana*, *Chlorella marina*, and *Saccharomyces cerevisiae*. Enrichment of *B. plicatilis* with squid oil contains 9-12% EPA and 2-3% DHA. The high content of EPA and DHA is caused the most squid lipids in the form of phospholipids which contain lots of essential fatty acids. Enrichment of *B. plicatilis* with a combination of fish oil and squid oil is assumed to increase the content of essential fatty acids needed in the development of grouper larvae. Some of them, namely fish oil and squid oil, are oils that are rich in EPA and DHA content. If fish oil and squid oil are added to enrich *B. plicatilis*, it is suspected that it can increase the EPA and DHA content to help accelerate the development of grouper larvae. This study aims to determine the development of the initial stage of the grouper abstraction by giving *Brochionus plicatilis* enriched with a combination of fish oil and squid oil.

Method

This method used an experimental method with the experimental design used t-test. The sample is divided into 2 treatments, namely P0 given *B. plicatilis* without enrichment and *Nannochloropsis sp.* and P1 given *B. plicatilis* by enriching the combination of fish oil and squid oil plus *Nannochloropsis sp.* The independent variable in this study was the administration of *B. plicatilis* enriched in a combination of fish oil and squid oil. The dependent variables in this study were the speed of development (width of the mouth opening, eye diameter, dorsalis spine length and ventral spine length) abstraction grouper larvae and growth in length (L) abstraction grouper larvae.

Research Material

The materials used in this study included D-0 grouper larva that was origin from Bali, Indonesia. Natural feed was the form of *B. plicatilis* and *Nannochloropsis sp.* Seawater of 30-31 ppt salinity as a medium for maintenance of grouper larvae. Chlorine, detergents, and chlorine were as treatment materials or sterilization tools and maintenance

media. Sodium thiosulphate function neutralized the chlorine or chlorine content in the device or media that has been treated. Checking the water with chlorine test, if yellow indicated the media is not neutral and if clear or no color indicated the media is neutral.

Enriching agents used included fish oil and squid oil. Duck egg yolk as an emulsifier which functions to keep oil grains suspended in water. Aquades was as dispersing media (Winarno, 1984). 5% formalin solution used to preserve (Kohno, 1998) and kill grouper larvae when observed through a microscope.

Manufacture of Fish Oil and Squid Oil Emulsions

Fish oil and squid oil cannot be given directly as an enrichment of *B. plicatilis* so it must be mixed with other ingredients such as duck egg yolk through an emulsion process. The ingredients needed were 10 mL of duck egg yolk, 40 mL of distilled water, 25 mL of fish oil, and 25 mL of squid oil. Then, the emulsion was stored in the refrigerator and can be used with thawing first.

Provision of *B. plicatilis*

The initial stage is filling of the structure given by the seeds of *Nannochloropsis sp.* with a density of 107 cells/mL. Then, fill 25% of seawater from the bottom of the tub. After that, *B. plicatilis* was spread with a density of 50 ind/mL. Harvesting of *B. plicatilis* from mass culture was carried out partially on the third day after stocking.

Enrichment of *B. plicatilis*

The density of harvested *B. plicatilis* is 150 ind/mL. The enrichment process of *B. plicatilis* with fish oil and squid oil emulsions at a dose of 0.5 ml/L enrichment media. Enrichment was carried out for 12 hours every day (Akbari *et al.*, 2011). In addition, *B. plicatilis* in the control treatment (P0) and enrichment treatment (P1) also added *Nannochloropsis sp.* with a density of $20-30 \times 10^6$ cells /mL (Redjeki, 1999). The administration of *Nannochloropsis sp.* served as *B. plicatilis* food for 12 hours of enrichment. This can keep *B. plicatilis* alive for 12 hours of the enrichment process.

Rearing Grouper Larvae

Stages in the maintenance of larvae include preparation of the container, feeding both natural and artificial, and water quality management media main-

tenance. Spread eggs as much as 500,000/tub. The grouper larvae are kept in larval rearing tanks with a capacity of 30 tons that have been sterilized.

There are two treatments given to *B. plicatilis* as grouper larvae feed in the control treatment (P0) *B. plicatilis* without enrichment. Treatment one (P1) *B. plicatilis* was enriched with emulsion of fish oil and squid oil as much as 0.5 mL/L of media yield of *B. plicatilis* for 12 hours.

The administration of *Nannochloropsis sp.* in grouper larval rearing tanks carried out since D2 with a density of $300-500 \times 10^5$ cells/mL. Apart from being a feed for *B. plicatilis*, *Nannochloropsis sp.* in grouper larval rearing tanks also acted as green water (Ismi *et al.*, 2012). The use of *Nannochloropsis sp.* as green water formed the color of the water to be dimmer, so that the grouper larvae do not cluster (Ismi, Asih and Kusumawati, 2014).

B. plicatilis was given when grouper larvae are D-2 to D-15. At the beginning of administration, the density of *B. plicatilis* given was 1-3 ind/mL. In D-3 to D-8 the density of *B. plicatilis* was maintained at 5 ind/mL, in larvae D-8 to D-15 the density of *B. plicatilis* was increased to 10-15 ind/mL. The frequency of administration of *B. plicatilis* was once a day at 08.00 a.m.

The grouper larval rearing tank was maintained static until D-7. Initially, water exchange was limited to only around 10% / day D-7 to D-12, to avoid sudden changes in water quality. Substitution of water gradually increased to 20% / day, when given artificial feed D-13 to D-15. From around D-12, grouper larvae droppings, dead grouper larvae, and food that accumulate at the bottom of the tank must be siphoned out at least once a day to maintain water quality. Initially, only one-quarter of the bottom of the tank was siphoned every day. This was done gradually increasing until all parts of the tank are siphoned every day.

Research Parameters

The main parameters in this study were the speed of development of mouth opening width, eye diameter (Glamuzina *et al.*, 1998), dorsalis spine length and ventral spine length, and specific growth rate of the grouper abbot larvae length. The total length was measured from the most anterior part of the mouth to the most posterior caudal fin (Glamuzina *et al.*, 1998). The mouth opened was calculated by the length of the lower jaw multiplied by $\tan (50^\circ)$

(Dabrowski and Bardega, 1984). The eye diameter was the longest diameter of the eye. Dorsal spine length measured from the base to the tip of the first radius of the dorsal fin (Moyle and Cech, 2004), whereas the length of the ventral spine was measured from the base to the tip of the first radius of the abdominal fin (Moyle and Cech, 2004).

Larvae samples were taken randomly using 3 dropper pipettes from each tub. Sampling was done every morning on D-3 to D-15 because grouper larvae generally consume *B. plicatilis* at that age. The larvae taken were placed on a concave object glass and turned off with 5% formalin to then be observed its development using a microscope equipped with an ocular micrometer at a magnification of 40 times.

Growth measurement aimed to determine the magnitude of larval growth during maintenance. Growth of absolute length larvae of grouper fish was calculated by measuring the total length of larvae at the end of maintenance minus the total length

of larvae at the beginning.

Supporting parameters in this study were observations of water quality media for maintenance of grouper larvae. Water quality parameters observed were temperature, DO, pH, and salinity. Observation of temperature, DO, pH and salinity was carried out three times a day at 07.00, 12.00, and 16.00.

Data Analysis

The observing results of the development speed of grouper abstraction larvae were explained descriptively. Growth data for absolute long grouper larvae were analyzed by Independent Samples T-Test by the SPSS of Windows program. Supporting data in the form of a range of water quality values for grouper larvae maintenance were revealed descriptively.

Results

The results of the grouper larvae growth for 15 days

Table 1. The Growth of Grouper Larvae

Larvae aged (D-)	Growth	
	P0	P1
D-1	New larvae hatch and cannot swim actively, they still have egg yolk and globule oil, the eyes have not been pigmented and the mouth not open yet.	New larvae hatch and cannot swim actively, they still have egg yolk and globule oil, the eyes have not been pigmented and the mouth not open yet.
D-3	The larvae begin to swim actively, egg yolk and globule oil are completely absorbed, the pectoral fins are formed, the eyes have pigmented and the mouth is open, the average eye diameter is 0.227 ± 0.006 mm.	The larvae begin to swim actively, egg yolk and globule oil are completely absorbed, the pectoral fins are formed, the eyes have pigmented and the mouth is open, the average eye diameter is 0.248 ± 0.015 mm.
D-4	Total length of 2.706 ± 0.048 mm.	Total length of 2.867 ± 0.101 mm.
D-5	The total length is 2.867 ± 0.101 mm.	The total length is 2.975 ± 0.023 mm.
D-6	The black spot of dorsal spina candidate has not been seen.	Spina dorsalis prospective black dots have been seen.
D-7	Black dots of dorsal spine candidates have been seen in the dorsal section.	Dorsal spine formed with a length of 0.033 ± 0.014 mm and ventral spine with a length of 0.425 ± 0.066 mm.
D-8	Dorsal spine formed with a length of 0.033 ± 0.014 mm and ventral spine with a length of 0.358 ± 0.014 mm.	Spina dorsalis has pigmented.
D-9	Spina dorsalis has pigmented.	The dorsal spina lengthens the dorsal section.
D-15	The total length of larvae increased to 4.872 ± 0.547 mm.	The total length of larvae increased to 5.792 ± 0.144 mm.
Average L (mm)	2.20 ± 0.43^a	3.27 ± 0.17^b

Note: Different superscripts on the same line showed differences ($P < 0.05$). The average L was measured at D3 - D15.

can be seen in Table 1. Results showed that grouper larvae given enrichment of *B. plicatilis* with L P1 of 3.27 ± 0.17 mm were longer than P0 (without enrichment) which was only $2, 20 \pm 0.43$ mm.

Water quality

Water quality parameters measured during the study were temperature, pH, DO, and salinity. The results of water quality measurements were carried out on each treatment as many as two replications every day at 07.00, 12.00 and 16.00. Data on water quality range for maintenance of grouper larvae during the study can be seen in Table 2.

The value of water quality media for maintenance of grouper larvae during the study was within the normal range for the development of grouper larvae. Water temperature during the study ranged from 29.0 °C - 31.3 °C. Water pH during the study ranged from 8.15 - 8.64. DO water during the study ranged from 5.4 to 6.8 ppm. Salinity of water during the study was worth 31 ppt.

Discussion

First day (D-1) grouper aberration larvae in the study has the following morphology: the body was still transparent, the skin has not been pigmented, the mouth has not opened, the eyes have not been segmented yet, the intestine was straight (not filled), the anus was still closed, the apical tip of the caudal fin was still rounded, pectoral fins, dorsal fins, anal fins, and ventral fins have not been differentiated. At this age, the larva grouper larvae still have egg yolks and oil bubbles that were used as a food source.

At the age of D-3 the yolk has been absorbed, the mouth can open, the stomach begins to contract, the anus has opened, and the eye pigment was clear. The eye pigment was very important for the grouper larvae to see and find food. Pigtail grouper larva eye pigments begin to form at the age of D-3 with different diameters at P0 and P1. Eye pigmentation

indicated that the grouper larvae were ready to search for prey / food. At this stage, the grouper larvae were in the first critical phase because the larvae undergo an adaptation process that came from the endogenous feeding phase to exogenous feeding. At this time, grouper larvae have to get food from the outside because the egg yolk has run out. This was consistent with previous research (Setyadi, 2008), that the critical period of fish larvae occurred in the transition period between the expiration of egg yolk and the time to start taking food from outside.

Apart from eye development, the development of the mouth opening was also very important in the development phase of the giant grouper larvae. The rapid development of mouth opening will affect the ability of giant grouper larvae to catch food. The size of the mouth opening was larger than the size of *B. plicatilis* which ranges from 0.08 to 0.12 mm, thus the larvae can catch its food because it matched the size of its mouth opening.

The dorsal spine and ventral spine on P1 form earlier on the seventh day (D-7) than on P0 that only forms on the eighth day (D-8). This was shown in Table 3, at the age of D-7, the dorsal spina P1 appeared to be prominent and formed earlier while the dorsal spina P0 was only formed at D-8. In addition, at the age of D-7, the ventral spine P1 has formed, whereas the ventral spina P0 was only formed at D-8. The formation of dorsal spine and ventral spine larvae of the grouper abstraction in this study was faster than in previous studies (Ch'ng and Senoo, 2008) where new dorsal spines formed at the age of D-9 while ventral spina was only seen at D-9 to D-10. The difference in the initial development of the dorsal spine and the ventral spina can be caused by differences in the nutrients contained in the feed given. The dorsal spine and ventral spine will reach a maximum length at the age of D-20 after hatching (Ch'ng and Senoo, 2008). After reaching the maximum length, the spine will reduce and turn into hard fin radii (Bulanin *et al.*,

Table 2. Data of Water Quality Enrichment Media of Grouper Larvae during Study

Parameter	Value		Standard value
	P0	P1	
Temperature (°C)	29.0 – 31	29.1 - 31.3	28-32
pH	8.15 - 8.64	8.28 - 8.57	7.5 – 8.6 (Ch'ng and Senoo, 2008)
DO (ppm)	5.4 - 6.8	5.6 - 6.7	5 – 6 (Chu <i>et al.</i> , 2016)
Salinity (ppt)	31	31	31 – 32 (Ch'ng and Senoo, 2008)

2018). The fingers of the hard fin on the dorsal fin and the abdominal fin are characteristic of fish from the family of serenades.

In the phase of grouper larvae where dorsal spines and ventral spines have formed, indicated that the larvae have entered the second critical phase. In this study, the second critical phase of the P1 grouper larvae occurred at the age of D-7, however in P0 it only occurred at D8. In this critical phase larvae already need feed with more complex nutritional content. This was consistent with study that during the growth of fish, especially in the younger phase requires more nutrient content in previous food (Effendi, 2004). The faster the dorsal and ventral spines were formed, the faster the larvae pass through the second critical phase, the faster the process of metamorphosis.

Growth was the increase in length or weight over time. Based on the result of the Independent-Samples, T Test analysis the absolute length (L) larvae growth of grouper revealed significantly different results. This examined that the enrichment of *B. plicatilis* with a combination of fish oil and squid oil influenced the growth of absolute length of the grouper larvae. The absolute length growth is illustrated in Table 3, where the absolute length growth P1 (with enrichment) was faster than P0 (without enrichment).

The enrichment of fish oil and squid oil combination has accelerated the growth of absolute length of the grouper larvae in P1 compared to P0. This happened because the enrichment has improved the nutritional quality of *B. plicatilis* consumed by the grouper larvae. These nutrients, especially EPA and DHA, can support faster metabolic processes and have an impact on better growth in larva grouper larvae.

Based on the results of the study, *B. plicatilis* with enriched combination of fish oil and squid oil have higher EPA and DHA content than *B. plicatilis* without enrichment. *B. plicatilis* enriched with a combination of fish oil and squid oil had a percentage of EPA and DHA respectively 15.60% and 4.56%, whereas *B. plicatilis* without enriching the percentage of EPA and DHA were lower, i.e. 12.92% and 2.26%. The higher EPA and DHA content in P1 was influenced by the addition of a combination of fish oil and squid oil emulsion given at the time of *B. plicatilis* enrichment.

Giant grouper was a hybrid grouper that results from a cross between tiger grouper (*Epinephelus*

fuscoguttatus) female and giant grouper (*Epinephelus lanceolatus*) male that live in sea waters. Fish that live in marine waters do not have the activity of the delta 5-desaturase enzyme as those in freshwater fish, so that marine fish were in dire need of long-chain polyunsaturated fatty acids such as EPA and DHA in their feed (Peng *et al.*, 2003).

Faster growth of grouper larvae in P1, due to the enrichment of *B. plicatilis* with a combination of fish oil and squid oil can increase the content of fatty acids, especially EPA and DHA needed in the development of grouper larvae. Fat that enters the body of the fish will be hydrolyzed by the lipase enzyme into fatty acids and glycerol (Afrianto and Liviawaty, 2005). In this hydrolysis process, lipases were aided by bile acids and lecithin. The process of hydrolysis occurs along the digestive tract, especially in the intestine. These fatty acids will be absorbed by the intestinal mucosal enterocytes and into the blood. Most of the absorbed fat was transported to the liver. Fat was converted into phospholipids and transported by blood to cells and tissues. These fatty acids, specifically EPA and DHA, were the largest structural component in the phospholipid membrane that regulated membrane fluidity and ion transport (Chapkin *et al.*, 2008). Hence, the existence of EPA and DHA can support the smooth metabolism of cells and tissues, thus it impacts on the rapid development of larva grouper larvae.

Other factors besides feed influenced the development of early stadia grouper larvae were water quality media maintenance. The condition of water quality of the maintenance media showed that conditions were still suitable for the maintenance of the grouper larvae. Water quality observed included temperature, pH, DO, and salinity.

Temperature of grouper larvae rearing media influences the process of hatching organ development in which the process of embryogenesis and early organogenesis occurs. Water temperature during rearing influenced the metabolic rate of aquatic animal metabolic processes that were also poikilothermic which also impacts on the development process of larval organs produced. The initial development of fish was strongly influenced by the temperature at which the eggs. The temperature during the study in both treatments ranged from 29.0 - 31.3 °C. This temperature was still in the range of temperatures that can be feasible in the development of bleach grouper larvae. This was consistent with the statement that the temperature 28–32 °C was the

water quality standard for grouper hatcheries.

The pH content in each treatment ranged from 8.15 to 8.64. This was consistent with the previous one, that the pH value for the maintenance of larva grouper larvae was 7.5 - 8.6 (Ch'ng and Senoo, 2008). Aquatic pH influences the pH of blood plasma which can negatively influence larval development. A low pH affected the oxygen consumption of grouper larvae in decomposing alkaline properties in water. This can be seen from the difficulty of the gills in the process of taking oxygen and interfering with the transportation of oxygen in the blood, causing metabolic processes to be disrupted (Saleh *et al.*, 2013).

Dissolved oxygen content during the study ranged from 5.4 - 6.7 ppm. This condition was very supportive for maintenance of larva grouper larvae. The measurement results revealed that the higher the temperature, the lower the dissolved oxygen (DO). This was because an increase in temperature causes an increase in the speed of metabolism and respiration of aquatic organisms, thus resulting in an increase in oxygen consumption which results in a decrease in the value of oxygen content in water. DO requirements in raising grouper larvae range from 5-6 ppm, so it can be said that the oxygen content during this study can meet the needs of dissolved oxygen needed by the grouper larvae (Chu, 2016).

Salinity was one of the determining factors in the development of grouper larvae. The ideal salinity for maintenance of larva grouper larvae was 31 - 32 ppt (Ch'ng and Senoo, 2008). During the study, there were no significant changes in salinity. Salinity of maintenance media water was 31 ppt.

Conclusion

Enrichment of *B. plicatilis* with a combination of fish oil and squid oil can accelerate the development of the grouper larvae larvae, especially on the development of dorsalis spina and ventral spina that formed earlier in D7 than the larva grouper larvae with the administration of *B. plicatilis* without enrichment newly formed in D8. The enrichment of *B. plicatilis* with the combination of fish oil and squid oil can accelerate the growth of absolute length of the grouper larvae by 3.27 ± 0.17 mm.

Acknowledgement

A high gratitude to Faculty of Fisheries and Marine,

Universitas Airlangga which has given permissions to the authors to use their laboratories for research purpose.

References

- Afrianto, E. and Liviawaty, E. 2005. *Pakan Ikan Dan Perkembangannya*. Yogyakarta: Kanisius.
- Akbary, P., Hosseini, S. A. and Imanpoor, M. R. 2011. Enrichment of *Artemia nauplii* with essential fatty acids and vitamin C: Effect on rainbow trout (*Oncorhynchus mykiss*) larvae performance. *Iranian Journal of Fisheries Sciences*. 10(4) : 557-569.
- Bulanin, U. *et al.* 2018. Perkembangan embrio dan penyerapan kuning telur larva ikan kerapu bebek', *Mangrove dan Pesisir*. III(3) : 16-23.
- Ch'ng, C. L. and Senoo, S. 2008. Egg and Larval Development of a New Hybrid Grouper, Tiger Grouper *Epinephelus fuscoguttatus* x Giant Grouper *Epinephelus lanceolatus*. *Aquaculture Science*. 56(4): 505-512.
- Chapkin, R. S. 2008. Bioactive dietary long-chain fatty acids: emerging mechanisms of action. *British Journal of Nutrition*. 100(6) : 1152. doi: 10.1017/S0007114508992576.
- Chu, I. K. C. 2016. Effect of salinity on embryonic development and hatching of hybrid grouper, *Epinephelus fuscoguttatus* x *epinephelus lanceolatus*. *AACL Bioflux*. 9(6) : 1278-1285.
- Dabrowski, K. and Bardega, R. 1984. Mouth size and predicted food size preferences of larvae of three cyprinid fish species. *Aquaculture*. 40 : 41-46.
- Effendi, I. 2004. *Pengantar Akuakultur*. Penebar Swadaya.
- Glamuzina, B. 1998. Egg and early larval development of laboratory reared dusky grouper. *Scientia Marina*. 62(4) : 373-378.
- Halver, J. E. and Hardy, R. W. 2002. *Fish Nutrition*. Academic Press.
- Ismi, S. 2012. *Cromileptes altivelis*. Secara Terkontrol', pp. 407-408.
- Ismi, S., Asih, Y. N. and Kusumawati, D. 2014. Peningkatan Produksi dan Kualitas Benih Kerapu dengan Program Hybridisasi Improvement of Seed Production and Quality of Grouper by Hybridization. *Jurnal Oseanologi Indonesia*. 1 (1) : 1-5.
- Kementrian Kelautan and Perikanan Republik Indonesia 2013. *Ditjen Perikanan Budidaya*. Available at: <https://kkp.go.id/artikel/1180-ditjen-perikanan-budidaya>.
- Kohno, H. 1998. Early Life History Features Influencing Larval Survival of Cultivated Tropical Finfish', in *Tropical Mariculture*. Elsevier, pp. 71-110. doi: 10.1016/B978-012210845-7/50003-9.
- Leaver, M. J. 2008. Towards fish lipid nutrigenomics: current state and prospects for fin-fish aquaculture. *Reviews in Fisheries Science*. Taylor & Francis,

- 16(sup1) : 73–94.
- Léger, P. 1986. The use and nutritional value of *Artemia* as a food source. *Oceanogr. Mar. Biol. Ann. Rev.* 24: 521–623.
- Moyle, P. B. and Cech, J. J. 2004. *Fishes: An Introduction to Ichthyology*. Pearson Prentice Hall.
- Peng, J. 2003. Polyunsaturated fatty acid profiles of whole body phospholipids and triacylglycerols in anadromous and landlocked Atlantic salmon (*Salmo salar* L.) fry. *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology*, 134(2), pp. 335–348.
- Purwaningsih, S., Salamah, E. and Dewantoro, R. 2014. Chemical Composition and Fatty Acids of Glodok Fish by High Thermal Processing. *Jurnal Pengolahan Hasil Perikanan Indonesia*. 17(2).
- Redjeki, S. 1999. Budidaya Rotifer (*Brachionus plicatilis*). *Jurnal Oseana*. 24 : 27-43.
- Saleh, J. H., Al-Zaidi, F. M. and Al-Faiz, N. A. 2013. Effect of pH on hatching and survival of Larvae of common carp *Cyprinus carpio* (Linnaeus, 1758). *Marsh Bulletin*. Basrah University, 8(1) : 58–64.
- Setyadi, I. 2008. Produksi Masal Larva Ikan Kerapu Pasir (*Epinephelus Corallicola*) dengan Ukuran Bak Berbeda', *Neptunus*, 14(1).
- Tocher, D. R. 2003. Metabolism and functions of lipids and fatty acids in teleost fish. *Reviews in Fisheries Science*. Taylor & Francis. 11(2) : 107–184.
- Watanabe, T. 1993. Importance of docosahexaenoic acid in marine larval fish. *Journal of the World Aquaculture Society*. Wiley Online Library, 24(2) : 152–161.
- Watanabe, T., Kitajima, C. and Fujita, S. 1983. Nutritional values of live organisms used in Japan for mass propagation of fish: a review. *Aquaculture*. Elsevier, 34(1–2) : 115–143.
- Winarno, F. G. 1984. *Kimia Pangan dan gizi*. P.T. Gramedia.
-
-