

5

by Mat Taib Mimi Iryanib

Submission date: 29-Dec-2020 04:28PM (UTC+0800)

Submission ID: 1481838623

File name: oro-RNA_interference_of_Hsp70_in_Artemia_franciscana_nauplii.pdf (1.18M)

Word count: 5974

Character count: 32665



1 RNA interference of Hsp70 in *Artemia franciscana* nauplii and its effect on morphology, growth, survival and immune response



Mat Taib Mimi Iryani^b, Thomas H. MacRae^c, Patrick Sorgeloos^{b,d},
 Tengku Sifzizul Tengku Muhammad^b, Muhd Danish-Daniel^b, Min Pau Tan^b,
 Woro Hastuti Satyantini^f, Mohd Effendy Abd Wahid^{b,e}, Jingfeng Sun^a, Aijun Lv^a,
 Yeong Yik Sung^{a,b,f,*}

^a Tianjin Key Lab of Aqua-ecology and Aquaculture, Fisheries College, Tianjin Agricultural University, Tianjin 300384, China

^b Institute of Marine Biotechnology, Universiti Malaysia Terengganu (UMT), 21030 Kuala Nerus, Malaysia

^c Department of Biology, Dalhousie University, Halifax, NS B3H 4R2, Canada

^d Lab of Aquaculture and Artemia Reference Center, Department of Animal Sciences and Aquatic Ecology, Campus Coupure, Coupure Links 653 - block F, B-9000 Gent, Belgium

^e School of Fisheries and Aquaculture Sciences, Universiti Malaysia Terengganu (UMT), 21030 Kuala Nerus, Malaysia

^f Department of Fish Health Management and Aquaculture, Faculty of Fisheries and Marine, Universitas Airlangga, 60115 Surabaya, Indonesia

ARTICLE INFO

Keywords:

Hsp70

Artemia franciscana

Nauplii

RNA interference (RNAi)

ABSTRACT

1 Brine shrimp *Artemia franciscana* release either encysted embryos via oviparous reproduction or motile nauplii via ovoviviparous reproduction. *Artemia* cysts contain an abundant amount of ArHsp21, ArHsp22, artemin, and p26, small heat shock proteins which regulate diapause and stress tolerance of this branchiopod crustacean. However, not much is known of the role of molecular chaperone, Hsp70 in this animal. In this study, the immunostimulatory role of Hsp70 in *A. franciscana* was investigated, work which included examining the function of this protein in embryo development, growth, survival and stress tolerance. Knockdown of Hsp70 was successfully performed by RNA interference (RNAi), with specific Hsp70 dsRNA construct delivered to female broods by microinjection to knockdown the protein in nauplii. The loss of Hsp70 neither affected the growth nor morphology as *Artemia* nauplii hatched normally and thereafter grew to adult within 28 days, observations similar to the control animals, the latter injected with GFP dsRNA. The survival of nauplii lacking Hsp70 appeared lower upon 28 days culture but the differences were insignificant when comparisons were made with the controls ($P > .05$). On other aspect, knockdown of Hsp70 reduced approximately 31% survival upon *Vibrio campbellii* challenge, indicating that Hsp70 assisted in the tolerance of *Artemia* nauplii during pathogen infection, perhaps through immune stimulation. Initial findings revealed that Hsp70 knockdown did not affect the expression of prophenoloxidase (proPO), a crucial immune protein of the shrimp innate immunity. By examining the morphology, growth, viability and molecular changes upon Hsp70 elimination, this study contributes to our understanding of the important roles of Hsp70 in *A. franciscana*, an important live food species used in aquaculture.

34

1. Introduction

The brine shrimp *Artemia* is a crustacean highly adapted to living in hypersaline waters of inland salt lakes, coastal lagoons and solar saltworks (Kappas et al., 2004; Persoone and Sorgeloos, 1980). *Artemia* occur in many colours, from white to pink, shadow or green with females appear pink or orange (King, 2013). Males are usually grey in color and distinguished from females based on several features, notably their anterior pincers and lack of an egg sac at sexual maturity (King,

2013). Adult *Artemia* are generally 1 cm long with females longer than males. Reproduction may proceed in two pathways: ovoviviparous or oviparous. Females release free swimming nauplii via ovoviviparous pathway, which then undergo a series of molts before reaching adulthood (Liang and MacRae, 1999). Under unfavorable or extremely critical environmental conditions, females produce a highly resistant encysted gastrula embryo, known as cyst via oviparous pathway.

Heat shock proteins (Hsp70) have been studied extensively and are associated with the protection of *Artemia* against abiotic and biotic

26

* Corresponding author at: Institute of Marine Biotechnology, Universiti Malaysia Terengganu (UMT), 21030 Kuala Nerus, Malaysia

E-mail address: yeong@umt.edu.my (Y.Y. Sung).

12

<https://doi.org/10.1016/j.aquaculture.2020.735012>

Received 1 August 2019; Received in revised form 14 January 2020; Accepted 23 January 2020

Available online 23 January 2020

0044-8486/© 2020 Published by Elsevier B.V.

stress. For example, exposure of *A. franciscana* nauplii to a non-lethal heat shock from 28 °C to 37 °C for 30 min with 6 h recovery enhances Hsp70 accumulation and promoted heat tolerance at 42 °C (Sung et al., 2008). The tolerance of *Artemia* against virulent *V. campbellii* and *V. proteolyticus* were enhanced, with survival increased two-fold when compared to the non-heated controls (Sung et al., 2007). As Hsp70 is involved in the immune response, the immunoregulatory roles of Hsp70 during shrimp disease and infection were examined. Hsp70 potentially function in shrimp immunity through activation of Toll like receptors (TLRs). As a ligand for TLRs, Hsp70 binds pathogen-associated molecular pattern (PAMP) molecules and modulates PAMP-induced Toll-like receptor (TLR) signaling, transferring signals to cells of the innate immune system and thereby promoting pathogen recognition and elimination (Sung et al., 2018; Sung and MacRae, 2011).

RNAi is a gene silencing technique in which dsRNA related in sequence to a specific mRNA causes degradation of that mRNA, enabling a specific gene to be silenced (Chopra et al., 2002). This powerful tool was first discovered during a study of the biochemical nature of inducers of gene silencing in the nematode *Caenorhabditis elegans*, mediated by dsRNA molecules (Aigner, 2006; Fire et al., 1998). To further examine the function of *Artemia* Hsp70 in stress tolerance and immune response, a protocol to knockdown this particular protein by way of RNAi was developed (Iryani et al., 2017). In this context, Hsp70 knockdown nauplii were produced by injection of Hsp70 dsRNA into the berried females of *Artemia* to eliminate both the Hsp70 mRNA and protein in nauplii released from these females. This protocol facilitated the study on the role of Hsp70 in the protection of *A. franciscana* against abiotic and biotic stressors. An established molecular techniques like RNAi makes *A. franciscana* an exceptional model organism to explore many basic studies such as biological evolution (Copf et al., 2006), gene function (King and MacRae, 2012) and stress tolerance (Iryani et al., 2017; Toxopeus et al., 2014). It was demonstrated through previous study that Hsp70 knockdown nauplii were less resistant to heat perturbation and pathogenic *Vibrio campbellii*, with survival percentage reduced approximately 41% and 34%, respectively in challenge assays (Iryani et al., 2017). The role and association of Spalt with Hox gene was studied using the RNAi, with results indicated that Spalt acts as a repressor of Hox gene expression and knocking down Spalt expression caused various segmental anomalies in *A. franciscana* (Copf et al., 2006). Gnotobiotic *A. franciscana* were used to investigate whether a phenolic compound, phloroglucinol is effective against *V. parahaemolyticus* and pretreatment of Hsp70-knockdown *Artemia* larvae with this compound indicated that phloroglucinol-induced Hsp70 mediates the survival of *Artemia* against *V. parahaemolyticus* (Kumar et al., 2018).

In this study, the induction of Hsp70 dsRNA into the egg sacs of *Artemia* females significantly reduced Hsp70 content in the released nauplii. Protein knockdown by dsRNA microinjection was successful in *Artemia* with caudal dsRNA (Copf et al., 2004), *C. pipiens* with catalase dsRNA (Sim and Denlinger, 2011), *Artemia* females with p26 dsRNA (King and MacRae, 2012) and *Artemia* females with AfrLEA1 dsRNA (Toxopeus et al., 2014). *Artemia* embryos with reduced amounts of Hsp70 developed normally, yielding nauplii similar morphology to those with normal amounts of Hsp70. Hsp70-knockdown nauplii however, displayed slower growth during the culture period. Hsp70 was absent in the nauplii but the expression resumed upon reaching juvenile and adult stage. The survival of *Artemia* lacking Hsp70 in the absence of infection was lower when compared to the control 28 days upon culture. This result suggested that Hsp70 plays crucial roles in cell maintenance, growth and survival of *Artemia*. Real time-PCR analysis revealed that there was no apparent difference between the expression levels of proPO with the control, whereas the expression level of Hsp70 is down-regulated. The number of proPO transcript was not elevated in *Artemia* without bacterial exposure. Upon *V. campbellii* infection, survival of nauplii lacking Hsp70 decreased, indicating that Hsp70 protects *Artemia* against *Vibrio* infection. This work makes fundamental but

important contributions to our understanding on the role of Hsp70 in *A. franciscana* and determining the Hsp70 function by evaluating effects of Hsp70 knockdown in nauplii could lead to the formulation of strategies to protect *Artemia* and perhaps other crustaceans against stress and disease.

2. Materials and methods

2.1. Culture of *A. franciscana*

A. franciscana cysts (INVE Aquaculture) were hatched in 30 mg/l of seawater with vigorous aeration and constant illumination for 24 to 48 h. Nauplii were collected with a 200 µm plankton net and grown to adult stage at 28 °C with constant aeration. Nauplii were fed daily with PKC Nutri+*, a formulated feed developed at Universiti Malaysia Terengganu (UMT) for use in *Artemia* biomass production. Adult *Artemia* were collected 28 days after for microinjection.

2.2. Preparation of Hsp70 dsRNA

pRSET-C plasmid (Invitrogen, Burlington, ON., Canada) containing *Artemia* Hsp70 cDNA were transformed into *E. coli* competent cells (Invitrogen, USA) and colonies grown on LB-Amp agar were cultured in LB-Amp broth at 37 °C prior to extraction. Hsp70 cDNA were harvested from 5 ml overnight cultures of *E. coli* using miniprep kit (PureLink™ Quick Plasmid Miniprep Kit, Invitrogen, USA). The Hsp70 cDNA was amplified using forward (5'-TAATACGAC TCACTATAGGATTCTCAA AGACAAGC-3') and reverse (5'-TAATAC GACTCACTAT GCATAGA GCITGGTAAT-3') primers, each specific for Hsp70 and containing the T7 promoter, TAATACGACTCACTATAGG, at the 5'-end (Bioline, UK). PCR was performed using Immix™ (Bioline, UK) at the manufacturer suggested concentration as follows: 5 min at 94 °C followed by 30 cycles of 94 °C for 30 s, 55 °C for 30 s and 72 °C for 1 min, followed by 10 min at 72 °C. The PCR product was resolved in 1.7% (w/v) agarose gels in 1 × TBE (20 mM TRIS, 10 mM acetic acid, 0.5 mM EDTA, pH 8.5) at 90 V for 45 min and stained with ethidium bromide (Sigma Aldrich, USA) prior to visualization via a Gel Doc™ XR + with Image Lab™ Software (BioRad, USA). The PCR product was used as template to generate dsRNA using the MEGAscript® RNAi kit (Ambion Applied Biosystems, USA). Green fluorescent protein (GFP) dsRNA was generated as control solution using forward (5'-TAATACGACTCA CTAAGGGAGACACATGAAGCAGCAGCAGCT-3') and reverse (5'-TAAT ACGACTCACTATAGGGAGAAAGTTCACCTTGATGCCGTTCC-3') primers, with PCR conditions as described above except annealing temperature was at 60 °C. The GFP cDNA used as template was cloned in the vector pEGFP-N1 (Clontech, Mountain View, CA, USA) (King and MacRae, 2012).

2.3. Injection of *A. franciscana* females with dsRNA

Adult *A. franciscana* females with unfertilized eggs sacs and highly possible to produce nauplii were chosen for dsRNA injection (Iryani et al., 2017). They were differentiated from females destined to produce cysts by the absence of a shell and (Liang and MacRae, 1999). dsRNA was mixed with 0.5% (w/v) phenol red in Dulbecco's phosphate buffered saline (DPBS) (Sigma Aldrich, USA). *Artemia* females were immobilized on cooled 3% (w/v) agar and Hsp70 dsRNA solution containing approximately 100 ng of dsRNA was injected into the egg sacs with a micromanipulator (InjectMan®, Eppendorf, Germany). Injected females were placed in 6 well plates and monitored for at least 2 h. Females exhibiting normal swimming, straight tails and dye retention were used in the experiment. Males were added to females 24 h post-injection and females were monitored with a stereomicroscope for fertilization, noted by the fusion of egg sacs (King and MacRae, 2012) and for the release of nauplii.

2.4. Detection of Hsp70 in nauplii, juvenile and adult *A. franciscana*

Protein were extracted from 40 nauplii released from *A. franciscana* females injected with either Hsp70 dsRNA or GFP dsRNA as described in previous study with minor modification (Sung, 2014). Sample was rinsed in cold distilled water and homogenized in cold Buffer K (150 mM sorbitol, 70 mM potassium gluconate, 5 mM MgCl₂, 5 mM NaH₂PO₄, 40 mM HEPES, pH 7.4) (Sung et al., 2007; Clegg et al., 2000) supplemented with Protease Inhibitor Cocktail (Sigma-Aldrich Inc-P8849, USA) as recommended by the manufacturer. The mixture was centrifuged at 4 °C for 5 min and the supernatant was transferred to a new tube. SDS polyacrylamide gel electrophoresis, aliquots of homogenate were combined with equal volumes of SDS sample buffer, vortexed and heated at 95 °C for 5 min (Laemmli, 1970). The samples were cooled and centrifuged at 2200 xg for 1 min. Fifteen (15) µl of each protein sample was applied per lane of 10% SDS polyacrylamide gels prior to electrophoresis at 120 V for 15 min and then at 150 V for 45 min. One gel was stained with Coomassie Biosafe (BioRad, USA) and the other transferred to polyvinylidene fluoride transfer membrane (BioRadImmuno-Blot™ PVDF, USA) for antibody probing. PVDF membrane were probed in mouse monoclonal antibody to Hsp70 (SMC-164D) (StressMarq, Canada), diluted 1:5000 in Tris (pH 7.6) for 2 h. Membrane was then washed in Trisaline (pH 7.6) prior to incubation in polyclonal HRP-conjugated goat anti-mouse IgG (Bioreagent-SAB-100 J) (Stressgen, Canada) diluted 1:5000 in Tris. For detection, antibody-reactive proteins were detected with 0.7 mM diaminobenzidine tetrahydrochloride dihydrate (DAB) used as substrate in association with 0.01% (v/v) H₂O₂ in 0.1 M Tris-HCL (pH 7.6). Detection of Hsp70 in either juvenile or adult stage was performed as described, with extracts from 1 whole animal used for SDS-PAGE and WB.

2.5. Densitometry analysis

For densitometry analysis, blots were scanned with a GS-800 calibrated densitometer (BioRad Laboratories, USA) and quantification was performed by measuring protein bands from the control and treatments with Quantity One software (BioRad Laboratories, USA). The amount of protein were interpreted as reflective density/mm², with the density value generated by the software.

2.6. Phenotypic modification, growth development and survival of nauplii upon knockdown

Nauplii released from females injected with Hsp70 dsRNA were observed with a stereomicroscope for 28 days to determine if they were morphologically normal compared to the nauplii released from females injected with control solution. Nauplii length was determined using a Nikon Profile Projector V-12B fixed with a Nikon Digital Counter SC-212 at 7 days intervals. Thirty nauplii were collected and measured each week.

Thirty nauplii released from females injected with dsRNA for either Hsp70 or GFP were incubated in seawater at 28 °C for 28 days. Nauplii were fed and observed daily to determine the number of surviving animals. Dead nauplii were removed after counting. The experiment was performed two times in triplicate.

2.7. Quantification of Hsp70 and ProPO mRNAs in Hsp70-knockdown nauplii

Total RNA were extracted from 20 nauplii using TRIsure™ reagent (Bioline, UK) by following the manufacturer's instructions with minor modification. Nauplii released from *A. franciscana* females injected with dsRNA specific to either Hsp70 or GFP were rinsed with ice cold distilled water, homogenized in 100 µl of TRIsure and the mixture incubated at room temperature for 5 min. Twenty (20) µl of chloroform were added to the sample, vortex vigorously for 15 s and incubated for

Table 1

Primers specific for Hsp70, Prophenoloxidase and β-actin.

Gene	Sequences of forward and reverse primers
Hsp70 (F)	5'-GATTCTCAAAGACAAGC-3'
Hsp70 (R)	5'-GCATAGAGCTTGGTAAT-3'
Prophenoloxidase (F)	5'-CGCTGGCATAAGCACATCGATG-3'
Prophenoloxidase (R)	5'-GTCATTCTCACTGTGAAAGC-3'
β-actin (F)	5'-GGTCGTGACTTGACGGACTATCT-3'
β-actin (R)	5'-AGCGGTGCCATTCTTGT-3'

2 min at room temperature. The samples were centrifuged at 12,000 xg for 15 min at 4 °C. The upper colorless aqueous phase which contains the RNA were transferred to a fresh tube prior to RNA isolation. First strand cDNA was synthesized using Tetro cDNA Synthesis Kit (Bioline, UK). The premix was prepared on ice (Mix 1), mixed gently by pipetting and incubated at 65 °C for 22 min before placed on ice for 2 min. The second mixture (Mix 2) was added and the sample was incubated at 42 °C for 45 min followed by a 15 min incubation at 70 °C.

qPCR was performed with 2 µl of cDNA (100 ng) using 2× SensiMix SYBR No-ROX Kit (Bioline, UK) in a CFX Connect™ System (Bio-Rad, USA) and primers for the immune related genes of *A. franciscana*, Hsp70 and prophenoloxidase (Baruah et al., 2015) with β-actin served as the house-keeping gene (Jiang et al., 2007) (Table 1). Amplification was performed at 95 °C for 10 min followed by 40 cycles of 95 °C for 5 s and 60 °C for 31 s. The cycle threshold (CT) values and fold difference for each immune-related genes was calculated according to the 2-ΔΔCt method provided by the CFX Manager Software (Bio-Rad, USA) (Clegg et al., 2000). Data generated in CFX Connect™ System were analyzed with the Bio-Rad CFX (Gene Expression Analysis) software and data were normalized against β-actin.

2.8. Tolerance of nauplii to *Vibrio campbellii*

Thirty nauplii from *Artemia* females injected with either Hsp70 dsRNA or control solution were transferred into Falcon tubes containing 30 ml of filtered, autoclaved seawater. Nauplii were challenged by immersion to *Vibrio campbellii* at 1 × 10⁸ cells/ml (Iryani et al., 2017) and vibrio tolerance was determined 24 h after challenge by counting actively swimming nauplii. The experiment was performed two times, each in triplicate.

3. Results

3.1. Hsp70 expression in different developmental stages of *A. franciscana*

As revealed by SDS gel electrophoresis and immunoprobings of western blots, Hsp70 was absent in *Artemia* nauplii. The administration of Hsp70 dsRNA by injection decreased Hsp70 accumulation in *Artemia* nauplii, indicating that the synthesis of Hsp70 in *Artemia* nauplii was obscured, an expected outcome when the dsRNA used were constructed and delivered appropriately. Though Hsp70 was undetectable in nauplii, the synthesis increased in juvenile and adult stages upon 28 days, revealing that knockdown of this protein was transient (Fig. 1). The Hsp70 content in juvenile *Artemia* produced from female broodstock receiving Hsp70 dsRNA was augmented 0.3 fold over time and the adults accumulated Hsp70 levels similar to those injected with GFP dsRNA (control) upon 28 days culture (Table 2).

3.2. Morphology of *Artemia* Lacking Hsp70

Microscopic observation of nauplii released from females injected with Hsp70 dsRNA displayed normal morphological structure, as with those released from control females (Fig. 2). It was noted that the growth of Hsp70 knockdown nauplii was slightly slower than the control, but the differences were insignificant ($P > .05$).




Fig. 1. Hsp70 expression in different growth stages of *A. franciscana*. Protein extracts prepared from different stages of *Artemia* were resolved by electrophoresis in SDS polyacrylamide gels and either stained with Coomassie brilliant blue or blotted to PVDF membrane and probed with antibody SMC-164. The secondary antibody was HRP-conjugated goat anti-mouse IgG. Antibody reactive protein was detected by DAB reagent. A & D, nauplii; B & E, juvenile; C & F, adult. 1, 13 kDa marker; 2, GFP dsRNA; 3, Hsp70 dsRNA. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

Table 2
Relative density of Hsp70 in different developmental stages of *A. franciscana*.

Nauplii	Adj. Vol. (RD)	Relative quantity
Control	2294.56	–
Hsp70 dsRNA	–	–
Juvenile	Adj. Vol. (RD)	Relative Quantity
Control	3124.56	1
Hsp70 dsRNA	890.80	0.3
Hsp70 dsRNA	–	–
Adult	Adj. Vol. (RD)	Relative Quantity
Control	1908.39	1
Hsp70 dsRNA	2143.09	1.1

3.3. Growth and survival of *A. franciscana* nauplii lacking Hsp70

Nauplii released from females receiving dsRNA for Hsp70 grown to adult and their morphology were similar to the control animals. The final length of Hsp70 knockdown *Artemia* was 11.3 ± 1.3 mm, whereas the control animals reached 12.1 ± 1.6 mm upon 28 days culture (Fig. 3). Approximately 43% of control nauplii (released from GFP dsRNA injected-females) survived 28 days upon culture whereas those lacking Hsp70 were maintained at 23% (Fig. 4).

3.4. Immune genes expression upon Hsp70 knockdown in nauplii

Hsp70 mRNA in nauplii released from *Artemia* females injected with Hsp70 dsRNA reduced approximately 4.2 fold. Prophenoloxidase gene transcripts in Hsp70 knockdown nauplii were augmented by 0.3 fold, but the differences were insignificant when compared to the controls, indicating that regulation of this immune gene was not affected by Hsp70 knockdown (Fig. 5).

3.5. *Vibrio* tolerance of Hsp70-knockdown nauplii

The 24 h survival of nauplii released from females injected with GFP dsRNA and subsequently challenged with 1×10^8 *V. campbellii*/ml was approximately 78%. In similar experimental conditions, the survival of nauplii lacking Hsp70 reduced to 31% (Fig. 6). The significance in

survival reduction strongly indicated that Hsp70 contributes to *V. campbellii* tolerance.

4. Discussion

RNAi was used in this study to elucidate Hsp70 function in *Artemia*, with particular emphasis on morphological changes, growth and survival upon Hsp70 knockdown in nauplii-destined embryos, work which included verification of prophenoloxidase expression, an important shrimp innate immune component to battle pathogens. Administration of Hsp70 dsRNA into the egg sacs specifically knocked down Hsp70 in *Artemia* females, an outcome consistent with our previous study (Iryani et al., 2017). *A. franciscana* nauplii with reduced amounts of Hsp70 developed normally, having morphology similar to those produced from females injected with GFP dsRNA, with neither malformation nor mutation noticed. Nauplii lacking Hsp70 grew to adult stage within 12 to 14 days upon given a standardized feeding regime with PKC-Nutri +*, suggesting that Hsp70 is unessential for *Artemia* growth at early developmental stages. Nevertheless, based on the outcome of our previous study, fertilization rate and the time used to release nauplii were slightly affected by the loss of Hsp70 (Iryani et al., 2017).

Further investigation into the Hsp70 content across different stages of *Artemia* revealed that this group of stress protein, initially undetected in nauplii stage, increased in juveniles and adults upon RNAi, indicating that Hsp70 knockdown with this method is transient. To compare, knockdown of p26 diapause-destined *Artemia* embryos lasted for at least four broods from the same female but the RNAi effect does not carry over into subsequent generations (King and MacRae, 2012). Various factors influence the efficiency of RNAi. A GFP knockdown with transfection of GFP-dsRNA into the fall armyworm *Spodoptera frugiperda* cells and subsequently infected with recombinant baculovirus carrying GFP gene revealed fluorescence inhibition of 60% and 30% at day 5 and 9, respectively (Flores-Jasso et al., 2004). The reduction in the inhibition efficiency might be due to the specificity, length and quantity of dsRNA, delivery method as well as target tissue site, developmental stage and host's characteristics (Yang et al., 2018). Interestingly, RNAi in the nematode *C. elegans* lasted for generations, and this effect was most probably due to the presence of RNA-directed RNA polymerases (RdRP), a crucial enzyme required in the RNA




Fig. 2. Morphology of *A. franciscana* nauplii lacking Hsp70. Microscopic observation using a stereomicroscope was employed for 28 days to determine if nauplii released from females injected with dsRNA for Hsp70 were morphologically normal compared to nauplii released from females injected with dsRNA for GFP. C, GFP dsRNA; T, Hsp70 dsRNA. C1 & T1, day 1; C7 & T7, day 7; C14 & T14, day 14; C21 & T21, day 21; C28 & T28, day 28.




Fig. 3. Knockdown of Hsp70 does not affect *A. franciscana* growth. Size of nauplii released from females injected with dsRNA for Hsp70 and GFP were determined at 7 days intervals for 28 days. Thirty nauplii were collected and measured each week for both knockdowns. Bars represent length of *Artemia*. The error bars represent the corresponding standard deviation (SD) with ($P > .05$).




Fig. 4. Hsp70 knockdown *Artemia* exhibited reduced survival. Nauplii released from females injected with dsRNA for Hsp70 and GFP were incubated at 28 °C and surviving (swimming) nauplii were counted for 28 days. The experiment was done two times for both knockdowns with 30 nauplii in each replicate. Lines represent survival percentage of *Artemia*. The error bars represent the corresponding standard deviation (SD) from three replicates with ($P > .05$).




Fig. 5. The effect of Hsp70 knockdown on gene expression of *A. franciscana* nauplii. The relative gene expression of Hsp70 and proPO genes was quantified by qRT-PCR in nauplii *Artemia*. The value at each point represents the fold change of mRNA expression (normalized against internal control β -actin) relative to the value of control. The error bars represent the corresponding standard deviation (SD) from three replicates independently prepared after knockdown. Asterisks denote statistically significant differences between treatments with ($P < .05$).

silencing pathway of many invertebrates (Fire et al., 1998).

In this study, *Artemia* with reduced amounts of Hsp70 appeared slightly shorter than the control in the early developmental phase. However, the total length upon reaching 28 days culture were




Fig. 6. Hsp70 protected *A. franciscana* nauplii against pathogenic *V. campbellii*. Nauplii released from females injected with either Hsp70 dsRNA or GFP dsRNA were exposed to 1×10^8 *Vibrio campbellii*/ml for 24 h. *Vibrio* tolerance was determined by counting surviving (swimming) nauplii. The experiment was done two times with 30 nauplii in each replicate. The bars in the graph represent survival percentages and the error bars represent the corresponding standard deviation (SD) from three replicates within each experiment. Asterisks denote statistically significant differences between treatments with ($P < .05$).

statistically indifferent than the controls. Knockdown of Hsp70 had no apparent effect on *Artemia* survival, though it was initially noted that the survival of *Artemia* lacking Hsp70 was lower than those accumulating normal amounts of the protein. Hsp70 are crucial in maintaining several biological processes such as protein folding and signaling which affect gene expression. Reducing the amount of Hsp70 decrease cellular protein folding efficiency and this condition may adversely affect the general well-being of *Artemia* and their tolerance against stress, resulting in low survival. Mortality was not observed in this study possibly because only one Hsp70 isotype was knocked down by the dsRNA. Several Hsp70 isotypes occur in *Artemia* and they may play a role in sustaining growth and survival (Jung et al., 2019). Hence, it is interesting to determine the number of Hsp70 isotypes in *Artemia* and examine their function(s) in stress tolerance.

Hsp70 was linked to several pathways of the vertebrate immune system. In *Artemia*, augmentation of Hsp70 by a non-lethal heat shock protected against *V. cholerae* and *V. proteolyticus* (Sung et al., 2008, 2007). Additionally, feeding *Artemia* with either *Artemia* Hsp70 or DnaK, a bacterial Hsp70, provided similar protection, boosting survival two-fold against *V. campbellii* challenge (Sung et al., 2009). The increase in proPO mRNA led to the assumption that this immune protein was associated with the enhanced tolerance (Baruah et al., 2011). ProPO typically confers protection by melanization and coagulation, two processes crucial in pathogen elimination in invertebrates, including brine shrimp *Artemia* (Norouzitallab et al., 2016). Specifically, the activation of the proPO system by pathogen pattern-recognition proteins (PPRPs), activates a serine proteinase cascade, eventually leading to the cleavage of the inactive proPO to the active PO that functions to produce the melanin and toxic reactive intermediates against invading pathogens (Amparyup et al., 2013). Using *Artemia* as a model organism, the link between Hsp70 and ProPO in this crustacean species was examined. As revealed by qRT-PCR, the number of ProPO mRNA in Hsp70-knockdown and control nauplii were insignificantly different, revealing that Hsp70 knockdown, under the experimental condition performed in this study, does not affect proPO expression. Interestingly, it was observed that the survival of nauplii containing Hsp70 upon 24 h challenge with *V. campbellii* was significantly higher than those lacking Hsp70, indicating that protection against pathogenic *Vibrio* in *Artemia* is perhaps due solely to Hsp70 accumulation and/or other immune proteins which were not examined in this study. In fact, many studies have revealed that Hsp70 accumulation boosts aquatic species tolerance against disease and infection other than *Artemia*. As one example, priming of adult mussels *P. viridis* with a 30 min NLHS at 38 °C followed

by 6 h recovery increased resistance to *V. alginolyticus*. The induction of Hsp70 in parallel with enhanced thermotolerance and improved protection against *V. alginolyticus*, suggested Hsp70 functions in *P. viridis* as a molecular chaperone and as a stimulator of the immune system (Aleng et al., 2015).

Whatever the outcome, this study represents the first to demonstrate the effect of Hsp70 knockdown on morphology, growth, survival and immune response of the brine shrimp *Artemia*. Considering the massive economic losses incurred by disease outbreak in the shrimp culture industry, studies focusing on Hsps have become increasingly popular because of their putative regulatory function in the immune system. Understanding the role of Hsp70 in *Artemia* and/or other shrimp species may assist in the formulation of strategies to boost stress and disease tolerance during aquaculture.

Acknowledgements

This work was partially supported by the Fundamental Research Grant Scheme (FRGS) 59331 and No. 59578 from the Ministry of Education, Malaysia and the 1000-Young Talent Professors Grant Scheme, China to YYS, a Natural Sciences and Engineering Research Council Discovery Grant, Canada to THM and Professorial Chair of Sultan Mizan Arch Fund No. 53264 to PS. MTMI was supported by MyBrain 15 Scholarship from the Ministry of Higher Education, Malaysia.

Declaration of Competing Interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

- Aigner, A., 2006. Gene silencing through RNA interference (RNAi) in vivo: strategies based on the direct application of siRNAs. *J. Biotechnol.* 124, 12–25.
- Aleng, N.A., Sung, Y.Y., MacRae, T.H., Abd Wahid, M.E., 2015. Non-lethal heat shock of the Asian green mussel, *Perna viridis*, promotes Hsp70 synthesis, induces thermotolerance and protects against *Vibrio* infection. *PLoS One* 10 (8), e0135603.
- Amparyup, P., Charoensapri, W., Tassanakajon, A., 2013. Prophenoloxidase system and its role in shrimp immune responses against major pathogens. *Fish Shellfish Immunol.* 34, 990–1001.
- Baruah, K., Ranjan, J., Sorgeloos, P., MacRae, T.H., Bossier, P., 2011. Priming the prophenoloxidase system of *Artemia franciscana* by heat shock proteins protects against *Vibrio campbellii* challenge. *Fish Shellfish Immunol.* 31, 134–141.
- Chopra, M., Pachuk, C., Satishchandran, C., Giordano, T., 2002. Using RNA interference to modulate gene expression. *TARGETS.* 1 (03), 1–7.
- Clegg, J.S., Susan, A., Jackson, N.V.H., Sorgeloos, P., 2000. Thermal resistance, developmental rate and heat shock proteins in *Artemia franciscana* from San Francisco Bay and southern Vietnam. *J. Exp. Mar. Biol. Ecol.* 252, 85–96.
- Copf, T., Schröder, R., Averof, M., 2004. Ancestral role of caudal genes in axis elongation and segmentation. *Proc. Natl. Acad. Sci. U. S. A.* 101, 17711–17715.
- Copf, T., Rabet, N., Averof, M., 2006. Knockdown of Spalt function by RNAi causes de-repression of Hox genes and homeotic transformations in the crustacean *Artemia franciscana*. *Dev. Biol.* 298, 87–94.
- Fire, A., Xu, S.Q., Montgomery, M.K., Kostas, M.K., Driver, S.E., Mello, C.C., 1998. Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*. *Nature.* 391, 806–811.
- Flores-Jasso, C.F., Valdes, V.J., Sampieri, A., Valadez-Graham, V., Recillas-Targa, F., Vaca, L., 2004. Silencing structural and nonstructural genes in baculovirus by RNA interference. *Virus Res.* 102, 75–84.
- Iryani, M.T.M., MacRae, T.H., Panchakshari, S., Tan, J., Bossier, P., Wahid, M.E.A., 2017. Knockdown of heat shock protein 70 (Hsp70) by RNAi reduces the tolerance of *Artemia franciscana* nauplii to heat and bacterial infection. *J. Exp. Mar. Biol. Ecol.* 487, 106–112. <https://doi.org/10.1016/j.jembe.2016.12.004>.
- Jiang, L., Hou, L., Zou, X., Zhang, R., Wang, J., Sun, W., Zhao, X., An, J., 2007. Cloning and expression analysis of p26 gene in *Artemia sinica*. *Acta Biochim. Biophys. Sin.* 39 (5), 351–358.
- Junprung, W., Norouzitallab, P., De Vos, S., Tassanakajon, A., Viet, D.N., Van Stappen, G., Bossier, P., 2019. Sequence and expression analysis of HSP70 family genes in *Artemia franciscana*. *Sci. Rep.* 9:839. <https://doi.org/10.1038/s41598-019-44884-y>.
- Kappas, I., Abatzopoulos, T.J., Van Hoa, N., Sorgeloos, P., Beardmore, J.A., 2004. Genetic and reproductive differentiation of *Artemia franciscana* in a new environment. *Mar. Biol.* 146, 103–117. <https://doi.org/10.1007/s00227-004-1420-9>.
- King, A., 2013. Elucidating the functions of proteins up-regulated during diapause in *Artemia franciscana* using RNAi. *FEBS J.* 280, 4761–4772.
- King, A.M., MacRae, T.H., 2012. The small heat shock protein p26 aids development of encysting *Artemia* embryos, prevents spontaneous diapause termination and protects against stress. *PLoS One* 7 (8), e43723. <https://doi.org/10.1371/journal.pone.0043723>.
- Kumar, V., Baruah, K., Nguyen, D.V., Smaghe, G., Vossen, E., Bossier, P., 2018. Phloroglucinol-mediated Hsp70 production in crustaceans: protection against *Vibrio parahaemolyticus* in *Artemia franciscana* and *Macrobrachium rosenbergii*. *Front. Immunol.* 9, 1091. <https://doi.org/10.3389/fimmu.2018.01091>.
- Laemmli, U.K., 1970. Cleavage of structural proteins during assembly of the head of bacteriophage T4. *Nature.* 227, 680–685.
- Liang, P., MacRae, T.H., 1999. The synthesis of a small heat shock/a-crystallin protein in *Artemia* and its relationship to stress tolerance during development. *Dev. Biol.* 207, 445–456.
- Norouzitallab, P., Baruah, K., Biswas, P., Vanrompay, D., Bossier, P., 2016. Probing the phenomenon of trained immunity in invertebrates during a transgenerational study, using brine shrimp *Artemia* as a model system. *Sci. Rep.* 6, 21166. <https://doi.org/10.1038/srep21166>.
- Persoon, G., Sorgeloos, P., 1980. General aspects of the ecology and biogeography of *Artemia*. In: Persoon, G., Sorgeloos, P., Roels, O., Jaspers, E. (Eds.), *The Brine Shrimp Artemia*. 3. Universa Press, Wetteren, pp. 3–24.
- Sim, C., Denlinger, D.L., 2011. Catalase and superoxide dismutase-2 enhance survival and protect ovaries during overwintering diapause in the mosquito *Culex pipiens*. *J. Insect Physiol.* 57, 628–634.
- Sung, Y.Y., 2014. Heat shock proteins: An alternative to control disease in aquatic organisms. *J. Mar. Sci.: Res. Dev.* 4, e126. <https://doi.org/10.4172/2155-9910.1000e126>.
- Sung, Y.Y., MacRae, T.H., 2011. Heat shock proteins and disease control in aquatic organisms. *J. Aquac. Res. Dev.* 52, 006. <https://doi.org/10.4172/2155-9546.52-006>.
- Sung, Y.Y., Van Damme, E.J.M., Sorgeloos, P., Bossier, P., 2007. Non-lethal heat shock protects gnotobiotic *Artemia franciscana* larvae against virulent *Vibrios*. *Fish Shellfish Immunol.* 22, 318–326.
- Sung, Y.Y., Pineda, C., MacRae, T.H., Sorgeloos, P., Bossier, P., 2008. Exposure of gnotobiotic *Artemia franciscana* larvae to abiotic stress promotes heat shock protein 70 synthesis and enhances resistance to pathogenic *Vibrio campbellii*. *Cell Stress Chaperones* 13, 59–66.
- Sung, Y.Y., Dhaene, T., Defoirdt, T., Boon, N., MacRae, T.H., Sorgeloos, P., Bossier, P., 2009. Ingestion of bacteria overproducing DnaK attenuates *Vibrio* infection of gnotobiotic *Artemia franciscana* larvae. *Cell Stress Chaperones* 14, 603–609.
- Sung, Y.Y., Rahman, N.A., Shazili, N.A.M., Chen, S.J., Lv, A.J., Sun, J.F., 2018. Non-lethal heat shock induces Hsp70 synthesis and promotes tolerance against heat, ammonia and metals in post-larvae of the white leg shrimp *Penaeus vannamei* (Boone, 1931). *Aquaculture.* 483, 21–26.
- Toxopeus, J., Warner, A.H., MacRae, T.H., 2014. Group 1 LEA proteins contribute to the desiccation and freeze tolerance of *Artemia franciscana* embryos during diapause. *Cell Stress Chaperones* 19, 939–948.
- Yang, D., Xu, X., Zhao, H., Yang, S., Wang, X., Zhao, D., Diao, Q., Hou, C., 2018. Diverse factors affecting efficiency of RNAi in honey bee viruses. *Front. Genet.* 9, 384. <https://doi.org/10.3389/fgene.2018.00384>.

ORIGINALITY REPORT

20%

SIMILARITY INDEX

15%

INTERNET SOURCES

15%

PUBLICATIONS

6%

STUDENT PAPERS

PRIMARY SOURCES

1	www.researchgate.net Internet Source	2%
2	Mohamed Omar El-Magsodi, Kartik Baruah, Parisa Norouzitallab, Peter Bossier, Patrick Sorgeloos, Gilbert Van Stappen. "Hydration/dehydration cycles imposed on Artemia cysts influence the tolerance limit of nauplii against abiotic and biotic stressors", Aquaculture International, 2015 Publication	1%
3	f1000.com Internet Source	1%
4	Submitted to Khulna University Student Paper	1%
5	academic.oup.com Internet Source	1%
6	www.longdom.org Internet Source	1%
7	Kartik Baruah, Parisa Norouzitallab, Linayati	

Linayati, Patrick Sorgeloos, Peter Bossier.
"Reactive oxygen species generated by a heat shock protein (Hsp) inducing product contributes to Hsp70 production and Hsp70-mediated protective immunity in *Artemia franciscana* against pathogenic vibrios", *Developmental & Comparative Immunology*, 2014

1%

Publication

8

portal.research.lu.se

Internet Source

1%

9

Wisarut Junprung, Premruethai Supungul, Anchalee Tassanakajon. "Litopenaeus vannamei heat shock protein 70 (LvHSP70) enhances resistance to a strain of *Vibrio parahaemolyticus*, which can cause acute hepatopancreatic necrosis disease (AHPND), by activating shrimp immunity", *Developmental & Comparative Immunology*, 2018

1%

Publication

10

Wisarut Junprung, Premruethai Supungul, Anchalee Tassanakajon. "HSP70 and HSP90 are involved in shrimp *Penaeus vannamei* tolerance to AHPND-causing strain of *Vibrio parahaemolyticus* after non-lethal heat shock", *Fish & Shellfish Immunology*, 2017

1%

Publication

11

Dung Nguyen Viet, Olivier Christiaens, Duy V.B.

1%

Le, Stephanie De Vos, Thomas H. MacRae, Guy Smagghe, Peter Bossier. "RNA interference in Artemia franciscana", Aquaculture, 2018

Publication

12

digitalarchive.worldfishcenter.org

Internet Source

1%

13

repositorio.uam.es

Internet Source

1%

14

www.nrcresearchpress.com

Internet Source

1%

15

Sung, Yeong Yik, Thomas H. MacRae, Patrick Sorgeloos, and Peter Bossier. "Stress response for disease control in aquaculture : Stress and disease in aquaculture", Reviews in Aquaculture, 2011.

Publication

1%

16

www.spandidos-publications.com

Internet Source

<1%

17

www.lifetechindia.com

Internet Source

<1%

18

aem.asm.org

Internet Source

<1%

19

spandidos-publications.com

Internet Source

<1%

- | | | |
|----|---|-----|
| 20 | Kartik Baruah, Jayant Ranjan, Patrick Sorgeloos, Peter Bossier. "Efficacy of heterologous and homologous heat shock protein 70s as protective agents to <i>Artemia franciscana</i> challenged with <i>Vibrio campbellii</i> ", <i>Fish & Shellfish Immunology</i> , 2010
Publication | <1% |
| 21 | M.M. Frankenberg, S.A. Jackson, J.S. Clegg. "The heat shock response of adult <i>Artemia franciscana</i> ", <i>Journal of Thermal Biology</i> , 2000
Publication | <1% |
| 22 | mediatum.ub.tum.de
Internet Source | <1% |
| 23 | doaj.org
Internet Source | <1% |
| 24 | www.scribd.com
Internet Source | <1% |
| 25 | Julie A. Crack. "Functional analysis of a small heat shock/alpha-crystallin protein from <i>Artemia franciscana</i> . Oligomerization and thermotolerance", <i>European Journal of Biochemistry</i> , 2/2002
Publication | <1% |
| 26 | Sinnasamy, S, N Mat Noordin, T H MacRae, M Ikhwanuddin bin Abdullah, P Bossier, M E bin Abdul Wahid, A Noriaki, and Y Y Sung. | <1% |

"Ingestion of food pellets containing Escherichia coli overexpressing the heat-shock protein DnaK protects Penaeus vannamei (Boone) against Vibrio harveyi (Baumann) infection", Journal of Fish Diseases, 2015.

Publication

27

Submitted to Mahidol University

Student Paper

<1%

28

Ahmed A. M. Mohamed. "A target-specific feeding toxicity of β 1 integrin dsRNA against diamondback moth, *Plutella xylostella*", Archives of Insect Biochemistry and Physiology, 12/2011

Publication

<1%

29

Submitted to Malaysia University of Science and Technology

Student Paper

<1%

30

Wisarut Junprung, Premruethai Supungul, Anchalee Tassanakajon. "Structure, gene expression, and putative functions of crustacean heat shock proteins in innate immunity", Developmental & Comparative Immunology, 2021

Publication

<1%

31

eccis.ub.ac.id

Internet Source

<1%

32

O'Connell, P.A.. "Characterization of the microtubule proteome during post-diapause

<1%

development of *Artemia franciscana*", BBA - Proteins and Proteomics, 200605

Publication

33

Kartik Baruah, Ho Phuong Pham Duy Phong, Parisa Norouzitallab, Tom Defoirdt, Peter Bossier. "The gnotobiotic brine shrimp (*Artemia franciscana*) model system reveals that the phenolic compound pyrogallol protects against infection through its prooxidant activity", *Free Radical Biology and Medicine*, 2015

Publication

34

Cheng, C.H.. "Expression and characterization of the JAK kinase and STAT protein from brine shrimp, *Artemia franciscana*", *Fish and Shellfish Immunology*, 201005/06

Publication

35

bmcdevbiol.biomedcentral.com

Internet Source

36

d-nb.info

Internet Source

37

digital.library.adelaide.edu.au

Internet Source

38

bmccancer.biomedcentral.com

Internet Source

39

L.H. Phuoc. "Virulence of luminescent and non-luminescent isogenic vibrios towards gnotobiotic

<1%

<1%

<1%

<1%

<1%

<1%

<1%

Artemia franciscana larvae and specific pathogen-free *Litopenaeus vannamei* shrimp", *Journal of Applied Microbiology*, 04/2009

Publication

40

Sagi, Amir, Rivka Manor, and Tomer Ventura. "Gene Silencing in Crustaceans: From Basic Research to Biotechnologies", *Genes*, 2013.

Publication

41

Zhuoran Han, Jingfeng Sun, Aijun Lv, Jian-An Xian, Yeong Yik Sung, Xueliang Sun, Xiucui Hu, Kezhi Xing. "Transcriptome profiling of immune-responsive genes in the intestine of *Cynoglossus semilaevis* Günther challenged with *Shewanella* algae", *Fish & Shellfish Immunology*, 2018

Publication

42

www.repositoriodigital.ipn.mx

Internet Source

43

dev.biologists.org

Internet Source

44

scialert.net

Internet Source

45

Eamy Nursaliza Yaacob, Parisa Norouzitallab, Bruno G. De Geest, Aline Bajek, Kristof Dierckens, Peter Bossier, Daisy Vanrompay. "Recombinant DnaK Orally Administered

<1%

<1%

<1%

<1%

<1%

<1%

Protects Axenic European Sea Bass Against
Vibriosis", *Frontiers in Immunology*, 2020

Publication

46

allie.dbcls.jp

Internet Source

<1%

47

www.jbc.org

Internet Source

<1%

48

Valerie J. Smith, Elisabeth A. Dyrzynda.

"Antimicrobial proteins: From old proteins, new tricks", *Molecular Immunology*, 2015

Publication

<1%

49

Emilyn E. Matsumura, Jared C. Nigg, Elizabeth M. Henry, Bryce W. Falk. " Development of a cricket paralysis virus-based system for inducing RNA interference-mediated gene silencing in ", Cold Spring Harbor Laboratory, 2020

Publication

<1%

50

"Heat Shock Proteins and Stress", Springer Science and Business Media LLC, 2018

Publication

<1%

51

archive.org

Internet Source

<1%

52

Yi-Hong Chen, Jian-Guo He. "Effects of environmental stress on shrimp innate immunity and white spot syndrome virus infection", *Fish & Shellfish Immunology*, 2019

<1%

Publication

Exclude quotes On

Exclude matches Off

Exclude bibliography On