

# The identification of plankton tropical status in the Wonokromo, Dadapan and Juanda extreme water estuary

*by* L A Sari

---

**Submission date:** 21-Jan-2020 02:51PM (UTC+0800)

**Submission ID:** 1244344449

**File name:** s\_in\_the\_Wonokromo,\_Dadapan\_and\_Juanda\_extreme\_water\_estuary.pdf (395.67K)

**Word count:** 2950

**Character count:** 16354

PAPER • OPEN ACCESS

3  
The identification of plankton tropical status in the Wonokromo, Dadapan and Juanda extreme water estuary

6  
To cite this article: L A Sari *et al* 2018 *IOP Conf. Ser.: Earth Environ. Sci.* **137** 012029

6  
View the [article online](#) for updates and enhancements.

Related content

- 3  
- [The biomass, abundance, and distribution of starfish \*Asterias\* sp. \(Echinodermata: Asteroidea\) in East Coast of Surabaya](#)  
N N Dewi, K T Pursetyo, L Aprilianitasari et al.
- 8  
- [The identification of plankton, water quality, blood cell, and histology in culture pond of tilapia \*Oreochromis niloticus\* which infected by viral nervous necrosis \(VNN\)](#)  
U Yanuhar, D T Rahayu, M Musa et al.
- 23  
- [Study of copper \(Cu\) contents in blood cockles \(\*Anadara\* sp.\) at Surabaya coastal waters](#)  
S Alfionita, K T Pursetyo and A M Sahidu

## The identification of plankton tropical status in the Wonokromo, Dadapan and Juanda extreme water estuary

L A Sari<sup>1</sup>, W H Satyantini<sup>1</sup>, A Manan<sup>1</sup>, K T Pursetyo<sup>1</sup> and N N Dewi<sup>1</sup>

<sup>1</sup> Department of Fish Health Management and Aquaculture, Faculty of Fisheries and Marine, Universitas Airlangga, Jl. Airlangga, Surabaya 60115, East Java, Indonesia

E-mail: luthfianaaprilianitas@gmail.com

**Abstract.** Wonokromo, Dadapan and Juanda estuaries are extreme waters located around Surabaya environment. This is because of a lot of organic material intake, which provided nutrients for plankton growth. In addition, the waters is also dynamic in reason of physico-chemical, geological and biological processes controlled by the tides and freshwater run-off from the river that empties into it. The objective of this study was to identify the presentation of plankton in extreme waters based on brightness and ammonia level. The study was conducted in January 2017. Three sampling locations were Wonokromo, Dadapan and Juanda estuaries. Each station consists of three points based on distances, which were 400, 700, and 1000 meters from the coastline. The brightness in Wonokromo, Dadapan, and Juanda environment was 60, 40, and 100 cm, respectively. The result of ammonia in Wonokromo, Dadapan, and Juanda estuary was 0.837, 0.626, and 0.396 mg/L, correspondingly. Nine classes of phytoplankton's were found in three locations (bacillariophyceae, dynophyceae, chlorophyceae, cyanophyceae, crysophyceae, euglenoidea, trebouxlophyceae, mediophyceae, and nitachiaceae) and five classes of zooplanktons (maxillopoda, hexanuplia, copepoda, malacostraca, and oligotrichea). The density of plankton in Wonokromo, Dadapan and Juanda environments, was 37.64, 63.80, and 352.85 cells/L, respectively.

### 1. Introduction

The fertility of waters can be seen from the existence of plankton. According to Sofarini [1], stated that the presence and density of phytoplankton is one indicator of the fertility of the aquatic environment. According to Hemraj *et al.* [2], also stated that planktons are a bio-indicator of the water conditions of coastal lagoons. This indicator is performed by measuring the concentration of the chlorophyll produced by plankton.

The plankton density in waters is influenced by a number of environmental parameters and physiological characteristics. The composition and plankton density change at various levels in response to changes in physical, chemical, and biological conditions. These include physical and chemical changes such as the intensity of light, dissolved oxygen, temperature stratification, and the availability of nutrients, nitrogen and phosphorus; while the biological aspect is the existence of animal predation activities, death, and decomposition [3].



The east coast of Surabaya is located in the northwestern part of Madura Strait, covering the area between Tambak Wedi village and the Dadapan river estuary (border area with Sidoarjo). The waters of the east coast of Surabaya can be regarded as extreme waters due to several factors. One of the factors is that there are approximately 10 rivers that empties into it [4]. These waters make the coastal waters of Surabaya's east coast an estuary area with an input of numerous organic material intake as nutrients for the growth of plankton. Moreover, the coastal waters of Surabaya's east coast can be considered as very dynamic waters because there are various physical-chemical, geological and biological processes controlled by the sea water tides and freshwater run-off from the rivers that empties into it.

According to Winarni *et al.* [5], have stated that the structure of communities in the east coast of Surabaya have led to the discovery of seven types of sea cucumbers with high to low distribution indexes, none very high. According to Affandi *et al.* [4], mentioned that there were four species of *kupang* (local name) and one *lorjuk* (local name) species in the east coast waters of Surabaya with a distribution that did not occur spontaneously or randomly but were closely related to preferences or selection of suitable habitats.

This research is a preliminary study to provide basic data and will be continued in several parts. The research aims to assess the extreme coastal waters of Surabaya with extreme water condition and high organic material which have dominance, diversity and plankton density. Extreme waters with good fertility will have a variety of species and good plankton density as well. Therefore, a study on the identification and fertility levels of plankton in the extreme waters in the east coast of Surabaya was carried out.

Seawater as public waters may at any time experience changes in conditions and quality caused by surrounding environmental factors, weather, climate as well as human activities conducted on these waters. Changes that occur in sea water will cause fluctuations in water quality such as temperature, salinity, water brightness (turbidity), degree of acidity (pH) and dissolved oxygen (DO). Changes that occur to water quality will influence the sustainability of aquatic organisms. Planktons are one of the most influential organisms in extreme water conditions [6].

Some types of plankton are able to survive and grow in extreme waters. Plankton is able to adapt to extreme environments. According to Seckback [6], explained that some algae are able to survive under conditions of low temperature, high temperature and high salinity waters. Tolerant organisms that can adapt to extreme waters will easier adapt to within the culture.

Research from Hemraj *et al.* [2], stated that coastal lagoons are characterized by extreme environments especially high salinity. This is led to community changing. Studies have been performed which have used plankton as an indicator of extreme environmental fluctuations. Two different communities of phytoplankton and zooplankton were identified, with salinity and nutrients being the main factors affecting species distribution. Polychaete and gastropod larvae showed positive indicators of high salinity fluctuations.

## 2. Methodology

The study was conducted in January 2017. The research included field observations and laboratory analysis. Sampling was carried out in the coastal waters of Surabaya on three stations, where each station consisted of three points. The identification of plankton samples were performed at the Dry Laboratory of the Faculty of Fisheries and Marine at Airlangga University, Surabaya and the water sample was measured there.


The research material used was plankton found in the Wonokromo, Jua and Dadapan waters. The research materials used were 4% formalin, lugol, walnut nutrient ( $\text{NaNO}_3$ ,  $\text{Na}_2\text{EDTA}$ ,  $\text{H}_3\text{BO}_3$ ,  $\text{NaH}_2\text{PO}_4$ ,  $2\text{H}_2\text{O}$ ,  $\text{FeCl}_3$ ,  $6\text{H}_2\text{O}$ ,  $\text{MnCl}_2$ ,  $4\text{H}_2\text{O}$ ), vitamin solution stock, stock of micro metal solution (BBPAP Jepara), sea water chlorine (Brataco), Na Thiosulphate (Brataco), alcohol (Brataco) paper, gloves and masks.

The equipment used in the research were: sample bottles, plastic clips, bottle insulation, gel ice, GPS, styrofoam, aerator set, aerator hose, cork, measuring cup, Erlenmeyer, dropper, volume pipette,

microscope, homogenizer, hand tally counter, autoclave American 25H, Haemocytometer, Sedgwick-Rafter, Spectrophotometer, Hettich EBA-20 centrifuge, hot plate stirrer, refractometer, pH paper, thermometer, ammoniac test kit, DO test kit, Ohaus PA 2102 digital scales, OHAUS digital scales, Analytical Balance PA413, TL 40 watts, cotton, water funnel, gauze and aluminum foil.

The plankton identification books used were: Freshwater Algae Identification, Enumeration and Use as Bioindicators [10]; Methods for the Study of Marine Benthos (Athlete 7 March) Ostracods, and Chaetognaths [16]; Algae [9]; Identifying Marine Phytoplankton [22]; and Plankton: A Guide to their Ecology and Monitoring for Water Quality [3].

Plankton and water samples were obtained from three stations in Wonokromo, Juanda and Dadapan waters. Each station consisted of 3 water sampling points at a distance of 200, 500, 800 meters from the coastal waters.



**Figure 1.** Research location and sampling site at Wonokromo, Dadapan and Juanda waters.

### 2.1. Sample collection

The sampling of plankton and water was done at each sampling location. Sampling of microalgae (phytoplankton) was carried out using a 10  $\mu\text{m}$  diameter plankton net. Zooplankton sampling was done using a 80  $\mu\text{m}$  diameter plankton net. First, 50 liters surface water was filtered by a plankton net in each station. Then the filtered water was put into a sample container. The ordinate sampling station was recorded. Bottles were labeled with information such as times, dates, points and sampling stations. Two sample bottles were collected at each sample site location, one bottle of sample was preserved by using lugol (sample for density calculation and plankton diversity), while the other contained a sample without additional preservatives but stored in a gelled ice coolbox (sample for plankton insulation) [3].

The identification of samples was performed at the laboratory of the Faculty of Fisheries and Marine at Airlangga University. Plankton analyses was determined by dripping 1 ml of sample water on the haemocytometer, then observed under the microscope with 100 enlargement. The checking of parameters of water samples such as Dissolved Oxygen (DO), temperature, brightness, current and pH were done directly at the sampling site.

### 2.2. Identification and Calculation of Plankton Density

The identification of the plankton is based on its morphological form observed by using a microscope [23]. The identification of zooplankton is based on [16] the calculation of plankton density using the formula from [20], while the Calculation of Dominance Index and Diversity Index was calculated using the formula from [19].



### 3. Result and Discussion

The research data stated that the extreme waters at the locations of Wonokromo, Dadapan and Juanda was caused by the condition of water brightness. The brightness at 3 stations were 60 cm at Wonokromo, 40 cm at Dadapan and 100 cm at Juanda waters. Ammonia at the three stations were measured at 0.837 mg/L in the waters of Wonokromo, 0.626 mg/L in Dadapan and 0.396 mg/L in Juanda.



**Figure 2.** Location of sample waters.

The development of phytoplankton is determined by the intensity of sunlight, temperature and nutrients. The high growth of phytoplankton may not always be beneficial for aquatic conditions; it can also cause a population explosion (blooming), which can produce harmful toxic substances.

The structure of the phytoplankton community is a collection of populations that live in a particular region or habitat that interconnect and interact or have a reciprocal relationship of a particular zone [19]; this includes the index of diversity, index dominance, uniformity index and species wealth index (Kamali, 2004). The Shannon-Wiener species diversity index is a mathematical calculation that describes the analysis of information about the number of individuals in each species, the number of species and the total of individuals within a community (Masson, 1981). The uniformity index (ETIsability) is a description of the uniformity of the individual distribution of the phytoplankton species in a community [19]. Simpson's dominance index represents the presence or absence of a dominant species in a community. The disappearance of the dominant species causes a change in the biotic community and its physical environment [19]. The richness index is used to find out the least amount of taxa and the concentration of biota in a community (Margalef, 1951 in Romimohtarto, 2001).

From the sampling sites, we found 10 classes of phytoplankton (bacillariophyceae, dynophyceae, chlorophyceae, cyanophyceae, crysophyceae, euglenoidea, trebouxlophyceae, mediophyceae, nitachiaceae) and 5 classes of zooplankton (maxillopoda, hexanuplia, copepoda, malacostraca, oligotrichea). The density of plankton were 37.64 cells/liter in Wonokromo, 63.80 cells/liter in Dadapan, and 352.85 cells/liter in Juanda waters.

Phytoplankton have chlorophyll that the function in photosynthesis to produce organic and oxygen in the water. However, certain phytoplankton may degrade the quality of the waters if the amount is excessive (blooming). The high populations of toxic phytoplankton in waters can cause negative consequences for aquatic ecosystems, such as reduced oxygen in the water that can cause the death of various aquatic creatures.

**Table 1.** Identification of Plankton Density at the Wonokromo, Juanda, and Dadapan locations.

Species of Plankton	Wonokromo			Juanda			Dadapan		
	400 m	700 m	1000 m	400 m	700 m	1000 m	400 m	700 m	1000 m
<i>Amphora</i>		x	x					x	x
<i>Amphidinium</i>	x				x			x	
<i>Bidulphia</i>							x	x	x
<i>Ceratium</i>			x						
<i>Chaetoceros</i>		x	x	x	x	x			
<i>Chromulina</i>			x	x	x	x			
<i>Coscinodiscus</i>		x	x	x	x	x	x		
<i>Cyclotella</i>	x		x	x			x	x	x
<i>Dynophysis</i>					x				x
<i>Dytilum</i>			x		x				x
<i>Euglena</i>									x
<i>Gymnodinium</i>									
<i>Lyngbia</i>	x				x			x	
<i>Naupli</i>			x	x					
<i>Navicula</i>	x		x	x	x	x			X
<i>Nitzchia</i>		x	x		x	x			
<i>Oocystis</i>	x	x							
<i>Oscillatoria</i>	x	x			x		x	x	X
<i>Parafavella</i>				x	x	x			
<i>Peridinium</i>				x				x	
<i>Planktoniella</i>			x	x	x	x			
<i>Pleurosigma</i>				x				x	
<i>Rhizosolenia</i>	x	x	x	x	x			x	
<i>Siola</i>		x							X
<i>Tabellaria</i>		x			x				
<i>Tintinopsis</i>						x			
<i>Thalassiothrix</i>									
<i>Skeletonema</i>			x	x	x	x			

Phytoplankton can be found throughout the mass of water from the surface to a depth where it is still possible for the intensity of sunlight to be used in the process of photosynthesis. This phytoplankton is the largest flora component of its role as a primary producer in a waters [19]

#### 4. Conclusion

Our gratitude goes out to the Faculty of Fisheries and Marine who have funded this research. We would also like to thank our team of researchers: both technicians in the laboratory and in the field. We hope that this research can be beneficial for future studies on this topic. The brightness in Wonokromo, Dadapan, and Juanda environment was 60, 40, and 100 cm, respectively. The result of ammonia in Wonokromo, Dadapan, and Juanda estuary was 0.837, 0.626, and 0.396 mg/L, correspondingly. Nine classes of phytoplankton's were found in three locations (bacillariophyceae, dynophyceae, chlorophyceae, cyanophyceae, crysophyceae, euglenoidea, trebouxlophyceae, mediophyceae, and nitachiaceae) and five classes of zooplanktons (maxillopoda, hexanuplia, copepoda, malacostraca, and oligotrichea). The density of plankton in Wonokromo, Dadapan and Juanda environments, was 37.64, 63.80, and 352, 85 cells/L, respectively.

#### 5. Refference

- [1] Sofarini D 2012 *Enviro Scientee* **8** 30-34

- [2] Hemraj DA, Hossain Q, Ye J, G Qin and SC Leterme 2017 *ECSA* **184** 102-114
- [3] Suthers IM and D Rissik 2008 Plankton: a guide to their ecology and monitoring for water quality *National Library of Australia Cataloguing-in-Publication* entry CSIRO Publishing p 273
- [4] Affandi MB, Irawan A, Soegianto and Sosmanida 2009 *J. Biol. Researches* **3B** 37–41
- [5] Harni D, M Affandi, ED Masithah and AN Kristanti 2014 *JPMIPA* **117**
- [6] Seckbach J 2007 Algae and Cyanobacteria in extreme environments (cellular origin life in extreme habitats and astrobiology) *Dordrecht Springer* p 786
- [7] Ahuja S 2013 Monitoring water quality pollution assessment analysis and remediation *Elsevier* p 374
- [8] Barnes RSK and KH Mann 1992 Fundamental of aquatic ecology second edition Blackwell Science p 271
- [9] Barsanti L and P Gualtieri 2014 Algae CRC Press Taylor and Francis p 344
- [10] Bellinger EG and DC Sigeo 2015 Freshwater algae : identification enumeration and use as bioindicator *Wileyblackwell* p 275
- [11] Boyd CE 2015. Water quality an introduction second edition *Springer International Publishing Switzerland* pp 14-16
- [12] Darley WM 1982 Algal biology: a physiological approach blackwell scientific publications. Pp 97-98
- [13] Dolan JR, DJS Montagnes, S Agatha, DW Coats and DK Stoecker 2013 The biology and ecology of tintinnid ciliates: models for marine *Wiley-Blackwell* p 319
- [14] Effendi H 2003 Water quality study for resource management and aquatic environments *Kanisius Yogyakarta* p 250
- [15] Eleftheriou 2013 Methods for the study of marine benthos *Wileyblackwell* p 502
- [16] Guglielmo L and A Ianora 1997 Atlas of marine zooplankton I straits of magellan amphipods, nauplii, mysids, ostracods and chaetognaths *Springer* p 263
- [17] Li Y and K Migliaccio 2011 Water quality concepts sampling and analyses *CRC Press* p 335
- [18] McVey JP 1983 *CRC Handbook of marine culture volume I* CRC Press Inc Boca Rton Florida pp 33- 36
- [19] Odum EP 1993 The third edition of the ecology base (*Yogyakarta: Gadjah Mada University Press*) p 697
- [20] Rice EW, RB Baird, AD Eaton and LS Clesceri 2012 Standard methods for the examination of water and wastewater 22nd edition *American Public Health Association American Water Works Association Water Environment Federation* 1496p
- [21] Soccodato A, F d'Ovidio, M Lévy, O Jahn, MJ Follows and SD Monte 2016 **29** 9–17
- [22] Tomas CR 1997 Identifying marine phytoplankton Printed in The United States of America p 875
- [23] Wehr JD, RG Sheath and JP Kociolek 2015 Freshwater algae of North America ecology and classification *Academic Press* is an imprint of Elsevier p 1045



# The identification of plankton tropical status in the Wonokromo, Dadapan and Juanda extreme water estuary

## ORIGINALITY REPORT

18%

SIMILARITY INDEX

9%

INTERNET SOURCES

8%

PUBLICATIONS

9%

STUDENT PAPERS

## PRIMARY SOURCES

1	<a href="https://repositorium.sdum.uminho.pt">repositorium.sdum.uminho.pt</a> Internet Source	2%
2	Jakob Wolf. "Can the religions contribute to fight climate change?", IOP Conference Series: Earth and Environmental Science, 2009 Publication	2%
3	<a href="http://www.ppjpi.unair.ac.id">www.ppjpi.unair.ac.id</a> Internet Source	1%
4	Deevesh A. Hemraj, Md A. Hossain, Qifeng Ye, Jian G. Qin, Sophie C. Leterme. "Plankton bioindicators of environmental conditions in coastal lagoons", Estuarine, Coastal and Shelf Science, 2017 Publication	1%
5	Submitted to Universitas Hasanuddin Student Paper	1%
6	<a href="https://repository.uhamka.ac.id">repository.uhamka.ac.id</a> Internet Source	1%

7	Submitted to University Of Tasmania Student Paper	1%
8	<a href="http://www.science.gov">www.science.gov</a> Internet Source	1%
9	Submitted to Heriot-Watt University Student Paper	1%
10	Viviana R. Lopes, Vitor M. Vasconcelos. "Planktonic and benthic cyanobacteria of European brackish waters: a perspective on estuaries and brackish seas", European Journal of Phycology, 2011 Publication	1%
11	Américo Montiel San Martín, Dieter Gerdes, Wolf E. Arntz. "Distributional patterns of shallow- water polychaetes in the Magellan region: a zoogeographical and ecological synopsis", Scientia Marina, 2007 Publication	1%
12	<a href="http://f1000research.com">f1000research.com</a> Internet Source	1%
13	<a href="http://revistas.unal.edu.co">revistas.unal.edu.co</a> Internet Source	<1%
14	Submitted to Bellarmine University Student Paper	<1%
15	<a href="http://skyros.locean-ipsl.upmc.fr">skyros.locean-ipsl.upmc.fr</a>	

Internet Source

<1%

16

Submitted to Trisakti University

Student Paper

<1%

17

Submitted to University of Nottingham

Student Paper

<1%

18

[www.ekolojidergisi.com](http://www.ekolojidergisi.com)

Internet Source

<1%

19

[hdl.handle.net](http://hdl.handle.net)

Internet Source

<1%

20

Submitted to South Bank University

Student Paper

<1%

21

Thushara Navodani Wickramaarachchi. "Rating curve based assessment of seasonal variability of sulfate in streamflow", Environmental Monitoring and Assessment, 2018

Publication

<1%

22

[docobook.com](http://docobook.com)

Internet Source

<1%

23

Submitted to Aston University

Student Paper

<1%

24

Binbin Liu, Feng Zhang, Xiaoxi Feng, Yongdi Liu, Xing Yan, Xiaojun Zhang, Linghua Wang, Liping Zhao. "Thauera and Azoarcus as

<1%

functionally important genera in a denitrifying quinoline-removal bioreactor as revealed by microbial community structure comparison", FEMS Microbiology Ecology, 2006

Publication

25

[onlinelibrary.wiley.com](https://onlinelibrary.wiley.com)

Internet Source

<1%

26

Maria Tereza Morais Pereira Souza Lobo, Paulo Sérgio Scalize, Cleber Nunes Kraus, Weliton José da Silva et al. "Biological index based on epiphytic diatom assemblages is more restrictive than the physicochemical index in water assessment on an Amazon floodplain, Brazil", Environmental Science and Pollution Research, 2020

Publication

<1%

27

[www.sjrbc.com](http://www.sjrbc.com)

Internet Source

<1%

28

Submitted to Hadlow College, Kent

Student Paper

<1%

29

Submitted to University of Edinburgh

Student Paper

<1%

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

