

Efficacy of an Education Session by Pharmacists for Patients With Asthma: Protocol and Design of a Randomized Controlled Trial

by Elida Zairina

Submission date: 02-Sep-2021 04:26PM (UTC+0800)

Submission ID: 1639984450

File name: C-10.JMIR.pdf (646.67K)

Word count: 4172

Character count: 23606

Protocol

Efficacy of an Education Session by Pharmacists for Patients With Asthma: Protocol and Design of a Randomized Controlled Trial

Elida Zairina¹, PhD; Gesnita Nugraheni¹, MSc; Gusti NV Achmad¹, MSc; Arie Sulistyari¹, MPharm; Yunita Nita¹, Dr; Arief Bakhtiar^{2,3,4}, MD; Muhammad Amin^{2,3,4}, MD, Prof Dr

¹Department of Pharmacy Practice, Faculty of Pharmacy, Universitas Airlangga, Surabaya, Indonesia

²Faculty of Medicine, Department of Pulmonology and Respiratory Medicine, Universitas Airlangga, Surabaya, Indonesia

³Department of Pulmonology, Dr Soetomo Hospital, Surabaya, Indonesia

⁴Department of Pulmonology, Universitas Airlangga Hospital, Surabaya, Indonesia

Corresponding Author:

Elida Zairina, PhD
Department of Pharmacy Practice
Faculty of Pharmacy
Universitas Airlangga
Dharmawangsa Dalam Surabaya
Surabaya, 60286
Indonesia
Phone: 62 31 5033710 ext 201
Fax: 62 31 5020514
Email: elida-z@ff.unair.ac.id

26

Abstract

Background: Asthma is a chronic disease that requires indefinite long-term therapy. Many approaches have been developed to enable people with asthma to live as normally as possible. In medication therapy management, pharmacists could play important roles in supporting the everyday life of asthmatic patients, such as by providing education therapy management to ensure that patients achieve optimal therapeutic outcomes. A good collaboration between health care practitioners and patients will produce a better system in terms of therapeutic management, which will lead to health care cost savings related to emergency visits. Although the Government has made various efforts to manage asthma in Indonesia, without commitment and support from both patients and health care professionals, the expected outcomes cannot be achieved.

Objective: This study aims to evaluate the effectiveness of an educational intervention provided by pharmacists compared with that of usual care.

Methods: A randomized controlled trial comparing usual care with an education session by pharmacists is underway. The intervention comprises a one-on-one education session of 60 minutes with a pharmacist comprising information regarding (1) asthma medication that has been used; (2) how to use asthma medication devices correctly; (3) asthma symptoms and how to prevent exacerbation of asthma; and (4) how to manage asthma triggers and environmental control measures. The primary outcome measure is change in asthma control, as measured using the Asthma Control Questionnaire. Secondary outcomes include changes in Asthma Quality of Life Questionnaire score, lung function, asthma-related health visits, days off from work or study, and oral corticosteroid use. Research assistants who are masked to the group allocation will collect outcome data at the baseline and every month for a 3-month period. Informed consent will be sought at enrollment and intention-to-treat analysis will be performed.

Results: This study was funded in January 2017 and ethical approval was obtained in June 2017. The enrollment was started in August 2017 and about 72 participants have been enrolled. First results are expected to be submitted for publication in 2019.

Conclusions: This is the first study to evaluate the effectiveness of a pharmacist-guided asthma education session compared with that of usual care in Indonesia. If it is proven effective, this intervention program could improve asthma self-management by patients, which may reduce risks of poorly controlled asthma. This intervention could also be implemented in addition to the current usual care for patients with asthma.

Trial Registration: Thai Clinical Trials Registry TCTR20171219001; <http://www.clinicaltrials.in.th/index.php?tp=regtrials&menu=trialsearch&smenu=fulltext&task=search&task2=view1&id=3068> (Archived by WebCite at <http://www.webcitation.org/73C5eKtv>)

International Registered Report Identifier (IRRID): DERR1-10.2196/10210

(*JMIR Res Protoc* 2018;7(12):e10210) doi: [10.2196/10210](https://doi.org/10.2196/10210)

KEYWORDS

asthma control; education session; pharmacist

16 Introduction

Asthma is a chronic disease that affects people of all ages worldwide. Uncontrolled asthma can restrict patients' daily activities and can place them at risk of death. According to the World Health Organization and Global Initiative for Asthma, as many as 300 million people worldwide, of different ages and races, are exposed to asthma; this number is predicted to increase to 400 million by 2025 [1]. Asthma is a health problem in both developed and developing countries [1-3]. Approximately 250,000 people die of asthma each year [3,4]. In 2007, asthma caused 3447 deaths in the United States—equivalent to approximately 9 people each day [5]. Death due to asthma is more common among adults than among children; it is also more common among women (2173) than among men (1274) [5]. In the United Kingdom, >10% of the adults exhibit asthma [6,7]. Although the proportion of population affected by asthma in Asian countries, including Indonesia, is lower than that in Europe or America, the proportion of elderly patients with asthma in Asian countries is quite high (1.3%-15.3%) [8]. Asthma can be controlled with appropriate therapeutic management. Patients with controlled asthma conditions can participate in normal activities and are not likely to experience fatal asthma symptoms [9].

Asthma is a chronic disease that requires indefinite long-term therapy. Various approaches are needed to optimize therapeutic treatments among asthma patients. Several studies in developed countries, such as the United States [10,11] and Australia [12-14], have showed that better health outcomes for asthma patients resulted from the following factors: adequate knowledge among asthma patients, regular monitoring of therapy, and high level of understanding among both health care professionals and patients regarding the disease management behavior. In Indonesia, asthma is one of the top 10 causes of morbidity and death, together with chronic bronchitis and emphysema [15]. In April 2007, observations in 5 provinces of Indonesia (North Sumatra, Central Java, East Java, West Kalimantan, and South Sulawesi) conducted by multiple chronic and degenerative disease subdivisions showed that, in general, asthma control efforts have not been effectively implemented; moreover, there is minimal availability of the equipment required for the diagnosis and management of asthma patients in health care facilities. In 1997, the prevalence of asthma in children was 12%; in 2008, results of the International Study of Asthma and Allergies in Childhood showed that asthma prevalence in 12-14-year-old children was 12.6% [16].

It is currently unknown how the implementation of pharmacist-guided education in asthma self-management might

affect asthma self-management in Indonesia as there has been limited research regarding the understanding and behavior of patients for the management of asthma. Therefore, the proposed study has been designed to determine the effectiveness of pharmacist-guided education for asthma patients in Indonesia. Because medications play important roles in successful management of chronic diseases, including asthma, the role of pharmacists' expertise is essential in the implementation of medication therapy management. Studies have shown that improved asthma control can be achieved if patients are involved in self-management, including self-monitoring of asthma symptoms or lung function, as well as when patients follow written asthma action plans while maintaining regular contact with their health care professionals [17].

This study aims to evaluate the efficacy of pharmacist-guided education sessions provided to patients with asthma compared with that of usual care. We hypothesize that the intervention group will demonstrate superior asthma control, as measured by changes in the Asthma Control Questionnaire (ACQ) scores, after 3 months from the baseline.

Methods

Study Setting

Participant recruitment is ongoing at the outpatient Departments of Pulmonology at Universitas Airlangga Hospital. The study has been approved by the human research ethics committee of Universitas Airlangga Hospital. All participants provide written informed consent at the time of enrollment.


15 Study Design

This study is designed as a prospective, single-blinded, randomized controlled trial (Thai Clinical Trials Registry # TCTR20171219001); outcome assessors will be masked to group allocation at follow-up assessments. The flow of participants with the expected number is illustrated in Figure 1. The total duration of the study is 3-6 months, depending on the timing of the first visit and patient enrollment in the study. Both groups will be followed up for 3 months; the outcomes will be compared at 1, 2, and 3 months from the baseline to evaluate the efficacy of the intervention.

Inclusion and Exclusion Criteria

Eligibility for this trial includes patients with asthma who have used any regular medications for asthma within the previous 12 months, who are 18 years of age or older, and are able to communicate in Indonesian. Those who are unable or contraindicated to demonstrate the lung function with spirometer will be excluded.

Figure 1. Flowchart of study participants.



Trial Recruitment

The following methods of identification and recruitment of participants will be used in this study: First, the doctors will identify all patients with asthma visiting the outpatient Department of Pulmonology at Universitas Airlangga Hospital on the date of each patient's clinic visit. The research assistants will approach potential participants and perform screening on the basis of the inclusion and exclusion criteria that were described earlier in this protocol. Then, participant explanatory statements and expression of interest forms will also be provided at the outpatient Department of Pulmonology at Universitas Airlangga Hospital, which has access to the relevant information. Potential participants will be asked to leave their contact details to allow one of the research assistants to contact them. Finally, if a patient agrees to participate, written informed consent will be sought.

Group Allocation

Recruited participants will be allocated to intervention or control group on a 1:1 basis. Allocation will be concealed using the sealed opaque envelope technique. Random blocks of 4 and 6 will be chosen, and random numbers will be generated using a random allocation software program [18] by an external researcher not involved in the study. Only this researcher will know the allocation sequence. At the time of recruitment, the researchers coordinating this study will open the numbered envelope and allocate each participant to the control (usual care) group or the intervention (education) group. The outcome assessors will be masked to participant group allocation during follow-up assessments.

30

Control and Intervention Group

Control: Usual Care Group

Participants allocated to the control group will receive usual medical care provided by the Department of Pulmonology and its health care professionals. This includes regular monthly visits, depending on each patient's asthma severity or complications. If, during follow-up, it becomes apparent that asthma control has deteriorated since the prior assessment (eg, using the asthma reliever ≥ 3 times per week or requiring an increased preventer dose), the participant and corresponding health professionals will be notified with the participant's permission.

Intervention: Education Group

Before the delivery of educational sessions to patients, pharmacists in this study will undergo training provided by a certified asthma educator (EZ). The trial evaluates an intervention involving pharmacists to deliver a *one-on-one* education session regarding (1) asthma medication that has been used; (2) how to use asthma medication devices correctly; (3) asthma symptoms and how to prevent exacerbation of asthma; and (4) how to manage asthma triggers and environmental control measures. A video that explains how properly use asthma medications with a variety of devices will be shown to the intervention group. The intervention group will also be provided with an asthma booklet that explains how to correctly use asthma medication and avoid asthma triggers. One of the researchers will contact participants' health care professionals if any medication changes or unscheduled asthma-related visits are needed.

A written asthma action plan, consistent with the Global Initiative for Asthma guidelines, has been translated into the Indonesian language and will be used to design a participant-specific treatment plan based on the information obtained at the baseline. The asthma action plan contains instructions regarding which medications to take when feeling well, how to recognize worsening asthma, what to do when symptoms are worsening, and what to do in the event of an acute attack, including a first aid plan. The flow of the study is described in Figure 2.

Outcome Measures

The primary outcome measure is change in asthma control, as measured by the Juniper ACQ [19]. Secondary outcomes include

changes in Juniper's Asthma Quality of Life Questionnaire score [20], Juniper's Adherence to Refills and Medications Scales scores, asthma-related health visits, days off from work or study related to asthma, and oral corticosteroid use.

Data Collection and Follow-Up


ACQ scores, Juniper's Asthma Quality of Life Questionnaire scores, Adherence to Refills and Medications Scale scores, asthma-related health visits, asthma-related days off from work study, oral corticosteroid use, and preventer or reliever use data will be collected at the baseline and at 1, 2, and 3 months from the baseline to allow comparisons. Identical data collection forms will be used for both groups. The assessors responsible for collecting outcome data at 1, 2, and 3 months will be masked to participant group allocation.

Sample Size

A sample size of 28 participants per arm using an estimated SD of 0.66 in ACQ scores will have 80% power (with 2-sided 5% significance level) to detect the minimal clinically important difference in the ACQ score of ≥ 0.5 between the groups [21,22]. To allow for 25% attrition, 41 participants will be recruited for each arm.

Data Analysis

The primary analysis will be performed in accordance with the intention-to-treat principle. The baseline characteristics of the 2 groups will be compared using Student's *t* test for normally distributed continuous variables, Mann-Whitney *U* test for nonnormally distributed continuous variables, and chi-square or Fisher's exact test (as appropriate) for categorical variables. Primary inferential analysis will be conducted using a mixed effects model for the intention-to-treat population. This model will include the treatment group and time as fixed effects, with an interaction between treatment and time to ascertain if the groups behave differently over time. Other demographic and clinical factors will be included as potential covariates in the mixed effects model. Comparisons will also be made of the following: the proportion of participants whose ACQ score improves >0.5 (minimal clinically important difference) over the study period, the proportion in whom asthma remains "not well controlled" (ACQ score ≥ 1.5), and those whose asthma is "well controlled" (ACQ score < 1.5) at each time point [23]. Secondary outcomes will be summarized using descriptive statistics; analyses will be performed using the methods described above.

Figure 2. Flowchart of the study.

Results

This study was funded in January 2017 and ethical approval was obtained in June 2017. The enrollment was started in August 2017 and is currently ongoing; we plan to complete the

enrollment by December 2018. About 72 participants have been enrolled in the trial. First results are expected to be submitted for publication in 2019.

Discussion

⁵This is the first study to evaluate the effectiveness of a pharmacist-guided asthma educational session compared with that of usual care in Indonesia. If it is proven effective, this intervention program could improve asthma self-management by patients, which may thus reduce the risk of poorly controlled asthma. This intervention could also be implemented along with the current usual care for patients with asthma. This trial is designed to evaluate an educational program for patients with asthma in order to enable better self-management for the control of their asthma. The individualized written asthma action plan designed for each patient provides clear guidelines in terms of actions to be taken in case of worsening asthma.

The proposed intervention has the potential to improve asthma outcomes by facilitating better asthma self-management. This ²³ translate to reduced health care costs in the form of fewer asthma-related unplanned medical and emergency department

visits. If the intervention is efficacious, this could potentially influence clinical practice and health policy.

In order to improve health outcomes of asthmatic patients, solutions include regular monitoring as well as education regarding medication use and compliance. Most chronic diseases require patients to undergo therapy for an indefinite duration to meet therapeutic goals. Pharmacists have an important role in ensuring patients achieve optimal therapeutic outcomes. A pharmacist should be aware of this role, which involves providing adequate information to patients with chronic diseases. By building collaborations among health care professionals, a robust system of therapeutic management for patients with asthma will be established, enabling improvement of health outcomes and potentially reducing health care costs. Multiple efforts have been initiated by the government to manage asthma in Indonesia; however, without support and commitment from both health care professionals and community, the desired results cannot be achieved.

Acknowledgments

The authors thank the Ministry of Research and Technology and Higher Degree of Indonesia. The authors also thank the Faculty of Pharmacy, Universitas Airlangga, and the Universitas Airlangga for the support and facilities provided during the study. The authors thank the Head of the Pharmacy Department at Universitas Airlangga Hospital, Dr Budi Suprapti, as well as staff members Dr Mahardian Rahmadi and Ms Dewi Wara Shinta for their help and support in developing the recruitment system. The authors would also like to thank Mr Alfian Nur Rosyd, MD, Ms Amalia Wardatul Firdaus, Ms Anggita Mirzautika, Ms Ishmatul Izza, Ms Kinanthi Pasissa Puspita, and Ms Putri Intan Pertiwi for their help in recruiting participants. This trial is supported by the Ministry of Research, Technology, and Higher Degree of Indonesia.

Authors' Contributions

EZ conceived the project idea with input from GN, GNVA, and YN. EZ further developed the trial with input from all other authors: AB, MA, AS, GN, GNVA, and YN. EZ is managing the trial, while GN is organizing research assistant rosters. EZ wrote the first draft of this manuscript and refined it on the basis of comments and feedback from all other authors.

Conflicts of Interest

None declared.

References

1. World Health Organization. 2018. Global status report on noncommunicable disease URL: <http://www.who.int/news-room/fact-sheets/detail/noncommunicable-diseases> [accessed 2018-10-24] [WebCite Cache ID 73NqTG38P]
2. Masoli M, Fabian D, Holt S, Beasley R, Global Initiative for Asthma (GINA) Program. The global burden of asthma: executive summary of the GINA Dissemination Committee report. *Allergy* 2004 May;59(5):469-478 [FREE Full text] [doi: [10.1111/j.1398-9995.2004.00526.x](https://doi.org/10.1111/j.1398-9995.2004.00526.x)] [Medline: [15080825](https://pubmed.ncbi.nlm.nih.gov/15080825/)]
3. Braman SS. The global burden of asthma. *Chest* 2006 Jul;130(1 Suppl):4S-12S. [doi: [10.1378/chest.130.1_suppl.4S](https://doi.org/10.1378/chest.130.1_suppl.4S)] [Medline: [16840363](https://pubmed.ncbi.nlm.nih.gov/16840363/)]
4. Kroegel C. Global Initiative for Asthma (GINA) guidelines: 15 years of application. *Expert Rev Clin Immunol* 2009 May;5(3):239-249. [doi: [10.1586/eci.09.1](https://doi.org/10.1586/eci.09.1)] [Medline: [20477002](https://pubmed.ncbi.nlm.nih.gov/20477002/)]
5. Xu J, Murphy SL, Kochanek KD, Arias E. Mortality in the United States, 2015. *NCHS Data Brief* 2016 Dec(267):1-8 [FREE Full text] [Medline: [27930283](https://pubmed.ncbi.nlm.nih.gov/27930283/)]
6. Netuveli G, Hurwitz B, Levy M, Fletcher M, Barnes G, Durham SR, et al. Ethnic variations in UK asthma frequency, morbidity, and health-service use: a systematic review and meta-analysis. *Lancet* 2005;365(9456):312-317. [doi: [10.1016/S0140-6736\(05\)17785-X](https://doi.org/10.1016/S0140-6736(05)17785-X)] [Medline: [15664226](https://pubmed.ncbi.nlm.nih.gov/15664226/)]
7. Netuveli G, Hurwitz B, Sheikh A. Ethnic variations in incidence of asthma episodes in England & Wales: national study of 502,482 patients in primary care. *Respir Res* 2005 Oct 21;6:120 [FREE Full text] [doi: [10.1186/1465-9921-6-120](https://doi.org/10.1186/1465-9921-6-120)] [Medline: [16242029](https://pubmed.ncbi.nlm.nih.gov/16242029/)]
8. Song W, Kang M, Chang Y, Cho S. Epidemiology of adult asthma in Asia: toward a better understanding. *Asia Pac Allergy* 2014 Apr;4(2):75-85 [FREE Full text] [doi: [10.5415/apallergy.2014.4.2.75](https://doi.org/10.5415/apallergy.2014.4.2.75)] [Medline: [24809012](https://pubmed.ncbi.nlm.nih.gov/24809012/)]

9. Global Initiative For Asthma. Global Strategy for Asthma Management and Prevention (2016 update) URL: https://ginasthma.org/wp-content/uploads/2016/04/GINA-2016-main-report_tracked.pdf [accessed 2018-10-23] [WebCite Cache ID 73NpvarNe]
10. Bunting BA, Cranor CW. The Asheville Project: long-term clinical, humanistic, and economic outcomes of a community-based medication therapy management program for asthma. *J Am Pharm Assoc* (2003) 2006;46(2):133-147. [Medline: [16602223](#)]
11. Isetts B, Schondelmeyer S, Artz M, Lenarz L, Heaton A, Wadd W, et al. Clinical and economic outcomes of medication therapy management services: the Minnesota experience. *J Am Pharm Assoc* (2003) 2008;48(2):203-11; 3 p following 211. [doi: [10.1331/JAPhA.2008.07108](#)] [Medline: [18359733](#)]
12. Armour CL, Lemay K, Saini B, Reddel HK, Bosnic-Anticevich SZ, Smith LD, et al. Using the community pharmacy to identify patients at risk of poor asthma control and factors which contribute to this poor control. *J Asthma* 2011 Nov;48(9):914-922. [doi: [10.3109/02770903.2011.615431](#)] [Medline: [21942306](#)]
13. Saini B, Krass I, Smith L, Bosnic-Anticevich S, Armour C. Role of community pharmacists in asthma - Australian research highlighting pathways for future primary care models. *Australas Med J* 2011;4(4):190-200 [FREE Full text] [doi: [10.4066/AMJ.2011.790](#)] [Medline: [23393510](#)]
14. Saini B, LeMay K, Emmerton L, Krass I, Smith L, Bosnic-Anticevich S, et al. Asthma disease management-Australian pharmacists' interventions improve patients' asthma knowledge and this is sustained. *Patient Educ Couns* 2011 Jun;83(3):295-302. [doi: [10.1016/j.pec.2011.05.001](#)] [Medline: [21621947](#)]
15. Dirjen PK. Direktorat Bina Farmasi Komunitas dan Klinik. Ditjen Bina Kefarmasian dan Alat Kesehatan. Departemen Kesehatan Republik Indonesia. 2007. Pharmaceutical Care untuk penyakit asma URL: <http://farmalkes.kemkes.go.id/> [accessed 2018-10-24] [WebCite Cache ID 73Nr1M94K]
16. Pfister M, Schaub MC, Watterson JG, Knecht M, Waser PG. Radioactive labeling and location of specific thiol groups in myosin from fast, slow and cardiac muscles. *Biochim Biophys Acta* 1975 Nov 20;410(1):193-209. [Medline: [73](#)]
17. Gibson P, Powell H, Coughlan J, Wilson A, Abramson M, Haywood P, et al. Self-management education and regular practitioner review for adults with asthma. *Cochrane Database Syst Rev* 2003(1):CD001117. [doi: [10.1002/14651858.CD001117](#)] [Medline: [12535399](#)]
18. Kim J, Shin W. How to do random allocation (randomization). *Clin Orthop Surg* 2014 Mar;6(1):103-109 [FREE Full text] [doi: [10.4055/cios.2014.6.1.103](#)] [Medline: [24605197](#)]
19. Juniper EF, O'Byrne PM, Guyatt GH, Ferrie PJ, King DR. Development and validation of a questionnaire to measure asthma control. *Eur Respir J* 1999 Oct;14(4):902-907 [FREE Full text] [Medline: [10573240](#)]
20. Juniper EF, Guyatt GH, Cox FM, Ferrie PJ, King DR. Development and validation of the Mini Asthma Quality of Life Questionnaire. *Eur Respir J* 1999 Jul;14(1):32-38 [FREE Full text] [Medline: [10489826](#)]
21. Zairina E, Abramson MJ, McDonald CF, Li J, Dharmasiri T, Stewart K, et al. Telehealth to improve asthma control in pregnancy: A randomized controlled trial. *Respirology* 2016 Dec;21(5):867-874. [doi: [10.1111/resp.12773](#)] [Medline: [27037722](#)]
22. Lim AS, Stewart K, Abramson MJ, Walker SP, Smith CL, George J. Multidisciplinary Approach to Management of Maternal Asthma (MAMMA): a randomized controlled trial. *Chest* 2014 May;145(5):1046-1054. [doi: [10.1378/chest.13-2276](#)] [Medline: [24522786](#)]
23. Juniper EF, Bousquet J, Abetz L, Bateman ED, GOAL Committee. Identifying 'well-controlled' and 'not well-controlled' asthma using the Asthma Control Questionnaire. *Respir Med* 2006 Apr;100(4):616-621 [FREE Full text] [doi: [10.1016/j.rmed.2005.08.012](#)] [Medline: [16226443](#)]

Abbreviations

ACQ: Asthma Control Questionnaire

Edited by G Eysenbach; submitted 24.02.18; peer-reviewed by E Arden-Close, A Gobbi; comments to author 11.05.18; revised version received 31.07.18; accepted 04.09.18; published 18.12.18

Please cite as:

Zairina E, Nugraheni G, Achmad GNV, Sulistyarini A, Nita Y, Bakhtiar A, Amin M

Efficacy of an Education Session by Pharmacists for Patients With Asthma: Protocol and Design of a Randomized Controlled Trial

JMIR Res Protoc 2018;7(12):e10210

URL: <https://www.researchprotocols.org/2018/12/e10210/>

doi: [10.2196/10210](#)

PMID: [30563816](#)

©Elida Zairina, Gesnita Nugraheni, Gusti NV Achmad, Arie Sulistyarini, Yunita Nita, Arief Bakhtiar, Muhammad Amin. Originally published in JMIR Research Protocols (<http://www.researchprotocols.org>), 18.12.2018. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in JMIR Research Protocols, is properly cited. The complete bibliographic information, a link to the original publication on <http://www.researchprotocols.org>, as well as this copyright and license information must be included.

Efficacy of an Education Session by Pharmacists for Patients With Asthma: Protocol and Design of a Randomized Controlled Trial

ORIGINALITY REPORT

16%

SIMILARITY INDEX

6%

INTERNET SOURCES

14%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- 1** Matteo Bonini, Omar S. Usmani. "Novel methods for device and adherence monitoring in asthma", *Current Opinion in Pulmonary Medicine*, 2018
Publication 3%
- 2** Bain, Emily, Kristen L Pierides, Vicki L Clifton, Nicolette A Hodyl, Michael J Stark, Caroline A Crowther, Philippa Middleton, and Emily Bain. "Interventions for managing asthma in pregnancy", *Cochrane Database of Systematic Reviews*, 2014.
Publication 2%
- 3** "Cell Biology and Immunology SIG 2", *Respirology*, 2015
Publication 1%
- 4** Konlawij Trongtrakul, Chookiat Feemuchang. "Prevalence and association of vitamin D deficiency and mortality in patients with

severe sepsis", International Journal of
General Medicine, 2017

Publication

5

Georgia Kourlaba, Petros Bakakos, Stelios Loukides, Katerina Vellopoulou, Argyro Solakidi, Nikos Maniadas. "The self-reported prevalence and disease burden of asthma in Greece", Journal of Asthma, 2018

Publication

1 %

6

S. M. van der Kooij. "Limited efficacy of conventional DMARDs after initial methotrexate failure in patients with recent-onset rheumatoid arthritis treated according to the Disease Activity Score", Annals of the Rheumatic Diseases, 2/9/2007

Publication

1 %

7

Elida Zairina, Kay Stewart, Michael J Abramson, Johnson George. "The effectiveness of non-pharmacological healthcare interventions for asthma management during pregnancy: a systematic review", BMC Pulmonary Medicine, 2014

Publication

1 %

8

www.pure.ed.ac.uk

Internet Source

1 %

9

Damilola Olufemi-Yusuf, Sophie Gabriel, Tatiana Makhinova, Lisa Guirguis. "'Being in Control of My Asthma Myself' Patient

1 %

Experience of Asthma Management: A Qualitative Interpretive Description", Pharmacy, 2018

Publication

10

"Poster Session Clinical", European Journal of Heart Failure, 2013.

Publication

<1 %

11

Rehan, V. K., J. Liu, R. Sakurai, and J. S. Torday. "Perinatal Nicotine-Induced Transgenerational Asthma", AJP Lung Cellular and Molecular Physiology, 2013.

Publication

<1 %

12

www.bmj.com

Internet Source

<1 %

13

eprints.nottingham.ac.uk

Internet Source

<1 %

14

inba.info

Internet Source

<1 %

15

Rose, Olaf, Corinna Schaffert, Kathrin Czarnecki, Hugo S. Mennemann, Isabel Waltering, Stefanie Hamacher, Moritz Felsch, Lena Herich, and Juliane Köberlein. "Effect evaluation of an interprofessional medication therapy management approach for multimorbid patients in primary care: a cluster-randomized controlled trial in

<1 %

community care (WestGem study protocol)",
BMC Family Practice, 2015.

Publication

16

corpus.ulaval.ca

Internet Source

<1 %

17

Karen B. Schmaling. "Asthma", Wiley, 2012

Publication

<1 %

18

M Arora, L A Harvey, J V Glinsky, H S Chhabra,
S Hossain, N Arumugam, P K Bedi, L
Lavrencic, A J Hayes, I D Cameron.

"Telephone-based management of pressure
ulcers in people with spinal cord injury in low-
and middle-income countries: a randomised
controlled trial", Spinal Cord, 2016

Publication

<1 %

19

Nicole W. Tsao, Amir Khakban, Louise
Gastonguay, Kathy Li, Larry D. Lynd, Carlo A.
Marra. "Perceptions of British Columbia
residents and their willingness to pay for
medication management services provided by
pharmacists", Canadian Pharmacists Journal /
Revue des Pharmaciens du Canada, 2015

Publication

<1 %

20

Richard N van Zyl-Smit, Matthias Krüll,
Christian Gessner, Yasuhiro Gon et al. "Once-
daily mometasone plus indacaterol versus
mometasone or twice-daily fluticasone plus
salmeterol in patients with inadequately

<1 %

controlled asthma (PALLADIUM): a randomised, double-blind, triple-dummy, controlled phase 3 study", The Lancet Respiratory Medicine, 2020

Publication

21

respiratory-research.biomedcentral.com

Internet Source

<1 %

22

Danielle Santos Alves, Valéria Cesário Times, Érika Maria Alves da Silva, Priscila Santos Alves Melo et al. "Advances in obstetric telemonitoring: a systematic review", International Journal of Medical Informatics, 2020

Publication

<1 %

23

Emily Bain, Kristen L Pierides, Vicki L Clifton, Nicolette A Hodyl, Michael J Stark, Caroline A Crowther, Philippa Middleton. "Interventions for managing asthma in pregnancy", Cochrane Database of Systematic Reviews, 2014

Publication

<1 %

24

Majed Almutairi, Khalid Al Sulaiman, Sultan Alenazi, Ramesh Vishwakarma, Ohoud Aljuhani. "Evaluation of standard dosing for selected broad spectrum Hydrophilic antibiotics in critically ill patients with Augmented renal clearance: An observational Study", Research Square Platform LLC, 2021

<1 %

25 [docslide.us](https://www.docslide.us) <1 %
Internet Source

26 ira.le.ac.uk <1 %
Internet Source

27 "Respiratory Infectious Diseases SIG: Poster Session", *Respirology*, 3/2008 <1 %
Publication

28 "Asthma, Allergic and Immunologic Diseases During Pregnancy", Springer Science and Business Media LLC, 2019 <1 %
Publication

29 Joanne M. Fuller, Keith K. Wong, Camilla Hoyos, Ines Krass, Bandana Saini. "Dispensing good sleep health behaviours not pills - a cluster-randomized controlled trial to test the feasibility and efficacy of pharmacist-provided brief behavioural treatment for insomnia", *Journal of Sleep Research*, 2016 <1 %
Publication

30 Mohit Arora, Lisa Anne Harvey, Alison Joy Hayes, Harvinder Singh Chhabra et al. "Effectiveness and cost-effectiveness of telephone-based support versus usual care for treatment of pressure ulcers in people with spinal cord injury in low-income and middle-income countries: study protocol for a

12-week randomised controlled trial", BMJ

Open, 2015

Publication

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

Efficacy of an Education Session by Pharmacists for Patients With Asthma: Protocol and Design of a Randomized Controlled Trial

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8
