

Pharmacy Practice

HOME ABOUT - IN PRESS CURRENT ARCHIVES AUTHORS FOR REVIEWERS

Home / Editorial Team

Editorial & Advisory Boards

Pharmacy Practice Editorial Board

David A. Holdford (associate editor) Virginia Commonwealth University (USA)

Shalom I. Benrimoj University of Granada (Spain)

Karen B. Farris University of Michigan (USA)

Martin Schulz
Freie Universitaet Berlin and ABDA – Federal Union of German Associations of Pharmacists (Germany)

Cassyano J. Correr Federal University of Parana (Brazil)

Fernanda S. Tonin Federal University of Parana (Brazil)

Jonathan Penm University of Sydney (Australia)

Meagen Rosenthal University of Mississippi (USA)

Carrie M. Blanchard University of North Carolina (USA)


Sarah Dineen-Griffin University of Technology Sydney (Australia)

Advisory Board

Marja Airaksinen, University of Halsiniki (Finland)
Simon Bell, University of South Australia (Australia)
Mana Cordina, University of Malifa (Malifa)
Souheil Hallit, Holy Spirit University of Kasilik (Jebanon)
Mohamed A. Hassali, University of Isasilik (Jebanon)
Mohamed A. Hassali, University of Growe (USA)
Carmel Hughes, Queen's University Belfost (UK)
Ulrich Jeade, University of Bortish Columbia (Canada)
Dawof Nau, University of Bortish Columbia (Canada)
Dawof Nau, University of Bortish Columbia (Canada)
Julie A. Patterson, Virginia Commonweath University (USA)
Bandana Saini, University of Sydney (Australia)
Anthony Serradino-Inglott, University of Malifa (Malifa)
Mary Tully, University of Manchester (UK)
Asim Ahmed Elnour, Al Ain University (UAE)

Pharmacy Practice

HOME ABOUT - IN PRESS CURRENT ARCHIVES AUTHORS FOR REVIEWERS Home / Archives / Vol. 17 No. 3 (2019); Jul-Sep □ jdi A transition from the BPharm to the PharmD degree in five selected countries Teerapom Supapsan, Bee Y. Low, Payom Wongpoowersk, Summans Moolasarn, Claire Anderson Abstract ● 4624 | pdf Downloads ≟ 1299 国 Page 1611 ORIGINAL RESEARCH □ pdf Role of community pharmacists in skin cancer screening: A descriptive study of skin cancer risk factors prevalence and photoprotection habits in Barcelona, Catalonia, Spain Joan F. Mir, Maria Estrada-Compmany, Anna Heredia, Cristina Rodríguez-Caba, Marta Alcalde, Natalia Espinosa, Illilana Monnier, Javiera Férez-Anker, Oriol Vélamos, Susana Puig, Núria Bosch, Josep Malvehy Abstract ூ 1330 | pdf Downloads ± 568 online appendix Downloads ± 0 ☐ Page 1455 **≥** par Biosimilars and implications for pharmacy practice: Ready or not, here they come! Joshua P. Crawford, Andrea I. Hobbs Abstract № 1609 | pdf Downloads 🕹 574 🖻 Page 1659 i par The Syrian refugee crisis in Jordan: a cross sectional pharmacist-led study assessing post-traumatic stress disorder. Iman A. Basheti, Shahnaz M. Ayasrah, Marian M. Basheti, Judeh Mahfuz, Betty Chaar Abstract © 1064 | pdf Downloads 去 440 online appendix Downloads 去 0 国 Page 1475 D jet Pharmacists and Medicare's Annual Wellness Visit: implications for pharmacy education and interprofessional primary care Kristin Zimmerman, Daniel Bluestein Abstract ூ 703 | pdf Download: ₺401 国 Page 1672



Pharmacy Practice

HOME ABOUT - IN PRESS CURRENT ARCHIVES AUTHORS FOR REVIEWERS

Home / Archives / Vol. 17 No. 3 (2019): Jul-Sep. / Original Research

Assessment of pharmacists' knowledge, attitude and practice in chain community pharmacies towards their current function and performance in Indonesia

Umi Athiyah http://orcid.org/2000-0003-4818-6847 Catur D. Setlawan http://orcid.org/2000-0002-5942-9947 Gesnifa Nugriaheni http://orcid.org/2000-0002-0791-6556 Elida Zairina http://orcid.org/2000-0003-5887-0846 Wahyu Ulani http://orcid.org/2000-0003-5887-0846 Andi Hermansyah http://orcid.org/2000-0002-9716-3126

Keywords

Professional Fractice, Pharmacies, Pharmaceutical Services, Fharmacists, Self Medication, Surveys and Questionnaires, Indonesia

Abstract

Background: The introduction of universal healthcare coverage in 2014 has affected the practice of community pharmacies in Indonesia. Studies regarding the practice of pharmacist in the chain community pharmacy extension in Indonesia are every limited. The chain community pharmacies in Indonesia are expected by the same management. The chain community pharmacies usually show better services compared to independent community pharmacies in Indonesia.

Objective: The study aimed to assess the knowledge, attitude and practice (KAP) of pharmacist working in chain community pharmacy towards their current function and performance in delivering pharmacy services

Methods: A cross-sectional study using questionnaires was conducted between January and March 2017 in KF, one of the largest chain community pharmacies in Indonesia. The total sampling method was used in the recruitment process. The data were analyzed using descriptive statistics, independent t-Test and one-way ANOVA. The KAP scores were assessed and categorized as "poort," moderate" and "good" based on the standardized scoring system.

Results: A total of 949 KF's pharmacists (100% response rate) were participated in the study. The majority of pharmacists showed a good score in terms of knowledge and attitude, which is in contrast to practice as majority only obtained a moderate score. Working experience, age and the availability of standard operating procedures (SOP) for both dispensing and self-medication services were found to be statistically significant (p<0.005) aspects to KAP of pharmacists in delivering pharmacy services.

Conclusions: This study identified several important aspects that could affect the KAP of pharmacists working in chain community pharmaciss in indonesia. Specific policies should be conceived to improve the competencies of pharmacist and to ensure the compliance with the SOP and standardization system within pharmacy sector.

D ndf

Aug 24, 2019

How to Cite

Athlyan U, Setiawan CD, Nugraheni G, Zairina E, Utami YV, Hermansyah A. Assessment of pharmacists' knowledge, attitude and practice in chain community pharmacies towards their current function and performance in Indonesia. Pharm Pract (Granada) [Internet]. 2019 Aug. 24 [cited 2023 Apr. 10];17]gi:1518. Available from: https://www.pharmacypractice.org/index.php/pp/arbide/view/1518

More Citation Formats

Issue

Vol. 17 No. 3 |2019|; Jul-Sep

Section

Original Research



This work is licensed under a Creative Commons Attribution-Non-Commercial-NoDerivatives 3.0 Unported License.

Original Research

Assessment of pharmacists' knowledge, attitude and practice in chain community pharmacies towards their current function and performance in Indonesia

Umi ATHIYAH, Catur D. SETIAWAN, Gesnita NUGRAHENI, Elida ZAIRINA, Wahyu UTAMI, Andi HERMANSYAH.

Received (first version): 6-Apr-2019

Accepted: 11-Aug-2019

Abstract

Background: The introduction of universal healthcare coverage in 2014 has affected the practice of community pharmacies in Indonesia. Studies regarding the practice of pharmacist in the chain community pharmacy setting in Indonesia are very limited. The chain community pharmacies in Indonesia are operated and controlled by the same management. The chain community pharmacies usually show better services compared to independent community pharmacies in Indonesia.

Published online: 24-Aug-2019

Objective: The study aimed to assess the knowledge, attitude and practice (KAP) of pharmacist working in chain community pharmacy towards their current function and performance in delivering pharmacy services

Methods: A cross-sectional study using questionnaires was conducted between January and March 2017 in KF, one of the largest chain community pharmacies in Indonesia. The total sampling method was used in the recruitment process. The data were analyzed using descriptive statistics, independent t-Test and one-way ANOVA. The KAP scores were assessed and categorized as "poor", "moderate" and "good" based on the standardized scoring system.

Results: A total of 949 KF's pharmacists (100% response rate) were participated in the study. The majority of pharmacists showed a good score in terms of knowledge and attitude, which is in contrast to practice as majority only obtained a moderate score. Working experience, age and the availability of standard operating procedures (SOP) for both dispensing and self-medication services were found to be statistically significant (p<0.005) aspects to KAP of pharmacists in delivering pharmacy services.

Conclusions: This study identified several important aspects that could affect the KAP of pharmacists working in chain community pharmacies in Indonesia. Specific policies should be conceived to improve the competencies of pharmacist and to ensure the compliance with the SOP and standardization system within pharmacy sector.

Kevwords

Professional Practice; Pharmacies; Pharmaceutical Services; Pharmacists; Self Medication; Surveys and Questionnaires; Indonesia

INTRODUCTION

Community pharmacy is currently undergoing a transformation from its traditional function as a supplier of medicines towards a health hub Internationally, the development of policy has changed the practice of community pharmacy significantly. The increasing pressure to provide effective, efficient and affordable care under constrained healthcare budget has encouraged policymakers and professional bodies to expand pharmacist's contribution within the healthcare system.² For example, a number of cognitive services such as medicine use reviews (MURs), medication therapy management for chronic illness and home medication reviews (HMRs) are now widely available in pharmacies in UK, USA, New Zealand, Australia and Canada. 3,4 Moreover, community pharmacists in some of these countries have also been involved in providing services beyond their conventional scope of practice e.g. health screening, vaccination and prescribing for minor ailments.^{5,6}

The need for pharmacists to change the current practice is crucial in Indonesia. The recent health care system changes marked by the introduction of universal health care coverage (Jaminan Kesehatan Nasional - JKN) in 2014 has created an opportunity for community pharmacist to be more involved in the primary care services.7 This could provide opportunity for pharmacist to expand their roles in community care. However, on the other hand, the preparedness of pharmacist to provide such roles has become a major concern. For example, whilst Indonesia has legislation that a community pharmacy can only be operated under pharmacist's responsibility and supervision, low pharmacist presence has been evident in the large proportion of pharmacies.^{8,9} The similar trend is noticeable in other low and middle-income countries (LMICs) reflecting an intractable problem towards pharmacist role expansion. 10

Since 2018, there were 26,658 community pharmacies in Indonesia with the large proportion (>90%) are independent pharmacies. ¹¹ The Indonesian government give no restriction in the ownership structure of pharmacy and allow corporations and non-pharmacists to own more than one pharmacy. ⁸ As a result, community pharmacy in

Umi ATHIYAH. Dr. MS. Dean & Professor. Faculty of Pharmacy, Airlangga University. Surabaya (Indonesia). umi-a@ff.unair.ac.id Catur Dian SETIAWAN. MKes. Lecturer. Faculty of Pharmacy, Airlangga University. Surabaya (Indonesia). catur-d-s@ff.unair.ac.id Gesnita NUGRAHENI. M. Pharm. Lecturer. Faculty of Pharmacy, Airlangga University. Surabaya (Indonesia). gesnita-n@ff.unair.ac.id Elida ZAIRINA. PhD. Associate Professor. Faculty of Pharmacy, Airlangga University Surabaya (Indonesia). elida-z@ff.unair.ac.id Wahyu UTAMI. Dr. MS. Associate Professor. Faculty of Pharmacy, Airlangga University. Surabaya (Indonesia). wahyu-u@ff.unair.ac.id Andi HERMANSYAH. PhD. Associate Professor. Faculty of Pharmacy, Airlangga University. Surabaya (Indonesia). andi-h@ff.unair.ac.id.



Indonesia ranges from small independent community pharmacies to a large network of chain community pharmacies, most of which are operated in private sectors. Whilst independent pharmacy is generally owned by individual or group of individuals, the chain community pharmacy model can be classified into three main categories. The first category is chain pharmacy operated under centralized management structures. This means the local pharmacy branches must firmly follow the policy and management system as regulated by the main company for running the business including in logistic and staffing. KF, Century and Guardian are some examples of chain pharmacies operated under this model. 12 The second category is chain pharmacy operated under franchise system. This type of pharmacy allows local branches to selfregulated and use brand identity for the pharmacy operation (e.g. K-24 pharmacy). 12 The last category is buying groups pharmacy. While the first two pharmacy models are operated under organized chain system and use a unified business name, the buying groups are formed by a number of individual pharmacies to collectively increase their purchasing power, thus obtaining cheaper prices than they would possibly do individually. Apart from purchasing purposes, this type of pharmacy is independently operated by the owner. 12

Community pharmacy in Indonesia has been known as the primary source for public to obtain medicines and pharmacy services. While the majority of pharmacies focused on the supply function of medicines and retail services, there is a small proportion provided extended services such as early detection and health management services as an adjunct to dispensing model. Typically, pharmacies providing such services are operating under chain model pharmacy. This is not surprising as these pharmacies generally have at least one pharmacist in duty to organize and supervise pharmacy operation. Arguably, this has made such pharmacies are in greater position to maximize opportunity within the universal health care era.

The evidence in other LMICs demonstrated that pharmacist working in chain pharmacies had a better understanding and attitude towards pharmaceutical care and provided more advanced services as compared to their colleagues working in individual pharmacies. 16 Nevertheless, there is a paucity of research investigating the performance of pharmacists working in chain community pharmacies in Indonesia. The corporatization of pharmacy in Indonesia on one hand has become a consolidated effort to provide better access and larger economics scale in the increased of competition and dynamic changes within pharmacy and healthcare sectors. 17 However, on the other hand, the forces of corporatization have been found to challenge the autonomy and professional rigor of pharmacists leading to moral dilemmas experienced by pharmacist regarding their professional roles. 18,19 Evidence has shown that the performances of independent pharmacies in Indonesia were below the minimum standards, however there were no evidence that the chain community or franchise model pharmacies gave the same results. 20 Therefore, this study aimed to assess the knowledge, attitude and practice of pharmacist working in chain community pharmacy towards their current function and performance in delivering pharmacy services.

METHODS

Study design and setting

A cross-sectional study was conducted between January and March 2017 using a self-developed and pre-validated questionnaire to assess the knowledge, attitude and practice (KAP) of chain community pharmacist working in KF, one of the largest chain pharmacy networks in Indonesia. All pharmacists signed the consent form and willing to participate were included in this study. The exclusion criteria included pharmacists who did not complete the survey responses.

Study instrument

The questionnaire was developed from the literatures regarding the pharmacists' work experiences and attitude towards pharmaceutical care services in Indonesia, also based on the discussion among research team and feedback from KF managerial members.^{21,22} questionnaire was developed in Indonesian language and it was consisted of five sections. Section one comprised four questions about the demographic profile including gender, age, educational level and years of working in KF. Section two (11 questions) inquiring about the pharmacy characteristics including the average number of patients, the availability of Standard Operating Procedure (SOP) and the number of workforces in pharmacy (pharmacists and non-pharmacists). Section 3, 4 and 5 focusing on Knowledge (24 questions), Attitude (34 questions) and Practice (26 questions), respectively.

Questions pertaining to KAP included concept of pharmaceutical care, services evaluation, patient expectation, understanding and compliance regarding drug therapy. A correct answer for each item of knowledge was given a score of 1, while wrong and unsure answer were marked as 0 (zero). The Likert scales were adopted in the section of attitude (four scales ranging from strongly agree to strongly disagree) and practice (four scales ranging from always to never), respectively. Highest score (4 points) reflected the most positive option and the lowest score (1 point) reflected the most negative option. The maximum score from each KAP aspect was used to rank the level of knowledge, attitude and practice, respectively. Subsequent analysis was conducted to classify the score as "poor" (low score), "moderate" (medium score) and "good" (high score). For example, if the respondent answered all questions correctly in Knowledge aspect, 24 scoring points were given. The total of 24 points were divided into three categories in which 0-8 points attributed to poor, 9-16 points attributed to moderate and 17-24 points attributed to good knowledge.

The representatives of the Faculty and KF examined and approved the face and content validity of the questionnaire. The questionnaire was then piloted to ten KF's pharmacists resulted in minor changes on the wording and questionnaire lay-out. The Internal consistency of the modified questionnaire was measured using Cronbach



https://doi.org/10.18549/PharmPract.2019.3.1518

Table 1. Characteristics of participants (N=949)				
· ·	· · · · · · · · · · · · · · · · · · ·			
Characteristics	n (%)			
Gender				
Male	390 (41.1)			
Female	559 (58.9)			
Age group (years)				
<u>≤</u> 30	694 (73.2)			
31-40	172 (18.1)			
41-50	58 (6.1)			
> 50	25 (2.6)			
Educational level				
Undergraduate + Pharmacist	915 (96.4)			
Master + Pharmacist	33 (3.5)			
Doctor + Pharmacist	1 (0.1)			
Working experience in KF (years)				
< 5	728 (76.7)			
5-10	111 (11.7)			
11-15	49 (5.2)			
> 15	61 (6.4)			

Alpha resulting coefficient 0.724 reflecting good internal consistency of the questionnaire (>0.70). Ethics approval was not deemed necessary as the study was part of internal audit to observe performance of pharmacists and pharmacy. However, each participant was required to provide signed consent prior to fill the questionnaire. In addition, participants received information about the nature of the study and procedure for completing the questionnaire.

Participants and data analysis

The study used a total sampling method to recruit all community pharmacists (n=949) working in 825 KF pharmacy outlets throughout the country (34 provinces). The questionnaire was distributed to each pharmacist through KF system in which the research team was provided access to collect the results. The data were analyzed using SPSS v22. The categorical variables were presented as frequencies and percentages while the continuous variables were reported as mean and standard deviation (SD). Independent t-Test and one-way ANOVA were used to compare means between KAP groups with pvalue < 0.05 was considered to be statistically significant. To address the objective of this study, five variables were analyzed as indicator for pharmacists' performance in delivering services with respect to their KAP, namely working experience, age, educational level and the availability of SOP for both dispensing and self-medication services. These variables were selected as they were intrinsic factors that contribute to the performance of individual pharmacist in delivering the service.²⁴

RESULTS

A total of 949 pharmacists returned the questionnaire resulting in 100% response rate. The majority of participants are female. Most of the participants are novice pharmacists with aged less than 30 years old with working experience in KF less than 5 years. Characteristics of the respondent are provided in Table 1.

The majority of pharmacies served less than 20 patients with prescriptions each day (Table 2). In contrast, the provision of self-medication was relatively high with 20-100 patients were treated as self-medication per day. As a

Table 2. Characteristics of KF pharmacies (N=949)					
Characteristics n (%)					
The average number of patients with prescriptions per day					
< 20	462 (48.7)				
20-50	278 (29.3)				
51-100	140 (14.8)				
> 100	69 (7.2)				
The availability of SOP for dispensing prescribed medicines					
Available	855 (91)				
Not available	94 (9)				
The average number of patients for self-medicat	. , ,				
< 20	80 (8.4)				
20-50	348 (36.7)				
51-100	339 (35.7)				
> 100					
The availability of SOP for self-medication servic	182 (19.2)				
l ·					
Available Net available	811 (85)				
Not available	138 (15)				
The average number of patients purchasing med	dical devices with				
prescriptions per day					
< 20	867 (91.4)				
20-50	63 (6.6)				
51-100	11 (1.2)				
> 100	8 (0.8)				
The number of pharmacists in each pharmacy	i				
1	813 (85.6)				
2	113 (11.9)				
3	12 (1.3)				
4	6 (0.6)				
5	3 (0.3)				
>5	2 (0.2)				
The number of pharmacy assistants in each pharmacy					
1	247 (26)				
2	406 (42.8)				
3	118 (12.4)				
4	97 (10.2)				
5	38 (4)				
>5	43 (4.5)				
The number of other pharmacy staffs (non-pharmacists and non-					
pharmacy assistants)					
1	819 (86.3)				
2	87 (9.2)				
3	23 (2.4)				
4	8 (0.8)				
5	3 (0.3)				
>5	9 (0.9)				
SOP = Standard operating procedure	•				

corporate chain pharmacy network, the availability of SOP for services is highly important. Although all pharmacies were expected to have SOP for delivering their services as required by the legislation, there was small yet significant proportion which deviated from such policy, both SOP for dispensing services (9%) and SOP for self-medication services (15%), respectively.²⁵

It was not surprising that majority of participants had good knowledge and attitude about their function and services they provided in pharmacy. However, the overall score for practice reflected a contradiction to this notion. The majority of participants only had moderate practice with relatively high proportion scored poor practice (15%). Table 3 shows the distribution of score for KAP and frequency of participant with particular level of KAP.

Table 4 displays the comparison between KAP and selected variables. Three statistically significant differences (p<0.05) were found namely (1) between attitude-practice and working experience, (2) between attitude-practice and age,



https://doi.org/10.18549/PharmPract.2019.3.1518

Table 3. Distribution of KAP score and frequency of participant with particular lev	el of knowledge, attitud	es and praction	e.
	Mean (SD)	Minimum	Maximum
Total knowledge score	22.73 <u>(</u> 1.77)	14	24
Total attitude score	119.99 <u>(</u> 11.98)	89	136
Total practice score	85.75 <u>(</u> 12.29)	47	104
	Poor	Moderate	Good
Number of participants with particular level of knowledge. n(%)	-	7 (0.7)	942 (99.3)
Number of participants with particular level of attitude. n(%)	-	92 (9.7)	857 (90.3)
Number of participants with particular level of practice. n(%)	146 (15.4)	803 (84.6)	-

and (3) between knowledge-attitude-practice and availability of SOP both for dispensing and self-medication services. There was no significant difference in terms of educational level with the KAP score.

With respect to working experience, it was found that pharmacists who worked for 5-10 years in KF had the lowest score for attitude and practice. Pharmacist aged of 31-40 had the lowest level of attitude, while the age above 50 years had the lowest score of practice. Moreover, pharmacies which did not have SOP resulted in the lowest score in three elements of KAP.

DISCUSSION

This study aimed to assess KAP of pharmacists working in Indonesian chain community pharmacy towards their function and performance in delivering pharmacy services. There were three main findings which should be part of concerns both for pharmacists as a profession and pharmacy stakeholders particularly the owners/managers of pharmacies in Indonesia. Firstly, it was observed that pharmacists who already worked for 5 to 10 years showed the lowest score in terms of attitude and practice. While this study was unable to explore the factors behind such result, pharmacists' burnout, job stress and dissatisfaction to work were reported to moderately increase in the later stage of work. The escalating pharmaceutical demand and dynamic changes in pharmacy practice have contributed to

the increase of pharmacists' workload and requirement for more interpersonal interaction with the patients.²⁷ A number of literatures mentioned that individuals who already worked for 10 years in particular setting tend to experience exhaustion and burnout feeling from their works. ²⁸⁻³⁰ Specific to KF, the low volumes of prescription in contrast to high volumes of self-medication consultations to some extent may have put pharmacist under constant pressure of conflict of interest affecting their professionalism. It is true that pharmacist can play unique role in self-medication for instance by facilitating individuals for appropriate self-care or medication. However, in environment where commercial-consumerism influences inflict with personal-professional decision like in pharmacy, such practice can be problematic. Several studies highlighted the lack of pharmacist ability to consistently and appropriately facilitate self-medication which may lead to adverse reaction and public risk e.g. drug resistance. 32,33 Such situation is without a doubt affecting pharmacists' behavior in delivering services which may become greater concern in the future if it goes unrecognized and untreated. This study implied that pharmacists are in a need of professional practice development to update, refresh and maintain their professional roles and competence. Such attempt may take in a number of forms ranging from an individual training and continuing education to a collective practice building such as teamwork exercise.

Madala		Knowledge		Attitude		Practice				
Variable N	N	Mean	Std. Error	Sig.	Mean	Std. Error	Sig.	Mean	Std. Error	Sig.
Working experience in KF (years)*										
< 5	728	22.780	0.063	0.336	120.0618	0.438	0.005 ⁺	86.520	0.441	0.015
5-10	111	22.522	0.192		116.3604	1.229		81.657	1.227	
11-15	49	22.551	0.331		121.7551	1.666		83.612	1.829	
> 15	61	22.721	0.195		124.4262	1.343		85.852	1.760	
Age (years)*										
<u>≤</u> 30	694	22.796	0.063	0.336	119.956	0.450	0.005 ⁺	86.448	0.455	0.015
31-40	172	22.564	0.154		118.546	0. 973		83.337	0.959	
41-50	58	22.620	0.240		125.034	1.262		85.982	1.671	
> 50	25	22.440	0.416		119.400	2.509		82.760	2.990	
Educational level*				-						
Undergraduate + pharmacist	915	22.732	0.058	0.772	119.875	0.396	0.229	85.617	0.403	0.119
Master + pharmacist	33	22.757	0.261		123.060	2.077		89.212	2.492	
Doctor + pharmacist**	1	24.000	1		130.000	-		101.000	-	
The availability of SOP for dispensing	g preso	ribed med	licines**	-						
Available	855	22,773	0.060	0.043 ⁺	120.366	0. 406	0.004 ⁺	86.349	0.414	<0.001
Not Available	94	22,383	0.190		116.638	1.286		80.383	1.312	
The availability of SOP for self-medi	cation	services**								
Available	811	22,812	0.061	0.001 ⁺	120.607	0.416	<0.001 ⁺	86.612	0.422	<0.001
Not Available	138	22,275	0.159		116.4058	1.029		80.739	1.070	

^{*}Using one-way ANOVA to compare means between three or more groups

^{*}Statistically significant (p<0.05)



^{**}Using Independent t Test to compare means between two independent groups

Secondly, age of pharmacists influences the attitude and practice in delivering care regardless their knowledge and educational level. Pharmacists in KF were found to have overall good knowledge and understanding about their function and performance in services. This is perhaps due to the fact that they have pursued adequate educational program and acquired basic and essential knowledge to become a pharmacist. Additionally, the corporate system in KF may have forged this knowledge through capacity building program.³⁴ However, pharmacists who fall into particular group of age had the lowest score for attitude (31-40 years old) and practice (>50 years old) which becomes a fact that should not be ignored by KF. It is not surprising that pharmacists in the late career or pension age may experience a diminished performance and function. Several studies have indicated that pharmacists who have been in the profession for 30 years may intend to leave the profession. 35,36 What might be greater concern is that early career person, 31-40 years as reflected in this study, also experienced a decreasing attitude towards their practice. This may a warranty to examine developmental process between career and life stage especially in KF. A study by Magola et al. indicated that early-career pharmacist was challenged with the same routine and responsibilities in the pharmacy as late-career pharmacist.³⁷ This may influence the transition of novice practitioners into practice and having negative implications for patient care. Pharmacists in this group of age are likely to find certain things – desire, expectation and motivation to deserve more – in work which are translated in their attitudes and intentions.³⁸ The ability to identify and response to such condition can be a winning program for a pharmacy particularly KF. What is perhaps interesting is the finding that pharmacist aged 41-50 years showed better overall KAP results as compared to the early and late career pharmacist. In terms of career management, this study argued that mid-career professionals may have developed resilience and adaptability, both intellectually and personally in viewing their career.³⁹ A number of studies underlined that mid-career level professionals have strong commitment and endorsement for learning, increasing autonomy and supporting others. 40-42 The key challenge, however, is to maximize such adaptive behavior to help pharmacists realize their career aspirations and hone their full potential constructively.

Thirdly, the availability of SOP significantly influenced the KAP of pharmacists. This study revealed that pharmacies that were not equipped with SOP for services proved to have the lowest level of three element of pharmacists' KAP. WHO and FIP have underlined the importance of having SOP as step-by-step guidance for good pharmacy practice. The findings in England and Wales showed that the availability of SOP has helped pharmacist in applying procedures to their everyday work. This is particularly the case for novice pharmacists which were relied on SOP as a guide to practice.⁴⁴ In Indonesia, pharmacies are required to have a number of SOP from the procurement, storage of medicines, dispensing and services to elimination of expired medicines.²⁵ This study adds to this notion that the availability of SOP can be an indicator for good pharmacy practice and therefore it is essential for the government and pharmacy stakeholders to ensure that SOP for service is available in every pharmacy. Learning from KF, a top down approach can be one way to encourage the availability of SOP. The fact that pharmacies involved in this study were operated under centralized management might explain why there was only small proportion of pharmacies without SOP in KF. It is important to note that centralized management system as developed by KF is likely to reduce variation and gap of performance between pharmacies which is common in pharmacies operating in LMICs. Alternatively, the government in LMICs can endorse the availability of SOP for example by including requirement to have SOP prior seeking approval and renewal of pharmacy permits which is currently the norm in Indonesia.

To the best of our knowledge, this is the first study examining KAP of pharmacists towards their function and performance in delivering pharmacy services in the setting of chain pharmacy in Indonesia. The findings of this study provoke deeper understanding about pharmacists' role and behavior and provide insight to the influence of a number of important aspects such as age, working experience and availability of SOP for services. This is similar to the findings of a study by Schafheutle et al. concluding that age and factors associated with workplace affected pharmacists' performance.²⁴ In addition, this study also highlights three main tasks for policymakers and pharmacy stakeholders in Indonesia. First, as a pharmacist continues to age and he/she constantly experiences pressure for change, there is significant need for training and competency development program to improve pharmacist's professional practice. Such training may comprise any unique approach to boost the cognitive, moral, emotional and social side of the pharmacist. Second, despite there are policies in place that requires pharmacy to have SOP, the evaluation for compliance to the SOP is unnoticed. This reflects more specific efforts should be made to ensure pharmacy compliance with the regulation and documentation system. Third, this study demonstrates that practice of pharmacist in chain community pharmacies is overrated despite the good level of knowledge and attitude as shown by the pharmacists. As the co-existence of chain and independent pharmacies is crucial to Indonesian healthcare system, it is imperative that there is a standardization system e.g. accreditation program to ensure that community pharmacy meets the minimum standard for practice. Such program is literally not available in Indonesian pharmacy legislation.

This study, however, is not without limitations. The administration of the questionnaire which is part of internal audit process may affect the integrity and honesty of the response. While the research team strived to ensure that participation in this study did not affect individual pharmacist *per se*, there is no guarantee that such method was effective to overcome bias in participants' response. Subsequently, despite the effort to measure pharmacist KAP based on five selected variables, this study did not analyze other variables such as the number of patients, number of pharmacists and other pharmacy characteristics which may or may not influence the KAP results. This is a call for further research to observe other variables using KAP approach.



https://doi.org/10.18549/PharmPract.2019.3.1518

CONCLUSIONS

This study examines pharmacist's function and performance working in chain pharmacy in Indonesia which can be an insight for policymakers and managers/owners in the context of LMICs. Using KAP approach, this study highlights that working experience, age and availability of SOP for services are significant aspects affecting pharmacists' function and performance. Evidence of poor practice were apparent indicating an imperative for improving pharmacist competence, ensuring compliance to SOP and standardization system in pharmacy.

ACKNOWLEDGEMENTS

The authors gratefully acknowledge and thank KF for collaborating and providing access to this research. We also thank participants of this study who have provided time and efforts for this study.

CONFLICT OF INTEREST

None Declared.

FUNDING

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

References

- 1. McMillan SS, Wheeler AJ, Sav A, King MA, Whitty JA, Kendall E, Kelly F. Community pharmacy in Australia: a health hub destination of the future. Res Social Adm Pharm. 2013;9(6):863-875. https://doi.org/10.1016/j.sapharm.2012.11.003
- 2. Scahill S, Harrison J, Carswell P, Shaw J. Health care policy and community pharmacy: implications for the New Zealand primary health care sector. N Z Med J. 2010;123(1317):41-51.
- 3. Patwardhan PD, Amin ME, Chewning BA. Intervention research to enhance community pharmacists' cognitive services: a systematic review. Res Social Adm Pharm. 2014;10(3):475-493. https://doi.org/10.1016/j.sapharm.2013.07.005
- 4. Benrimoj S, Feletto E, Gastelurrutia MÁ, Martínez-Martínez F, Faus Dáder MJ. A holistic and integrated approach to implementing cognitive pharmaceutical services. Ara Pharmaceutica. 2010;51(2):69-87.
- Hunt BD, Hiles SL, Chauhan A, Ighofose C, Bharakhada N, Jain A, Davies MJ, Khunti K. Evaluation of the Healthy LifeCheck programme: a vascular risk assessment service for community pharmacies in Leicester city, UK. J Public Health (Oxf). 2013;35(3):440-446. https://doi.org/10.1093/pubmed/fdt017
- 6. Laliberté MC, Damestoy N, Perreault S, Lalonde L. The role of community pharmacists in health promotion and prevention. Can Pharm J. 2012;145(4):S6.
- 7. Hermansyah A, Sainsbury E, Krass I. Investigating the impact of the universal healthcare coverage programme on community pharmacy practice. Health Soc Care Community. 2018;26(2):e249-e260. https://doi.org/10.1111/hsc.12506
- 8. Ministry of Health Indonesia. Permenkes No. 9 tahun 2017 tentang Apotek (Community Pharmacy Decree). Jakarta: Ministry of Health Indonesia; 2017.
- 9. Hermansyah A, Pitaloka D, Sainsbury E, Krass I. Prioritising recommendations to advance community pharmacy practice. Res Social Adm Pharm. 2018;14(12):1147-1156. https://doi.org/10.1016/j.sapharm.2018.02.003
- 10. Miller R, Goodman C. Performance of retail pharmacies in low- and middle-income Asian settings: a systematic review. Health Policy Plan. 2016;31(7):940-953. https://doi.org/10.1093/heapol/czw007
- 11. Ministry of Health Indonesia. Recapitulation of Community Pharmacy in Indonesia (2018) Jakarta: Ministry of Health Indonesia. Available at: http://apif.binfar.depkes.go.id/grafik-apotek.php (accessed Jul 29, 2019).
- Ken Research Private Ltd. Indonesia Pharmacy Retail Market Outlook to 2022 by Pharmacies & Drugstores and By Region (Central Java, East Java, West Java, North Sumatra, Jakarta and Others). Indonesia: Ken Research Private Limited, 2018 ID: 4586463.
- 13. Puspitasari H, Faturrohmah A, Hermansyah A. Do Indonesian community pharmacy workers respond to antibiotics requests appropriately? Trop Med Int Health. 2011;16(7):840-846. https://doi.org/10.1111/j.1365-3156.2011.02782.x
- Hermansyah A, Sainsbury E, Krass I. Community pharmacy and emerging public health initiatives in developing Southeast Asian countries: a systematic review. Health Soc Care Community. 2016;24(5):e11-e22. https://doi.org/10.1111/hsc.12289
- 15. Andayani TM, Satibi S. Pharmacy practice in Indonesia. In: Fathelrahman A, Ibrahim MI, Wertheimer AI, ed. Pharmacy practice in Developing Countries: achievements and challenges. London: Academic Press; 2016.
- Xi X, Huang Y, Lu Q, Ung COL, Hu H. Community pharmacists' opinions and practice of pharmaceutical care at chain pharmacy and independent pharmacy in China. Int J Clin Pharm. 2019;41(2):478-487. https://doi.org/10.1007/s11096-019-00802-w
- 17. Bush J, Langley CA, Wilson KA. The corporatization of community pharmacy: Implications for service provision, the public health function, and pharmacy's claims to professional status in the United Kingdom. Res Social Adm Pharm. 2009;5(4):305-318. https://doi.org/10.1016/j.sapharm.2009.01.003
- 18. Kruijtbosch M, Göttgens-Jansen W, Floor-Schreudering A, van Leeuwen E, Bouvy ML. Moral dilemmas of community pharmacists: a narrative study. Int J Clin Pharm. 2018;40(1):74-83. https://doi.org/10.1007/s11096-017-0561-0
- 19. Bodreau C. The Corporatization of Retail Pharmacy: Changes in Efficacy, Reliability, and Expedience. 2015.
- Hermansyah A, Sukorini AI, Setiawan CD, Priyandani Y. The conflicts between professional and non professional work of community pharmacists in Indonesia. Pharm Pract (Granada). 2012;10(1):33-39. https://doi.org/10.4321/s1886-36552012000100006



- 21. Athiyah U. Model Asuhan Kefarmasian Pelayanan Obat yang diresepkan dan pengaruhnya terhadap Perilaku Apoteker dalam Asuhan Kefarmasian (The model of pharmaceutical care services for prescribed medicines and its impact to pharmacists' behavior in delivering pharmaceutical care services). Surabaya: Airlangga University; 2007.
- 22. Suhartono, Athiyah U, Utami W. Analisis Hubungan Profesionalisme Apoteker dengan Praktek Asuhan Kefarmasian: Studi pada Kasus Terapi Diabetes di Apotek Wilayah Kabupaten Sidoarjo (Analysing the correlation between professionalism of pharmacist and pharmaceutical care services: the case of diabetes mellitus therapy in community pharmacy in Sidoarjo). Jurnal Ilmu Kefarmasian Indonesia. 2017;13(2):166-73.
- 23. Wassenaar DR, Mamotte N. Ethical issues and ethics reviews in social science research. In: Leach MM, Stevens MJ, Korkut Y, Lindsay G, Ferrero A, eds.The Oxford handbook of international psychological ethics. Oxford: OUP; 2012.
- 24. Schafheutle EI, Seston EM, Hassell K. Factors influencing pharmacist performance: a review of the peer-reviewed literature. Health Policy. 2011;102(2-3):178-192. https://doi.org/10.1016/j.healthpol.2011.06.004
- 25. Ministry of Health Indonesia. Permenkes No. 73 tahun 2016 tentang Standar Pelayanan Kefarmasian di Apotek (Standard of pharmacy services in community pharmacy). Jakarta: Ministry of Health Indonesia; 2016.
- 26. McCann L, Hughes CM, Adair CG, Cardwell C. Assessing job satisfaction and stress among pharmacists in Northern Ireland. Pharm World Sci. 2009;31(2):188-194. https://doi.org/10.1007/s11096-008-9277-5
- Cooksey JA, Knapp KK, Walton SM, Cultice JM. Challenges To The Pharmacist Profession From Escalating Pharmaceutical Demand. Health Aff (Millwood). 2002;21(5):182-188.
- 28. Calgan Z, Aslan D, Yegenoglu S. Community pharmacists' burnout levels and related factors: an example from Turkey. Int J Clin Pharm. 2011;33(1):92-100. https://doi.org/10.1007/s11096-010-9461-2
- 29. Bruce SP. Recognizing stress and avoiding burnout. Curr Pharm Teach Learn. 2009;1(1):57-64.
- Gaither CA, Kahaleh AA, Doucette WR, Mott DA, Pederson CA, Schommer JC. A modified model of pharmacists' job stress: The role of organizational, extra-role, and individual factors on work-related outcomes. Res Social Adm Pharm. 2008;4(3):231-243. https://doi.org/10.1016/j.sapharm.2008.04.001
- 31. World Health Organization. The Role of the pharmacist in self-care and self-medication: report of the 4th WHO Consultative Group on the Role of the Pharmacist, The Hague, The Netherlands, 26-28 August 1998. Geneva: WHO; 1998
- 32. Bennadi D. Self-medication: A current challenge. J Basic Clin Pharm. 2013;5(1):19. https://doi.org/10.4103%2F0976-0105.128253
- 33. Jain S, Malvi R, Purviya JK. Concept of self medication: A review. Int J Pharm Biol Arch. 2011;2(3):831-836.
- 34. Budiningsih I, Soehari TD, Ahmad M. Increase Competency Through Training Interventions. Int J Appl Bus Econ Res. 2017;15(6):249-266.
- 35. Mak VSL, March GJ, Clark A, Gilbert AL. Why do Australian registered pharmacists leave the profession? a qualitative study. Int J Clin Pharm. 2013;35(1):129-137. https://doi.org/10.1007/s11096-012-9720-5
- Seston E, Hassell K, Ferguson J, Hann M. Exploring the relationship between pharmacists' job satisfaction, intention to quit the profession, and actual quitting. Res Social Adm Pharm. 2009;5(2):121-132. https://doi.org/10.1016/j.sapharm.2008.08.002
- 37. Magola E, Willis SC, Schafheutle EI. What can community pharmacy learn from the experiences of transition to practice for novice doctors and nurses? A narrative review. Int J Pharm Pract. 2018;26(1):4-15. https://doi.org/10.1111/ijpp.12349
- 38. Gaither CA. Job satisfaction and intention to leave the profession: Should we care? Res Social Adm Pharm. 2009;5(2):91-93. https://doi.org/10.1016/j.sapharm.2009.04.001
- 39. Bimrose J, Hearne L. Resilience and career adaptability: Qualitative studies of adult career counseling. J Vocat Behav. 2012;81(3):338-344. http://doi.org/10.1016/j.jvb.2012.08.002
- 40. Bimrose J, Brown A. Mid-career progression and development: The role for career guidance and counseling. Handbook of career development: Springer: New York; 2014.
- 41. Griffiths V, Thompson S, Hryniewicz L. Landmarks in the professional and academic development of mid-career teacher educators. Eur J Teach Educ. 2014;37(1):74-90. https://doi.org/10.1080/02619768.2013.825241
- 42. Crawford L, Morris P, Thomas J, Winter M. Practitioner development: From trained technicians to reflective practitioners. Int J Project Manag. 2006;24(8):722-733. https://doi.org/10.1016/j.iiproman.2006.09.010
- 43. WHO-FIP. Annex 8: Joint FIP/WHO guidelines on good pharmacy practice: standards for quality of pharmacy services. World Health Organization, 2010 0512-3054 Contract No.: Report.
- 44. Thomas CE, Phipps DL, Ashcroft DM. When procedures meet practice in community pharmacies: qualitative insights from pharmacists and pharmacy support staff. BMJ Open. 2016;6(6):e010851. https://doi.org/10.1136/bmjopen-2015-010851
- 45. Lu H, Eriksson T, Wiffen P. Evidence based pharmacy for developing countries. Eur J Hosp Pharm. 2015;22(2):66-72. http://doi.org/10.1136/ejhpharm-2015-000643



Viz Tools

Help

About Us

Country Rankings

Home Journal Rankings

Pharmacy Practice 3

COUNTRY	SUBJECT AREA AND CATEGORY	PUBLISHER	H-INDEX
Universities and research institutions in Spain Media Ranking in Spain	Health Professions Pharmacy Pharmacology, Toxicology and Pharmaceutics Pharmaceutical Science	Centro de Investigaciones y Publicaciones Farmaceuticas	28
PUBLICATION TYPE	ISSN	COVERAGE	INFORMATION
Journals	1885642X, 18863655	2006-2021	Homepage How to publish in this journal journal@pharmacypractice.org


SCOPI

Pharmacy Practice is a free full-text peer-reviewed journal with a scope on pharmacy practice. Pharmacy Practice is published quarterly. Pharmacy Practice does not charge and will never charge any publication fee or article processing charge (APC) to the authors. The current and future absence of any article processing charges (APCs) is signed in the MoU with the Center for Pharmacy Practice Innovation (CPPI) at Virginia Commonwealth University (VCU) School of Pharmacy. Pharmacy Practice is the consequence of the efforts of a number of colleagues from different Universities who belief in collaborative publishing: no one pays, no one receives. Although focusing on the practice of pharmacy, Pharmacy Practice covers a wide range of pharmacy activities, among them and not being comprehensive, clinical pharmacy, pharmaceutical care, social pharmacy, pharmacy education, process and outcome research, health promotion and education, health informatics, pharmacoepidemiology, etc.

1 dari 4 27/03/2023, 02.27







Metrics based on Scopus® data as of April 2022

S Sara 3 months ago

How much does it cost to publish in the magazine in dollars?

reply

2 dari 4 27/03/2023, 02.27



Dear Sara,

Melanie Ortiz 3 months ago

Thank you for contacting us.

We suggest you visit the journal's homepage or contact the journal's editorial staff , so they could inform you more deeply.

Best Regards, SCImago Team

S sayed 7 months ago

I submit this request to you in order to obtain support in publishing, as the magazine is classified as open access, but it requires publishing expenses



Melanie Ortiz 7 months ago

SCImago Team

Dear Sayed, thank you very much for your comment. We suggest you contact the journal's editorial staff so they could inform you more deeply. Best Regards, SCImago Team

Temesgen Mulugeta 1 year ago

IS Pharmacy practice journal indexing in Scopus and PubMed discontinued after 2022? It says the Scopus coverage years:from 2006 to 2021. I could not find articles indexed in Scopus



Melanie Ortiz 1 year ago

SCImago Team

Dear Temesgen,

Thank you for contacting us.

SJR is a portal with scientometric indicators of journals indexed in Elsevier/Scopus. Unfortunately, we cannot help you with your request referring to the index status. We suggest you consult Scopus database (see the current status of the journal) or the mentioned database for further information.

We suggest you contact Scopus support for further details here: https://service.elsevier.com/app/answers/detail/a_id/14883/kw/scimago/supporthub

Best Regards, SCImago Team

A ANDREW GNANAPRAKASAM J 3 years ago

i need to do publications

reply



SCImago Team

Dear Andrew, thank you very much for your comment. Sorry to tell you that SCImago Journal & Country Rank is not a journal. SJR is a portal with scientometric indicators of journals indexed in Elsevier/Scopus. We suggest you to look for author's instructions/submission guidelines in the journal's website. Best Regards, SCImago Team

Avinash Kumar Singh 4 years ago

The page for the journal is to be more attractive and must be advertised. Pharmacy Practices research should not be published in nor practice journals.

reply

Leave a comment

Name

Email

(will not be published)

Submit

The users of Scimago Journal & Country Rank have the possibility to dialogue through comments linked to a specific journal. The purpose is to have a forum in which general doubts about the processes of publication in the

3 dari 4 27/03/2023, 02.27 journal, experiences and other issues derived from the publication of papers are resolved. For topics on particular articles, maintain the dialogue through the usual channels with your editor.

Developed by:

Powered by:





Follow us on @ScimagoJR

Scimago Lab, Copyright 2007-2022. Data Source: Scopus®

EST MODUS IN REBUS

Horatio (Satire 1,1,106)

Cookie settings

Cookie policy

4 dari 4 27/03/2023, 02.27