



Journal of Advanced Pharmaceutical Technology & Research

An Official Publication of Society of Pharmaceutical Scientists in Research
Volume 10, Issue 11, November 2019
ISSN 2090-0722

JAPTR



Editorial Board

Editor-in-Chief

Dr. Upendra Nagaich

M.Pharm., Ph.D., FSPER
 Professor & Research Coordinator,
 Amity Institute of Pharmacy, Amity University,
 Noida (U.P.) India
 &
 National Secretary,
 Society of Pharmaceutical Education & Research (SPER), India

Executive Editors

Dr. Paul Wan Sia Heng

Department of Pharmacy,
 National University of Singapore,
 Singapore

Dr. Jaya Pandey

School of Studies in Pharmaceutical Sciences,
 Jiwaji University,
 Gwalior (M.P.) India

Associate Editor

Dr. Vandana Chaudhary

School of Pharmacy,
 Pharmacy Australia Centre of Excellence,
 The University of Queensland,
 Brisbane, Australia

Consultant Editor

Prof. (Dr.) Laszlo Endrenyi

Professor Emeritus
 Department of Pharmacology & Toxicology
 University of Toronto, Toronto, Canada

Section Editors (Alphabetical Order)

Dr. Ikoni Ogaji

Dept. of Pharmaceutics & Pharmaceutical Technology,
 Faculty of Pharmaceutical Sciences,
 University of Jos, Nigeria

Dr. Ke Li

J. David Gladstone Institutes,
 San Francisco, California, USA

Editorial Board Members (In Alphabetical Order)

Prof. (Dr.) A. N. Kalia

Vice-President, Society of Pharmacognosy,
 Professor & Head
 I.S.F. College of Pharmacy,
 Moga (Punjab), India

Dr. Abdul Wahab

Department of Pharmacy
 Kohat University of Science & Technology
 Kohat, KPK, Pakistan


Prof. (Dr.) Arun Dharmarajan

Professor
 School of Biomedical Sciences,
 Faculty of Health Sciences,

Feedback

[Subscribe](#)

[About the journal](#)

[Most popular articles](#) 

[Join as Reviewer](#) 

[Email alerts](#) 

[Recommend this journal to your library](#) 

Curtin University,
Perth, Australia

Prof. (Dr.) Ashoke Kumar Ghosh

Vice-Chancellor
IFTM University,
Moradabad (Uttar Pradesh), India

Prof. (Dr.) B. Mishra

Professor & Head
Department of Pharmaceutics,
Indian Institute of Technology,
Banaras Hindu University,
Varanasi (Uttar Pradesh), India

Prof. B. G. Shivananda

Secretary, APTI
Bangalore (Karnataka), India

Prof. (Dr.) Bharat B. Aggarwal

Professor of Cancer Medicine &
Chief, Cytokine Research Laboratory
The University of Texas
Texas, USA

Dr. Bhaswat Sundar Chakraborty

Senior VP & Chairman, R&D Core Committee
Cadila Pharmaceuticals Ltd.
Ahmedabad (Gujarat), India

Prof. (Dr.) C. K. Kokate

National Advisor, Society of Pharmacognosy,
Vice-Chancellor
KLE University,
Belgaum (Karnataka), India

Prof. (Dr.) Cedric Pearce

Adjunct Professor
University of North Carolina
CEO, Mycosynthetix Inc.
North Carolina, USA
[Associate Editor, Journal of Natural Products]

Prof. (Dr.) Ciddi Veeresham

Professor
University College of Pharmaceutical Sciences,
Kakatiya University,
Warangal (Andhra Pradesh), India

Dr. Dâmaris Silveira

Universidade de Brasilia
Brasilia, Brazil

Prof. (Dr.) Dilipkumar Pal

Professor
Dept. of Pharmaceutical Sciences,
Guru Ghasidas Vishwavidyalaya,
Bilaspur (Chhattisgarh), India

Prof. (Dr.) Divakar Goli

Associate Secretary, IPA
Secretary, Education Division, IPA
Professor & Principal
Acharya & B.M. Reddy College of Pharmacy,
Bangalore (Karnataka), India

Prof. (Dr.) G. K. Dash

Professor
Universiti Kuala Lumpur
Perak Darul Ridzuan, Malaysia

Prof. (Dr.) H. S. Buttar

Adjunct Professor,
School of Medicine,
University of Ottawa,
Ottawa, Canada

Kaushik Desai

Gen. Secretary, IPA
Immediate Vice President, IPA
Mumbai (Maharashtra), India

Prof. (Dr.) Mainul Haque

Professor
Universiti Sultan Zainal Abidin
Kuala Terengganu, Terengganu, Malaysia

Dr. Muhammad Akhlaq

University of Auckland
Auckland, New Zealand

Prof. (Dr.) N. Udupa

Professor & Research Director [Health Sciences]
Manipal University,
Manipal (Karnataka), India

Prof. (Dr.) Pornsak Sriamornsak

Professor,
Department of Pharmaceutical Technology &
Pharmaceutical Biopolymer Group (PBiG),
Faculty of Pharmacy, Silpakorn University,
Nakhon Pathom, Thailand

Dr. Rakesh Kumar Sharma

Scientist 'F'
Director,
Defence Food Research Laboratory (DFRL),
Mysuru (Karnataka), India

Prof. (Dr.) Roop K. Khar

Former Professor
Department of Pharmaceutics,
Jamia Hamdard (Hamdard University)
New Delhi, India

Prof. (Dr.) Roopa Karki

Professor & Head
Department of Pharmaceutics,
Acharya & BM Reddy College of Pharmacy,
Bangalore (Karnataka), India

Prof. (Dr.) S. D. Tonpay

Former Professor & Head
Govt. Gajra Raja Medical College,
Gwalior (Madhya Pradesh), India

Prof. (Dr.) S. H. Ansari

Professor & Head,
Department of Pharmacognosy & Phytochemistry,
Jamia Hamdard (Hamdard University)
New Delhi, India

Prof. (Dr.) S. H. Mishra

Professor Emeritus
Faculty of Pharmacy,
The M. S. University of Baroda,
Vadodara (Gujarat), India

Prof. (Dr.) Satish Sardana

Immediate Past President, APTI (Haryana)
Director,
Amity Institute of Pharmacy,
Amity University,
Manesar (Haryana), India

Prof. (Dr.) T. V. Narayana

President, IPA
Director
Vikas Institute of Pharmaceutical Sciences
Rajahmundry (Andhra Pradesh), India

Prof. (Dr.) V. K. Dixit

President, Society of Pharmacognosy,
Former Professor
Department of Pharmaceutical Sciences,
Dr. H. S. Gour University,
Sagar (M.P.), India


Prof. (Dr.) Y. K. Agrawal

Director
Institute of Research & Development,
Gujarat Forensic Sciences University,

[Sitemap](#) | [What's New](#) [Feedback](#) | [Copyright and Disclaimer](#) | [Privacy Notice](#)

© Journal of Advanced Pharmaceutical Technology & Research | Published by Wolters Kluwer - Medknow
Online since 1st November, 2010

[Editorial and Ethics Policies](#)

 [Open Access](#) [View mobile site](#)
ISSN: Print -2231-4040, Online - 0976-2094



Home | About JAPTR | Editorial board | Search | Ahead of print | Current issue | Archives | Submit article | Instructions | Subscribe | Advertise | Contacts | Login

Users Online: 1207     

Table of Contents



July-September 2021

Volume 12 | Issue 3

Page Nos. 215-309

Online since Friday, July 16, 2021

Accessed 19,377 times.

PDF access policy

Journal allows immediate open access to content in HTML + PDF

EPub access policy

allows EPub access only to users from **INDIA**, developing countries and paid subscribers.

 View issue as eBook

 Issue citations

 Issue statistics

 RSS

Feedback

Subscribe

Next Issue

Previous Issue

Most popular articles 

Join as Reviewer 

Email alerts 

Recommend this journal to your library 


Show all abstracts

Show selected abstracts

Export selected to ▼

Add to my list

REVIEW ARTICLE




Efficacy and safety of potential vaccine candidates against coronavirus disease 2019: A systematic review p. 215

Subodh Kumar, Manoj Kumar Saurabh, Vikas Maharshi

DOI:10.4103/japtr.JAPTR_229_20

[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [Sword Plugin for Repository]^{Beta}

ORIGINAL ARTICLES




Anticancer, antioxidant, and antibacterial activities of the methanolic extract from *Sphagneticola trilobata* (L.) J. F pruski leaves p. 222

Vivi Mardina, Syafruddin Ilyas, Halimatussakdiah Halimatussakdiah, Tisna Harmawan, Masitta Tanjung, Faridah Yusof

DOI:10.4103/japtr.JAPTR_131_21

[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [Sword Plugin for Repository]^{Beta}




The effect of rice husk nanosilica hydroxyl compound on dentin biominerization p. 227

Iffi Aprillia, Endang Suprastiwi, Aryo Megantoro, Luh Putu Trisna, Budi Utami, Sarmayana Yana

DOI:10.4103/japtr.JAPTR_227_21

[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [Sword Plugin for Repository]^{Beta}




Antimutagenic activity of nanoparticles of *Rhabdophora pinnata* leaves in mice using micronucleus assay p. 232

Masfria Masfria, Marianne Marianne, Yade Metri Permata, Steven Octavio, Sri Mulyani

DOI:10.4103/japtr.JAPTR_380_20

[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [Sword Plugin for Repository]^{Beta}




In silico estrogen receptor alpha antagonist studies and toxicity prediction of *Melia azedarach* leaves bioactive ethyl acetate fraction p. 236

Martha Ervina, Mohammad Rizki Fadhil Pratama, Hadi Poerwono, Juni Ekowati, Retno Widyowati, Katsuyoshi Matsunami, Sukardiman

DOI:10.4103/japtr.JAPTR_198_21

[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [Sword Plugin for Repository]^{Beta}




Performance and drug deposition of kappa-carrageenan microspheres encapsulating ciprofloxacin HCl: Effect of polymer concentration p. 242

Dewi Melani Hariyadi, Tutiek Purwanti, Dinda Mauldyia, Cindy Alicia Estherline, Esti Hendradi, Mahardian Rahmadi

DOI:10.4103/japtr.JAPTR_197_21

[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [Sword Plugin for Repository]^{Beta}

Encapsulation mechanism of α -mangostin by β -cyclodextrin: Methods of molecular docking and molecular dynamics p. 250




Dian Triwahyuningtyas, Sandra Megantara, Tai Tze Hong, Muhammad Yusuf, Muchtaridi Muchtaridi
DOI:10.4103/japtr.JAPTR_298_20

[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [Sword Plugin for Repository]^{Beta}

Phytochemical screening, antimalarial activities, and genetic relationship of 16 indigenous Thai Asteraceae medicinal plants: A combinatorial approach using phylogeny and ethnobotanical bioprospecting in antimalarial drug discovery p. 254

Desy Liana, Kanchana Rungsihirunrat
DOI:10.4103/japtr.JAPTR_238_21


[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [Sword Plugin for Repository]^{Beta}



In silico identification of natural products from *Centella asiatica* as severe acute respiratory syndrome coronavirus 2 main protease inhibitor p. 261

Putu Gita Maya, Widyaswari Mahayasih, Harizal, Herman, Islamudin Ahmad
DOI:10.4103/japtr.JAPTR_297_20


[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [Sword Plugin for Repository]^{Beta}



Development and validation of an automated solid-phase extraction-LC-MS/MS method for the bioanalysis of fluoxetine in human plasma p. 267

Ishtiaque Ahmad, Zabih Ullah, Mohd Ibrahim Khan, Abdullah Khaloofa Alahmar, Mohd Faiyaz Khan
DOI:10.4103/japtr.JAPTR_308_20

[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [Sword Plugin for Repository]^{Beta}



The color improvement of postdebonding white spot lesions after fluoride and casein phosphopeptide-amorphous calcium phosphate application p. 274

Haru Setyo Anggani, Putri Arifiani, Erwin Siregar
DOI:10.4103/japtr.JAPTR_203_21


[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [Sword Plugin for Repository]^{Beta}



3,4-dimethoxychalcone novel ultraviolet-A-protection factor in conventional sunscreen cream p. 279

Erlina Fatmasari, Abdul Karim Zulkarnain, Rina Kuswahyuning
DOI:10.4103/japtr.JAPTR_89_21


[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [Sword Plugin for Repository]^{Beta}



The interaction of alpha-mangostin and its derivatives against main protease enzyme in COVID-19 using in silico methods p. 285

Syahrul Hidayat, Faisal Maulana Ibrahim, Kelvin Fernando Pratama, Muchtaridi Muchtaridi
DOI:10.4103/japtr.JAPTR_299_20


[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [Sword Plugin for Repository]^{Beta}



Standardization of *Leonurus sibiricus* L. aerial part and capillary electrophoresis quantitative analysis of its leonurine content p. 291

Maneewan Suwatronnakorn, Somchai Issaravanich, Chanida Palanuvej, Nijsiri Ruangrungsi
DOI:10.4103/japtr.JAPTR_243_21


[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [Sword Plugin for Repository]^{Beta}



Working goal of Brazilein sappan wood as a candidate for SARS-coV-2 antivirus drug against spike (S) glycoprotein, papain-like proteinase, and main protease: In silico study p. 298

Entuy Kurniawan, Dwi Kihariyani
DOI:10.4103/japtr.JAPTR_222_21


[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [Citations (1)] [Sword Plugin for Repository]^{Beta}



Characterization and acute oral toxicity of concentrated minerals of Pamekasan Madura seawater p. 305

Syafika Alaydrus, Abd Kakhar Umar, Sriwidodo Sriwidodo, Ajeng Diantini, Nasrul Wathoni, Riezki Amalia
DOI:10.4103/japtr.JAPTR_250_20

[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [Sword Plugin for Repository]^{Beta}



Performance and drug deposition of kappa-carrageenan microspheres encapsulating ciprofloxacin HCl: Effect of polymer concentration

Dewi Melani Hariyadi,
Tutiek Purwanti, Dinda Maulydia,
Cindy Alicia Estherline,
Esti Hendradi,
Mahardian Rahmadi¹

Departments of Pharmaceutical
Sciences and ¹Pharmacy Practice,
Faculty of Pharmacy, Universitas
Airlangga, Surabaya, Indonesia

J. Adv. Pharm. Technol. Res.

ABSTRACT

It has been known that in respiratory disease, antibiotic is selected for respiratory diseases or lung infections and this research focused on ciprofloxacin HCl as a model. The aim was to evaluate the effect of kappa-carrageenan polymer concentrations on characteristics, release, and drug deposition in the lung. Ciprofloxacin HCl-carrageenan microspheres were produced with kappa carrageenan (0.75%, 0.50%, and 0.25%) as polymer and KCl (1.5%) as crosslinker. Physical characteristics were included morphology, size, moisture content, swelling index, mucoadhesivity, drug loading, entrapment efficiency, and yield. Freeze-dried microspheres were inhaled by animal, and drug deposition was observed. Results showed that dried, smooth, and spherical microspheres of size of 1.34 to 1.70 μm and loading of 15.63% to 38.72%. Entrapment efficiency and yield were 25.38%–51.61% and 52.53%–63.19%, respectively. Mucoadhesivity was 0.0059–0.0096 kg force, and release in 24 h was 74.38%–81.02%. Release kinetics demonstrated Higuchi mechanism. Increasing carrageenan concentration affected size, loading, and efficiency but did not influence adhesivity, yield, and release. Higher amount of polymer caused the lower deposit on the lungs. Respirable size of ciprofloxacin HCl-kappa carrageenan microspheres was successfully achieved target site and prolonged residence time in lungs.

Key words: Ciprofloxacin HCl, deposition, kappa carrageenan, lung delivery, microspheres, release, respiratory disease

INTRODUCTION

Respiratory tract infection is an infectious respiratory disease that occurs in both upper and lower respiratory tract. This can cause a wide spectrum of diseases, ranging

from mild-to-severe infection to death, which is influenced by causative pathogens, environmental, and host factors. Bacteria that can cause infection in the human respiratory tract are *Pseudomonas aeruginosa* and *Staphylococcus aureus*.^[1] *Staphylococcus aureus* contained in the respiratory tract will enter the lungs causes irritation of the respiratory tract from the pharynx to the lungs.^[2] Antibiotics can be used to treat respiratory infections, and their use refers to the results of sputum culture.^[3]

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: WKHLRPMedknow_reprints@wolterskluwer.com

How to cite this article: Hariyadi DM, Purwanti T, Maulydia D, Estherline CA, Hendradi E, Rahmadi M. Performance and drug deposition of kappa-carrageenan microspheres encapsulating ciprofloxacin HCl: Effect of polymer concentration. *J Adv Pharm Technol Res* 2021;12:242-9.

Address for correspondence:

Dr. Dewi Melani Hariyadi,
Department of Pharmaceutical Sciences, Faculty of Pharmacy,
Universitas Airlangga, Surabaya, Indonesia.
E-mail: dewi-m-h@ff.unair.ac.id

Submitted: 19-Feb-2021

Revised: 31-Mar-2021

Accepted: 20-May-2021

Published: 16-Jul-2021

Access this article online

Quick Response Code:



Website:

www.japtr.org

DOI:

10.4103/japtr.JAPTR_197_21

Ciprofloxacin is a fluoroquinolone class antibiotic that is active both against Gram-negative and some Gram-positive bacteria.^[4] Ciprofloxacin can be used by oral twice a day. However, only 0.5%–5% can enter the systemic circulation and of these levels, only 10% can reach target organ. Thus, delivery system is needed by changing the route or modifying the release to increase efficacy and reduce side effects of antibiotics.^[5]

Inhalation drug experiences rapid absorption because the wide of the absorption surface, avoiding first-pass metabolism in the liver, can be administered directly into the bronchi, and the lesser amount of drug. Advantages of this require a lower dose along with minimal drug side effects.^[1] Microspheres are one of the delivery systems which can deliver large amounts of drug and slower clearance in the lungs, therefore, it can increase effectiveness.

Microsphere is a small spherical particle (1 μm to 1000 μm). Mechanism of drug release is mediated by diffusion, swelling, and degradation of the matrix.^[6] For the inhalation route, the particle size ranges from 1 μm to 5 μm . If it is greater than 5 μm , the particles will only reach the oropharynx, whereas if it is smaller than 1 μm , the particles will come out with expired air.^[7] Spherical has good flow properties that allow drug to reach respiratory tract. Microencapsulation can increase drug bioavailability and sustained release to target therapy.^[8] Release is influenced by various factors.^[9] Hariyadi *et al.* entrapped ciprofloxacin HCl in alginate microspheres using CaCl_2 .^[10] Hariyadi *et al.* also investigated ciprofloxacin-carrageenan microspheres with high concentration of drug and low concentration of KCl,^[11] however, the entrapment efficiency was less than 30%. Kinetics and micromeritics of those microspheres were also studied (unpublished paper).

Carrageenan has been used for encapsulation of drug and pharmaceutical purposes.^[12-15] Carrageenan has different types which produce distinct gel and solution for specific texturing applications.^[15]

This study encapsulated ciprofloxacin HCl using kappa carrageenan produced by ionic gelation method and administered to mice using nose-only inhalation method as modification of Kaur's method, and the lung tissue was collected using the cryotome technique.^[16] The animals were observed for drug deposition. Carrageenan-ciprofloxacin HCl microspheres were designed with 0.75%, 0.50%, and 0.25% polymer; 5% maltodextrin; and 1.5% KCl using aerosolization technique. This research investigated the effect of polymer concentration on characteristics includes size, morphology, entrapment efficiency, loading, yield, mucoadhesive, and release. Drug deposition in the lungs was also observed.

MATERIALS AND METHODS

Materials

Ciprofloxacin HCl (Interbath); Kappa-Carrageenan (Danisco Cultor); KCl (Solvay Chemicals International); Na_2HPO_4 p.a (Merck); KH_2PO_4 p.a (Merck); NaCl p.a (Merck); KCl p.a (Merck); Maltodextrin (PT Bratachem); HCl p.a (Merck); Aquademineralisata (PT. Bratachem).

Microspheres preparations

Microspheres were made by aerosolization with modification, by spraying solution with a spray diameter of 35 μm , distance of 8 cm, and pressure of 40 psi.^[10,11] Formula is shown in Table 1. Ciprofloxacin-carrageenan solution was sprayed on the KCl solution and stirred for 2 h at 1000 rpm. Microspheres were washed for 6 min by centrifugation at 2500 rpm and freeze dried for 24 h at -80°C .

Evaluation of microspheres

Particle size and morphology

Size was characterized by optical microscope and software with a magnification of $\times 100$. For morphology, scanning electron microscope (SEM) was used.

Entrapment efficiency and drug loading

Entrapment efficiency was calculated from ciprofloxacin contained in microspheres, as below equation:

$$\% \text{ EE} = \frac{\text{ciprofloxacin mass in the microspheres}(\text{mg})}{\text{ciprofloxacin mass in the formula}(\text{mg})} \times 100\% .^{[9]}$$

Loading was calculated from the total microspheres obtained compared to the total weight of microspheres.

In vitro release

Ciprofloxacin release from microspheres was carried out in phosphate-buffered saline (PBS) pH 7.4 ± 0.05 at 37°C . Sample was shaken at 100 rpm. A 5.0 ml of sample was taken from the PBS at 10 min up to 1440 min, and 5 ml media was replaced in each interval. Sample was filtered using 0.45 μm millipore paper. Cumulative ciprofloxacin HCl released from microspheres was obtained by below equation:

$$\% \text{ Ciprofloxacin HCl cumulative} = \frac{\text{Ciprofloxacin mass in the 100 ml of media}(\text{mg})}{\text{ciprofloxacin mass}(\text{mg})} \times 100\%$$

Release profile was shown by the value of b (slope) of the regression equation.

Mucoadhesive examination

Mucoadhesivity was performed using TA-XT stage plus texture analyzer (Stable Micro Systems, Godalming, UK).

Drug deposition and lung irritation

Wistar rats which met inclusions criterias were used, including male rats, 150-200 grams, healthy and no defects

or wounds on their body. Minimum number of animals in each group was 6 rats. Animal experiments were done according to the animal handling protocols and have passed the ethical clearance by Ethical committee of Universitas Airlangga with Reference No. 038/HRECC.FODM/II/2019.

Drugs administration to animal

Ciprofloxacin dosage inhalation for animals is 20 mg/Kg BW.^[17] Microspheres given to rats were calculated from drug loadings. Administration of microspheres by nose only exposure were applied using modified tool as same principle as DPI tool [Figure 1]. The apparatus was adopted from Kaur.^[16] Microspheres were flown and inhaled by rats through their nose.^[16]

Drug deposition examination

Microspheres were labeled with rhodamine B and were inhaled by previous method. Rats were euthanized and lungs were taken. Lungs were rinsed with normal saline then sliced longitudinally horizontally with a thickness of 6 um using a cryotom (Tissue-Tek Cyro3, Sakura) at a temperature of -59°C at the Anatomical Pathology Laboratory, Integrated Central Surgical Building at Soetomo Hospital. The cryostat preparations were stored at -15°C-20°C until further analysis. Drug deposition observation was carried out using Fluorescence Microscope (FS × 100, Olympus) with a magnification of ×42 and an exposure time of 1/1.2 ms and using a red filter.

Data analysis

For physical characteristics and drug release, statistical

data analysis was performed using one-way ANOVA with a confidence degree of 95% ($\alpha = 0.05$). Drug deposition was observed using a fluorescent microscope and intensity was compared. Data analysis of the intensity quantitatively used imageJ software and compared statistically with the Kruskal-Wallis nonparametric analysis.

RESULTS AND DISCUSSIONS

Particle size and morphology of microspheres

Particle size of all formulas was obtained below 5 µm as shown in Figure 2. This size corresponds to the desired size to achieve alveoli site. If it is greater than 5 µm, particles will only reach the oropharynx, whereas if it is smaller than 1 µm, the particles will come out with expired air.^[18]

Based on statistical analysis, the increase in polymer concentration showed a significant difference to the particle size.

Morphological observations of the microspheres using SEM can be seen in Figure 3. It was showed that formulas formed spherical microspheres and smooth surfaces.

Yield

Microspheres' yield was seen in Figure 4 and showed at about 52.53%–63.19%.

Table 1: Ciprofloxacin HCl-carrageenan microspheres

Formula	Components			
	Ciprofloxacin (%)	Kappa-carrageenan (%)	KCl (%)	Maltodextrin (%)
F1	0.5	0.75	1.5	5
F2	0.5	0.50	1.5	5
F3	0.5	0.25	1.5	5




Figure 1: Inhalation Drug Apparatus [16]: (a) A tapered poly(propylene) centrifuge tube of capacity 15ml that formed the powder fluidization or aerosol generation chamber; (b) Flexible, C-Flex tubing to introduce a turbulent fluidizing air stream into the chamber; (c) a rubber pipette bulb that provided the source of turbulent air when pressed and released.




Figure 2: Size of ciprofloxacin HCl-carrageenan microspheres




Figure 4: Microspheres yield

Entrapment efficiency and drug loading

Entrapment efficiency of microspheres and drug loading was presented in Figure 5.

F1 had the highest entrapment efficiency and drug loading. This was due to the highest concentration of F1 polymer. Increasing polymer concentration increased entrapment and drug loading which will cause high polymer solution viscosity and produce large droplets and sizes and cross-linking points on the polymer chain increased to bind drugs so that entrapment efficiency and drug loading increased.^[19,20] Meanwhile, F3 had the smallest entrapment efficiency and drug loading because due to smallest polymer concentration caused low viscosity and results in small droplets and a reduced cross-link point on the polymer chain. Thus, small efficiency and drug loading were found. Addition of maltodextrin may cause small entrapment efficiency and loading due to ability of maltodextrin to form hydrogen bonds which can force the intermolecular bonds between K⁺ with sulfate groups and carrageenan polymer oxygen, thereby causing a decrease in cross-link point of the polymer.^[9]

In vitro release study

Release of all microspheres formula is shown in Figure 6.




Figure 3: Morphological of ciprofloxacin HCl-carrageenan microspheres Formula 1 (a), Formula 2 (b), and Formula 3 (c) with a magnification of $\times 5000$




Figure 5: Entrapment efficiency and drug loading of ciprofloxacin-carrageenan microspheres

Cumulative drug release up to 1440 min from F1 to F3 was 71% to 77.96%. This was in line with the expectation of slow-release modification. Statistical showed that there was no significant difference in the cumulative regardless of each formula. This could be because differences in concentration between formulas of 0.25% still did not make a significant difference at the use of 1.5% KCl. Thus, it is recommended to optimize other factors that may influence the release. Based on statistical analysis, there was no significant difference toward all slopes.

Release kinetics

Release kinetics model showed that kinetics follows zero order with the highest r value [Table 2]. Zero-order kinetics model stated that drug release occurs through a solid gel layer which has a constant thickness with respect to the time associated with the swelling process of carrageenan and erosion rate of the gel layer.^[21] In consequence, sustain release of microspheres in carrageenan matrix can be applied.

F1, F2, F3, and F4 showed no significant differences between amount of polymer and release rate. This could be because of the small differences in concentration used did not make a difference.

Table 2: Coefficient of the relation (*R*) of the kinetics model

Formula	Order 0 (<i>R</i> ²)	Order 1 (<i>R</i> ²)	Higuchi (<i>R</i> ²)	Korsmeyer-Peppas (<i>R</i> ²)
F1	0.9994	0.9940	0.9723	0.9359
F2	0.9996	0.9855	0.9777	0.9372
F3	0.9996	0.9900	0.9770	0.9370




Figure 6: Release of ciprofloxacin HCl from microspheres

Mucoadhesivity of microspheres

Mucoadhesivity of microspheres formula is shown in Figure 7.

F1 had the highest mucoadhesive compared to others, however, no significant difference between formulas. The most optimized formula was formula with high carrageenan because formula provided high drug loading and entrapment efficiency, spherical, small, and smooth surface microspheres which were not cause irritation to the respiratory tract.

Drug deposition of microspheres

Observations of drug deposition in the trachea at 1st and 4th h can be observed in Figure 8. It can be seen that intensity was higher in Formula 3 compared to others in the 1st h. For the 4th h, intensity became more even in Formulas 1 and 2. For Formula 3, intensity was fainter than the 1st h [Table 3]. It was indicated that there was a significant difference in drug deposition in trachea.

Observations of drug deposition in bronchioles at the first and the 4th h can be seen in Figure 9.

It can be seen that intensity was greater in Formula 3 compared to others [Figure 10]. For the 4th h, intensity became more even in Formulas 1 and 2. In Formula 3, intensity was fainter than 1st h [Table 4]. It was indicated that there was a significant difference in drug deposition.

Observations of drug deposition in the alveoli at the 1st and the 4th h can be seen in Figure 11.

Highest intensity in alveoli at 1 h was Formula 3. For the 4th h, highest intensity was Formula 1 and decreased in Formula 3. On the other hand, in Formula 3, intensity




Figure 7: Mucoadhesivity of microspheres

was fainter than the 1st h [Table 5]. It was indicated that a significant difference in drug deposition was found. Based on all results, the distribution of microspheres intensity in rat lungs at the one and 4 h was seen clearly.

Drug inhalation in the form of DPI uses nose-only method because simple no need for anesthesia, no surgery, and quite effective.^[12] Observation of deposition using fluorescence microscopy was a sensitive method.^[21]

Results of F1 and F2 observation, it can be seen that at 1st h, the red luminescence weak in the trachea, bronchi, and alveoli. In F3, the red luminescence was higher and had spread to the lower lungs. Furthermore, an increase in polymer concentration showed significant difference in the intensity. This may occur because with a concentration of 0.75% and 0.5%, microspheres hold release of rhodamine, therefore, at the 1st h, only a little rhodamine was released and red luminescence was only visible around the walls of the trachea and bronchioles.

Increase in polymer concentration caused larger size, which means thicker matrix and more polymers that entrap the drug produced longer drug release. Therefore, at the 1st h, formula with smallest polymer concentration produced more luminescence than formula with higher concentration.

At the 4th h, F3 luminescence had started to fade and intensity was the lowest, while in F1 and F2, the intensity was higher than 1st h. This may occur because a lot of rhodamine was released from microspheres enter into the lower lungs and alveoli.

The ideal size range for inhalation system is 1–5 μm, therefore, all formulas were included as particles




Figure 8: Observation of ciprofloxacin HCl-carrageenan microspheres in the trachea at 1st h of Formula 1 (a), Formula 2 (b), Formula 3 (c) and 4th h of Formula 1 (d), Formula 2 (e), Formula 3 (f)




Figure 9: Observation of ciprofloxacin HCl-carrageenan microspheres in rat bronchioles at the 1st h of Formula 1 (a), Formula 2 (b), Formula 3 (c) and the 4th h of Formula 1 (d), Formula 2 (e), Formula 3 (f)

which capable of entering the lower respiratory system. Microspheres were able to deliver drugs to the lower lungs. Further research is needed to determine whether ciprofloxacin reaches the lungs with sufficient concentration levels to kill bacteria.

CONCLUSION

Increased polymer concentration carrageenan kappa influenced physical characteristics of the ciprofloxacin-kappa carrageenan microspheres but had no effect on mucoadhesive, release rate, and release profile. The use of carrageenan was capable of delivering microspheres to the lower lung. Carrageenan concentration affected drug deposition in the lungs. The higher the carrageenan levels, the slower the drug deposition in the lungs. It is suggested to evaluate more quantitatively the ciprofloxacin HCl concentration in the blood that is separated from the microspheres.

Table 3: Intensity in the trachea at 1st and 4th h

Formula	Mean intensity±SD	
	1 st h	4 th h
F1	7.4056±0.908	12.085±0.1589
F2	11.944±1.934	24.932±6.084
F3	40.083±6.879	5.354±0.717

SD: Standard deviation

Table 4: Intensity in the bronchioles at 1st and 4th h

Formula	Mean intensity±SD (intensity unit)	
	1 st h	4 th h
F1	8.414±0.911	31.407±7.378
F2	19.7024±3.4765	25.973±5.913
F3	35.1182±3.894	9.458±1.994

SD: Standard deviation

Acknowledgment

Authors would like to thank to the Directorate of General Higher Education (DRPM DIKTI) for grant and the Faculty




Figure 10: Observation of ciprofloxacin HCl-carrageenan microspheres in rat bronchulus at the 1st h of Formula 1 (a), Formula 2 (b), Formula 3 (c) and the 4th h of Formula 1 (d), Formula 2 (e), Formula 3 (f) with a ×42 magnification red filter fluoresce




Figure 11: Observation of ciprofloxacin HCl-carrageenan microspheres in the alveoli of rats at the 1st h of Formula 1 (a), Formula 2 (b), Formula 3 (c) and 4th h of Formula 1 (d), Formula 2 (e), Formula 3 (f)

Table 5: Intensity in the alveoli at 1st and 4th h

Formula	Mean intensity unit±SD	
	1 st h	4 th h
F1	4.212±1.325	21.487±6.541
F2	4.169±1.079	16.992±1.813
F3	15.376±2.520	8.251±2.824

SD: Standard deviation

of Pharmacy Universitas Airlangga for research support and facilities.

Financial support and sponsorship

This study was financially supported by DRPM DIKTI for grant.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

1. Ghosh Dastidar D, Saha S, Chowdhury M. Porous microspheres: Synthesis, characterisation and applications in pharmaceutical and medical fields. *Int J Pharm* 2018;548:34-48.
2. English JP, Dang W, and Zhao Z. Fabrication of Controlled Delivery Devices. *Encyclopedia of controlled drug delivery Volume 1 & 2* United State of America: Wiley Intersciences Publication 1999. p. 356.
3. Dhakar CR, Maurya SD, Sagar B, Bhagat S, Prajapati SK, Jain CP. Variables influencing the drug entrapment efficiency of microspheres: A pharmaceutical review. *Scholars Res Libr* 2010;2:102-16.
4. Flume PA, VanDevanter DR. Clinical applications of pulmonary delivery of antibiotics. *Adv Drug Deliv Rev* 2015;85:1-6.
5. Jin M, Zheng Y, Hu Q. Preparation and characterization of bovine serum albumin alginate/chitosan microspheres for oral administration. *Asian J Pharm Sci* 2009;4:216-8.
6. Joshi S, Patel P, Lin S, Mada PL. Development of crosslinked

- alginate spheres by ionotropic gelation technique for controlled release of naproxen oral. *Asian J Pharm* 2012;7:134-14.
7. Katzung BG, Trevor AJ. *Basic Clinical and Pharmacology* 12th ed. New York: Mc Graw Hill Companies Inc.; 2012. p. 753-6.
 8. Knowles MR, Gilligan PH, Boucher RC. *Cystic Fibrosis*. New York: Churchill Livingstone; 2000. p. 767-72.
 9. Li L, Ni R, Shao Y, Mao S. Carrageenan and its applications in drug delivery. *Carbohydr Polym* 2014;103:1-11.
 10. Liew CV, Chan LW, Ching AL, Heng PW. Evaluation of sodium alginate as drug release modifier in matrix tablets. *Int J Pharm* 2006;309:25-37.
 11. Ramteke KH, Jadhav VB, Dhole SN. Microspheres: As carriers used for novel drug delivery system. *J Pharm* 2012;2:44-8.
 12. Peñalver DA, Alemán A, Montero MP, Gómez-Guillen MC. Entrapment of natural compounds in spray-dried and heat-dried iota-carrageenan matrices as functional ingredients in surimi gels. *Food Function* 2021;12:2137-47.
 13. Fauzi MA, Hendradi E, Pudjiastuti P, Widodo RT. Analysis of dissolution of salicylamide from carrageenan based hard-shell capsules: A study of the drug-matrix interaction, Indones. *J Chem* 2021;21:148-56.
 14. Gonzalez CA, Sosnik A, Kalmar J, De Marco I, Erkey C, Concheiro A, *et al.* Aerogels in drug delivery: From design to application. *J Control Release* 2021;332:40-63.
 15. Fu M, Wu Y, Zhang Q, Xue Z, Geng C, Xia Y. Study on the mechanical and flame retardant properties of a novel carrageenan fiber-CAF-K, Al, *Eur Polymer J* 2021;150:110408.
 16. Kaur J, Muttill P, Verma RK, Kumar K, Yadav AB, Sharma R, *et al.* A hand-held apparatus for “nose-only” exposure of mice to inhalable microparticles as a dry powder inhalation targeting lung and airway macrophages. *Eur J Pharm Sci* 2008;34:56-65.
 17. Rondhianto R, Kurniawati D, Vidiany AK. Effective Cough and Deep Breath Decreases the Colonization of *Staphylococcus Aureus* in Secret of Post Surgery Patients with General Aenesthesia in Soebandi Jember Hospital. *Nurse Line J* 2016;1:151-8.
 18. Wichelhaus TA, Böddinghaus B, Besier S, Schäfer V, Brade V, Ludwig A. Biological cost of rifampin resistance from the perspective of *Staphylococcus aureus*. *Antimicrob Agents Chemother* 2002;46:3381-5.
 19. Hariyadi DM, Hendradi E. Optimization performance and physical stability of ciprofloxacin hcl-ca alginate microspheres: Effect of different concentration of alginate and CaCl₂. *Int J Drug Deliv Technol* 2018;10:89-94.
 20. Hariyadi DM, Hendradi E, Sharon N. Development of carrageenan polymer for encapsulation of ciprofloxacin HCl: *In vitro* characterization. *Int J Drug Deliv Technol* 2019;9:89-93.
 21. Liu C, Shi J, Dai Q, Yin X, Zhang X, Zheng A. *In-vitro* and *in-vivo* evaluation of ciprofloxacin liposomes for pulmonary administration. *Drug Dev Ind Pharm* 2015;41:272-8.



UNIVERSITAS AIRLANGGA FACULTY OF DENTAL MEDICINE
HEALTH RESEARCH ETHICAL CLEARANCE COMMISSION

ETHICAL CLEARANCE CERTIFICATE

Number : 038/HRECC.FODM/II/2019

Universitas Airlangga Faculty Of Dental Medicine Health Research Ethical Clearance Commission has studied the proposed research design carefully, and therefore, shall herewith certify that the research entitled :

"PENGARUH KONSENTRASI POLIMER TERHADAP DRUG DEPOSITION MIKROSFER CIPROFLOXACIN HCI-KARAGENAN DI PARU TIKUS"

Principal Researcher : CINDY ALICIA ESTHERLINE

Unit/Institution/Place of Research : - Fakultas Farmasi Universitas Airlangga
Surabaya

CERTIFIED TO BE ETHICALLY CLEARED



February 11, 2019
Chairman,

Prok. Dr. M. Rubianto, drg., MS, Sp.Perio(K)
Official No.195009081978021001



[i](#) [x](#)

Complexity Research Papers

Join Leading Researchers in the Field of Complexity. Publish Your Next Research With Us

Hindawi

Open

Journal of Advanced Pharmaceutical Technology and Research

COUNTRY	SUBJECT AREA AND CATEGORY	PUBLISHER	H-INDEX
India 	Pharmacology, Toxicology and Pharmaceutics Pharmaceutical Science	Wolters Kluwer Medknow Publications	38

PUBLICATION TYPE	ISSN	COVERAGE	INFORMATION
Journals	01105558, 09762094	2010-2021	Homepage How to publish in this journal editor@japtr.org

SCOPE

Journal of Advanced Pharmaceutical Technology & Research (JAPTR) is an Official Publication of Society of Pharmaceutical Education & Research™. It is an international journal published Quarterly. Journal of Advanced Pharmaceutical Technology & Research (JAPTR) is available in online and print version. It is a peer reviewed journal aiming to communicate high quality original research work, reviews, short communications, case report, Ethics Forum, Education Forum and Letter to editor that contribute significantly to further the scientific knowledge related to the field of Pharmacy i.e. Pharmaceutics, Pharmacology, Pharmacognosy, Pharmaceutical Chemistry. Articles with timely interest and newer research concepts will be given more preference.



Complexity Research Papers


Join Leading Researchers in the Field of Complexity. Publish Your Next Research With Us Hindawi



Quartiles

FIND SIMILAR JOURNALS ?




<p>1 Brazilian Journal of Pharmaceutical Sciences BRA</p> <p>41% similarity</p>	<p>2 Saudi Pharmaceutical Journal SAU</p> <p>40% similarity</p>	<p>3 DARU, Journal of Pharmaceutical Sciences GBR</p> <p>39% similarity</p>	<p>4 Asian Journ Pharmaceut IND</p> <p>3 s</p>
-----------------------------------------------------------------------------------------------------------	-------------------------------------------------------------------------------------------	-------------------------------------------------------------------------------------------------------	--------------------------------------------------------------------------



Complexity Research Papers

Join Leading Researchers in the Field of Complexity. Publish Your Next Research With Us Hindawi





Journal of Advanced Pharmaceutical Technology...

Q3 Pharmaceutical Science
best quartile

SJR 2021 0.31

powered by scimagojr.com


Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scimaç
```

SCImago Graphica

Explore, visually communicate and make sense of data with our [new data visualization tool](#).



Metrics based on Scopus® data as of April 2022



Loading comments...

Complexity Research Papers

Join Leading Researchers in the Field of Complexity. Publish Your Next Research With Us

Complexity Research Papers

Join Leading Researchers in the Field of Complexity. Publish Your Next Research With Us Hindawi



Developed by:



Powered by:



Follow us on @ScimagoJR

Scimago Lab, Copyright 2007-2022. Data Source: Scopus®

EST MODUS IN REBUS

Horatio (Satire 1,1,106)

[Edit Cookie Consent](#)



Complexity Research Papers

Join Leading Researchers in the Field of Complexity. Publish Your Next Research
With Us Hindawi

