

Differentiation of osteoblasts: the links between essential transcription factors

by Mahardian Rahmadi

Submission date: 01-Apr-2023 02:57PM (UTC+0800)

Submission ID: 2052757051

File name: oblasts_the_links_between_essential_transcription_factors_1.pdf (2.73M)

Word count: 17319

Character count: 99468



Differentiation of osteoblasts: the links between essential transcription factors

Junaidi Khotib, Honey Dzikri Marhaeny, Andang Miatmoko, Aniek Setiya Budiadin, Chrismawan Ardianto, Mahardian Rahmadi, Yusuf Alif Pratama & Muhammad Tahir

To cite this article: Junaidi Khotib, Honey Dzikri Marhaeny, Andang Miatmoko, Aniek Setiya Budiadin, Chrismawan Ardianto, Mahardian Rahmadi, Yusuf Alif Pratama & Muhammad Tahir (2022): Differentiation of osteoblasts: the links between essential transcription factors, Journal of Biomolecular Structure and Dynamics, DOI: [10.1080/07391102.2022.2148749](https://doi.org/10.1080/07391102.2022.2148749)

To link to this article: <https://doi.org/10.1080/07391102.2022.2148749>



Published online: 24 Nov 2022.



Submit your article to this journal [↗](#)




View related articles [↗](#)



View Crossmark data [↗](#)



Differentiation of osteoblasts: the links between essential transcription factors

Junaidi Khotib^a, Honey Dzikri Marhaeny^a, Andang Miatmoko^b, Aniek Setiya Budiati^a, Chrismawan Ardianto^a, Mahardian Rahmadi^a, Yusuf Alif Pratama^a and Muhammad Tahir^c 

^aDepartment of Pharmacy Practice, Faculty of Pharmacy, Universitas Airlangga, Surabaya, Indonesia; ^bDepartment of Pharmaceutical Science, Faculty of Pharmacy, Universitas Airlangga, Surabaya, Indonesia; ^cDepartment of Pharmaceutical Science, Kulliyah of Pharmacy, International Islamic University Malaysia, Pahang, Malaysia

Communicated by Ramaswamy H. Sarma

ABSTRACT

Osteoblasts are cells derived from mesenchymal stem cells (MSCs) in the bone marrow, are cells responsible for bone formation and remodeling. The differentiation of osteoblasts from MSCs is triggered by the expression of specific genes, which are subsequently controlled by pro-osteogenic pathways. Mature osteoblasts then differentiate into osteocytes and are embedded in the bone matrix. Dysregulation of osteoblast function can cause inadequate bone formation, which leads to the development of bone disease. Various key molecules are involved in the regulation of osteoblastogenesis, which are transcription factors. Previous studies have heavily examined the role of factors that control gene expression during osteoblastogenesis, both in vitro and in vivo. However, the systematic relationship of these transcription factors remains unknown. The involvement of ncRNAs in this mechanism, particularly miRNAs, lncRNAs, and circRNAs, has been shown to influence transcriptional factor activity in the regulation of osteoblast differentiation. Here, we discuss nine essential transcription factors involved in osteoblast differentiation, including Runx2, Osx, Dlx5, β -catenin, ATF4, Ihh, Satb2, and Shn3. In addition, we summarize the role of ncRNAs and their relationship to these essential transcription factors in order to improve our understanding of the transcriptional regulation of osteoblast differentiation. Adequate exploration and understanding of the molecular mechanisms of osteoblastogenesis can be a critical strategy in the development of therapies for bone-related diseases.

ARTICLE HISTORY

Received 30 May 2022
Accepted 12 November 2022

KEYWORDS

Bone development; bone remodeling; osteoblast differentiation; essential transcription factors; molecular mechanisms; transcriptional regulation; bone disorders; ncRNAs

Introduction

Bone is a metabolically active organ that is dynamic in maintaining its strength and integrity through the actions of osteoblasts and osteoclasts (El-Ganzuri et al., 2016; Shahi et al., 2017). In vertebrates, bone formation (ossification) occurs in the craniofacial intramembrane and endochondral bones in other parts of the skeletal system. Endochondral ossification is the replacement of cartilage with mineralized bone affected by chondrocyte differentiation in the central cartilage anlagen. This is followed by the invasion of perichondrial osteoblast progenitors, osteoclasts, vascular endothelial cells, and hematopoietic cells into hypertrophic cartilage. Dense mesenchymal progenitor cells differentiate into osteoblasts and form bone directly during intramembranous ossification (Berendsen & Olsen, 2015; laquinta et al., 2019).

Bone tissue has an amazing ability to repair itself and generally heals through regeneration. Under homeostatic conditions, the balance between bone formation (mediated by osteoblasts) and bone resorption (mediated by osteoclasts) is tightly regulated without major changes in net bone mass or mechanical strength, a process known as bone

remodeling (Kim et al., 2020). Osteoclasts degrade bone by secreting polarized proteolytic enzymes, such as cathepsin K, and acids, such as HCl, which dissolve collagen and matrix proteins during bone resorption. Meanwhile, osteoblasts generate an extracellular collagen matrix with specific properties that will be mineralized following hydroxyapatite ($\text{Ca}_5(\text{PO}_4)_3(\text{OH})$) crystal deposition (Kim et al., 2020; Shahi et al., 2017; Zhang, 2010). Once this balance is disrupted, abnormal bone remodeling occurs, resulting in bone deformities and a variety of bone diseases (Chan et al., 2021).

Osteoblast differentiation, also known as osteoblastogenesis, is a major component of bone formation due to the initial very rapid cell proliferation followed by extracellular matrix maturation and mineralization (Huang et al., 2007; Shahi et al., 2017). Classically, osteoblast differentiation is governed by a complex activity involving signal transduction and transcriptional regulation of gene expression (Huang et al., 2007). Runx2 has been identified as the master regulatory switch in osteoblast differentiation, with Osx acting as the 'downstream' regulator of Runx2 (Baldini et al., 2009; Komori, 2019). Furthermore, Dlx5, β -catenin, ATF4, Ihh, Satb2, and Shn3 are other essential transcription factors known to be involved in osteoblast differentiation (Bialek et al., 2004;

CONTACT Junaidi Khotib  junaidi-k@ff.unair.ac.id  Department of Pharmacy Practice, Faculty of Pharmacy, Universitas Airlangga, Nanizar Zaman Joenoes Building, Mulyorejo, Surabaya, 60115, Indonesia.

© 2022 Informa UK Limited, trading as Taylor & Francis Group




Figure 1. Schematic representation of multilineage mesenchymal stem cell (MSC) differentiation. Adipocyte, osteoblasts, and chondrocyte are all part of multilineage differentiation. The progression of osteoblast differentiation toward a mature cell phenotype will result in bone formation.

Lee et al., 2003a; Long et al., 2004; Okamoto et al., 2014; Shim et al., 2013; Tang et al., 2011; Yang & Karsenty, 2004). The discovery of new molecules, known as ncRNAs, that control transcription of osteoblast differentiation and function has recently opened up new avenues to understanding bone pathogenesis (Aurilia et al., 2021; Beermann et al., 2016). Thus, a better understanding of the regulatory mechanisms of osteoblast differentiation provides valuable opportunities for preventing or treating bone-related diseases.

This review looks at recent advances in the regulation of signaling and transcription in osteoblast differentiation. We also discuss about the utilization of these novel molecular players in this mechanism for future clinical applications.

Mesenchymal stem cells in osteoblast differentiation

Stem cells are cells with specific functions that can renew themselves, varying potentials, and differentiate into multiple lineages. Mesenchymal stem cells (MSCs) are stem cells that develop from mesoderm (Ullah et al., 2015). MSCs were first isolated by Friedenstein et al. in the bone marrow and described as adherent cells capable of forming fibroblastic colonies (Friedenstein et al., 1970). Extensive MSC evaluations in recent decades have revealed that MSCs can be isolated from a variety of locations throughout the body. Furthermore, MSCs are multipotent due to their ability to differentiate into specific functional cells such as osteoblasts, adipocytes, or chondrocytes in response to specific factors and signaling cascades in the microenvironment (Knight & Hankenson, 2013; Pino et al., 2012). MSCs are also reported to be capable of expressing CD73, CD90, and CD105, as well as having a low level of expression of surface molecules CD11b, CD14, CD19, CD34, CD45, CD79 α , and human leukocyte antigen-related D antigen (HLA-DR) (Hu et al., 2018). MSCs can be found in the bone compartment in the bone marrow, periosteum, and endosteum, as well as thin layers of connective tissue on the bone surface and the bone itself. They are also a major source of cellular renewal during bone repair. The

capacity of MSCs to differentiate into functional osteoblasts is regulated by osteoblast-specific transcription factors that trigger osteoblast commitment and differentiation, as shown in Figure 1 (Capulli et al., 2014; Hu et al., 2018; Zhang, 2010). The roles of each essential transcription factor involved in osteoblast differentiation are shown in Table 1.

Osteoprogenitor cells

Osteoprogenitor cells (OPCs), also known as osteoblast progenitors (preosteoblasts), are bone stem cells that help with tissue formation and bone repair. OPCs are more common during bone development and can activate a multifunctional stage for bone reconstruction. OPCs can be found in the endosteum, the periosteum's cellular layer, and the osteogenic cell layer (Nahian & Davis, 2021). A large number of OPCs can also be found in bone marrow stromal cells, which are multidirectional. Periosteum and bone marrow-derived osteoprogenitors differentiate directly into osteogenic bone without involving other inducers. These properties of OPCs are known as determined OPCs (DOPCs). While OPCs are found only in pathological situations, such as heterotopic ossification and fracture repair, they are derived from undifferentiated mesenchymal cells found throughout the body. These OPCs can differentiate into osteoblasts via cartilage osteogenesis, hence the name osteoprogenitor-induced OPCs (Qiu et al., 2019). Alpha smooth muscle actin (α SMA) has recently been identified as a marker of OPCs in bone and periodontium, as well as a progenitor of osteochondral in the periosteum that contributes to fracture healing (Matthews et al., 2014). SMA-expressing osteoprogenitors have also been proven to improve site-specific periosteal osteoblast differentiation induced by mechanical loading (Matthews et al., 2020).

OPCs have been shown to divide, propagate, and differentiate further into functionally specialized cells. The majority of OPCs differentiate as osteoblasts during bone development. OPCs attach to the bone surface and are known as inactive osteoblasts

Table 1. Essential transcription factors in osteoblast differentiation.

Model		Model	
No.	Transcription factors	Role	In vivo In vitro
1.	Runx2	Regulates osteoprogenitor proliferation and functions as a master switch in the induction of osteoblast differentiation.	Knockout of Runx2 causes an increased proliferation of osteoblast progenitors.
2.	Osx	Promotes the differentiation of preosteoblasts into osteoblasts, the maturation and function of osteocytes, and cartilage resorption.	Silencing Osx decreases the regulation of Col-X expression, Dlx5, and ALP.
3.	Dlx5	Promotes osteoblast proliferation and differentiation, and is involved in the later stages of osteogenesis.	Dlx5 is a direct transcription activator of Runx2.
4.	β -catenin	An important component in canonical Wnt signaling.	Cells with low β -catenin activity enter the chondrocyte lineage.
5.	ATF4	Regulates terminal osteoblast differentiation as well as chondrocyte proliferation and differentiation during skeletal development.	Knockout of ATF4 shows severe cell proliferative defects and increases cell apoptosis.
6.	lhh	Promotes MSC differentiation and is involved in bone formation.	Knockdown of lhh causes inhibition of osteoblast growth, increased apoptosis, termination of the cell cycle, decreased ALP activity and deposition of osteoblast minerals.
7.	Satb2	Multifunctional determinants of craniofacial pattern and osteoblast differentiation.	Knockout of Satb2 shows a decrease in BSP and OCN regulation.
8.	Shn3	Negative regulator of Runx2.	Knockout of Shn3 causes an increase in the number of terminalized nodules as well as the expression of BSP, OCN, Osx, and ATF4.
9.	Twist1	Inhibits osteoblast differentiation.	Twist-1 haploinsufficiency leads to increased regulation of osteogenic markers, including ALP and COL1A1.

Col-X, X-type collagen.

45

References

Kawane et al., 2018; Komori, 2019
 Zhang et al., 2008a; Zhou et al., 2010; Martin et al., 2011; Omoteyama & Takagi, 2010
 Holleville et al., 2007; Samee et al., 2008
 Day et al., 2005; Hill et al., 2005
 Yang et al., 2004; Yang & Karsenty, 2004; Wang et al., 2009; Zhang et al., 2008b; Zhang et al., 2019
 Nakamura et al., 1997; Deng et al., 2017; Long et al., 2004; Wang et al., 2012; Yang et al., 2015
 Dobrev et al., 2006; Mouillé et al., 2022
 Jones et al., 2006
 Bialek et al., 2004; Quarto et al., 2015; Zhang et al., 2014

Knockout of Runx2 causes osteoblast and oscillation defects and inhibits chondrocyte maturation.
 Osx inactivation leads to the cessation of osteoblast differentiation and new bone formation.
 Dlx5-null showed a significant decrease in bone volume, proliferation and differentiation of osteoblasts, and potentially induced osteoclastogenesis.
 The knockdown of β -catenin inhibits osteoblast differentiation and induces chondrocyte differentiation.
 Ablation of ATF4 causes growth defects and osteogenesis as well as a reduction in bone mass.
 ATF4 deficiency causes delayed bone formation during embryonic development and low postnatal bone mass.
 lhh deficiency causes decreased proliferation and maturation of chondrocytes as well as failure of osteoblast development in endochondral bone.
 Knockout of Satb2 causes craniofacial disability as well as defects in differentiation and function of osteoblasts.
 Double heterozygous mutants, Satb2/Runx2 and Satb2/ATF4, showed defects in bone formation.
 Knockout of Shn3 causes an increase in bone time phenotype and bone formation.
 Twist-1 hypoinsufficiency causes Saethre-Chotzen syndrome.

53
 18
 22
 58
 44
 18
 3
 6




Figure 2. The Wnt and BMP signaling cascades. Both can be activated via canonical and noncanonical pathways, causing osteoblastic transcription factors to be expressed.

in mature bone (Qiu et al., 2019). The recruitment of OPCs initiates the process of new bone formation via a complex and highly regulated interaction between signaling from the systemic and local biomechanical and biophysical environments. This signaling regulates the activation and maturation of OPCs into osteoblasts, which produce and form the extracellular bone matrix (Ibrahim et al., 2016). Age can have a significant impact on OPC regenerative capacity. The accumulation of marrow adipose tissue (MAT) in the bone marrow cavity due to aging contributes to a pathological process that interferes with the maintenance of proper bone tissue repair and the hematopoietic system, increasing the risk of fractures and complications (Ambrosi et al., 2017). Furthermore, estrogen deficiency contributes to increased bone resorption, which results in bone loss (Manolagas, 2000; Syed et al., 2008). The estrogen receptor α (ER α) in osteoblast progenitors has been shown to promote bone formation on the cortex's periosteal surface and prevent resorption on the endocortical surface (Almeida et al., 2013).

Several studies utilizing OPCs as therapy for bone defects have been developed due to their promising role in bone development and healing. Autogenous OPC transplantation into a porous calcium phosphate scaffold can improve mandibular segmental defect repair (Schliephake et al., 2001). In another study, exogenous murine MC3T3-E1 OPCs with a high propensity for osteoblast differentiation demonstrated the ability to migrate systemically to the femoral bone defect and accelerate bone healing (Gibson et al., 2012). Thus, additional research on this topic may provide a potential therapeutic alternative for bone repair in clinical practice.

Osteoblasts

Osteoblasts are mononuclear cuboid cells derived from OPCs in the bone marrow that differentiate from MSCs. Osteoblasts are responsible for bone formation. This cell is distinguished from mesenchymal progenitor cells at the site of membrane and endochondral bone formation (Kobayashi et al., 2008; Ponzetti

& Rucci, 2021). Osteoblasts occupy 4–6% of the total resident cells in bone and play an important role in the fulfilment and maintenance of bone mass along with osteoclasts and osteocytes. In addition, osteoblasts can also differentiate into osteocytes (Capulli et al., 2014; Rutkovskiy et al., 2016). Osteoblasts can site and secrete bone matrix and contribute to bone mineralization to regulate the balance of calcium and phosphate ions in bone formation. Once the OPCs differentiate into osteoblasts, they will be followed by the secretion of collagen I to form osteoid, followed by the precipitation of calcium and phosphorus salts from the blood by osteoblasts and the formation of bonds with osteoid for the mineralization of bone tissue. Furthermore, presence of estrogen receptors in osteoblasts promotes an increase in the number of osteoblasts, which leads to an increase in collagen production. ALP, an enzyme involved in bone mineralization and an early marker of osteoblast differentiation, is also produced by osteoblasts. Increased ALP expression is associated with osteoblast differentiation (Bassi et al., 2011).

The key role of signalling pathways in osteoblast differentiation

The differentiation of osteoblasts from MSCs is triggered by the expression of specific genes, which are subsequently controlled by pro-osteogenic pathways. The wingless-related integration site (Wnt)/ β -catenin and bone morphogenetic proteins (BMPs) pathways are the main pathways that play an important role in promoting MSC's commitment to osteo/chondroprogenitor cells in the initial steps of osteoblastogenesis. Figure 2 illustrates schematically how the Wnt and BMP signaling pathways regulate osteoblast differentiation.

Wnt signaling pathway in osteoblast differentiation

The wingless-related integration site (Wnt) signaling pathway is divided into two parts: a canonical pathway that mediates

signaling via β -catenin stabilization, which is involved in increased bone formation and a noncanonical pathway that works independently of β -catenin, which plays a role in regulation of cell migration and polarity during embryogenesis (Kim et al., 2013; Nemoto et al., 2012). Canonical Wnt, such as Wnt3a and Wnt10b, binds to Frizzled (Fzd) and low-density lipoprotein receptor-associated protein 5/6 (Lrp5/6) to inhibit glycogen synthase kinase-3 β (GSK-3 β) activations, an enzyme that phosphorylates β -catenin, causing ubiquitination and degrade. This inhibition induces the accumulation of β -catenin in the target cell, which results in translocation into the nucleus. β -catenin will initiate the transcription of the target gene through its interaction with members of the T-cell family factor/lymphoid enhancer factor (Tcf/Lef). Meanwhile, noncanonical Wnt, such as Wnt5a, binds to the Fzd, Ror1/2 or Ryk receptor complexes. In addition, some Wnt antagonists, such as dickkopf-1 (DKK-1), sclerostin (SOST), kremen, and others, can inhibit this signaling.

Multipotential cells, as previously stated, can differentiate into osteoblasts and adipocytes. The balance of adipogenic and osteoblastogenic components via Wnt/ β -catenin signaling is thought to be a determinant of the differences in outcomes of these mesenchymal precursor cells. Previous research has shown that ectopic expression of Wnt10b, a subfamily of canonical Wnt pathway, suppresses the expression of adipogenic transcription factors CCAAT/enhancer-binding protein α (C/EBP α) and peroxisome proliferator-activated receptor γ (PPAR γ) in ST2 cells. Following this condition, osteoblastogenic transcription factors are activated, as evidenced by increased regulation of Runx2, Osx, and Dlx5. Meanwhile, in Wnt10b-expressing ST2 cells, partially forced expression of C/EBP α or PPAR γ promotes lipid accumulation while decreasing mineralization. Thus, in Wnt/ β -catenin signaling, C/EBP α or PPAR γ repression is required to direct precursor cells into osteoblasts (Kang et al., 2007).

Furthermore, Okamoto et al. (2014) stated that good bone formation requires cooperation between Wnt5a-induced non-canonical signaling, a subfamily of Wnt noncanonical pathways, and Wnt/ β -catenin signaling. Wnt5a has also been implicated in osteoblast differentiation. Through upregulation of Lrp 5/6 expression in osteoblast cell lineages, Wnt5a can suppress PPAR γ and increase Wnt/ β -catenin signaling (Nemoto et al., 2012; Okamoto et al., 2014). Wnt5a deficiency in osteoblast lineage cells reduces Lrp5/6 expression, lowering the sensitivity of canonical Wnt ligands such as Wnt3a and Wnt10b. This condition interferes with osteoblast differentiation while increasing adipocyte differentiation (Okamoto et al., 2014). TAZ, a Hippo pathway transcription factor, is also known to induce osteoblastogenesis and suppress canonical Wnt signaling semiconductor adipogenesis (Okamoto et al., 2014; Zarka et al., 2022). Thus, during osteoblast differentiation, Wnt5a can play a role in increasing Wnt/ β -catenin and Wnt/TAZ signaling by upregulating Lrp5/6 (Okamoto et al., 2014). The receptor tyrosine kinase-like orphan receptor 2 (Ror2) has been known as the Wnt5a receptor or co-receptor. Nemoto et al. (2012) has proven that the Wnt5a/Ror2 signaling pathway is involved in BMP-2-mediated osteoblast differentiation in Smad-independent

ways. The suppression of Wnt5a/Ror2 expression resulted in the suppression of osteoblast differentiation marker gene expression, specifically ALP and OCN, induced by BMP-2 (Nemoto et al., 2012).

BMP signaling in osteoblast differentiation


BMPs are members of the transforming growth factor β (TGF- β) superfamily that play a role in regulating osteoblast differentiation and inducing bone formation. During development, genetic interventions in the BMP gene cause a variety of extracellular and bone abnormalities (Wang et al., 2017). BMP signaling is mediated by BMP receptors type I (BMPR-I) and type II (BMPR-II) (BMPR-II). Both are serine-threonine receptors that are required for BMP signaling. BMPR-I is consisted of three receptors: BMPR type IA (BMPR-IA), which binds BMP-2 and BMP-4 efficiently; BMPR type IB (BMPR-IB), which binds BMP-4 and BMP-7 efficiently; and activin type I receptor (ActRI), which binds activins, proteins from TGF- β /BMP family members, and BMP-7 (Nohno et al., 1995; Chen et al., 2012). Meanwhile, type II receptors include BMPR-II, which binds to BMP-4 and BMP-7, and activin type II receptors (ActRII) and ActRIIb, which bind to activin and BMP-7 (Rosenzweig et al., 1995; Yamaguchi et al., 2008). Unlike the TGF- β receptor, BMPR-I binds BMP directly without the involvement of BMPR-II, which only binds BMPR-I on the extracellular N-terminus. Furthermore, BMPR-I has a GS domain, which is a cytoplasmic juxta-membrane area made up of glycine and serine that serves as a site for phosphorylation of serine and threonine after the receptor binds to a ligand, activating BMPR-II. This dynamic interaction directs downstream BMP signals via BMP-specific Smad (Smad 1, 5, or 8) or p38 MAPK. Activated receptor kinases, in collaboration with other co-factors, regulate the transcription of specific target genes by forming heterodimeric complexes with nuclear Smad4 (Kim et al., 2017; Chen et al., 2012; Yamaguchi et al., 2008).

Based on its role, BMP-2 has been proven to promote the expression of Runx2, Osx, and osteoblast differentiation markers (ALP, OCN, and type I collagen) in a variety of cells (Yamaguchi et al., 2008; Ogasawara et al., 2004). Furthermore, BMP-2 and BMP-4 are involved in the formation of bone nodules (Wada et al., 1998). The application of recombinant human bone morphogenetic protein-2 (rhBMP-2) therapy has shown promising results both pre-clinically and clinically. rhBMP-2 has the ability to stimulate bone repair and regeneration (Chen et al., 2012; Ueyama et al., 2021). BMP-7 is well-known for its osteogenic activity (Chen et al., 2019; Lavery et al., 2009). It was recently discovered that immature BMP-7, also known as bone-forming peptide-2 (BFP-2), has higher osteogenic activity than mature BMP-7 and induces bone formation in vitro and in vivo (Kim et al., 2017).

The osteoblast differentiation regulation, essential transcription factors involved, and their links

The osteoblast differentiation begins with the commitment of MSCs into osteoblast lineage progenitor cells, later known as preosteoblasts. Preosteoblasts then undergo proliferation,

The role of essential transcription factors in the regulation of osteoblast differentiation



The involvement of non-coding RNAs in the regulation of osteoblast differentiation




Figure 3. Regulation of osteoblast differentiation by essential transcription factors. ncRNAs (miRNAs, lncRNAs, and circRNAs) are novel players that can influence the molecular regulation of osteoblast differentiation by targeting essential transcription factors.

extracellular matrix (ECM) secretion, matrix maturation, and matrix mineralization, which are complexly regulated by various transcription factors (Figure 3). Runx2 is a transcription factor that is required for osteoprogenitor proliferation and osteoblast differentiation. Runx2 expression is known to be low in MSCs, but it increases throughout cell proliferation and then decreases in maturing osteoblasts. Regulation of Runx2 expression in osteoprogenitors requires Ihh at the stage of MSC commitment formation into preosteoblasts (Amarasekara et al., 2021; Nakashima & De Crombrughe, 2003; Shimoyama et al., 2007). In regard, the upstream of Runx2 is involved in controlling the early stages of osteoblast differentiation, including Twist-1 and Satb2. Twist-1, through its physical interaction with Runx2, functions as a switch that blocks Runx2 function, delaying osteoblast differentiation and preventing premature osteoblast formation. Twist-1 expression is also downregulated when osteogenesis begins. Satb2 regulates the expression of BSP and OCN, the osteoblast-forming components during the cell differentiation phase, as well as inhibiting Hoxa2, a bone formation-inhibiting gene (Bialek et al., 2004; Liu & Lee, 2013).

Following cell commitment, preosteoblasts proliferate and express OPN, fibronectin, collagen, and TGF- β 1 receptors (Rutkovskiy et al. 2016). The proliferation phase is regulated in order to induce maturation during osteogenesis (Vimalraj et al., 2015). Runx2 expression is still increasing at this stage, promoting Osx to mediate osteoblast commitment and differentiation. Moreover, Dlx5 is present to aid in early osteoblast differentiation and advanced stages of osteogenesis. Dlx5 has been shown to influence the expression of Osx, ALP, OCN, and BSP (Holleville et al., 2007; Samee et al., 2008). Following the proliferation phase, the formed osteoblasts begin to express the bone matrix protein gene at various levels depending on cell maturation. ATF4 is also involved in indirect interactions with Runx2 during cell maturation to increase OCN expression, a marker of terminal osteoblast differentiation (Xiao et al., 2005). Along with this, Satb2 also acts as a mediator to increase the synergy of both actions (Dobrev et al., 2006). Mature osteoblasts secrete COL1A1, a key component of ECM, as well as ALP, which aids in ECM maturation. This process is then followed by matrix mineralization. This step is triggered after osteoblasts

Table 2. Essential osteoblastic markers.

No.	Marker	Role	References
1.	ALP	ECM maturation.	Stein and Lian, 1993
2.	BSP-II	Promotes mineralization by regulating the formation of hydroxyapatite crystals.	Gordon et al., 2007; Kim et al., 1994; Lin et al., 2020
3.	COL1A1	ECM main constituents.	Stein and Lian, 1993
4.	OCN	Terminal osteoblast differentiation markers, which regulate calcium metabolism and promote mineral deposition in ECMs.	Xiao et al., 2005
5.	OPN	Increases MSC proliferation capacity in a dose-dependent manner and promotes bone formation and mineralization.	Lin et al., 2020

bind to the existing matrix via integrin $\beta 1$, forming a single layer that is linked to cadherin. Furthermore, cells secreted the matrix by expressing OPN, OCN, and BSP, as well as maintaining ALP and COL1A1 expression (Table 2) (Huang et al., 2007; Stein & Lian, 1993). Runx2 protein levels decreased at the end of this stage, which could be regulated by Shn3, an adapter that induces Runx2 degradation via ubiquitination (Jonason et al., 2009; Jones et al., 2006; Shim et al., 2013). Additionally, osteoblasts that have completed their roles in bone homeostasis will undergo apoptosis, becoming bone-lining cells, or terminally differentiate osteocytes (Nakashima & De Crombrughe, 2003; Amarasekara et al., 2021).

The following are reviews of the essential transcription factors involved in osteoblast differentiation:

Runx-related transcription factor 2

Runx-related transcription factor 2 (Runx2) is a transcription factor that is required for osteogenesis and is responsible for activating osteoblast differentiation marker genes. Runx2 specifically increases the expression of osteoblastogenic markers such as ALP, BSP-II, collagen1 α 1 chain (Col1A1), OCN, and OPN, which leads to osteoblast commitment (Vimalraj et al., 2015; Ponzetti & Rucci, 2021). Furthermore, Runx2 regulates osteoblast progenitor proliferation by inducing fibroblast growth factor receptor (FgfR)-2 and Fgf3 expression. Both promote proliferation by activating the mitogen-activated protein kinase (MAPK) pathway (Kawane et al., 2018; Komori, 2019).

Osterix

Osterix (Osx) or Sp7 is an osteoblast-specific transcription factor that is involved in preosteoblast differentiation into osteoblasts and bone formation. Osx has a proline-rich region (PRR) transactivation domain near the N-terminus of the protein and a three-type DNA binding domain C2H2 zinc finger near the C-terminus with motifs similar to Sp1, Sp3, and Sp4 (Zhang et al., 2008a). Osx may play a role in the bone microenvironment during osteogenesis (Liu et al., 2020). As a downstream of Runx2, Osx is expressed specifically in osteoblasts of all endochondral bones and membranes, as well as at low levels in pre-hypertrophic

chondrocytes (Zhang et al., 2010; Tang et al., 2011). Zhou et al. (2010) demonstrated that inactivating Osx during and after birth resulted in the cessation of osteoblast differentiation and new bone formation in mice. Furthermore, Osx is required for cartilage resorption, maturation, and osteocyte function. During the bone formation, osteoblasts are known to store osteoid, which are unmineralized matrixes containing type I collagen. In this case, osteocytes participate in the osteoid mineralization process by regulating extracellular matrix mineralization and Fgf23 production by bone via Dmp1 and Phex, gene products that are highly expressed in normal osteoblasts and osteocytes (Martin et al., 2011; Zhou et al., 2010). Osteocyte anomalies in $Osx^{postnatal}$ mutants revealed defects in the mineralization process due to decreased Dmp1 and Phex expression. Furthermore, terminalized cartilage resorption also defects $Osx^{postnatal}$ mutants due to a significant decrease in osteoclast density (Zhou et al., 2010).

Distal-less homeobox 5

Distal-less homeobox 5 (Dlx5) is a proliferation and early osteoblast differentiation driver that also influences the later stages of osteogenesis. This transcription factor is expressed specifically in osteogenic lineage cells, such as chondroprogenitor cells (Samee et al., 2008). Dlx5 can induce osteoblast differentiation from endochondral and membrane ossifying bone (Erceg et al., 2003). Furthermore, in vitro studies have reported that Dlx5 acts as a direct transcription activator of Runx2 by binding to the P1 promoter, which is the transcriptional regulator of the Runx2-II isoform. Dlx5 expression is specifically induced by BMP signaling pathways such as BMP-2 or BMP-4 stimulation (Holleville et al., 2007; Samee et al., 2008). Earlier studies have shown that the Dlx5 modulation of osteoblast differentiation mediates Runx2 expression via BMP-2 stimulation (Lee et al., 2003a; Holleville et al., 2007). Dlx5 has also been shown to promote ALP and Osx via a Runx2-independent pathway. Osx expression was also reduced in $Dlx5^{-/-}$ osteoblast cultures (Holleville et al., 2007; Samee et al., 2008). Osx, as previously stated, is a downstream of Runx2 (Samee et al., 2008; Lee et al., 2003b). However, Dlx5 directly controls OCN and BSP transcription in in vitro studies due to a significant decrease in $Dlx5^{-/-}$ osteoblast culture (Samee et al., 2008).

β -catenin

β -catenin is an important component that transduces canonical Wnt signaling to determine the direction of mesenchymal progenitor differentiation, regardless of regional location or oscillation mechanism. Activation of β -catenin leads to increase oscillation and suppression of chondrocyte formation. In the meantime, its inactivation inhibits osteoblast differentiation during intramembranous and endochondral ossification and promotes chondrocyte differentiation (Day et al., 2005; Hill et al., 2005; Hu et al., 2005). Furthermore, β -catenin/TCF1 has been shown to increase Runx2 promoter expression and activity, causing osteoprogenitor cells to differentiate into preosteoblasts. β -catenin, along with Osx, regulates the differentiation of preosteoblasts into immature osteoblasts (Gaur et al., 2005).

Activating transcription factor 4

Activating transcription factor 4 (ATF4) is a leucine basic zipper (bZip) transcription factor from the ATF/cAMP family response element-binding protein (CREB). The ATF4 gene appears to be expressed in variety of cells during development and embryonic life. However, the accumulation of ATF4 protein is strongly inversely related to its gene expression because ATF4 protein is degraded in most cells, except osteoblasts, via ubiquitination mediated by β -TrCP1, ligase ubiquitin E3 (Yang & Karsenty, 2004; Zhang et al., 2019). ATF4 is required for terminal osteoblast differentiation via OCN activation, as well as regulation of chondrocyte proliferation and differentiation during skeletal development via *Ihh* activation (Wang et al., 2009; Yang & Karsenty, 2004; Zhang et al., 2019). ATF4 ablation has been linked to severe osteopenia, impaired terminal osteoblast differentiation, and decreased OCN expression and type I collagen production in mice (Yang et al., 2004). Previous studies demonstrate that ATF4 acts as a specific activator of osteocalcin-specific element 1 (OSE1) in an RSK2-dependent manner, and that it indirectly associates with Runx2 to increase OCN expression, which eventually leads to terminal osteoblast differentiation (Xiao et al., 2005; Yang et al., 2004). *Satb2* mediates the synergistic action of both. Transcription factor general IIag (TFIIAg) has also been reported to increase OCN expression via interactions with Runx2 and ATF4 (Dobrev et al., 2006; Yu et al., 2008). Furthermore, Tominaga et al. (2008) stated that CCAAT/enhancer-binding proteins (C/EBPs), a bZip protein family, are responsible for increasing OCN promoter activity via a heterodimeric bond with ATF4 on OSE1. C/EBP β is expressed in osteoblastic cells, and its regulation becomes more active during osteoblast differentiation. The heterodimerization also facilitates it to collaborate with Runx2 (Tominaga et al., 2008). Xiao et al. (2005) demonstrated that the physical interaction between C/EBP β and Runx2 promotes OCN promoter gene expression. Thus, C/EBP β promotes the formation of complexes and associations between ATF4 and Runx2 in order to encourage OCN expression during terminal osteoblast differentiation (Tominaga et al., 2008). Moreover, ATF4 also plays a role in the regulation of

chondrocyte proliferation and differentiation during skeletal formation by involving *Ihh* transcription and signaling. ATF4 overexpression in mutant chondrocytes restored osteoblastic marker gene (OCN and BSP) expression in developing bone, according to studies using the ATF4^{-/-};COL2A1-ATF4 mouse model, in which ATF4 was expressed in chondrocytes selectively in an ATF4-null background. This is followed by correction of the bone elongation defect as well as improvement in decreased *Ihh* expression and *Hh* signaling. As a result, ATF4 is involved in the autonomic role of chondrocytes in growth plate development and may also be involved in osteogenesis regulation during postnatal bone development and remodeling (Wang et al., 2009, 2012).

Indian hedgehog

The transcription factor Indian hedgehog (*Ihh*) is a *Drosophila hedgehog* (*Hh*) mammalian homologue that is important for osteoblast differentiation and bone formation. *Ihh* acts as the primary regulator in the longitudinal growth and development of the endochondral skeleton. It is primarily expressed by peritrophic chondrocytes that have recently exited the cell cycle and send signals to proliferative chondrocytes to divide and perichondrial mesenchymal cells to differentiate into osteoblasts. The dysfunctional regulation of *Hh* signaling causes problems with bone homeostasis and development, as well as the onset of several bone diseases such as progressive heteroplasia and osseous dysplasia (Long et al., 2004; Wang et al., 2012; Yang et al., 2015). *In vivo* studies revealed that *Ihh* gene deficiency reduced chondrocyte proliferation and maturation, as well as failed osteoblast development in endochondral bone (St-Jacques et al., 1999). In an *in vitro* study using MC3T3-E1 osteoblast cells, *Ihh* expression increases ALP activity through cooperation with BMP-2 (Nakamura et al., 1997). Meanwhile, knocking out *Ihh* causes an increase in apoptosis, cell cycle termination in the G1 to S phases in osteoblasts, as well as a decrease in ALP activity and osteoblast mineral deposition, which are associated with the TGF- β /Smad and OPG/RANKL signaling pathways (Deng et al., 2017). Furthermore, *Hh* proteins, including *Ihh*, exert biological effects via their receptor components, patched (PTCH) and smoothened (Smo). *Ihh* binding to PTCH activates Smo and transduces signals in the cytoplasm via the fusion of intracellular signal molecules and transcription factors from the Gli family with the zinc finger domain. Gli2 and Gli3 are direct *Ihh* signaling mediators. *Ihh*/Gli2 signaling promotes mesenchymal cell differentiation in osteoblastogenesis by regulating expression and stimulating Runx2 osteoblastogenic function. Gli2 physically increases Runx2 expression and Runx2 osteogenic activity (Shimoyama et al., 2007). Meanwhile, Gli3 functions as a transcriptional repressor of the *Hh* target gene (Hilton et al., 2005).

Special at-rich sequence binding protein 2

Special AT-rich sequence binding protein 2 (*Satb2*) is a gene derived from a family of special AT-rich binding proteins that bind nuclear matrix attachment regions (MARS), an AT-rich

DNA sequence involved in gene transcription regulation. MAR affects the organization of eukaryotic chromosomes by structurally defining the boundary of the chromatin domain and increasing the ability of enhancers to work over long distances. *Satb2* is known to be located in the poor region of the 2q32-q33 gene, and its coding transcript consists of 11 exons and 191 kb of genomic DNA (FitzPatrick et al., 2003; Dobreva et al., 2006). *Satb2* is thought to increase expression of gene differentiation-specific type cells by regulating chromatin recurrence in MAR (Dobreva et al., 2006). *Satb2* has been proven to be a multifunctional determinant of craniofacial pattern and osteoblast differentiation. Activation of *Satb2* by *Osx* overexpression is responsible for MSC differentiation into osteoblasts by targeting inhibition of several Hox genes (*Hoxa13*, *Hoxa2*, and *Hoxb2*) in osteoblasts. Hox is a gene that regulates branchial arch patterns by directly recognizing MAR-like sequences (Dobreva et al., 2006; Mouillé et al., 2022). Furthermore, *Satb2* also targets *BSP* as an initial marker for osteoblast differentiation and *OCN* as a terminal marker for osteoblast differentiation (Dobreva et al., 2006). *BSP* is the main structural protein of the bone matrix that promotes osteoblast differentiation and thus increases the production of terminalized matrix (Gordon et al., 2007; Kim et al., 1994). *Satb2* directly binds to the *BSP* promoter region associated with the three osteoblast-specific element sequences, according to ChIP and EMSA analysis of fully-differentiated osteoblasts (Dobreva et al., 2006; Kim et al., 1994). Meanwhile, *OCN*, which is involved in bone matrix mineralization, is the second most abundant protein in bone after collagen. *OCN* is highly expressed in matured osteoblasts, which initiate the bone formation (Li et al., 2016; Rutkovskiy et al., 2016). Existing literature has shown that *Runx2* and *ATF4*, transcription factors that promote mineralization at different stages of the bone formation process, play a role in *OCN* regulation. In their functional synergy, *Runx2* and *ATF4* have indirect interactions. *Satb2* acts as a synergistic activation mediator of *Runx2* and *ATF4* in this regard because double heterozygous mutant mice, *Satb2/Runx2* and *Satb2/ATF4*, showed defects in bone formation. Thus, *Satb2* physically interacts with *Runx2* and *ATF4*, increasing their transactivation function; incorporates the specific-transcription factor at the *OCN* promoter, encouraging *OCN* expression indirectly (Bidwell et al., 1993; Dobreva et al., 2006; Ducky & Karsenty, 1995).

Schnurri-3

Schnurri-3 (Shn3) is a large zinc finger protein that plays an important role in embryogenesis as a cofactor for Decapentaplegic signaling (Dpp), a *Drosophila* homolog of the BMP/TGF- β signaling pathway. *Shn3* is one of three *Drosophila* Shn mammalian homologs that act as essential regulators of bone formation, regulating osteoblast activity (Jones et al., 2006). *Shn3* knockout mice had a higher bone tenure phenotype due to increased synthetic osteoblast activity and bone formation. In osteoblasts, a multimeric complex composed of *Runx2*, *Shn3*, and the Nedd4 family's E3 ubiquitin ligase *WWP1* can inhibit *Runx2* function. This is

because *Shn3* promotes *Runx2* poly-ubiquitination and proteasome-dependent degradation via *WWP1*. Therefore, the absence of *Shn3* in osteoblasts causes an increase in *Runx2* protein levels, followed by an increase in *Runx2* transcriptional activity and target gene, which increases extracellular matrix mineralization during the bone formation process (Jones et al., 2006). Furthermore, *Shn3* mediates interaction and inhibition of ERK activity in the Wnt signaling pathway in osteoblasts. In vivo studies revealed that knockdown in this section causes abnormal ERK activation, resulting in osteoblast hyperactivity and bone development problems (Shim et al., 2013).

Twist-associated protein 1

Twist-related protein 1 (Twist-1) is a basic helix-loop-helix (bHLH) transcription factor that acts as an anti-osteogenic and osteogenesis initiator (Komaki et al., 2007; Lee et al., 1999; Zhang et al., 2014). In humans and mice, heterozygous loss of *Twist-1* functions causes Saethre-Chotzen syndrome, which is characterized by craniosynostosis, a condition caused by premature osteoblast differentiation in the skull (Quarto et al., 2015). *Twist-1* has been shown to suppress osteoblast differentiation by inhibiting *Runx2* function. The decreasing *Twist-1* gene expression triggers osteoblast differentiation through increased expression of *Runx2* downstream in vivo. *Twist-1* overexpression also inhibits osteoblast differentiation while having no effect on *Runx2* expression (Bialek et al., 2004; Zhang et al., 2014). Similar to *Runx2*, *Twist-1* also inhibits *ATF4* function without interfering with *ATF4* protein levels (Danciu et al., 2012). In vitro experiments with C3H10T1/2 cells revealed that decreased *Twist-1* gene expression resulted in increased *ALP* and *COL1A1* expression. Moreover, *Twist-1* has been shown to modulate *Fgfr2* expression, activating ERK1/2 and PI3K signaling during osteoblastogenesis (Guenou et al., 2005; Miraoui et al., 2010). *Twist-1* and *Twist-2* haploinsufficient mouse models revealed that hereditary expression of *Fgfr2* and *Fgfr1-4* causes a decrease in bone formation, proliferative disorders, and osteoprogenitor differentiation (Huang et al., 2014).

Noncoding RNAs and their interplay with essential transcription factors: an advanced regulation concept for osteoblast differentiation

Noncoding RNAs (ncRNAs) are functional RNA molecules that do not have the ability to encode proteins, so they were initially considered 'evolutionary garbage'. However, emerging evidence has established the role of ncRNAs as potent and multifunctional regulators in all biological processes, including transcriptional regulation of osteoblast differentiation (Beermann et al., 2016; Aurilia et al., 2021). In parallel, these studies have discovered an association between ncRNA expression and disease progression in humans, including bone-related diseases. MicroRNAs (miRNAs), small interfering RNAs (siRNAs), PIWI-interacting RNAs (piRNAs), long noncoding RNAs (lncRNAs), circular RNAs (circRNAs), and other ncRNAs have been identified due to advances in RNA-Seq

Table 3. miRNAs involved in osteoblast differentiation regulation.

No.	miRNAs	Therapeutic prospects	Roles	References
1.	miR 23a–27a–24-2	Bone formation	Inhibits bone formation by targeting Runx2 and Satb2.	Hassan et al., 2010
2.	miR-39a and -b	Bone loss	Promotes osteoblast differentiation by decreasing the expression of Wnt signaling antagonists (DKK-1, Kremen2, and sFRP2) and AKT/ β -catenin (PTEN).	Kapinas et al., 2010; Xia et al., 2020
3.	miR-214	Osteoporosis	Inhibits osteoblast differentiation by targeting ATF4.	Wang et al., 2013
4.	miR-34b and -c	Skeletogenesis (embryogenesis and postnatally)	Inhibits osteoblast terminal differentiation by targeting Satb2.	Wei et al., 2012
5.	miR-31	Bone formation	Inhibits osteoblast differentiation by reversing Osx expression and suppressing the level of the Satb2 protein.	Baglio et al., 2013; Deng et al., 2013; Xie et al., 2014
6.	miR-322	Bone formation	Enhances BMP-2 response and Osx expression by targeting Tob2.	Gómez et al., 2013
7.	miR-203 and miR-320b	Bone formation	Inhibits BMP-2 stimulates osteoblast differentiation by targeting Dlx5, which inhibits the roles of Runx2 and Osx.	Laxman et al., 2016
8.	miR-27a	Bone formation	Inhibits Osx expression and attenuates Satb2-induced osteoblast differentiation.	Gong et al., 2016
9.	miR-467g	Bone formation	Inhibits osteoblast differentiation by targeting Runx2 and Ihh signaling.	Kureel et al., 2017
10.	miR-376c-3p	Skeletal abnormalities	Inhibits osteoblast proliferation and differentiation through Twist-1 regulation.	Camp et al., 2018
11.	miR-145	Adolescent idiopathic scoliosis	Disrupt osteoblast and osteocyte function through upregulation of β -catenin expression.	Zhang et al., 2018
12.	miR-26b	Osteoporosis and osteoarthritis	Promotes osteoblast differentiation by regulating β -catenin.	Hu et al., 2019; Yang et al., 2022
13.	miR-103	Osteoporosis	Inhibits osteoblast proliferation and differentiation by targeting Satb2.	Lv et al., 2020
14.	miR-92a-3p	Fracture healing	Inhibits IBSP expression and promotes osteoblast differentiation via the PI3K/AKT signaling pathway.	Hu et al., 2021

sFRP2, secreted frizzled related protein 2; PTEN, phosphatase and tensin homolog; IBSP, integrin binding sialoprotein.

(Lekka & Hall, 2018; Li et al., 2020, 2021). Here, we will discuss three ncRNAs that have been hot topics in the development of new therapeutic targets for bone-related diseases: miRNA, lncRNA, and circRNA. Figure 3 depicts the involvement of these ncRNAs in osteoblast differentiation.

MicroRNA

MicroRNAs (miRNAs) are a type of short noncoding RNA that contains 20–22 nucleotides. MiRNAs do not encode proteins, but they regulate the levels of other proteins, particularly at the post-transcriptional level, by lowering messenger RNA (mRNA) levels or inhibiting translation by binding to the 3'UTR of the target mRNA (Inose et al., 2009; Wang et al., 2020). Interestingly, Davis and Hata (2009) reported that miRNAs regulate approximately one-third of human genes. Evidence of miRNA involvement in regulating osteoblast differentiation and bone formation has been studied continuously over the last two decades. Wnt and BMP are the main signaling pathways in osteoblast differentiation, as previously described, and miRNAs have been shown to target both.

Understanding the role of miRNAs in osteoblastogenesis will provide important therapeutic insights. Previous research has found that miRNAs play a role in the regulation of osteoblast differentiation (Table 3). Several miRNAs, including miR-322, miR-27a, miR-26b, and miRNA-92a-3p, act as positive regulators of osteoblast differentiation. MiR 23a27a24-2, miR-214, miR-34b and -c, miR-31, miR-203, miR-320b, miR-467g,

miR-376c-3p, miR-145, and miR-103 are examples of negative regulators.

Long noncoding RNA

Long noncoding (lnc) RNA (lncRNAs) is a diverse class of transcripts that is 200–10,000 nucleotides longer than other types of ncRNAs (Nardocci et al., 2018; Silva et al., 2019). This type of ncRNA is widely transcribed in the nucleus by RNA polymerase II via 5' capping, 3' poly-A tail addition, and RNA splicing (Aurilia et al., 2021). lncRNAs are poorly conserved among mammalian species but play critical roles in transcriptional and post-transcriptional regulation, mRNA translation control, and chromatin structure regulation (Nardocci et al., 2018). Furthermore, lncRNAs can act as endogenous RNAs (ceRNAs) for miRNA 'sponges' and influence miRNA expression, thereby reducing the regulatory effect of miRNAs on miRNA targets (Thomson & Dinger, 2016; Xiao et al., 2017).

Unlike miRNAs, lncRNAs can be folded into complex secondary and higher-order structures to improve target recognition (Huang et al., 2015). These findings point to the role in the emergence and development of a disease. Several recent studies have found that lncRNAs play a role in osteoblast differentiation and act as ceRNAs targeting downstream miRNAs (Table 4). H19, TUG1, MEG3, and MALAT1 are lncRNAs that act as positive regulators of osteoblast differentiation. H19, ANCR, ODIR1, AK045490, HOTAIR, UCA1, Xist,

Table 4. lncRNAs involved in osteoblast differentiation regulation.

No.	lncRNAs	ceRNAs	Effects	Therapeutic prospects	References
1.	lncRNA H19	miR-141 and miR-22	<ul style="list-style-type: none"> Positive regulator of osteoblast differentiation by negatively regulates TGF-β1 with miR-675, thus increased Runx2 expression. H19 is negatively regulated by miR-675-5p via the formation of a feedback loop. 	Osteoporosis	Huang et al., 2015; Liang et al., 2016
2.	lncRNA ANCR	miR-19b-3p	Negative regulator of osteoblast differentiation.	PMOP	Xiaoling et al., 2020
3.	lncRNA ODIR1	Not mentioned.	Negative regulator of osteogenic differentiation.	PMOP	Cai et al., 2019
4.	lncRNA AK045490	Not mentioned.	Negative regulator of osteoblast differentiation.	Bone formation	He et al., 2019
5.	lncRNA MEG3	Not mentioned.	Positive regulator of osteoblast differentiation.	Osteoporosis	Li et al., 2019a
6.	lncRNA MALAT1	miR-204	Positive regulator of osteoblast differentiation.	Fracture healing	Li et al., 2019b
7.	lncRNA TUG1	miR-30	Positive regulator of osteoblast differentiation.	CAVD	Yi et al., 2019;
8.	lncRNA HOTAIR	Not mentioned.	Negative regulator of osteoblast differentiation and bone formation.	mineralized bone matrix.	Xiao et al., 2017
9.	lncRNAUCA1	Not mentioned.	Negative regulator of osteoblast differentiation.	Bone formation	Liu et al., 2019
10.	lncRNA Xist	miR-19a-3p	Negative regulator of osteoblast differentiation.	Osteoporosis	Shen et al., 2019
11.	lncRNA DANCR	Not mentioned.	Negative regulator of osteoblast differentiation.	Osteoporosis	Zhang et al., 2019
12.	lncRNA AK039312 and AK079370	miR-199b-5p	Negative regulator of osteoblast differentiation.	Osteoporosis	Chen et al., 2020

ANCR, anti-differentiation noncoding RNA; ODIR1, osteogenic differentiation inhibitory regulator 1; TUG1, taurine up-regulated gene 1; HOTAIR, HOX transcript antisense RNA;UCA1, urothelial carcinoma associated 1; DANCR, differentiation antagonizing nonprotein coding RNA; PMOP, postmenopausal osteoporosis; CAVD, calcific aortic valve disease.

Table 5. circRNAs involved in osteoblast differentiation regulation.

No.	circRNAs	Roles	ceRNAs	Effects	Therapeutic prospects	References
1.	circRNA_AFF4	Positive regulator of osteoblast differentiation.	miR-7223-5p	Activating the PI3K-AKT signaling pathway.	Fracture healing	Mi et al., 2019
2.	hsa_circ_0074834	Positive regulator of osteoblast differentiation.	miR-942-5p	Promotes osteoblast differentiation and repair of bone defects. ¹⁰⁴	Bone nonunion	Ouyang et al., 2019
3.	circSIPA1L1	Positive regulator of osteoblast differentiation.	miR-617	Promotes osteogenesis by targeting Smad3.	Bone defect	Ge et al., 2020
4.	hsa_circ_0076690	Positive regulator of osteoblast differentiation.	miR-152	Enhances osteoblast differentiation by targeting Runx2.	Osteoporosis	Han et al., 2020
5.	circ_0024097	Positive regulator of osteoblast differentiation. ³⁹	miR-376b-3p	Activating the Wnt/ β -catenin signaling pathway by targeting YAP1 ⁸⁸	Osteoporosis	Huang et al., 2020
6.	circRNA124534	Positive regulator of osteoblast differentiation.	miR-496	Modulates osteoblast differentiation by engaging the Wnt/ β -catenin signaling pathway. ¹⁰⁹	Bone formation	Ji et al., 2020
7.	circ_0076906	Positive regulator of osteoblast differentiation.	miR-1305	Promotes osteoblast differentiation by competing with OGN, resulting in increased Runx2 and OCN expression.	Osteoporosis	Wen et al., 2020
8.	circRNA_25487	Negative regulator of osteoblast differentiation.	miR-1344-3p	Inhibits bone repair by upregulating p21 expression.	TONFH	Zhang et al., 2020
9.	circRNA_TGFBR2	Negative regulator of osteoblast differentiation.	miR-25-3p	Positively regulates Twist1, which inhibits the formation of calcified nodules; ALP activity, as well as Runx2 and OPN expression.	CAVD	Yu et al., 2021
10.	circRNA-vglI3	Positive regulator of osteoblast differentiation.	miR-326-5p	Promotes osteoblast differentiation of ADSCs via Itga5.	Nonhealing bone defects	Zhang et al., 2021
11.	circ_0000020	Positive regulator of osteoblast differentiation.	miR-142-5p	Regulates BMP2 expression, thereby increasing the expression of Runx2, Osx, OCN, OPN, ALP activity, and mineral accumulation.	Osteoporosis	Zhou et al., 2021
12.	circStag1	Positive regulator of osteoblast differentiation.	Not mentioned.	Regulated HuR-mediated Wnt signaling.	PMOP	Chen et al., 2022
13.	circ_0019693	<ul style="list-style-type: none"> Positive regulator of osteoblast differentiation. Circ_0019693 overexpression increases ALP activity, RUNX2, OPN and OCN expression, and promotes angiogenesis. 	miR-942-5p	Regulates PCP4.	Osteoporosis	He et al., 2022
14.	circ_0003204	Negative regulator of osteoblast differentiation.	miR-370-3p	Inhibits osteoblast differentiation by promoting HDAC4 expression.	Bone defect	Yu et al., 2022

YAP1, yes-associated protein 1; OGN, osteoglycin; TONFH, trauma-induced osteonecrosis of femoral head; PMOP, postmenopausal osteoporosis; CAVD, calcific aortic valve disease; TGFBR, transforming growth factor beta receptor 2; Itga5, integrin 5; HuR, human antigen R; PCP4, purkinje cell protein 4.

DANCR, AK039312, and AK079370 are examples of negative regulators.

Circular RNA

Circular RNAs (circRNAs) are a type of ncRNA that is being studied extensively as therapeutic targets and biomarkers. CircRNA has a covalently closed ring structure and is an endogenous biomolecule that lacks a 5' end cap or 3' poly(A) tail, making it highly stable and resistant to RNase R and other exonucleases, with a mean half-life of more than 48 hours (Barrett & Salzman, 2016; Chen et al., 2021). CircRNAs are found exclusively in the cytoplasm, but some can also be found in the nucleus, particularly circRNAs containing introns. Like lncRNA, circRNA also acts as a miRNA sponge.

Furthermore, circRNA regulates gene transcription and translation, modifies alternative splicing, and can interact with regulatory RNA-binding protein (RBP) (Chen et al., 2021; Patil et al., 2020). Recent research has revealed that circRNAs play an important role in bone disease by regulating osteoblast differentiation. Table 5 shows some examples of circRNA's role in osteogenic regulation. CircRNAs that promote osteoblast differentiation include AFF4, has_circ_0074834, circSIPA1L1, has_circ_0076690, circ_0024097, circRNA124534, circ_0076906, circRNA-vgll3, circ_0000020, circStag1, and circ_0019693. CircRNA 25487, circRNA TGFBR2, and circ_0003204, on the other hand, inhibited osteoblast differentiation.

Prospect for the future

Rapid technological advances have created a plethora of new therapeutic targets for bone-related diseases. In recent years, studies on the involvement of ncRNAs in osteoblast differentiation among the classical mechanisms were reported to influence the activity of transcription factors in regulating physiological and pathological processes. According to this evidence, there is a growing interest in ncRNA-based therapies for bone diseases. Various exogenous ncRNA delivery systems to target sites have also been investigated. Several base scaffolds and carriers are being developed for delivery of ncRNAs, including liposomes, hydrogels, exosomes, synthetic and natural nanoparticles, nanofibers, and microspheres (Balagangadharan et al., 2018; Guan et al., 2021; Li et al., 2021). These findings suggest a new challenge in the discovery and development of therapeutic strategies for bone-related diseases, as well as a promising opportunity in controlling bone regeneration. Notably, it was discovered that microchannel porous hydroxyapatite scaffolds interfere with miRNA expression (Jiajun et al., 2020). More intriguingly, the combination of ncRNAs and specific scaffold biomaterials, such as hydroxyapatite, is expected to not only improve the osteogenic performance of the scaffolds but also act as drugs for bone-related diseases (Damiani & El-Messeiri, 2021; Khotib et al., 2021; Pan et al., 2021). Unfortunately, studies on the potential role of ncRNAs in the mechanism of osteoblast differentiation have not yet covered all bone-related

diseases. The research available is limited to common cases like osteoporosis, fractures, and osteoarthritis. Meanwhile, it has not been widely investigated in other cases, such as Paget's disease of bone (PDB), periodontitis, osteogenic imperfecta, osteosarcoma, etc.

As an example, consider PDB. It is a bone regeneration disorder characterized by excessive osteoclastic bone resorption followed by an increase in osteoblastic activity to compensate for bone remodeling (Nebeker, Valenzuela & Pietschmann, 2017). Despite the fact that PDB is the second most common bone disease after osteoporosis, recent reports indicate a global decline in prevalence and severity, which may be due to changes in environmental and lifestyle factors. Finally, these conditions lead to a decrease in PDB diagnoses (Michou & Orcel, 2019; Gennari et al., 2019). Current research on PDB therapeutic targets has focused on osteoclast-related pathways, such as the receptor activator of nuclear factor- κ B ligand (RANKL). Surprisingly, Marshall et al. (2009) and Yavropoulou et al. (2012) described the role of osteoblasts in PDB. SOST and DKK-1 levels of antagonists of the Wnt signaling pathway were found to be higher in PDB patients than in healthy controls. By contrast, Idolazzi et al. (2017) and Werner de Castro et al. (2019) reported that serum levels of SOST and DKK-1 in PDB patients were comparable to healthy subjects. On the other hand, the role of the Wnt/ β -catenin signaling system in the pathogenesis of PDB cannot be denied.

Likewise, periodontitis is a chronic bacterial (*Porphyromonas gingivalis*)-related inflammation of the soft tissues that support tooth structure. The significant proportion of the variation in periodontitis severity is thought to be due to genetic factors (Sayad et al., 2020). In this case, osteoblasts are functionally important cells that, along with periodontal ligament stem cells (PDLSCs), contribute to the physiological function of periodontal tissues and participate in periodontal regeneration (Yu et al., 2017). Several recent studies have reported the role of ncRNAs in the regulation of osteogenic differentiation on gene expression in human PDLSCs (Cuevas-González et al., 2021; Santonocito et al., 2021; Sayad et al., 2020). Anyway, more detailed reports on the various roles of ncRNAs in osteoblast differentiation and their potential effects in this area are still lacking.

Overall, further exploration into the roles of major signaling pathways, essential transcription factors, and ncRNAs in the molecular mechanisms of osteoblast differentiation is urgently needed in order to develop better therapeutic strategies for bone-related diseases.

Conclusion

Studies into the molecular mechanisms of osteoblast differentiation have greatly evolved. Previously, classical mechanisms thought that essential transcription factors were present to control MSC differentiation and commitment to osteogenesis. Surprisingly, in recent years, the use of high-throughput sequencing technology in conjunction with bioinformatics analysis has successfully identified the involvement of ncRNAs in bone regeneration, gaining insights in

98 orthopaedic and endodontic fields. This review described advances in the understanding of molecular mechanisms of osteoblast differentiation involving the roles of essential transcription factors and ncRNAs. Both interactions form a regulatory complex that controls gene expression. These findings provide exciting and valuable information in the identification of novel molecular players, paving the way for the future development of therapeutic agents as well as biomarkers for diagnostic and beneficial follow-up procedures in the treatment of bone-related diseases.

Disclosure statement

25 The authors declare no conflict of interest in the publication of this manuscript.

Funding

This work was supported by the Ministry of Education, Culture, Research, and Technology of the Republic of Indonesia with PMDSU scheme No. 891/UN3.15/PT/2022.

ORCID

Muhammad Tahir  <http://orcid.org/0000-0002-1463-3090>

Authors' contributions

Junaedi Khotib, Andang Miatmoko, and Aniek Setiya Budiati: Conceptualization and supervising manuscript. Honey Dzikri Marhaeny: collecting literature data, visualization, writing and editing original draft preparation. Chrismawan Ardianto, Mahardian Rahmadi, and Muhammad Tahir: methodology, reviewing, and validating manuscript. Yusuf Alif Pratama: collecting literature data.

References

- Almeida, M., Iyer, S., Martin-Millan, M., Bartell, S. M., Han, L., Ambrogini, E., Onal, M., Xiong, J., Weinstein, R. S., Jilka, R. L., O'Brien, C. A., & Manolagas, S. C. (2013). Estrogen receptor- α signaling in osteoblast progenitors stimulates cortical bone accrual. *The Journal of Clinical Investigation*, 123(1), 394–404. <https://doi.org/10.1172/JCI65910>
- Amarasekara, D. S., Kim, S., & Rho, J. (2021). Regulation of osteoblast differentiation by cytokine networks. *International Journal of Molecular Sciences*, 22(6), 2851. <https://doi.org/10.3390/ijms22062851>
- Ambrosi, T. H., Scialdone, A., Graja, A., Gohlke, S., Jank, A. M., Bocian, C., Woelk, L., Fan, H., Logan, D. W., Schürmann, A., Saraiva, L. R., & Schulz, T. J. (2017). Adipocyte Accumulation in the bone marrow during obesity and aging impairs stem cell-based hematopoietic and bone regeneration. *Cell Stem Cell*, 20(6), 771–784.e6. <https://doi.org/10.1016/j.stem.2017.02.009>
- Aurilia, C., Donati, S., Palmieri, G., Miglietta, F., Iantomasi, T., & Brandi, M. L. (2021). The involvement of long non-coding RNAs in bone. *International Journal of Molecular Sciences*, 22(8), 3909. <https://doi.org/10.3390/ijms22083909>
- Baglio, S. R., Devescovi, V., Granchi, D., & Baldini, N. (2013). MicroRNA expression profiling of human bone marrow mesenchymal stem cells during osteogenic differentiation reveals Osterix regulation by miR-31. *Gene*, 527(1), 321–331. <https://doi.org/10.1016/j.gene.2013.06.021>
- Balagandharan, K., Viji Chandran, S., Arumugam, B., Saravanan, S., Devanand Venkatasubbu, G., & Selvamurugan, N. (2018). Chitosan/nano-hydroxyapatite/nano-zirconium dioxide scaffolds with miR-590-5p for bone regeneration. *International Journal of Biological Macromolecules*, 111, 953–958. <https://doi.org/10.1016/j.ijbiomac.2018.01.122>
- Baldini, N., Cenni, E., Ciapetti, G., Granchi, D., & Savarino, L. (2009). Bone repair and regeneration. In J. A. Planell (Ed.), *Bone Repair Biomaterials* (pp. 69–105). Woodhead Publishing Limited. <https://doi.org/10.1533/9781845696610.1.69>
- Barrett, S. P., & Salzman, J. (2016). Circular RNAs: analysis, expression and potential functions. *Development (Cambridge, England)*, 143(11), 1838–1847. [10.1242/dev.128074](https://doi.org/10.1242/dev.128074)
- Bassi, A., Gough, J., Zakikhani, M., & Downes, S. (2011). Bone tissue regeneration. In L. A. Bosworth & S. Downes (Eds.), *Electrospinning for tissue regeneration* (pp. 93–110). Woodhead Publishing Limited. <https://doi.org/10.1533/9780857092915.2.93>
- Beermann, J., Piccoli, M. T., Viereck, J., & Thum, T. (2016). Non-coding RNAs in development and disease: Background, mechanisms, and therapeutic approaches. *Physiological Reviews*, 96(4), 1297–1325. <https://doi.org/10.1152/physrev.00041.2015>
- Berendsen, A. D., & Olsen, B. R. (2015). Bone development. *Bone*, 80, 14–18. [10.1016/j.bone.2015.04.035](https://doi.org/10.1016/j.bone.2015.04.035)
- Bialek, P., Kern, B., Yang, X., Schrock, M., Sosic, D., Hong, N., Wu, H., Yu, K., Ornitz, D. M., Olson, E. N., Justice, M. J., & Karsenty, G. (2004). A twist code determines the onset of osteoblast differentiation. *Developmental Cell*, 6(3), 423–435. [https://doi.org/10.1016/S1534-5807\(04\)00058-9](https://doi.org/10.1016/S1534-5807(04)00058-9)
- Bidwell, J. P., Van Wijnen, A. J., Fey, E. G., Dworetzky, S., Penman, S., Stein, J. L., Lian, J. B., & Stein, G. S. (1993). Osteocalcin gene promoter-binding factors are tissue-specific nuclear matrix components. *Proceedings of the National Academy of Sciences of the United States of America*, 90(8), 3162–3166. <https://doi.org/10.1073/pnas.90.8.3162>
- Cai, N., Li, C., & Wang, F. (2019). Silencing of lncRNA-ANCR promotes the osteogenesis of osteoblast cells in postmenopausal osteoporosis via targeting EZH2 and RUNX2. *Yonsei Medical Journal*, 60(8), 751–759. <https://doi.org/10.3349/ymj.2019.60.8.751>
- Camp, E., Pribadi, C., Anderson, P. J., Zannettino, A., & Gronthos, S. (2018). miRNA-376c-3p mediates TWIST-1 inhibition of bone marrow-derived stromal cell osteogenesis and can reduce aberrant bone formation of TWIST-1 haploinsufficient calvarial cells. *Stem Cells and Development*, 27(23), 1621–1633. <https://doi.org/10.1089/scd.2018.0083>
- Capulli, M., Paone, R., & Rucci, N. (2014). Osteoblast and osteocyte: Games without frontiers. *Archives of Biochemistry and Biophysics*, 561(May), 3–12. <https://doi.org/10.1016/j.abb.2014.05.003>
- Chan, W., Tan, Z., To, M., & Chan, D. (2021). Regulation and role of transcription factors in osteogenesis. *International Journal of Molecular Sciences*, 22(11), 5445. <https://doi.org/10.3390/ijms22115445>
- Chen, G., Deng, C., & Li, Y.-P. (2012). TGF- β and BMP signaling in osteoblast differentiation and bone formation. *International Journal of Biological Sciences*, 8(2), 272–288. [10.7150/ijbs.2929](https://doi.org/10.7150/ijbs.2929)
- Chen, F., Bi, D., Cheng, C., Ma, S., Liu, Y., & Cheng, K. (2019). Bone morphogenetic protein 7 enhances the osteogenic differentiation of human dermal-derived CD105+ fibroblast cells through the Smad and MAPK pathways. *International Journal of Molecular Medicine*, 43(1), 37–46. <https://doi.org/10.3892/ijmm.2018.3938>
- Chen, G., Long, C., Wang, S., Wang, Z., Chen, X., Tang, W., He, X., Bao, Z., Tan, B., Zhao, J., Xie, Y., Li, Z., Yang, D., Xiao, G., & Peng, S. (2022). Circular RNA circStag1 promotes bone regeneration by interacting with HuR. *Bone Research*, 10(1), 32. <https://doi.org/10.1038/s41413-022-00208-x>
- Chen, G., Tang, W., Wang, S., Long, C., He, X., Yang, D., & Peng, S. (2021). Promising diagnostic and therapeutic circRNAs for skeletal and chondral disorders. *International Journal of Biological Sciences*, 17(5), 1428–1439. <https://doi.org/10.7150/ijbs.57887>
- Chen, S., Li, Y., Zhi, S., Ding, Z., Huang, Y., Wang, W., Zheng, R., Yu, H., Wang, J., Hu, M., Miao, J., & Li, J. (2020). lncRNA Xist regulates osteoblast differentiation by sponging miR-19a-3p in aging-induced osteoporosis. *Aging and Disease*, 11(5), 1058–1068. <https://doi.org/10.14336/AD.2019.0724>
- Cuevas-González, M. V., Suaste-Olmos, F., García-Calderón, A. G., Tovar-Carrillo, K. L., Espinosa-Cristóbal, L. F., Nava-Martínez, S. D., Cuevas-González, J. C., Zambrano-Galván, G., Saucedo-Acuña, R. A., &

- Donohue-Cornejo, A. (2021). Expression of MicroRNAs in periodontal disease: A systematic review. *BioMed Research International*, 2021, 2069410. <https://doi.org/10.1155/2021/2069410>
- Damiati, L. A., & El-Messeiry, S. (2021). An overview of RNA-based scaffolds for osteogenesis. *Frontiers in Molecular Biosciences*, 8, 682581. <https://doi.org/10.3389/fmolb.2021.682581>
- Danciu, T. E., Li, Y., Koh, A., Xiao, G., McCauley, L. K., & Franceschi, R. T. (2012). The basic helix loop helix transcription factor Twist1 is a novel regulator of ATF4 in osteoblasts. *Journal of Cellular Biochemistry*, 113(1), 70–79. <https://doi.org/10.1002/jcb.23329>
- Davis, B. N., & Hata, A. (2009). Regulation of MicroRNA Biogenesis: A miRiad of mechanisms. *Cell Communication and Signaling*, 7(1), 18. <https://doi.org/10.1186/1478-811X-7-18>
- Day, T. F., Guo, X., Garrett-Beal, L., & Yang, Y. (2005). Wnt/beta-catenin signaling in mesenchymal progenitors controls osteoblast and chondrocyte differentiation during vertebrate skeletogenesis. *Developmental Cell*, 8(5), 739–750. <https://doi.org/10.1016/j.devcel.2005.03.016>
- Deng, A., Zhang, H., Hu, M., Liu, S., Gao, Q., Wang, Y., & Guo, C. (2017). Knockdown of Indian hedgehog protein induces an inhibition of cell growth and differentiation in osteoblast MC3T3-E1 cells. *Molecular Medicine Reports*, 16(6), 7987–7992. <https://doi.org/10.3892/mmr.2017.7669>
- Deng, Y., Wu, S., Zhou, H., Bi, X., Wang, Y., Hu, Y., Gu, P., & Fan, X. (2013). Effects of a miR-31, Runx2, and Satb2 regulatory loop on the osteogenic differentiation of bone mesenchymal stem cells. *Stem Cells and Development*, 22(16), 2278–2286. <https://doi.org/10.1089/scd.2012.0686>
- Dobrev, G., Chahrouh, M., Dautzenberg, M., Chirivella, L., Kanzler, B., Fariñas, I., Karsenty, G., & Grosschedl, R. (2006). SATB2 is a multifunctional determinant of craniofacial patterning and osteoblast differentiation. *Cell*, 125(5), 971–986. <https://doi.org/10.1016/j.cell.2006.05.012>
- Ducy, P., & Karsenty, G. (1995). Two distinct osteoblast-specific cis-acting elements control expression of a mouse osteocalcin gene. *Molecular and Cellular Biology*, 15(4), 1858–1869. <https://doi.org/10.1128/MCB.15.4.1858>
- El-Ganzuri, M. A., Ahmed, R. R., & Bastawy, E. M. (2016). Regulatory Mechanisms of Bone Development and Function. *Annals of Cytology and Pathology*, 1(1), 005–017. <https://doi.org/10.17352/acp.000002>
- Erceg, I., Tadić, T., Kronenberger, M. S., Marijanović, I., & Lichtler, A. C. (2003). Dlx5 regulation of mouse osteoblast differentiation mediated by avian retrovirus vector. *Croatian Medical Journal*, 44(4), 407–411.
- FitzPatrick, D. R., Carr, I. M., McLaren, L., Leek, J. P., Wightman, P., Williamson, K., Gautier, P., McGill, N., Hayward, C., Firth, H., Markham, A. F., Fantes, J. A., & Bonthron, D. T. (2003). Identification of SATB2 as the cleft palate gene on 2q32-q33. *Human Molecular Genetics*, 12(19), 2491–2501. <https://doi.org/10.1093/hmg/ddg248>
- Friedenstein, A. J., Chailakhjan, R. K., & Lalykina, K. S. (1970). The development of fibroblast colonies in monolayer cultures of guinea-pig bone marrow and spleen cells. *Cell and Tissue Kinetics*, 3(4), 393–403. <https://doi.org/10.1111/j.1365-2184.1970.tb00347.x>
- Gámez, B., Rodríguez-Carballo, E., Bartrons, R., Rosa, J. L., & Ventura, F. (2013). MicroRNA-322 (miR-322) and its target protein Tob2 modulate Osterix (Ox) mRNA stability. *The Journal of Biological Chemistry*, 288(20), 14264–14275. <https://doi.org/10.1074/jbc.M112.432104>
- Gaur, T., Lengner, C. J., Hovhannisyann, H., Bhat, R. A., Bodine, P. V., Komm, B. S., Javed, A., van Wijnen, A. J., Stein, J. L., Stein, G. S., & Lian, J. B. (2005). Canonical WNT signaling promotes osteogenesis by directly stimulating Runx2 gene expression. *The Journal of Biological Chemistry*, 280(39), 33132–33140. <https://doi.org/10.1074/jbc.M500608200>
- Ge, X., Li, Z., Zhou, Z., Xia, Y., Bian, M., & Yu, J. (2020). Circular RNA SIPA1L1 promotes osteogenesis via regulating the miR-617/Smad3 axis in dental pulp stem cells. *Stem Cell Research & Therapy*, 11(1), 364. <https://doi.org/10.1186/s13287-020-01877-3>
- Gennari, L., Rendina, D., Falchetti, A., & Merlotti, D. (2019). Paget's Disease of Bone. *Calcified Tissue International*, 104(5), 483–500. <https://doi.org/10.1007/s00223-019-00522-3>
- Gibon, E., Batke, B., Jawad, M. U., Fritton, K., Rao, A., Yao, Z., Biswal, S., Gambhir, S. S., & Goodman, S. B. (2012). MC3T3-E1 osteoprogenitor cells systemically migrate to a bone defect and enhance bone healing. *Tissue Engineering. Part A*, 18(9-10), 968–973. <https://doi.org/10.1089/ten.TEA.2011.0545>
- Gong, Y., Lu, J., Yu, X., & Yu, Y. (2016). Expression of Sp7 in Satb2-induced osteogenic differentiation of mouse bone marrow stromal cells is regulated by microRNA-27a. *Molecular and Cellular Biochemistry*, 417(1–2), 7–16. <https://doi.org/10.1007/s11010-016-2709-y>
- Gordon, J. A., Tye, C. E., Sampaio, A. V., Underhill, T. M., Hunter, G. K., & Goldberg, H. A. (2007). Bone sialoprotein expression enhances osteoblast differentiation and matrix mineralization in vitro. *Bone*, 41(3), 462–473. <https://doi.org/10.1016/j.bone.2007.04.191>
- Guan, S., Zhang, Z., & Wu, J. (2022). Non-coding RNA delivery for bone tissue engineering: Progress, challenges, and potential solutions. *iScience*, 25(8), 104807. <https://doi.org/10.1016/j.isci.2022.104807>
- Guenou, H., Kaabeche, K., Mée, S. L., & Marie, P. J. (2005). A role for fibroblast growth factor receptor-2 in the altered osteoblast phenotype induced by Twist haploinsufficiency in the Saethre-Chotzen syndrome. *Human Molecular Genetics*, 14(11), 1429–1439. <https://doi.org/10.1093/hmg/ddi152>
- Han, S., Kuang, M., Sun, C., Wang, H., Wang, D., & Liu, Q. (2020). Circular RNA hsa_circ_0076690 acts as a prognostic biomarker in osteoporosis and regulates osteogenic differentiation of hBMSCs via sponging miR-152. *Aging*, 12(14), 15011–15020. <https://doi.org/10.18632/aging.103560>
- Hassan, M. Q., Gordon, J. A., Beloti, M. M., Croce, C. M., van Wijnen, A. J., Stein, J. L., Stein, G. S., & Lian, J. B. (2010). A network connecting Runx2, SATB2, and the miR-23a~27a~24-2 cluster regulates the osteoblast differentiation program. *Proceedings of the National Academy of Sciences of the United States of America*, 107(46), 19879–19884. <https://doi.org/10.1073/pnas.1007698107>
- He, S., Yang, S., Zhang, Y., Li, X., Gao, D., Zhong, Y., Cao, L., Ma, H., Liu, Y., Li, G., Peng, S., & Shuai, C. (2019). LncRNA ODIR1 inhibits osteogenic differentiation of hUC-MSCs through the FBXO25/H2BK120ub/H3K4me3/OSX axis. *Cell Death & Disease*, 10(12), 947. <https://doi.org/10.1038/s41419-019-2148-2>
- He, W., Shi, X., Guo, Z., Wang, H., Kang, M., & Lv, Z. (2022). Circ_0019693 promotes osteogenic differentiation of bone marrow mesenchymal stem cell and enhances osteogenesis-coupled angiogenesis via regulating microRNA-942-5p-targeted purkinje cell protein 4 in the development of osteoporosis. *Bioengineered*, 13(2), 2181–2193. <https://doi.org/10.1080/21655979.2021.2023982>
- Hill, T. P., Später, D., Taketo, M. M., Birchmeier, W., & Hartmann, C. (2005). Canonical Wnt/beta-catenin signaling prevents osteoblasts from differentiating into chondrocytes. *Developmental Cell*, 8(5), 727–738. <https://doi.org/10.1016/j.devcel.2005.02.013>
- Hilton, M. J., Tu, X., Cook, J., Hu, H., & Long, F. (2005). Ihh controls cartilage development by antagonizing Gli3, but requires additional effectors to regulate osteoblast and vascular development. *Development (Cambridge, England)*, 132(19), 4339–4351. <https://doi.org/10.1242/dev.02025>
- Holleville, N., Matéos, S., Bontoux, M., Bollerot, K., & Monsoro-Burq, A. (2007). Dlx5 drives Runx2 expression and osteogenic differentiation in developing cranial suture mesenchyme. *Developmental Biology*, 304(2), 860–874. <https://doi.org/10.1016/j.ydbio.2007.01.003>
- Hu, H., Hilton, M. J., Tu, X., Yu, K., Ornitz, D. M., & Long, F. (2005). Sequential roles of Hedgehog and Wnt signaling in osteoblast development. *Development (Cambridge, England)*, 132(1), 49–60. <https://doi.org/10.1242/dev.01564>
- Hu, H., Zhao, C., Zhang, P., Liu, Y., Jiang, Y., Wu, E., Xue, H., Liu, C., & Li, Z. (2019). miR-26b modulates OA induced BMSC osteogenesis through regulating GSK3β/β-catenin pathway. *Experimental and Molecular Pathology*, 107, 158–164. <https://doi.org/10.1016/j.yexmp.2019.02.003>
- Hu, L., Liu, J., Xue, H., Panayi, A. C., Xie, X., Lin, Z., Wang, T., Xiong, Y., Hu, Y., Yan, C., Chen, L., Abududilibaier, A., Zhou, W., Mi, B., & Liu, G. (2021). miRNA-92a-3p regulates osteoblast differentiation in patients with concomitant limb fractures and TBI via IBS/Pi3K-AKT inhibition. *Molecular Therapy. Nucleic Acids*, 23, 1345–1359. <https://doi.org/10.1016/j.omtn.2021.02.008>

- Hu, L., Yin, C., Zhao, F., Ali, A., Ma, J., & Qian, A. (2018). Mesenchymal stem cells: Cell fate decision to osteoblast or adipocyte and application in osteoporosis treatment. *International Journal of Molecular Sciences*, 19(2), 360. <https://doi.org/10.3390/ijms19020360>
- Huang, W., Yang, S., Shao, J., & Li, Y. P. (2007). Signaling and transcriptional regulation in osteoblast commitment and differentiation. *Frontiers in Bioscience*, 12(8–12), 3068–3092. <https://doi.org/10.2741/2296>
- Huang, Y., Meng, T., Wang, S., Zhang, H., Mues, G., Qin, C., Feng, J. Q., D'Souza, R. N., & Lu, Y. (2014). Twist1- and Twist2-haploinsufficiency results in reduced bone formation. *PLoS One*, 9(6), e99331. <https://doi.org/10.1371/journal.pone.0099331>
- Huang, Y., Xiao, D., Huang, S., Zhuang, J., Zheng, X., Chang, Y., & Yin, D. (2020). Circular RNA YAP1 attenuates osteoporosis through up-regulation of YAP1 and activation of Wnt/ β -catenin pathway. *Biomedicine & Pharmacotherapy = Biomedicine & Pharmacotherapie*, 129, 110365. <https://doi.org/10.1016/j.biopha.2020.110365>
- Huang, Y., Zheng, Y., Jia, L., & Li, W. (2015). Long noncoding RNA H19 promotes osteoblast differentiation via TGF- β 1/Smad3/HDAC signaling pathway by deriving miR-675. *Stem Cells (Dayton, Ohio)*, 33(12), 3481–3492. <https://doi.org/10.1002/stem.2225>
- Iaquinta, M. R., Mazzoni, E., Bononi, I., Rotondo, J. C., Mazziotta, C., Montesi, M., Sprio, S., Tampieri, A., Tognoni, M., & Martini, F. (2019). Adult stem cells for bone regeneration and repair. *Frontiers in Cell and Developmental Biology*, 7, 268. <https://doi.org/10.3389/fcell.2019.00268>
- Ibrahim, A., Bulstrode, N. W., Whitaker, I. S., Eastwood, D. M., Dunaway, D., & Ferretti, P. (2016). Nanotechnology for stimulating osteoprogenitor differentiation. *The Open Orthopaedics Journal*, 10, 849–861. <https://doi.org/10.2174/1874325001610010849>
- Idolazzi, L., Fassio, A., Tripi, G., Braga, V., Viapiana, O., Adami, G., Rossini, M., & Gatti, D. (2017). Circulating Dickkopf-1 and sclerostin in patients with Paget's disease of bone. *Clinical Rheumatology*, 36(4), 925–928. <https://doi.org/10.1007/s10067-016-3497-1>
- Inose, H., Ochi, H., Kimura, A., Fujita, K., Xu, R., Sato, S., Iwasaki, M., Sunamura, S., Takeuchi, Y., Fukumoto, S., Saito, K., Nakamura, T., Siomi, H., Ito, H., Arai, Y., Shinomiya, K., & Takeda, S. (2009). A microRNA regulatory mechanism of osteoblast differentiation. *Proceedings of the National Academy of Sciences of the United States of America*, 106(49), 20794–20799. <https://doi.org/10.1073/pnas.0909311106>
- Ji, F., Pan, J., Shen, Z., Yang, Z., Wang, J., Bai, X., & Tao, J. (2020). The circular RNA circRNA124534 promotes osteogenic differentiation of human dental pulp stem cells through modulation of the miR-496/ β -catenin pathway. *Frontiers in Cell and Developmental Biology*, 8, 230. <https://doi.org/10.3389/fcell.2020.00230>
- Jiajun, Z., Wei, Q., Lijuan, H., Jing, R., Chunhui, L., Pairan, P., Jie, H., & Yandong, M. (2020). miRNA expression profiling of osteogenic differentiation of bone marrow mesenchymal stem cells induced by microchannel porous hydroxyapatite scaffold. *Chinese Journal of Tissue Engineering Research*, 24(13), 1989–1995. <https://doi.org/10.3969/j.issn.2095-4344.2053>
- Jonason, J. H., Xiao, G., Zhang, M., Xing, L., & Chen, D. (2009). Post-translational regulation of Runx2 in bone and cartilage. *Journal of Dental Research*, 88(8), 693–703. <https://doi.org/10.1177/0022034509341629>
- Jones, D. C., Wein, M. N., Oukka, M., Hofstaetter, J. G., Glimcher, M. J., & Glimcher, L. H. (2006). Regulation of adult bone mass by the zinc finger adapter protein Schnurri-3. *Science (New York, N.Y.)*, 312(5777), 1223–1227. <https://doi.org/10.1126/science.1126313>
- Kang, S., Bennett, C. N., Gerin, I., Rapp, L. A., Hankenson, K. D., & Macdougald, O. A. (2007). Wnt signaling stimulates osteoblastogenesis of mesenchymal precursors by suppressing CCAAT/enhancer-binding protein alpha and peroxisome proliferator-activated receptor gamma. *The Journal of Biological Chemistry*, 282(19), 14515–14524. <https://doi.org/10.1074/jbc.M700030200>
- Kapinas, K., Kessler, C., Ricks, T., Gronowicz, G., & Delany, A. M. (2010). miR-29 modulates Wnt signaling in human osteoblasts through a positive feedback loop. *The Journal of Biological Chemistry*, 285(33), 25221–25231. <https://doi.org/10.1074/jbc.M110.116137>
- Kawane, T., Qin, X., Jiang, Q., Miyazaki, T., Komori, H., Yoshida, C. A., Matsuura-Kawata, V., Sakane, C., Matsuo, Y., Nagai, K., Maeno, T., Date, Y., Nishimura, R., & Komori, T. (2018). Runx2 is required for the proliferation of osteoblast progenitors and induces proliferation by regulating Fgfr2 and Fgfr3. *Scientific Reports*, 8(1), 13551. <https://doi.org/10.1038/s41598-018-31853-0>
- Khotib, J., Gani, M. A., Budiati, A. S., Lestari, M., Rahadiansyah, E., & Ardianto, C. (2021). Signaling pathway and transcriptional regulation in osteoblasts during bone healing: Direct involvement of hydroxyapatite as a biomaterial. *Pharmaceuticals (Basel, Switzerland)*, 14(7), 615. <https://doi.org/10.3390/ph14070615>
- Kim, H. K., Lee, J. S., Kim, J. H., Seon, J. K., Park, K. S., Jeong, M. H., & Yoon, T. R. (2017). Bone-forming peptide-2 derived from BMP-7 enhances osteoblast differentiation from multipotent bone marrow stromal cells and bone formation. *Experimental & Molecular Medicine*, 49(5), e328. <https://doi.org/10.1038/emm.2017.40>
- Kim, J. H., Liu, X., Wang, J., Chen, X., Zhang, H., Kim, S. H., Cui, J., Li, R., Zhang, W., Kong, Y., Zhang, J., Shui, W., Lamplot, J., Rogers, M. R., Zhao, C., Wang, N., Rajan, P., Tomal, J., Statz, J., ... He, T.-C. (2013). Wnt signaling in bone formation and its therapeutic potential for bone diseases. *Therapeutic Advances in Musculoskeletal Disease*, 5(1), 13–31. <https://doi.org/10.1177/1759720X12466608>
- Kim, J. M., Lin, C., Stavre, Z., Greenblatt, M. B., & Shim, J. H. (2020). Osteoblast-osteoclast communication and bone homeostasis. *Cells*, 9(9), 2073. <https://doi.org/10.3390/cells9092073>
- Kim, R. H., Shapiro, H. S., Li, J. J., Wrana, J. L., & Sodek, J. (1994). Characterization of the human bone sialoprotein (BSP) gene and its promoter sequence. *Matrix Biology*, 14(1), 31–40. [https://doi.org/10.1016/0945-053X\(94\)90027-2](https://doi.org/10.1016/0945-053X(94)90027-2)
- Knight, M. N., & Hankenson, K. D. (2013). Mesenchymal stem cells in bone regeneration. *Advances in Wound Care*, 2(6), 306–316. <https://doi.org/10.1089/wound.2012.0420>
- Kobayashi, Y., Maeda, K., & Takahashi, N. (2008). Roles of Wnt signaling in bone formation and resorption. *Japanese Dental Science Review*, 44(1), 76–82. <https://doi.org/10.1016/j.jdsr.2007.11.002>
- Komaki, M., Karakida, T., Abe, M., Oida, S., Mimori, K., Iwasaki, K., Noguchi, K., Oda, S., & Ishikawa, I. (2007). Twist negatively regulates osteoblastic differentiation in human periodontal ligament cells. *Journal of Cellular Biochemistry*, 100(2), 303–314. <https://doi.org/10.1002/jcb.21038>
- Komori, T. (2019). Regulation of proliferation, differentiation and functions of osteoblasts by Runx2. *International Journal of Molecular Sciences*, 20(7), 1694. <https://doi.org/10.3390/ijms20071694>
- Kureel, J., John, A. A., Dixit, M., & Singh, D. (2017). MicroRNA-467g inhibits new bone regeneration by targeting Ihh/Runx-2 signaling. *The International Journal of Biochemistry & Cell Biology*, 85, 35–43. <https://doi.org/10.1016/j.biocel.2017.01.018>
- Lavery, K., Hawley, S., Swain, P., Rooney, R., Falb, D., & Alaoui-Ismaili, M. H. (2009). New insights into BMP-7 mediated osteoblastic differentiation of primary human mesenchymal stem cells. *Bone*, 45(1), 27–41. <https://doi.org/10.1016/j.bone.2009.03.656>
- Laxman, N., Mallin, H., Nilsson, O., & Kindmark, A. (2016). miR-203 and miR-320 Regulate Bone Morphogenetic Protein-2-Induced Osteoblast Differentiation by Targeting Distal-Less Homeobox 5 (Dlx5). *Genes*, 8(1), 4. <https://doi.org/10.3390/genes8010004>
- Lee, M. H., Kim, Y. J., Kim, H. J., Park, H. D., Kang, A. R., Kyung, H. M., Sung, J. H., Wozney, J. M., Kim, H. J., & Ryoo, H. M. (2003a). BMP-2-induced Runx2 expression is mediated by Dlx5, and TGF-beta 1 opposes the BMP-2-induced osteoblast differentiation by suppression of Dlx5 expression. *The Journal of Biological Chemistry*, 278(36), 34387–34394. <https://doi.org/10.1074/jbc.M211386200>
- Lee, M. H., Kwon, T. G., Park, H. S., Wozney, J. M., & Ryoo, H. M. (2003b). BMP-2-induced Osterix expression is mediated by Dlx5 but is independent of Runx2. *Biochemical and Biophysical Research Communications*, 309(3), 689–694. <https://doi.org/10.1016/j.bbrc.2003.08.058>
- Lee, M. S., Lowe, G. N., Strong, D. D., Wergedal, J. E., & Glackin, C. A. (1999). TWIST, a basic helix-loop-helix transcription factor, can regulate the human osteogenic lineage. *Journal of Cellular Biochemistry*,

- 75(4), 566–577. [https://doi.org/10.1002/\(sici\)1097-4644\(19991215\)75:4<566::aid-jcb3>3.0.co;2-0](https://doi.org/10.1002/(sici)1097-4644(19991215)75:4<566::aid-jcb3>3.0.co;2-0)
- Lekka, E., & Hall, J. (2018). Noncoding RNAs in disease. *FEBS Letters*, 592(17), 2884–2900. [10.1002/1873-3468.13182](https://doi.org/10.1002/1873-3468.13182) 29972883
- Li, D., Tian, Y., Yin, C., Huai, Y., Zhao, Y., Su, P., Wang, X., Pei, J., Zhang, K., Yang, C., Dang, K., Jiang, S., Miao, Z., Li, M., Hao, Q., Zhang, G., & Qian, A. (2019a). Silencing of lncRNA AK045490 promotes osteoblast differentiation and bone formation via β -catenin/TCF1/Runx2 signaling axis. *International Journal of Molecular Sciences*, 20(24), 6229. <https://doi.org/10.3390/ijms20246229>
- Li, H., Zheng, Q., Xie, X., Wang, J., Zhu, H., Hu, H., He, H., & Lu, Q. (2021). Role of exosomal non-coding RNAs in bone-related diseases. *Frontiers in Cell and Developmental Biology*, 9, 811666. <https://doi.org/10.3389/fcell.2021.811666>
- Li, J., Zhang, H., Yang, C., Li, Y., & Dai, Z. (2016). An overview of osteocalcin progress. *Journal of Bone and Mineral Metabolism*, 34(4), 367–379. <https://doi.org/10.1007/s00774-015-0734-7>
- Li, X. G., Liu, S. C., Qiao, X. F., Kong, Y., Liu, J. G., Peng, X. M., Wang, Y. X., & Abdulkarim Mohammed Al-Mohana, R. A. (2019b). LncRNA MEG3 promotes proliferation and differentiation of osteoblasts through Wnt/ β -catenin signaling pathway. *European Review for Medical and Pharmacological Sciences*, 23(11), 4521–4529. https://doi.org/10.26355/eurev_201906_18027
- Li, Y., Shan, G., Teng, Z. Q., & Wingo, T. S. (2020). Editorial: Non-coding RNAs and human diseases. *Frontiers in Genetics*, 11, 523. <https://doi.org/10.3389/fgene.2020.00523>
- Liang, W. C., Fu, W. M., Wang, Y. B., Sun, Y. X., Xu, L. L., Wong, C. W., Chan, K. M., Li, G., Waye, M. M., & Zhang, J. F. (2016). H19 activates Wnt signaling and promotes osteoblast differentiation by functioning as a competing endogenous RNA. *Scientific Reports*, 6, 20121. <https://doi.org/10.1038/srep20121>
- Lin, X., Patil, S., Gao, Y. G., & Qian, A. (2020). The bone extracellular matrix in bone formation and regeneration. *Frontiers in Pharmacology*, 11, 757. <https://doi.org/10.3389/fphar.2020.00757>
- Liu, Q., Li, M., Wang, S., Xiao, Z., Xiong, Y., & Wang, G. (2020). Recent advances of Osterix transcription factor in osteoblast differentiation and bone formation. *Frontiers in Cell and Developmental Biology*, 8, 601224. <https://doi.org/10.3389/fcell.2020.601224>
- Liu, S. C., Sun, Q. Z., Qiao, X. F., Li, X. G., Yang, J. H., Wang, T. Q., Xiao, Y. J., & Qiao, J. M. (2019). LncRNA TUG1 influences osteoblast proliferation and differentiation through the Wnt/ β -catenin signaling pathway. *European Review for Medical and Pharmacological Sciences*, 23(11), 4584–4590. https://doi.org/10.26355/eurev_201906_18035
- Liu, T. M., & Lee, E. H. (2013). Transcriptional regulatory cascades in Runx2-dependent bone development. *Tissue Engineering, Part B, Reviews*, 19(3), 254–263. <https://doi.org/10.1089/ten.TEB.2012.0527>
- Long, F., Chung, U.-i., Ohba, S., McMahon, J., Kronenberg, H. M., & McMahon, A. P. (2004). Ihh signaling is directly required for the osteoblast lineage in the endochondral skeleton. *Development (Cambridge, England)*, 131(6), 1309–1318. <https://doi.org/10.1242/dev.01006>
- Lv, H., Yang, H., & Wang, Y. (2020). Effects of miR-103 by negatively regulating SATB2 on proliferation and osteogenic differentiation of human bone marrow mesenchymal stem cells. *PLoS One*, 15(5), e0232695. <https://doi.org/10.1371/journal.pone.0232695>
- Manolagas, S. C. (2000). Birth and death of bone cells: basic regulatory mechanisms and implications for the pathogenesis and treatment of osteoporosis. *Endocrine Reviews*, 21(2), 115–137. [10.1210/edrv.21.2.0395](https://doi.org/10.1210/edrv.21.2.0395) 10782361
- Marshall, M. J., Evans, S. F., Sharp, C. A., Powell, D. E., McCarthy, H. S., & Davie, M. W. (2009). Increased circulating Dickkopf-1 in Paget's disease of bone. *Clinical Biochemistry*, 42(10–11), 965–969. <https://doi.org/10.1016/j.clinbiochem.2009.04.007>
- Martin, A., Liu, S., David, V., Li, H., Karydis, A., Feng, J. Q., & Quarles, L. D. (2011). Bone proteins PHEX and DMP1 regulate fibroblastic growth factor Fgf23 expression in osteocytes through a common pathway involving FGF receptor (FGFR) signaling. *The FASEB Journal*, 25(8), 2551–2562. <https://doi.org/10.1096/fj.10-177816>
- Matthews, B. G., Grcevic, D., Wang, L., Hagiwara, Y., Roguljic, H., Joshi, P., Shin, D. G., Adams, D. J., & Kalajzic, I. (2014). Analysis of α SMA-labeled progenitor cell commitment identifies notch signaling as an important pathway in fracture healing. *Journal of Bone and Mineral Research*, 29(5), 1283–1294. <https://doi.org/10.1002/jbmr.2140>
- Matthews, B. G., Wee, N., Widjaja, V. N., Price, J. S., Kalajzic, I., & Windahl, S. H. (2020). α SMA osteoprogenitor cells contribute to the increase in osteoblast numbers in response to mechanical loading. *Calcified Tissue International*, 106(2), 208–217. <https://doi.org/10.1007/s00223-019-00624-y>
- Mi, B., Xiong, Y., Chen, L., Yan, C., Endo, Y., Liu, Y., Liu, J., Hu, L., Hu, Y., Sun, Y., Cao, F., Zhou, W., & Liu, G. (2019). CircRNA AFF4 promotes osteoblast cells proliferation and inhibits apoptosis via the Mir-7223-5p/PIK3R1 axis. *Aging*, 11(24), 11988–12001. <https://doi.org/10.18632/aging.102524>
- Michou, L., & Orcel, P. (2019). Has Paget's bone disease become rare? *Joint Bone Spine*, 86(5), 538–541. <https://doi.org/10.1016/j.jbspin.2019.01.015>
- Miraoui, H., Severe, N., Vaudin, P., Pagès, J. C., & Marie, P. J. (2010). Molecular silencing of Twist1 enhances osteogenic differentiation of murine mesenchymal stem cells: Implication of FGFR2 signaling. *Journal of Cellular Biochemistry*, 110(5), 1147–1154. <https://doi.org/10.1002/jcb.22628>
- Mouillé, M., Rio, M., Breton, S., Piketty, M. L., Afenjar, A., Amiel, J., Capri, Y., Goldenberg, A., Francannet, C., Michot, C., Mignot, C., Perrin, L., Quelin, C., Van Gils, J., Barcia, G., Pingault, V., Maruani, G., Koumakis, E., & Comier-Daire, V. (2022). SATB2-associated syndrome: Characterization of skeletal features and of bone fragility in a prospective cohort of 19 patients. *Orphanet Journal of Rare Diseases*, 17(1), 100. <https://doi.org/10.1186/s13023-022-02229-5>
- Nahian, A., & Davis, D. D. (2021). Histology, osteoprogenitor cells. In: *StatPearls [Internet]*. StatPearls Publishing. <https://www.ncbi.nlm.nih.gov/books/NBK559160/>
- Nakamura, T., Aikawa, T., Iwamoto-Enomoto, M., Iwamoto, M., Higuchi, Y., Pacifici, M., Kinto, N., Yamaguchi, A., Noji, S., Kurisu, K., Matsuya, T., & Maurizio, P. (1997). Induction of osteogenic differentiation by hedgehog proteins. *Biochemical and Biophysical Research Communications*, 237(2), 465–469. <https://doi.org/10.1006/bbrc.1997.7156>
- Nakashima, K., & De Crombrugge, B. (2003). Transcriptional mechanisms in osteoblast differentiation and bone formation. *Trends in Genetics*, 19(8), 458–466. [https://doi.org/10.1016/S0168-9525\(03\)00176-8](https://doi.org/10.1016/S0168-9525(03)00176-8)
- Nardocci, G., Carrasco, M. E., Acevedo, E., Hodar, C., Meneses, C., & Montecino, M. (2018). Identification of a novel long noncoding RNA that promotes osteoblast differentiation. *Journal of Cellular Biochemistry*, 119(9), 7657–7666. <https://doi.org/10.1002/jcb.27113> 29806713
- Nebot Valenzuela, E., & Pietschmann, P. (2017). Epidemiology and pathology of Paget's disease of bone- a review. *Wiener Medizinische Wochenschrift (1946)*, 167(1-2), 2–8. <https://doi.org/10.1007/s10354-016-0496-4> 27600564
- Nemoto, E., Ebe, Y., Kanaya, S., Tsuchiya, M., Nakamura, T., Tamura, M., & Shimauchi, H. (2012). Wnt5a signaling is a substantial constituent in bone morphogenetic protein-2-mediated osteoblastogenesis. *Biochemical and Biophysical Research Communications*, 422(4), 627–632. <https://doi.org/10.1016/j.bbrc.2012.05.039>
- Nohno, T., Ishikawa, T., Saito, T., Hosokawa, K., Noji, S., Wolsing, D. H., & Rosenbaum, J. S. (1995). Identification of a human type II receptor for bone morphogenetic protein-4 that forms differential heteromeric complexes with bone morphogenetic protein type I receptors. *The Journal of Biological Chemistry*, 270(38), 22522–22526. <https://doi.org/10.1074/jbc.270.38.22522> 7673243
- Ogasawara, T., Kawaguchi, H., Jinno, S., Hoshi, K., Itaka, K., Takato, T., Nakamura, K., & Okayama, H. (2004). Bone morphogenetic protein 2-induced osteoblast differentiation requires Smad-mediated down-regulation of Cdk6. *Molecular and Cellular Biology*, 24(15), 6560–6568. <https://doi.org/10.1128/MCB.24.15.6560-6568.2004>
- Okamoto, M., Udagawa, N., Uehara, S., Maeda, K., Yamashita, T., Nakamichi, Y., Kato, H., Saito, N., Minami, Y., Takahashi, N., & Kobayashi, Y. (2014). Noncanonical Wnt5a enhances Wnt/ β -catenin signaling during osteoblastogenesis. *Scientific Reports*, 4, 4493. <https://doi.org/10.1038/srep04493>

- Omoteyama, K., & Takagi, M. (2010). The effects of Sp7/Osterix gene silencing in the chondroprogenitor cell line, ATDC5. *Biochemical and Biophysical Research Communications*, 403(2), 242–246. <https://doi.org/10.1016/j.bbrc.2010.11.023>
- Ouyang, Z., Tan, T., Zhang, X., Wan, J., Zhou, Y., Jiang, G., Yang, D., Guo, X., & Liu, T. (2019). CircRNA hsa_circ_0074834 promotes the osteogenesis-angiogenesis coupling process in bone mesenchymal stem cells (BMSCs) by acting as a ceRNA for miR-942-5p. *Cell Death & Disease*, 10(12), 932. <https://doi.org/10.1038/s41419-019-2161-5>
- Pan, T., Song, W., Xin, H., Yu, H., Wang, H., Ma, D., Cao, X., & Wang, Y. (2021). MicroRNA-activated hydrogel scaffold generated by 3D printing accelerates bone regeneration. *Bioactive Materials*, 10, 1–14. <https://doi.org/10.1016/j.bioactmat.2021.08.034>
- Patil, S., Dang, K., Zhao, X., Gao, Y., & Qian, A. (2020). Role of LncRNAs and CircRNAs in bone metabolism and osteoporosis. *Frontiers in Genetics*, 11, 584118. <https://doi.org/10.3389/fgene.2020.584118>
- Pino, A. M., Rosen, C. J., & Rodríguez, J. P. (2012). In Osteoporosis, differentiation of mesenchymal stem cells (MSCs) improves bone marrow adipogenesis. *Biological Research*, 45(3), 279–287. <https://doi.org/10.4067/s0716-97602012000300009>
- Ponzetti, M., & Rucci, N. (2021). Osteoblast differentiation and signaling: Established concepts and emerging topics. *International Journal of Molecular Sciences*, 22(13), 6651. <https://doi.org/10.3390/ijms22136651>
- Qiu, Z.-Y., Cui, Y., & Wang, X.-M. (2019). Natural bone tissue and its biomimetic. In X.-M. Wang, Z.-Y. Qiu, & H. Cui (Eds.), *Mineralized collagen bone graft substitutes* (pp. 1–22). Woodhead Publishing Limited. <https://doi.org/10.1016/b978-0-08-102717-2.00001-1>
- Quarto, N., Senarath-Yapa, K., Renda, A., & Longaker, M. T. (2015). TWIST1 silencing enhances in vitro and in vivo osteogenic differentiation of human adipose-derived stem cells by triggering activation of BMP-ERK/FGF signaling and TAZ upregulation. *Stem Cells (Dayton, Ohio)*, 33(3), 833–847. <https://doi.org/10.1002/stem.1907>
- Rosenzweig, B. L., Imamura, T., Okadome, T., Cox, G. N., Yamashita, H., Ten Dijke, P., Heldin, C. H., & Miyazono, K. (1995). Cloning and characterization of a human type II receptor for bone morphogenetic proteins. *Proceedings of the National Academy of Sciences of the United States of America*, 92(17), 7632–7636. [10.1073/pnas.92.17.7632](https://doi.org/10.1073/pnas.92.17.7632)
- Rutkovskiy, A., Stensløyken, K. O., & Vaage, I. J. (2016). Osteoblast differentiation at a glance. *Medical Science Monitor Basic Research*, 22, 95–106. <https://doi.org/10.12659/msmbr.901142>
- Samee, N., Geoffroy, V., Marty, C., Schiltz, C., Vieux-Rochas, M., Levi, G., & de Vemejoul, M. C. (2008). Dlx5, a positive regulator of osteoblastogenesis, is essential for osteoblast-osteoclast coupling. *The American Journal of Pathology*, 173(3), 773–780. <https://doi.org/10.2353/ajpath.2008.080243>
- Santonocito, S., Polizzi, A., Palazzo, G., & Isola, G. (2021). The emerging role of microRNA in periodontitis: Pathophysiology, clinical potential and future molecular perspectives. *International Journal of Molecular Sciences*, 22(11), 5456. <https://doi.org/10.3390/ijms22115456>
- Sayad, A., Mirzajani, S., Gholami, B., Razzaghi, P., Ghafouri-Fard, S., & Taheri, M. (2020). Emerging role of long non-coding RNAs in the pathogenesis of periodontitis. *Biomedicine & Pharmacotherapy = Biomedecine & Pharmacotherapie*, 129, 110362. <https://doi.org/10.1016/j.biopha.2020.110362>
- Schliephake, H., Knebel, J. W., Aufderheide, M., & Tauscher, M. (2001). Use of cultivated osteoprogenitor cells to increase bone formation in segmental mandibular defects: An experimental pilot study in sheep. *International Journal of Oral and Maxillofacial Surgery*, 30(6), 531–537. <https://doi.org/10.1054/ijom.2001.0164>
- Shahi, M., Peymani, A., & Sahmani, M. (2017). Regulation of bone metabolism. *Reports of Biochemistry & Molecular Biology*, 5(2), 73–82.
- Shen, J. J., Zhang, C. H., Chen, Z. W., Wang, Z. X., Yang, D. C., Zhang, F. L., & Feng, K. H. (2019). LncRNA HOTAIR inhibited osteogenic differentiation of BMSCs by regulating Wnt/ β -catenin pathway. *European Review for Medical and Pharmacological Sciences*, 23(17), 7232–7246. https://doi.org/10.26355/eurev_201909_18826
- Shim, J. H., Greenblatt, M. B., Zou, W., Huang, Z., Wein, M. N., Brady, N., Hu, D., Charron, J., Brodtkin, H. R., Petsko, G. A., Zaller, D., Zhai, B., Gygi, S., Glimcher, L. H., & Jones, D. C. (2013). Schnurri-3 regulates ERK downstream of WNT signaling in osteoblasts. *The Journal of Clinical Investigation*, 123(9), 4010–4022. <https://doi.org/10.1172/JCI69443>
- Shimoyama, A., Wada, M., Ikeda, F., Hata, K., Matsubara, T., Nifuji, A., Noda, M., Amano, K., Yamaguchi, A., Nishimura, R., & Yoneda, T. (2007). Ihh/Gli2 signaling promotes osteoblast differentiation by regulating Runx2 expression and function. *Molecular Biology of the Cell*, 18(7), 2411–2418. <https://doi.org/10.1091/mbc.e06-08-0743>
- Silva, A. M., Moura, S. R., Teixeira, J. H., Barbosa, M. A., Santos, S. G., & Almeida, M. I. (2019). Long noncoding RNAs: A missing link in osteoporosis. *Bone Research*, 7, 10. <https://doi.org/10.1038/s41413-019-0048-9>
- St-Jacques, B., Hammerschmidt, M., & McMahon, A. P. (1999). Indian hedgehog signaling regulates proliferation and differentiation of chondrocytes and is essential for bone formation. *Genes & Development*, 13(16), 2072–2086. [10.1101/gad.13.16.2072](https://doi.org/10.1101/gad.13.16.2072)
- Stein, G. S., & Lian, J. B. (1993). Molecular mechanisms mediating proliferation/differentiation interrelationships during progressive development of the osteoblast phenotype. *Endocrine Reviews*, 14(4), 424–442. <https://doi.org/10.1210/edrv-14-4-424>
- Syed, F. A., Oursler, M. J., Hefferan, T. E., Peterson, J. M., Riggs, B. L., & Khosla, S. (2008). Effects of estrogen therapy on bone marrow adipocytes in postmenopausal osteoporotic women. *Osteoporosis International*, 19(9), 1323–1330. <https://doi.org/10.1007/s00198-008-0574-6>
- Tang, W., Li, Y., Osimiri, L., & Zhang, C. (2011). Osteoblast-specific transcription factor Osterix (Osx) is an upstream regulator of Satb2 during bone formation. *The Journal of Biological Chemistry*, 286(38), 32995–33002. <https://doi.org/10.1074/jbc.M111.244236>
- Thomson, D. W., & Dinger, M. E. (2016). Endogenous microRNA sponges: Evidence and controversy. *Nature Reviews. Genetics*, 17(5), 272–283. <https://doi.org/10.1038/nrg.2016.20>
- Tominaga, H., Maeda, S., Hayashi, M., Takeda, S., Akira, S., Komiya, S., Nakamura, T., Akiyama, H., & Imamura, T. (2008). CCAAT/enhancer-binding protein beta promotes osteoblast differentiation by enhancing Runx2 activity with ATF4. *Molecular Biology of the Cell*, 19(12), 5373–5386. <https://doi.org/10.1091/mbc.e08-03-0329>
- Ueyama, H., Ohta, Y., Imai, Y., Suzuki, A., Sugama, R., Minoda, Y., Takaoka, K., & Nakamura, H. (2021). Topical co-administration of zoledronate with recombinant human bone morphogenetic protein-2 can induce and maintain bone formation in the bone marrow environment. *BMC Musculoskeletal Disorders*, 22(1), 94. <https://doi.org/10.1186/s12891-021-03971-w>
- Ullah, I., Subbarao, R. B., & Rho, G. J. (2015). Human mesenchymal stem cells - current trends and future prospective. *Bioscience Reports*, 35(2), e00191. <https://doi.org/10.1042/BSR20150025>
- Vimalraj, S., Arumugam, B., Miranda, P. J., & Selvamurugan, N. (2015). Runx2: Structure, function, and phosphorylation in osteoblast differentiation. *International Journal of Biological Macromolecules*, 78, 202–208. <https://doi.org/10.1016/j.ijbiomac.2015.04.008>
- Wada, Y., Kataoka, H., Yokose, S., Ishizuya, T., Miyazono, K., Gao, Y. H., Shibasaki, Y., & Yamaguchi, A. (1998). Changes in osteoblast phenotype during differentiation of enzymatically isolated rat calvaria cells. *Bone*, 22(5), 479–485. [https://doi.org/10.1016/S8756-3282\(98\)00039-8](https://doi.org/10.1016/S8756-3282(98)00039-8)
- Wang, C. G., Hu, Y. H., Su, S. L., & Zhong, D. (2020). LncRNA DANCR and miR-320a suppressed osteogenic differentiation in osteoporosis by directly inhibiting the Wnt/ β -catenin signaling pathway. *Experimental & Molecular Medicine*, 52(8), 1310–1325. <https://doi.org/10.1038/s12276-020-0475-0>
- Wang, W., Lian, N., Li, L., Moss, H. E., Wang, W., Perrien, D. S., Elefteriou, F., & Yang, X. (2009). Atf4 regulates chondrocyte proliferation and differentiation during endochondral ossification by activating Ihh transcription. *Development (Cambridge, England)*, 136(24), 4143–4153. <https://doi.org/10.1242/dev.043281>
- Wang, W., Lian, N., Ma, Y., Li, L., Gallant, R. C., Elefteriou, F., & Yang, X. (2012). Chondrocytic Atf4 regulates osteoblast differentiation and function via Ihh. *Development (Cambridge, England)*, 139(3), 601–611. <https://doi.org/10.1242/dev.069575>
- Wang, X., Guo, B., Li, Q., Peng, J., Yang, Z., Wang, A., Li, D., Hou, Z., Lv, K., Kan, G., Cao, H., Wu, H., Song, J., Pan, X., Sun, Q., Ling, S., Li, Y.,

- Zhu, M., Zhang, P., ... Li, Y. (2013). miR-214 targets ATF4 to inhibit bone formation. *Nature Medicine*, 19(1), 93–100. <https://doi.org/10.1038/nm.3026>
- Wei, J., Shi, Y., Zheng, L., Zhou, B., Inose, H., Wang, J., Guo, X. E., Grosschedl, R., & Karsenty, G. (2012). miR-34s inhibit osteoblast proliferation and differentiation in the mouse by targeting SATB2. *The Journal of Cell Biology*, 197(4), 509–521. <https://doi.org/10.1083/jcb.201201057>
- Wen, J., Guan, Z., Yu, B., Guo, J., Shi, Y., & Hu, L. (2020). Circular RNA hsa_circ_0076906 competes with OGN for miR-1305 binding site to alleviate the progression of osteoporosis. *The International Journal of Biochemistry & Cell Biology*, 122, 105719. <https://doi.org/10.1016/j.biocel.2020.105719>
- Werner de Castro, G. R., Buss, Z., Rosa, J. S., Facchin, B. M., & Fröde, T. S. (2019). Evaluation of bone metabolism biomarkers in Paget's disease of bone. *Cureus*, 11(5), e4791. <https://doi.org/10.7759/cureus.4791>
- Xia, T., Dong, S., & Tian, J. (2020). miR-29b promotes the osteogenic differentiation of mesenchymal stem cells derived from human adipose tissue via the PTEN/AKT/ β -catenin signaling pathway. *International Journal of Molecular Medicine*, 46(2), 709–717. <https://doi.org/10.3892/ijmm.2020.4615>
- Xiao, G., Jiang, D., Ge, C., Zhao, Z., Lai, Y., Boules, H., Phimpilai, M., Yang, X., Karsenty, G., & Franceschi, R. T. (2005). Cooperative interactions between activating transcription factor 4 and Runx2/Cbfa1 stimulate osteoblast-specific osteocalcin gene expression. *The Journal of Biological Chemistry*, 280(35), 30689–30696. <https://doi.org/10.1074/jbc.M500750200>
- Xiao, X., Zhou, T., Guo, S., Guo, C., Zhang, Q., Dong, N., & Wang, Y. (2017). LncRNA MALAT1 sponges miR-204 to promote osteoblast differentiation of human aortic valve interstitial cells through up-regulating Smad4. *International Journal of Cardiology*, 243, 404–412. <https://doi.org/10.1016/j.ijcard.2017.05.037>
- Xiaoling, G., Shuaibin, L., & Kailu, L. (2020). MicroRNA-19b-3p promotes cell proliferation and osteogenic differentiation of BMSCs by interacting with lncRNA H19. *BMC Medical Genetics*, 21(1), 11. <https://doi.org/10.1186/s12881-020-0948-y>
- Xie, Q., Wang, Z., Bi, X., Zhou, H., Wang, Y., Gu, P., & Fan, X. (2014). Effects of miR-31 on the osteogenesis of human mesenchymal stem cells. *Biochemical and Biophysical Research Communications*, 446(1), 98–104. <https://doi.org/10.1016/j.bbrc.2014.02.058>
- Yamaguchi, M., Goto, M., Uchiyama, S., & Nakagawa, T. (2008). Effect of zinc on gene expression in osteoblastic MC3T3-E1 cells: enhancement of Runx2, OPG, and osteocalcin mRNA expressions. *Molecular and Cellular Biochemistry*, 312(1–2), 157–166. <https://doi.org/10.1007/s11010-008-9731-7>
- Yang, J., Andre, P., Ye, L., & Yang, Y. Z. (2015). The Hedgehog signalling pathway in bone formation. *International Journal of Oral Science*, 7(2), 73–79. <https://doi.org/10.1038/ijos.2015.14>
- Yang, J., Xu, Y., Xue, X., Zhang, M., Wang, S., & Qi, K. (2022). MicroRNA-26b regulates BMSC osteogenic differentiation of TMJ subchondral bone through β -catenin in osteoarthritis. *Bone*, 162, 116448. <https://doi.org/10.1016/j.bone.2022.116448>
- Yang, X., & Karsenty, G. (2004). ATF4, the osteoblast accumulation of which is determined post-translationally, can induce osteoblast-specific gene expression in non-osteoblastic cells. *The Journal of Biological Chemistry*, 279(45), 47109–47114. <https://doi.org/10.1074/jbc.M410010200>
- Yang, X., Matsuda, K., Bialek, P., Jacquot, S., Masuoka, H. C., Schinke, T., Li, L., Brancorsini, S., Sassone-Corsi, P., Townes, T. M., Hanauer, A., & Karsenty, G. (2004). ATF4 is a substrate of RSK2 and an essential regulator of osteoblast biology; implication for Coffin-Lowry Syndrome. *Cell*, 117(3), 387–398. [https://doi.org/10.1016/S0092-8674\(04\)00344-7](https://doi.org/10.1016/S0092-8674(04)00344-7)
- Yavropoulou, M. P., van Lierop, A. H., Hamdy, N. A., Rizzoli, R., & Papapoulos, S. E. (2012). Serum sclerostin levels in Paget's disease and prostate cancer with bone metastases with a wide range of bone turnover. *Bone*, 51(1), 153–157. <https://doi.org/10.1016/j.bone.2012.04.016>
- Yi, J., Liu, D., & Xiao, J. (2019). LncRNA MALAT1 sponges miR-30 to promote osteoblast differentiation of adipose-derived mesenchymal stem cells by promotion of Runx2 expression. *Cell and Tissue Research*, 376(1), 113–121. <https://doi.org/10.1007/s00441-018-2963-2>
- Yin, C., Tian, Y., Yu, Y., Li, D., Miao, Z., Su, P., Zhao, Y., Wang, X., Pei, J., Zhang, K., & Qian, A. (2021). Long noncoding RNA AK039312 and AK079370 inhibits bone formation via miR-199b-5p. *Pharmacological Research*, 163, 105230. <https://doi.org/10.1016/j.phrs.2020.105230>
- Yu, C., Wu, D., Zhao, C., & Wu, C. (2021). CircRNA TGFBR2/MiR-25-3p/TWIST1 axis regulates osteoblast differentiation of human aortic valve interstitial cells. *Journal of Bone and Mineral Metabolism*, 39(3), 360–371. <https://doi.org/10.1007/s00774-020-01164-4>
- Yu, L., Xia, K., Zhou, J., Hu, Z., Yin, X., Zhou, C., Zou, S., & Liu, J. (2022). circ_0003204 regulates the osteogenic differentiation of human adipose-derived stem cells via miR-370-3p/HDAC4 axis. *International Journal of Oral Science*, 14(1), 30. <https://doi.org/10.1038/s41368-022-00184-2>
- Yu, M., Wang, L., Ba, P., Li, L., Sun, L., Duan, X., Yang, P., Yang, C., & Sun, Q. (2017). Osteoblast progenitors enhance osteogenic differentiation of periodontal ligament stem cells. *Journal of Periodontology*, 88(10), e159–e168. <https://doi.org/10.1902/jop.2017.170016>
- Yu, S., Jiang, Y., Galson, D. L., Luo, M., Lai, Y., Lu, Y., Ouyang, H. J., Zhang, J., & Xiao, G. (2008). General transcription factor IIA-gamma increases osteoblast-specific osteocalcin gene expression via activating transcription factor 4 and runt-related transcription factor 2. *The Journal of Biological Chemistry*, 283(9), 5542–5553. <https://doi.org/10.1074/jbc.M705653200>
- Zarka, M., Hay, E., & Cohen-Solal, M. (2022). YAP/TAZ in bone and cartilage biology. *Frontiers in Cell and Developmental Biology*, 9, 788773. <https://doi.org/10.3389/fcell.2021.788773>
- Zhang, C. (2010). Transcriptional regulation of bone formation by the osteoblast-specific transcription factor Osx. *Journal of Orthopaedic Surgery and Research*, 5(1), 37. <https://doi.org/10.1186/1749-799x-5-37>
- Zhang, C., Cho, K., Huang, Y., Lyons, J. P., Zhou, X., Sinha, K., McCrear, P. D., & de Crombrughe, B. (2008a). Inhibition of Wnt signaling by the osteoblast-specific transcription factor Osterix. *Proceedings of the National Academy of Sciences of the United States of America*, 105(19), 6936–6941. <https://doi.org/10.1073/pnas.0710831105>
- Zhang, D., Ni, N., Wang, Y., Tang, Z., Gao, H., Ju, Y., Sun, N., He, X., Gu, P., & Fan, X. (2021). CircRNA-vgl3 promotes osteogenic differentiation of adipose-derived mesenchymal stem cells via modulating miRNA-dependent integrin $\alpha 5$ expression. *Cell Death and Differentiation*, 28(1), 283–302. <https://doi.org/10.1038/s41418-020-0600-6>
- Zhang, J., Chen, H., Leung, R., Choy, K. W., Lam, T. P., Ng, B., Qiu, Y., Feng, J. Q., Cheng, J., & Lee, W. (2018). Aberrant miR-145-5p/ β -catenin signal impairs osteocyte function in adolescent idiopathic scoliosis. *FASEB Journal*, 32(12), 6537–6549. <https://doi.org/10.1096/fj.201800281>
- Zhang, R. F., Liu, J. W., Yu, S. P., Sun, D., Wang, X. H., Fu, J. S., & Xie, Z. (2019). LncRNA UCA1 affects osteoblast proliferation and differentiation by regulating BMP-2 expression. *European Review for Medical and Pharmacological Sciences*, 23(16), 6774–6782. https://doi.org/10.26355/eurrev_201908_18715
- Zhang, X. W., Zhang, B. Y., Wang, S. W., Gong, D. J., Han, L., Xu, Z. Y., & Liu, X. H. (2014). Twist-related protein 1 negatively regulated osteoblastic transdifferentiation of human aortic valve interstitial cells by directly inhibiting runt-related transcription factor 2. *The Journal of Thoracic and Cardiovascular Surgery*, 148(4), 1700–1708.e1. <https://doi.org/10.1016/j.jtcvs.2014.02.084>
- Zhang, X., Yu, S., Galson, D. L., Luo, M., Fan, J., Zhang, J., Guan, Y., & Xiao, G. (2008b). Activating transcription factor 4 is critical for proliferation and survival in primary bone marrow stromal cells and calvarial osteoblasts. *Journal of Cellular Biochemistry*, 105(3), 885–895. <https://doi.org/10.1002/jcb.21888>
- Zhang, Y., Jia, S., Wei, Q., Zhuang, Z., Li, J., Fan, Y., Zhang, L., Hong, Z., Ma, X., Sun, R., He, W., Wang, H., Liu, Y., & Li, W. (2020). CircRNA_25487 inhibits bone repair in trauma-induced osteonecrosis of femoral head by sponging miR-134-3p through p21. *Regenerative Therapy*, 16, 23–31. <https://doi.org/10.1016/j.reth.2020.12.003>
- Zhou, R., Miao, S., Xu, J., Sun, L., & Chen, Y. (2021). Circular RNA circ_0000020 promotes osteogenic differentiation to reduce osteoporosis via sponging microRNA miR-142-5p to up-regulate Bone

Morphogenetic Protein BMP2. *Bioengineered*, 12(1), 3824–3836. <https://doi.org/10.1080/21655979.2021.1949514>
Zhou, X., Zhang, Z., Feng, J. Q., Dusevich, V. M., Sinha, K., Zhang, H., Darnay, B. G., & de Crombrugge, B. (2010). Multiple functions of

Osterix are required for bone growth and homeostasis in postnatal mice. *Proceedings of the National Academy of Sciences of the United States of America*, 107(29), 12919–12924. <https://doi.org/10.1073/pnas.0912855107>

Differentiation of osteoblasts: the links between essential transcription factors

ORIGINALITY REPORT

19%

SIMILARITY INDEX

14%

INTERNET SOURCES

15%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1	dokumen.pub Internet Source	1%
2	harvest.usask.ca Internet Source	1%
3	api.research-repository.uwa.edu.au Internet Source	1%
4	link.springer.com Internet Source	<1%
5	www.wjgnet.com Internet Source	<1%
6	"Cartilage", Springer Science + Business Media, 2016 Publication	<1%
7	www.academia.dk Internet Source	<1%
8	Linghui Hu, Wei Wu, Jun Zou. "环状RNA: 骨相关疾病的典型生物标志物", Journal of Zhejiang University-SCIENCE B, 2022 Publication	<1%

9	e-century.us Internet Source	<1 %
10	Chong Yin, Ye Tian, Dijie Li, Yang Yu et al. "Long non-coding RNA Lnc-DIF inhibits bone formation by sequestering miR-489-3p", iScience, 2022 Publication	<1 %
11	ir.vanderbilt.edu Internet Source	<1 %
12	era.ed.ac.uk Internet Source	<1 %
13	"Cell Signaling and Growth Factors in Development", Wiley, 2005 Publication	<1 %
14	theses.ncl.ac.uk Internet Source	<1 %
15	"Dental Stem Cells", Springer Nature, 2016 Publication	<1 %
16	"Mineralized Tissues in Oral and Craniofacial Science", Wiley, 2012 Publication	<1 %
17	etheses.whiterose.ac.uk Internet Source	<1 %
18	pingpdf.com Internet Source	<1 %

19	"Cell Engineering and Regeneration", Springer Science and Business Media LLC, 2020 Publication	<1 %
20	digitallibrary.usc.edu Internet Source	<1 %
21	epdf.pub Internet Source	<1 %
22	www2.mdpi.com Internet Source	<1 %
23	Toshihisa Komori. "Regulation of osteoblast differentiation by transcription factors", Journal of Cellular Biochemistry, 2006 Publication	<1 %
24	www.research.manchester.ac.uk Internet Source	<1 %
25	Dijie Li, Chaofei Yang, Chong Yin, Fan Zhao et al. "LncRNA, Important Player in Bone Development and Disease", Endocrine, Metabolic & Immune Disorders - Drug Targets, 2020 Publication	<1 %
26	Qiuling Zhang, Yifei Long, Liangyu Jin, Chenghao Li, Jie Long. "Non-coding RNAs regulate the BMP/Smad pathway during osteogenic differentiation of stem cells", Acta Histochemica, 2023 Publication	<1 %

27

Hichem Miraoui, Nicolas Severe, Pascal Vaudin, Jean-Christophe Pagès, Pierre J. Marie. "Molecular silencing of Twist1 enhances osteogenic differentiation of murine mesenchymal stem cells: Implication of FGFR2 signaling", *Journal of Cellular Biochemistry*, 2010

Publication

<1 %

28

Jianfeng Ping, Laifeng Li, Yongqiang Dong, Xudong Wu, Xiaogang Huang, Bin Sun, Bin Zeng, Fangming Xu, Wenqing Liang. "The role of long non - coding RNAs and circular RNAs in bone regeneration: Modulating miRNAs function", *Journal of Tissue Engineering and Regenerative Medicine*, 2022

Publication

<1 %

29

www.utupub.fi

Internet Source

<1 %

30

"RUNX Proteins in Development and Cancer", Springer Science and Business Media LLC, 2017

Publication

<1 %

31

Qadir, Abdul S., Soyoun Um, Heesu Lee, Kyunghwa Baek, Byoung Moo Seo, Gene Lee, Gwan-Shik Kim, Kyung Mi Woo, Hyun-Mo Ryoo, and Jeong-Hwa Baek. "miR-124 Negatively Regulates Osteogenic

<1 %

Differentiation and In Vivo Bone Formation of Mesenchymal Stem Cells : miR-124 inhibits osteogenic differentiation", Journal of Cellular Biochemistry, 2014.

Publication

32

livrepository.liverpool.ac.uk

Internet Source

<1 %

33

Dwight A. Towler, Rene St. Arnaud. "Use of Cultured Osteoblastic Cells to Identify and Characterize Transcriptional Regulatory Complexes", Elsevier BV, 2002

Publication

<1 %

34

orca.cf.ac.uk

Internet Source

<1 %

35

udspace.udel.edu

Internet Source

<1 %

36

www.frontiersin.org

Internet Source

<1 %

37

Danzhou Yang, Laurence H. Hurley. "Structure of the Biologically Relevant G-Quadruplex in The c-MYC Promoter", Nucleosides, Nucleotides and Nucleic Acids, 2006

Publication

<1 %

38

Egawa, Shiro, Shinichirou Miura, Hitoshi Yokoyama, Tetsuya Endo, and Koji Tamura. "Growth and differentiation of a long bone in

<1 %

limb development, repair and regeneration",
Development Growth & Differentiation, 2014.

Publication

39

Marcela Angelica De la Fuente-Hernandez,
Miguel Angel Sarabia-Sanchez, Jorge
Melendez-Zajgla, Vilma Maldonado-Lagunas.
"Role of IncRNAs into Mesenchymal Stromal
Cell Differentiation", American Journal of
Physiology-Cell Physiology, 2022

Publication

40

Takuo Kubota. "Wnt signaling in bone
metabolism", Journal of Bone and Mineral
Metabolism, 05/2009

Publication

41

escholarship.org

Internet Source

42

hdl.handle.net

Internet Source

43

www.mdpi.com

Internet Source

44

www.tandfonline.com

Internet Source

45

Akira Yamaguchi, Toshihisa Komori, Tatsuo
Suda. "Regulation of Osteoblast
Differentiation Mediated by Bone
Morphogenetic Proteins, Hedgehogs, and
Cbfa1", Endocrine Reviews, 2000

<1 %

<1 %

<1 %

<1 %

<1 %

<1 %

<1 %

46

smartech.gatech.edu

Internet Source

<1 %

47

"Transcriptional and Translational Regulation of Stem Cells", Springer Science and Business Media LLC, 2013

Publication

<1 %

48

d-nb.info

Internet Source

<1 %

49

Tong Ming Liu, Eng Hin Lee. "Transcriptional Regulatory Cascades in Runx2-Dependent Bone Development", Tissue Engineering Part B: Reviews, 2013

Publication

<1 %

50

dr.library.brocku.ca

Internet Source

<1 %

51

eprints.soton.ac.uk

Internet Source

<1 %

52

Fatemeh Ranjbarnejad, Mozafar Khazaei, Alireza Shahryari, Fatemeh Khazaei, Leila Rezakhani. "Recent advances in gene therapy for bone tissue engineering", Journal of Tissue Engineering and Regenerative Medicine, 2022

Publication

<1 %

53

Patsy Gomez-Picos, Katie Ovens, B. Frank Eames. "Limb Mesoderm and Head

<1 %

Ectomesenchyme Both Express a Core Transcriptional Program During Chondrocyte Differentiation", *Frontiers in Cell and Developmental Biology*, 2022

Publication

54

Wanjin Tang, Yang Li, Lindsey Osimiri, Chi Zhang. "Osteoblast-specific Transcription Factor Osterix (Osx) Is an Upstream Regulator of Satb2 during Bone Formation", *Journal of Biological Chemistry*, 2011

Publication

<1 %

55

hub.hku.hk

Internet Source

<1 %

56

zidapps.boku.ac.at

Internet Source

<1 %

57

Saki, Najmaldin, Majid Farshdousti Hagh, Esmaeil Mortaz, and Abdolreza Ardehshiry Lajimi. "Does DNA Methylation Plays a Critical Role in Osteoblastic Differentiation of Mesenchymal Stem Cells (MSCs)?", *Iranian Red Crescent Medical Journal*, 2013.

Publication

<1 %

58

espace.curtin.edu.au

Internet Source

<1 %

59

jmhg.springeropen.com

Internet Source

<1 %

60

Internet Source

<1 %

61

Carrie S. Soltanoff, Shuying Yang, Wei Chen, Yi-Ping Li. "Signaling Networks that Control the Lineage Commitment and Differentiation of Bone Cells", *Critical Reviews™ in Eukaryotic Gene Expression*, 2009

Publication

<1 %

62

Gengyan Liu, Jia Luo, Zhengguang Wang, Yong Zhou, Yong Li. "CircZNF367 suppresses osteogenic differentiation of human bone marrow mesenchymal stromal/stem cells via reducing HuR-mediated mRNA stability of LRP5", *Human Cell*, 2022

Publication

<1 %

63

Jingwei Wang, Yutao Cui, He Liu, Shaorong Li, Shouye Sun, Hang Xu, Chuangang Peng, Yanbing Wang, Dankai Wu. "MicroRNA-loaded biomaterials for osteogenesis", *Frontiers in Bioengineering and Biotechnology*, 2022

Publication

<1 %

64

Toshihisa Komori. "Runx2, an inducer of osteoblast and chondrocyte differentiation", *Histochemistry and Cell Biology*, 2018

Publication

<1 %

65

dev.biologists.org

Internet Source

<1 %

66	edoc.hu-berlin.de Internet Source	<1 %
67	elifesciences.org Internet Source	<1 %
68	jme.endocrinology-journals.org Internet Source	<1 %
69	kclpure.kcl.ac.uk Internet Source	<1 %
70	matin.gatech.edu Internet Source	<1 %
71	prod--journal.elifesciences.org Internet Source	<1 %
72	tel.archives-ouvertes.fr Internet Source	<1 %
73	www.escholar.manchester.ac.uk Internet Source	<1 %
74	"ASBMR 26th Annual Meeting 1001-1222", Journal of Bone and Mineral Research, 2004 Publication	<1 %
75	"Primer on the Metabolic Bone Diseases and Disorders of Mineral Metabolism", Wiley, 2008 Publication	<1 %
76	Anny W. Robert, Bruna H. Marcon, Bruno Dallagiovanna, Patrícia Shigunov. "Adipogenesis, Osteogenesis, and	<1 %

Chondrogenesis of Human Mesenchymal Stem/Stromal Cells: A Comparative Transcriptome Approach", *Frontiers in Cell and Developmental Biology*, 2020

Publication

77

Carmen Lanzillotti, Monica De Mattei, Chiara Mazziotta, Francesca Taraballi et al. "Long Non-coding RNAs and MicroRNAs Interplay in Osteogenic Differentiation of Mesenchymal Stem Cells", *Frontiers in Cell and Developmental Biology*, 2021

Publication

<1 %

78

Minghua Hu, Xi Zhu, Heng Yuan, Haifeng Li, Hongchun Liao, Shijie Chen. "The function and mechanism of the miR - 210 - 3p/KRAS axis in bone marrow - derived mesenchymal stem cell from patients with osteoporosis", *Journal of Tissue Engineering and Regenerative Medicine*, 2021

Publication

<1 %

79

Shiyao Guan, Zhen Zhang, Jun Wu. "Non-coding RNA delivery for bone tissue engineering: progress, challenges and potential solutions", *iScience*, 2022

Publication

<1 %

80

Tiantian He, Wei Liu, Lihua Cao, Ying Liu, Zi Zou, Yancheng Zhong, Haihua Wang, Yuqing Mo, Shuping Peng, Cijun Shuai. "CircRNAs and

<1 %

LncRNAs in osteoporosis", Differentiation, 2020

Publication

81

Xiaochen Fan, Ashley J. Waardenberg, Madeleine Demuth, Pierre Osteil et al. "TWIST1 homodimers and heterodimers orchestrate lineage-specific differentiation", Cold Spring Harbor Laboratory, 2019

Publication

<1 %

82

Yao Weng, Ziyi Wang, Yoko Fukuhara, Airi Tanai et al. " drives calcium signaling toward osteoblast differentiation: A bioinformatics - oriented study ", BioFactors, 2021

Publication

<1 %

83

biolres.biomedcentral.com

Internet Source

<1 %

84

clonerresources.com

Internet Source

<1 %

85

core.ac.uk

Internet Source

<1 %

86

doi.org

Internet Source

<1 %

87

ebin.pub

Internet Source

<1 %

88

josr-online.biomedcentral.com

Internet Source

<1 %

89	mafiadoc.com Internet Source	<1 %
90	onlinelibrary.wiley.com Internet Source	<1 %
91	repo.lib.jfn.ac.lk Internet Source	<1 %
92	repository.up.ac.za Internet Source	<1 %
93	rupress.org Internet Source	<1 %
94	trepo.tuni.fi Internet Source	<1 %
95	www.doria.fi Internet Source	<1 %
96	www.iconceptpress.com Internet Source	<1 %
97	www.pubmedcentral.nih.gov Internet Source	<1 %
98	www.repository.cam.ac.uk Internet Source	<1 %
99	Anna K. Coussens, Ian P. Hughes, Christopher R. Wilkinson, C. Phillip Morris et al. "Identification of genes differentially expressed by prematurely fused human	<1 %

sutures using a novel in vivo – in vitro approach", Differentiation, 2008

Publication

100

B. Arumugam, K. Balagangadharan, N. Selvamurugan. "Syringic acid, a phenolic acid, promotes osteoblast differentiation by stimulation of Runx2 expression and targeting of Smad7 by miR-21 in mouse mesenchymal stem cells", Journal of Cell Communication and Signaling, 2018

Publication

<1 %

101

Epigenetics of Aging, 2010.

Publication

<1 %

102

Pavitra K. Jadaun, Shuang Zhang, Marijke Koedam, Jeroen Demmers, Suvro Chatterjee, Johannes P. Leeuwen, Bram C. Eerden. "Inhibition of hypoxia - induced Mucin 1 alters the proteomic composition of human osteoblast - produced extracellular matrix, leading to reduced osteogenic and angiogenic potential", Journal of Cellular Physiology, 2021

Publication

<1 %

103

Prem Swaroop Yadav, Yingzi Yang. "Lineage Determination of Chondrocytes and Osteoblasts: From the Perspective of Cellular and Molecular Biology", Elsevier BV, 2020

Publication

<1 %

- | | | |
|-----|--|------|
| 104 | Suryaji Patil, Kai Dang, Xin Zhao, Yongguang Gao, Airong Qian. "Role of LncRNAs and CircRNAs in Bone Metabolism and Osteoporosis", Frontiers in Genetics, 2020
Publication | <1 % |
| 105 | Yang Yang, Wang Yujiao, Wang Fang, Yuan Linhui, Guo Ziqi, Wei Zhichen, Wang Zirui, Wang Shengwang. "The roles of miRNA, IncRNA and circRNA in the development of osteoporosis", Biological Research, 2020
Publication | <1 % |
| 106 | "Bone and Development", Springer Science and Business Media LLC, 2010
Publication | <1 % |
| 107 | "Metabolism and Epigenetic Regulation: Implications in Cancer", Springer Science and Business Media LLC, 2022
Publication | <1 % |
| 108 | Chuan Wang, Haiqing Liao, Zhengguo Cao. "Role of Osterix and MicroRNAs in Bone Formation and Tooth Development", Medical Science Monitor, 2016
Publication | <1 % |
| 109 | Cinzia Aurilia, Simone Donati, Gaia Palmi, Francesca Miglietta, Teresa Iantomasi, Maria Luisa Brandi. "The Involvement of Long Non- | <1 % |

Coding RNAs in Bone", International Journal of Molecular Sciences, 2021

Publication

110 Essentials of Mesenchymal Stem Cell Biology and Its Clinical Translation, 2013. <1 %
Publication

111 Fang, Sijie, Yuan Deng, Ping Gu, and Xianqun Fan. "MicroRNAs Regulate Bone Development and Regeneration", International Journal of Molecular Sciences, 2015. <1 %
Publication

112 Hironori Hojo, Shinsuke Ohba, Fumiko Yano, Ung-il Chung. "Coordination of chondrogenesis and osteogenesis by hypertrophic chondrocytes in endochondral bone development", Journal of Bone and Mineral Metabolism, 2010 <1 %
Publication

113 Youqiang Sun, Vincent Kuek, Yuhao Liu, Jennifer Tickner, Yu Yuan, Leilei Chen, Zhikui Zeng, Min Shao, Wei He, Jiake Xu. "MiR-214 is an important regulator of the musculoskeletal metabolism and disease", Journal of Cellular Physiology, 2018 <1 %
Publication

Exclude quotes On

Exclude matches Off

Exclude bibliography On