

The Role of Rice Hull Liquid Smoke in the Traumatic Ulcer Healing

by Ira Arundina

Submission date: 02-Aug-2022 11:07AM (UTC+0800)

Submission ID: 1877944315

File name: of_Rice_Hull_Liquid_Smoke_in_the_Traumatic_Ulcer_Healing_2.pdf (1.26M)

Word count: 4260

Character count: 22558

The Role of Rice Hull Liquid Smoke in the Traumatic Ulcer Healing

Ira Arundina¹ Indeswati Diyatri¹ Tuti Kusumaningsih¹ Meircurius Dwi Condro Surboyo² Elita Monica³ Novitasari Mira Afanda³

¹Department of Oral Biology, Faculty of Dental Medicine, Universitas Airlangga, Surabaya, Indonesia

²Department of Oral Medicine, Faculty of Dental Medicine, Universitas Airlangga, Surabaya, Indonesia

³Undergraduate Student, Faculty of Dental Medicine, Universitas Airlangga, Surabaya, Indonesia

Address for correspondence Dr. Ira Arundina, drg., M.Si, Department of Oral Biology, Faculty of Dental Medicine, Universitas Airlangga, Jln. Prof. Dr. Moestopo 47 Surabaya, Indonesia 60132 (e-mail: arundinafkg@yahoo.com).

Eur J Dent

Abstract

Keywords

- ▶ traumatic ulcer
- ▶ fibroblast
- ▶ macrophage
- ▶ lymphocyte
- ▶ rice hull
- ▶ liquid smoke
- ▶ IL-6 and TGF- β

Objective The aim of this study was to prove the role of rice hull liquid smoke (RH-LS) on lymphocytes, macrophages, fibroblasts, interleukin 6 (IL-6), and transforming growth factor β (TGF- β) expression during traumatic ulcer healing.

Materials and Methods The RH-LS was obtained from the pyrolysis process. Traumatic ulcers were made 10 mm along the labial fornix incisive inferior of Wistar rat using a round stainless-steel blade. In control group, traumatic ulcers were treated using sterile water, and meanwhile in experimental group were treated using RH-LS once a day for 3, 5, and 7 days. After treatment, animal was terminated and their labial fornix incisive inferior tissues were biopsy and stained using hematoxylin and eosin staining to determine lymphocytes, macrophages, and fibroblasts. The IL-6 and TGF- β expressions were analyzed used immunohistochemistry staining.

Result The lymphocytes, macrophages, and fibroblasts were higher in the RH-LS group for 3-, 5-, and 7-day treatment ($p < 0.05$). The IL-6 expression was higher only in the 5-day treatment, and the TGF- β expression was higher in the 3- and 7-day treatment.

Conclusion The RH-LS able to accelerated the traumatic ulcer healing by increasing the number of lymphocytes, macrophages, fibroblasts, IL-6, and TGF- β expression.

Introduction

Traumatic ulcer is a common lesion in the oral cavity, around 83.6% of people have ever experienced it.¹ The prevalence of traumatic ulcers varies around 3 to 24% worldwide.² Ulcer is characterized by loss of the epithelium, causing a chief complaint of pain.^{3,4} The current topical treatment of traumatic ulcers, such as benzydamine hydrochloride in some special cases like in the diabetic condition, does not promote healing⁵ that is why an alternative treatment is needed.

The carbonization of wood at high temperature and without oxygen produces liquid smoke, then the smoke is condensed at a room temperature, and this whole of this

process is called pyrolysis.^{6,7} Liquid smoke compositions are obtained from the degradation of wood components through pyrolysis.^{7,8} The degradation of lignin generates phenolic compounds that have several beneficial effects in wound healing including traumatic ulcer.^{5,8}

Rice (*Oryza sativa L*) is the staple food for people all around the world, especially in Asia including Indonesia.⁹ Postharvesting rice process produces 20% rice hull as its byproduct,⁹ and rice hull can be processed through pyrolysis to become rice hull liquid smoke (RH-LS). RH-LS contains phenolic compounds, such as guaiacol, 2,6-dimethoxyphenols, phenol, and 4-ethyl-2-methoxyphenol (2-EMP) have antioxidant effect which bind to reactive oxygen species (ROS), inhibit the production of

DOI <https://doi.org/10.1055/s-0040-1714445>
ISSN 1305-7456.

©2020 Dental Investigation Society

License terms



pro-inflammatory cytokines, and inhibit the activation of nuclear factor kappa β (NF- κ B).⁹⁻¹¹ The inhibition of NF- κ B makes the polarization of anti-inflammatory macrophage (M2) more dominant compared with the pro-inflammatory macrophage (M1).¹² M2 with the help of lymphocyte secrete growth factors, such as fibroblast growth factor-2 (FGF-2), transforming growth factor β (TGF- β), and chemokine ligand-18 (CCL-18). Those growth factors accelerate the proliferation of fibroblast and collagen which are needed in the healing process.^{9,13-17}

Despite all the promising finding in the literatures mentioned above, RH-LS has a promising potential as treatment to accelerate the traumatic ulcer healing. Therefore, an *in vivo* research on the effect of RH-LS using macrophage, lymphocytes, fibroblast, IL-6, and TGF- β expressions as the indicators of the healing of traumatic ulcer needs to be conducted.

Materials and Methods

Pyrolysis and Distillation of Liquid Smoke

The rice hull (*Oryza sativa* L) were collected at the Center of Rice Processing in Tumpang village. The pyrolysis process was conducted in the Research and Development of Forest Products Laboratory, Bogor, Indonesia (ISO number 17025:2008). First of all, total 1,760 g of rice hull were air-dried at room temperature.

A kerosene pump stove was used as the heater of the pyrolysis furnace, as well as 30 cm in diameter and 40 cm in height encircling reactor was used. The reactor is connected to the cooling tubes by pipeline for condensing the fumes and generating the liquid smoke. The air-dried rice hull was put inside the furnace and then the furnace was closed. After that, the setting of the condenser was done, and cold water was used to stream the cooling tube. At a temperature of 400°C, the pyrolysis was performed for 4.5 hours with a heating rate of 3.33°C.¹⁸ The formula below was used to calculate the yield of the liquid smoke (%):¹⁸:

$$\text{yield (\%)} = \frac{\text{liquid smoke (g)}}{\text{rice hull (g)}} \times 100$$

The liquid smoke was then stored for 48 hours and after that the filtration process using Whatman 52 (Whatman 52, Cat No 1452, GE Healthcare Life Science, Singapore) was done. After that, a distillation process at the temperature of 120 to 150°C was done to purificate the liquid smoke.¹⁹ The yield of distilled liquid smoke derived from rice hull (%) was calculated using the following formula¹⁸:

$$\text{liquid smoke (\%)} = \frac{\text{condensed liquid smoke (ml)}}{\text{liquid smoke (ml)}} \times 100$$

Animals

Ethics approval was obtained from the Ethical Clearance of Health Experiment Committee, Faculty of Dental Medicine, Airlangga University, Surabaya under registered-number 132/HRECC.FODM/IV/2019.

This research was comprised of 2-month-old 30 male Wistar rats with a weight range of 120 to 160 g. Animal Testing Laboratory, Department of Biochemistry, Faculty of Medicine, Airlangga University, Surabaya was chosen as the place where the treatment and experiment of animals were done. Collective cages equipped with artificial light with a cycle of 12 hours light/12 hours dark at room temperature (27°C) were used to house Wistar rat in which free access of water and standard diet were given.

Traumatic Ulcer Induction

Wistar rats were put under anesthesia using a ketamine/xylazine cocktail, after that, at the labial fornix incisive inferior a 10 mm-long incision were made using a round stainless steel blade to induce traumatic ulcers.²⁰ They were confirmed if a yellowish-white ulcer with an erythematous halo appear after 24 hours.⁵

Wistar rats were randomly assigned to a control group, and an experimental group after the traumatic ulcers were confirmed. The intraoral dropping method was used for the topical treatment of RH-LS and sterile water. In the control group, the traumatic ulcers were treated topically using sterile water; meanwhile, in the experimental group, they were treated topically using RH-LS with a dosage of 20 μ L/20 gr once a day for 3, 5, and 7 days.¹⁸

Fibroblast, Macrophage, and Lymphocyte Counting on Traumatic Ulcer

After being treated for 3, 5, and 7 days in each group, the rats were terminated and their labial fornix incisive inferior tissues were subjected to biopsy before a hematoxylin and eosin staining to determine the number of fibroblasts, macrophages, and lymphocytes. While the TGF- β (TGF- β mouse monoclonal, Santacruz biotechnology) and IL-6 (IL-6 mouse monoclonal, Santacruz biotechnology) expression was analyzed using immunohistochemistry. All measurement was done using light microscope (Nikon H600L microscope; Nikon, Japan) with a magnification of \times 400 at five fields of view with single blind operator.

Statistical Analysis

The data were shown in the form of mean \pm standard deviation ($X \pm$ SD) for each group and each measurement. Then, the data were analyzed using independent *t*-test and the significance was set at $p < 0.05$.

Result

Pyrolysis and Distillation of Liquid Smoke

The pyrolysis process of 1,760 g rice hull was performed with the final temperature 400°C for 4.5 hours and then the liquid smoke was distilled at 120 to 150°C (–Table 1).

Table 1 The yield of pyrolysis and distillation of liquid smoke

| Component | % |
|--------------|-------|
| Liquid smoke | 34.82 |
| distilled | 87 |

The Number of Macrophages on Traumatic Ulcer

The histopathology of traumatic ulcer after topical treatment of RH-LS can be seen in **Fig. 1**. The number of macrophages after topical treatment of RH-LS for 3 days was higher than control (7.20 ± 0.84 ; $p = 0.005$). The topical treatment of RH-LS for 5 and 7 days was higher than control ($p = 0.004$ and $p = 0.022$, respectively; **Fig. 2A**).

The Number of Lymphocytes on Traumatic Ulcer

The number of lymphocytes after topical treatment of RH-LS for 3 and 5 days was higher than control ($p = 0.001$

and $p = 0.000$, respectively), but there is no difference in the treatment for 7 days ($p = 0.108$; **Fig. 2B**).

The Number of Fibroblasts on Traumatic Ulcer

The number of fibroblasts after topical treatment of RH-LS for 3, 5, and 7 days was higher than control ($p = 0.032$, $p = 0.001$, and $p = 0.025$, respectively; **Fig. 2C**).

The Interleukin-6 Expression on Traumatic Ulcer

The IL-6 expression after topical treatment of RH-LS for 5 days was higher than control ($p = 0.001$). The topical treatment of




Fig. 1 The histopathology of traumatic ulcer tissue. The lymphocytes (blue arrow), macrophages (black arrow), and fibroblasts (green arrow) cells were analyzed using Hematoxylin and Eosin staining. Magnification at $\times 400$.




Fig. 2 The number of lymphocytes (A), macrophages (B), and fibroblasts (C). The same character on the top of column, indicated different each group (t -test: $p < 0.05$). The interleukin-6 expression on traumatic ulcer.

RH-LS for 3 and 7 days showed no differences with control groups ($p = 0.088$ and $p = 0.293$, respectively; **Fig. 3** and **Fig. 4A**).

The Transforming Growth Factor-β Expression on Traumatic Ulcer

The TGF-β expression after topical treatment of RH-LS for 3 and 7 days was higher than control ($p = 0.034$ and $p = 0.019$, respectively). The topical treatment of RH-LS for 5 days showed no differences with control groups ($p = 0.059$; **Fig. 5** and **Fig. 4B**).

Discussion

RH-LS is a liquid smoke that produced from rice hull through the pyrolysis process. The process of pyrolysis includes different response forms to be specific decomposition, polymerization, oxidation, and condensation. The reactions that occur during pyrolysis are the decomposition of structures such as cellulose, hemicellulose, and lignin. In the pyrolysis process, compounds are formed such as carbonyl, acids, and phenols. In addition, it also produces carcinogenic compounds such as polycyclic aromatic hydrocarbons (PAHs). PAHs are formed naturally resulting from incomplete combustion, and this

compound is formed at pyrolysis temperatures ranging from 500 to 900°C. PAH formation rate is also influenced by the source. Some PAH compounds such as benzo(a)pyrene²¹ is a carcinogenic substance and can cause damage to amino acids and vitamins.²² One way to separate and eliminate unwanted components such as benzo(a)pyrene is by purification. One of the purifications methods is distillation to remove that compound.²³

RH-LS in this study was made with a pyrolysis temperature of 400°C because the products can be produced optimally at temperature of 400 to 550°C. Temperature 400°C produced more phenol, ferulic acid, and guaiacol than the temperature of 800°C.²⁴

The treatment of RH-LS to the traumatic ulcer showed a higher number of lymphocytes, macrophages, fibroblasts for 3, 5, and 7 days. Phenolic compounds which are contained in RH-LS, such as EMP and guaiacol, might have the role to increase them. EMP and guaiacol has antioxidant abilities that can increase proliferation lymphocytes, macrophages, and fibroblasts. The antioxidant ability is related to ability of inhibit the formation of ROS. Elevated ROS will cause oxidative stress in lymphocytes leading them to apoptosis; hence, with the inhibition of ROS, the number of lymphocytes will be increased.²⁵ Lymphocytes play a role




Fig. 3 The interleukin-6 expression on traumatic ulcer tissue with immunohistochemistry staining. Magnification at ×400.




Fig. 4 The interleukin-6 expression (A) and transforming growth factor-β expression (B). The same character on the top of column, indicated different each group (t-test: $p < 0.05$).




Fig. 5 Transforming growth factor- β expression on traumatic ulcer tissue with immunohistochemistry staining. Magnification at $\times 400$.

in releasing lymphokines, epidermal growth factor, basic fibroblast growth factor,²⁶ that influence the recruitment of macrophages to the wound and in the end promote the fibroblast proliferation.²⁷

The phenolic compound in RH-LS able to interfere the I κ B kinase (IKK) and NF- κ B signaling pathway by inhibiting the ROS and resulting inhibition of cytokines production, such as tumor necrosis factor (TNF)- α , IL-6, and IL-1 β .²⁸ The previous research has proved that RH-LS can reduce the pro-inflammatory cytokine, the TNF- α , by inhibiting the activation of NF- κ B through the inhibition of IKK activity.²⁹ Other research with liquid smoke from coconut shell also confirmed that liquid smoke able to decrease the NF- κ B and TNF- α expression.¹⁸ The result of this study is different with two previous studies. The pro-inflammatory cytokines, IL-6 expression, is higher in the treatment of RH-LS for 5-day treatment. Even the IL-6 is pro-inflammatory cytokines, its need in the traumatic ulcer healing.³⁰ IL-6 is released by M2 macrophages along with growth factors, such as TGF- β , platelet-derived growth factor, and basic fibroblast growth factor, which is the higher TGF- β expression also observed after treated with RH-LS for 3-day and 5-day treatment. That IL-6 and TGF- β have contributed in the fibroblast proliferation and collagen production.^{15,17} The increase of IL-6 expression promoted the fibroblasts proliferation and extracellular matrix production, including procollagen and fibronectin.³¹ This higher IL-6 and TGF- β expression in this study strengthen the previous study that liquid smoke able increased the fibroblast,³² and collagen formation in oral ulcer healing.⁵ The higher level of IL-6 and TGF- β can accelerate wound closure, granulation tissue formation, neovascularization, and re-epithelialization.^{33,34}

The number of macrophages and lymphocytes is significant increased during treatment with RH-LS in this research. The bidirectional interaction between lymphocytes and macrophages is well established, and the polarization paradigm of macrophages is derived from their interaction with lymphocytes. Recruitment of macrophages plays a important

role in healing process of traumatic ulcers.³⁵ Macrophages play a role in the wound healing process of M1 polarization in the initial stages of wound healing and the properties of M2 polarization expression during the later stages of wound healing. M1 macrophages mediate tissue damage and have a proinflammatory effect, while M2 macrophages function in cleansing debris, induction of angiogenesis, and tissue remodeling.³⁶ The inhibition of NF- κ B results a switch from M1 polarization to M2 polarization. This condition make the M2 becomes more dominant, for produce the anti-inflammatory cytokine.³⁶

The dominant M2 polarization results in the more abundant release of anti-inflammatory cytokines and growth factors, one of them is TGF- β which is proved to be increased after the application of RH-LS. In every phase of wound healing, TGF- β takes an important role to suppress the inflammatory response and to promote the formation of granulation tissue,^{37,38} promoting fibroblast migration and promoting wound healing.^{39,40} In the current study, it was found that RH-LS increased the TGF- β expression. The increase of TGF- β 1 promoted the fibroblast proliferation which can be seen in this study that there was an increase of fibroblast after RH-LS topical application. The ability to increase fibroblast of TGF- β plays an important role to accelerate wound healing. On the 5-day treatment, TGF- β expression found not significant compared with the control. TGF- β secretion starts in the early proliferative phase, then the secretion keeps on going, and reaches its peak on the 7 day.^{41,42} That is why the significant differences were found on the 3 days and the 7 days after RH-LS treatment.

Based on the results of the study, there was a significant difference between the number of macrophages in the RH-LS treatment. This is due to the ability of RH-LS in inhibiting the production of ROS and inhibiting the production of pro-inflammatory cytokines such as TNF- α to accelerating the inflammatory phase.⁴³ The inhibition of ROS production resulted in the inhibition of NF- κ B which later resulted in a switch from M1 polarization to M2 polarization and makes

M2 more dominant. The dominant M2 increases IL-6, TGF- β 1, PDGF, FGF-2, and IL-10 secretion. It can be seen in this study that IL-6 and TGF- β 1 increased after RH-LS treatment, which later contributes in wound healing by increasing the fibroblast. From the mechanism that is explained above, topical application of RH-LS to traumatic ulcer can increase the infiltration of lymphocytes, macrophages, increase TGF- β 1, and IL-6 expression and later increase the fibroblasts. This condition accelerates the inflammatory phase and indicates a faster traumatic ulcer healing.

Conclusion

The topical application of RH-LS increased the infiltration of macrophages, lymphocytes, IL-6, and TGF- β expression and resulted in the increase of fibroblasts number in traumatic ulcer. Therefore, it accelerated the healing of traumatic ulcer.

31

Funding

This work was supported by Ministry of Higher Education 2020 with grant number 607/UN3.14/PT/2020.

Conflict of Interest

None declared.

References

- Puspitasari D, Apriasari ML. Analysis of traumatic ulcer healing time under the treatment of the Mauli banana (*Musa acuminata*) 25% stem extract gel. *Padjadjaran J Dent* 2017;29(1):21–25
- Apriasari ML, Rahayu RP, Ernawati D. Mauli banana stem extract application increased expression of NF- κ B in traumatic ulcer healing. *Research* 2018;67(32):67–70
- Mortazavi H, Safi Y, Baharvand M, Rahmani S. Diagnostic features of common oral ulcerative lesions: an updated decision tree. *Int J Dent* 2016;2016:7278925
- Ali SA, Saudi HI. An expert system for the diagnosis and management of oral ulcers ScienceDirect: an expert system for the diagnosis and management of oral ulcers. *Tanta Dent J* 2015;11(1):42–46
- Surboyo MDC, Arundina I, Rahayu RP. Increase of collagen in diabetes-related traumatic ulcers after the application of liquid smoke coconut shell. *Dent J* 2017;71(32):71–75 (Majalah Kedokt Gigi)
- Oramahi HA, Diba F. Maximizing the production of liquid smoke from bark of durio by studying its potential compounds. *Procedia Environ Sci* 2013;17:60–69
- Wagiman FX, Ardiansyah A, Witjaksono. Activity of coconut-shell liquid-smoke as an insecticide on the rice brown planthopper (*Nilaparvata lugens*) *J Agric Biol Sci* 2014;9(9):293–296
- Ramakrishnan S, Moeller P. Liquid smoke. *Product Hardwood Pyrolysis* 2002;47(1):366–367
- Kim SP, Yang JY, Kang MY, Park JC, Nam SH, Friedman M. Composition of liquid rice hull smoke and anti-inflammatory effects in mice. *J Agric Food Chem* 2011;59(9):4570–4581
- Hayden MS, Ghosh S. Regulation of NF- κ B by TNF family cytokines. Vol. 26, *Seminars in Immunology*. Elsevier Ltd; 2014; 253–266
- Lingappan K. NF- κ B in Oxidative Stress. *Curr Opin Toxicol* 2018;7:81–86
- Zhou Y, Zhang T, Wang X, et al. Curcumin modulates macrophage polarization through the inhibition of the toll-like receptor 4 expression and its signaling pathways. *Cell Physiol Biochem* 2015;36(2):631–641
- Townsend CM, Beauchamp RD, Evers BM, Mattox KL. *Sabiston Textbook of Surgery: The Biological Basis of Modern Surgical Practice*. Shock. 20th edition. Elsevier Ltd; 2001;15(5):408
- Chistiakov DA, Myasoedova VA, Revin VV, Orekhov AN, Bobryshev YV. The impact of interferon-regulatory factors to macrophage differentiation and polarization into M1 and M2. *Immunobiology* 2018;223(1):101–111
- Ploeger DT, Hoesper NA, Schipper M, Koerts JA, de Rond S, Bank RA. Cell plasticity in wound healing: paracrine factors of M1/M2 polarized macrophages influence the phenotypical state of dermal fibroblasts. *Cell Commun Signal* 2013;11(1):29
- Akiyama M, Yasuoka H, Yoshimoto K, Takeuchi T. CC-chemokine ligand 18 is a useful biomarker associated with disease activity in IgG4-related disease. *Ann Rheum Dis* 2018;77(9):1386–1387
- Bi Y, Chen J, Hu F, Liu J, Li M, Zhao L. M2 Macrophages as a potential target for atherosclerosis treatment. *Neural Plast* 2019. Available at: <https://www.hindawi.com/journals/np/2019/6724903/>.
- Surboyo MDC, Arundina I, Rahayu RP, Mansur D, Bramantoro T. Potential of Distilled Liquid Smoke Derived from Coconut (*Cocos nucifera* L) Shell for Traumatic Ulcer Healing in Diabetic Rats. *Eur J Dent* 2019;13(2):271–279
- Desniorita M. The effect of adding liquid smoke powder to shelf life of sauce. *Int J Adv Sci Eng Inf Technol* 2015;5(6):457–459
- Surboyo MDC, Emawati DS, Arundina I, Rahayu RP. Oral ulcer healing after treatment with distilled liquid smoke of coconut shell on diabetic rats. *J Krishna Inst Med Sci Univ* 2019;8(2):70–79
- Lingbeck JM, Cordero P, O'Bryan CA, Johnson MG, Ricke SC, Crandall PG. Functionality of liquid smoke as an all-natural antimicrobial in food preservation. *Meat Sci* 2014;97(2):197–206
- Anggraini SPA, Yuniningsih S. Liquid smoke purification process for benzo (A) pyrene levels lowering on food safety. *J Agric Food Tech* 2013;3(12):1–4
- Lombok JZ, Setiaji B, Trisunaryati W, Wijaya K. Effect of pyrolysis temperature and distillation on character of coconut shell liquid smoke. Available at: <http://www.journalajst.com/sites/default/files/1576.pdf>. Accessed June 19, 2020
- Sukharnikov IY, Esengaraev E, Turgenbaev AT, Iskakov MS. Production, properties and use of the rice hulls pyrolysis organic product as a disinfectant. *World J Vet Med Eng* 2015;1(1):1–8
- Belikov AV, Schraven B, Simeoni L. T cells and reactive oxygen species. *J Biomed Sci* 2015;22(1):85
- Istiati NI, Surjono I, Surboyo MDC. Role of lactoferrin fibroblast growth factor 2 and vascular endothelial growth factor in gingival wounds. *J Krishna Inst Med Sci Univ* 2019;8(3):38–45
- Suryadi IA, Asmarajaya A, Maliawan S. *Proses Penyembuhan dan Penanganan Luka*. Ilmu Penyakit Bedah; 2012 ;1–19
- Limtrakul P, Yodkeeree S, Pitchakarn P, Punfa W. Suppression of Inflammatory Responses by Black Rice Extract in RAW 264.7 Macrophage Cells via Downregulation of NF- κ B and AP-1 Signaling Pathways. *Asian Pac J Cancer Prev* 2015;16(10):4277–4283
- Friedman M. Rice brans, rice bran oils, and rice hulls: composition, food and industrial uses, and bioactivities in humans, animals, and cells. *J Agric Food Chem* 2013;61(45):10626–10641

- 30 Luig M, Kluger MA, Goerke B, et al. Inflammation-induced IL-6 functions as a natural brake on macrophages and limits GN. *J Am Soc Nephrol* 2015;26(7):1597–1607
- 31 Choy E. Interleukin-6 as a multifunctional regulator: inflammation, immune response, and fibrosis. *Cold Spring Harb Perspect Biol* 2017;2(2):1–5
- 32 Tarawan VM, Mantilidewi KI, Dhini IM, Radhiyanti PT, Sutedia E. Coconut shell liquid smoke promotes burn wound healing. *J Evid Based Complementary Altern Med* 2017;22(3):436–440
- 33 Barrientos S, Stojadinovic O, Golinko MS, Brem H, Tomic-Canic M. Growth factors and cytokines in wound healing. *Wound Repair Regen* 2008;16(5):585–601
- 34 Gallucci RM, Sloan DK, Heck JM, Murray AR, O'Dell SJ. Interleukin 6 indirectly induces keratinocyte migration. *J Invest Dermatol* 2004;122(3):764–772
- 35 Surboyo MDC, Mahdani FY, Ernawati DS, Sarasati A, Rezkita F. The macrophage responses during diabetic oral ulcer healing by liquid coconut shell smoke: an immunohistochemical analysis. *Eur J Dent* 2020;1–5 DOI: 10.1055/s-0040-1712776
- 36 Okizaki S, Ito Y, Hosono K, et al. Suppressed recruitment of alternatively activated macrophages reduces TGF- β 1 and impairs wound healing in streptozotocin-induced diabetic mice. *Biomed Pharmacother* 2015;70:317–325
- 37 Wang L, Qin W, Zhou Y, et al. Transforming growth factor β plays an important role in enhancing wound healing by topical application of povidone-iodine. *Sci Rep* 2017;7(1):991
- 38 Liu Y, Li Y, Li N, et al. TGF- β 1 promotes scar fibroblasts proliferation and transdifferentiation via up-regulating MicroRNA-21. *Sci Rep* 2016;6:32231
- 39 Liarte S, Bernabé-García Á, Nicolás FJ. Role of TGF- β in skin chronic wounds: a keratinocyte perspective. *Cells* 2020;9(2):306
- 40 Surboyo MDC, Mahdani FY, Savitri DS, et al. Number of macrophages and transforming growth factor β expression in Citrus limon L. Tlekung peel oil-treated traumatic ulcers in diabetic rats. *Trop J Pharm Res* 2019;18(7):1427–1433
- 41 Hendriati L, Hamid IS, Widodo T, Wandasari C, Risata PM. Effect of egg white gel against burn healing on white rat (*Rattus novergicus*) *J ILMU KEFARMASIAN Indones [Internet]* 2018;16(2):231
- 42 Savari R, Shafiei M, Galehdari H, Kesmati M. Expression of VEGF and TGF- β genes in skin wound healing process induced using phenytoin in male rats. *Jundishapur J Heal Sci* 2019;1–5 : 10.5812/jjhs.86041
- 43 Zduńska K, Dana A, Kolodziejczak A, Rotsztejn H. Antioxidant properties of ferulic acid and its possible application. *Skin Pharmacol Physiol* 2018;31(6):332–336

The Role of Rice Hull Liquid Smoke in the Traumatic Ulcer Healing

ORIGINALITY REPORT

20%

SIMILARITY INDEX

13%

INTERNET SOURCES

15%

PUBLICATIONS

2%

STUDENT PAPERS

PRIMARY SOURCES

- | | | |
|---|---|----|
| 1 | jppres.com Internet Source | 2% |
| 2 | Submitted to iGroup Student Paper | 1% |
| 3 | Diah Savitri Ernawati, Meircurius Dwi Condro Surboyo, Nurina Febriyanti Ayuningtyas, Ayu Anggraini Broto Nagoro. "Role of Inflammatory Cell Responses in Stimulating Fibroblasts in Diabetic Oral Ulcer after Treatment with Liquid Smoke of Coconut Endocarp: A Histological Assessment", European Journal of Dentistry, 2020 Publication | 1% |
| 4 | test.dovepress.com Internet Source | 1% |
| 5 | Shin-ichiro Okizaki, Yoshiya Ito, Kanako Hosono, Kazuhito Oba et al. "Suppressed recruitment of alternatively activated macrophages reduces TGF- β 1 and impairs wound healing in streptozotocin-induced | 1% |

diabetic mice", Biomedicine & Pharmacotherapy, 2015

Publication

6

Jody M. Lingbeck, Paola Cordero, Corliss A. O'Bryan, Michael G. Johnson, Steven C. Ricke, Philip G. Crandall. "Functionality of liquid smoke as an all-natural antimicrobial in food preservation", Meat Science, 2014

Publication

1 %

7

"Therapeutic Dressings and Wound Healing Applications", Wiley, 2020

Publication

1 %

8

L.R. Lockett-Chastain. "Interleukin (IL)-6 modulates transforming growth factor- β expression in skin and dermal fibroblasts from IL-6-deficient mice", British Journal of Dermatology, 08/2009

Publication

1 %

9

www.biolifesas.org

Internet Source

1 %

10

Yasuo Morimoto, Tohru Tsuda, Hiroshi Yamato, Takako Oyabu et al. "COMPARISON OF GENE EXPRESSION OF CYTOKINES mRNA IN LUNGS OF RATS INDUCED BY INTRATRACHEAL INSTILLATION AND INHALATION OF MINERAL FIBERS", Inhalation Toxicology, 2008

Publication

1 %

| | | |
|----|---|------|
| 11 | www.phcogres.com Internet Source | 1 % |
| 12 | Ruslin Hadanu, Daniel Ambrosius Nicholas Apituley. "Volatile Compounds Detected in Coconut Shell Liquid Smoke through Pyrolysis at a Fractioning Temperature of 350-420 °C", Makara Journal of Science, 2016 Publication | <1 % |
| 13 | sabre.lib.sussex.ac.uk Internet Source | <1 % |
| 14 | www.painphysicianjournal.com Internet Source | <1 % |
| 15 | www.frontiersin.org Internet Source | <1 % |
| 16 | Nirawati Pribadi, Dwita Budiarti, Hendy Jaya Kurniawan, Ira Widjiastuti. "The NF-kB and Collagen Type 1 Expression in Dental Pulp after Treated Calcium Hydroxide Combined with Propolis", European Journal of Dentistry, 2021 Publication | <1 % |
| 17 | link.springer.com Internet Source | <1 % |
| 18 | Mendel Friedman. "Rice Brans, Rice Bran Oils, and Rice Hulls: Composition, Food and Industrial Uses, and Bioactivities in Humans, | <1 % |

Animals, and Cells", Journal of Agricultural and Food Chemistry, 2013

Publication

19

Zhang, Fengmei, Yuguo Ren, Peng Liu, Yufeng Ren, and Debao Wang. "Expression of TGF- β 1 and miRNA-145 in patients with diabetic foot ulcers", Experimental and Therapeutic Medicine, 2016.

Publication

<1 %

20

iris.unige.it

Internet Source

<1 %

21

pubmed.ncbi.nlm.nih.gov

Internet Source

<1 %

22

Mohammed Ahmed Aljuanid, Huda Rashad Qaid, Dur Muhammad Lashari, Rini Devijanti Ridwan et al. "Nano-emulsion of mangosteen rind extract in a mucoadhesive patch for periodontitis regenerative treatment: An in vivo study", Journal of Taibah University Medical Sciences, 2022

Publication

<1 %

23

f1000research.com

Internet Source

<1 %

24

jkimsu.com

Internet Source

<1 %

25

journal.uwks.ac.id

Internet Source

<1 %

26

www.ratoc.com

Internet Source

<1 %

27

"Abstracts of the 2nd International Conference (quote)Strategies in Tissue Engineering(quote)", *Cytotherapy*, 5/1/2006

Publication

<1 %

28

Abbas Shapouri - Moghaddam, Saeed Mohammadian, Hossein Vazini, Mahdi Taghadosi et al. "Macrophage plasticity, polarization, and function in health and disease", *Journal of Cellular Physiology*, 2018

Publication

<1 %

29

Hasan Ashari Oramahi, Farah Diba, Nurhaida. "New Bio Preservatives from Lignocelluloses Biomass Bio-oil for Anti Termites *Coptotermes Curvignathus Holmgren*", *Procedia Environmental Sciences*, 2014

Publication

<1 %

30

Roghaye Savari, Mohammad Shafiei, Hamid Galehdari, Mahnaz Kesmati. "Expression of VEGF and TGF- β Genes in Skin Wound Healing Process Induced Using Phenytoin in Male Rats", *Jundishapur Journal of Health Sciences*, 2019

Publication

<1 %

31

oamjms.eu

Internet Source

<1 %

| | | |
|----|---|------|
| 32 | vtechworks.lib.vt.edu Internet Source | <1 % |
| 33 | core.ac.uk Internet Source | <1 % |
| 34 | genesandnutrition.biomedcentral.com Internet Source | <1 % |
| 35 | ir.lib.uwo.ca Internet Source | <1 % |
| 36 | jeccr.biomedcentral.com Internet Source | <1 % |
| 37 | www.herbmedpharmacol.com Internet Source | <1 % |
| 38 | www.science.gov Internet Source | <1 % |
| 39 | Meircurius Dwi Condro Surboyo, Fatma Yasmin Mahdani, Diah Savitri Ernawati, Andari Sarasati, Fianza Rezkita. "The Macrophage Responses during Diabetic Oral Ulcer Healing by Liquid Coconut Shell Smoke: An Immunohistochemical Analysis", European Journal of Dentistry, 2020 Publication | <1 % |
| 40 | Yusnaini, Edi Suryanto, Said Hasan, Angela Wulansari, Eka Kusuma Dewi. "The Characteristics of Volatile Compounds of | <1 % |

Kenari (*Canarium indicum* L.) Shell Liquid Smoke", IOP Conference Series: Earth and Environmental Science, 2021

Publication

41

Elly Munadziroh, Muhammad Genadi Askandar, Anita Yuliati, Meircurius Dwi Condro Surboyo, Wan Himratul Aznita Wan Harun. "The effect of secretory leukocyte protease inhibitor amnion membrane on incisional wound healing", *Journal of Oral Biology and Craniofacial Research*, 2022

Publication

<1 %

42

FDI World Dental Federation. "P098AWDC 2015 – Abstract book - 23 September 2015 - FREE COMMUNICATION SESSIONS 21–40 and POSTER SESSIONS 17–32", *International Dental Journal*, 2015

Publication

<1 %

43

Rongguo He, Hui Yin, Baohong Yuan, Tao Liu, Li Luo, Ping Huang, Liangcheng Dai, Kang Zeng. "IL-33 improves wound healing through enhanced M2 macrophage polarization in diabetic mice", *Molecular Immunology*, 2017

Publication

<1 %

44

ÖZLEM ONAY. " Production of Bio-Oil from Biomass: Slow Pyrolysis of Rapeseed () in a Fixed-Bed Reactor ", *Energy Sources*, 2003

Publication

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

The Role of Rice Hull Liquid Smoke in the Traumatic Ulcer Healing

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7
