

 BMC

# Journal of Biological Engineering

ISSN 1546-1718

Part of Springer Nature

[Search](#)[Explore journals](#)  
[Menu](#)[Get published](#)[About BMC](#)[Login](#)

# Journal of Biological Engineering

[Home](#) [About](#) [Articles](#) [Submission Guidelines](#)[About](#) ▾[Contact](#)[Editorial Board](#)

## Editorial Board

### Editors-in-Chief

Prof Raj R Rao, *University of Arkansas, USA*

Prof Jeong-Yeol Yoon, *The University of Arizona, USA*

### Founding Editor-in-Chief

Prof Mark Riley, *University of Nebraska, USA*

### Past Editor-in-Chief

Prof Mark Eiteman, *University of Georgia, USA*

### Regional Editors

Prof Alfonso Jaramillo, *University of Warwick, UK*

Prof Sang Yup Lee, *Korea Advanced Institute of Science and Technology, South Korea*

### Associate Editors

Prof Ke Cheng, *North Carolina State University, USA*

Prof Marko Dolinar, *University of Ljubljana, Slovenia*

Prof Gyoo Yeol Jung, *Pohang University of Science and Technology, South Korea*

Prof Michael Ladisch, *Purdue University, USA*

Prof Min Lee, *University of California, Los Angeles, USA*

Prof Iram Mondaca Fernandez, *Instituto Tecnológico de Sonora, Mexico*

Prof ZhiYong Qian, *Sichuan University, China*

Prof Jixun Zhan, *Utah State University, USA*

## **Editorial Board**

Prof Wilfred Chen, *University of Delaware, USA*

Prof Hyun Jung Chung, *Korea Advanced Institute of Science and Technology, South Korea*

Prof Paul Freemont, *Imperial College London, UK*

Prof Sharon Gerecht, *Johns Hopkins University, USA*

Prof Man Bock Gu, *Korea University, South Korea*

Prof Joseph Irudayaraj, *Purdue University, USA*

Prof Katy Kao, *Texas A&M University, USA*

Prof Deok-Ho Kim, *Johns Hopkins University, USA*

Prof Jin-Woo Kim, *University of Arkansas, USA*

Prof Yeu-Chun Kim, *Korea Advanced Institute of Science and Technology, South Korea*

Prof Heinz Köppl, *TU Darmstadt, Germany*

Prof Gaetan Laroche, *University of Laval, Canada*

Prof Dali Li, *East China Normal University, China*

Prof Mahitosh Mandal, *Indian Institute of Technology - Kharagpur, India*

Prof John March, *Cornell University, USA*

Prof Robert Mitchell, *Ulsan National Institute of Science and Technology, South Korea*

Prof Tom Richard, *Penn State University, USA*

Prof Beatrix Suess, *Technical University of Darmstadt, Germany*

Prof Volker Wendisch, *Bielefeld University, Germany*

Prof Phillip C Wright, *University of Southampton, UK*

Prof Hu Yang, *Missouri University of Science and Technology, USA*

Prof Liju Yang, *North Carolina Central University, USA*

Prof An-Ping Zeng, *Technical University of Hamburg-Harburg, Germany*

Submit manuscript

---

[Editorial Board](#)

---

[Manuscript editing services](#)

---

[Instructions for Editors](#)

---

[Sign up for article alerts and news from this journal](#)

---

ISSN: 1754-1611 (electronic)

---

 **Journal of Biological Engineering**

ISSN: 1754-1611

### Contact us

Submission enquiries: [john.aceron@springernature.com](mailto:john.aceron@springernature.com)

General enquiries: [info@biomedcentral.com](mailto:info@biomedcentral.com)

---

[Read more on our blogs](#)

[Policies](#)

[Support and Contact](#)

[Receive BMC newsletters](#)

[Accessibility](#)

[Leave feedback](#)

[Manage article alerts](#)

[Press center](#)

[Careers](#)

[Language editing for authors](#)

[Scientific editing for authors](#)

Follow BMC



By using this website, you agree to our [Terms and Conditions](#), [California Privacy Statement](#), [Privacy statement](#) and [Cookies](#) policy. [Manage cookies/Do not sell my data](#) we use in the preference centre.

---

**SPRINGER NATURE**

© 2023 BioMed Central Ltd unless otherwise stated. Part of [Springer Nature](#).



Search

Explore journals  
Menu

Get published

About BMC

Login



# Journal of Biological Engineering

[Home](#) [About](#) [Articles](#) [Submission Guidelines](#)

About ▾

[Contact](#)

[Editorial Board](#)

## Editorial Board

### Editors-in-Chief

Prof Raj R Rao, *University of Arkansas, USA*

Prof Jeong-Yeol Yoon, *The University of Arizona, USA*

### Founding Editor-in-Chief

Prof Mark Riley, *University of Nebraska, USA*

### Past Editor-in-Chief

Prof Mark Eiteman, *University of Georgia, USA*

### Regional Editors

Prof Alfonso Jaramillo, *University of Warwick, UK*

Prof Sang Yup Lee, *Korea Advanced Institute of Science and Technology, South Korea*

### Associate Editors

Prof Ke Cheng, *North Carolina State University, USA*

Prof Marko Dolinar, *University of Ljubljana, Slovenia*

Prof Gyoo Yeol Jung, *Pohang University of Science and Technology, South Korea*

Prof Michael Ladisch, *Purdue University, USA*

Prof Min Lee, *University of California, Los Angeles, USA*

Prof Iram Mondaca Fernandez, *Instituto Tecnológico de Sonora, Mexico*

Prof ZhiYong Qian, *Sichuan University, China*

Prof Jixun Zhan, *Utah State University, USA*

## **Editorial Board**

Prof Wilfred Chen, *University of Delaware, USA*

Prof Hyun Jung Chung, *Korea Advanced Institute of Science and Technology, South Korea*

Prof Paul Freemont, *Imperial College London, UK*

Prof Sharon Gerecht, *Johns Hopkins University, USA*

Prof Man Bock Gu, *Korea University, South Korea*

Prof Joseph Irudayaraj, *Purdue University, USA*

Prof Katy Kao, *Texas A&M University, USA*

Prof Deok-Ho Kim, *Johns Hopkins University, USA*

Prof Jin-Woo Kim, *University of Arkansas, USA*

Prof Yeu-Chun Kim, *Korea Advanced Institute of Science and Technology, South Korea*

Prof Heinz Köppl, *TU Darmstadt, Germany*

Prof Gaetan Laroche, *University of Laval, Canada*

Prof Dali Li, *East China Normal University, China*

Prof Mahitosh Mandal, *Indian Institute of Technology - Kharagpur, India*

Prof John March, *Cornell University, USA*

Prof Robert Mitchell, *Ulsan National Institute of Science and Technology, South Korea*

Prof Tom Richard, *Penn State University, USA*

Prof Beatrix Suess, *Technical University of Darmstadt, Germany*

Prof Volker Wendisch, *Bielefeld University, Germany*

Prof Phillip C Wright, *University of Southampton, UK*

Prof Hu Yang, *Missouri University of Science and Technology, USA*

Prof Liju Yang, *North Carolina Central University, USA*

Prof An-Ping Zeng, *Technical University of Hamburg-Harburg, Germany*

Submit manuscript



# Journal of Biological Engineering



[Volumes and issues](#) > Volume 17, issue 1

Search within journal

## Volume 17, issue 1, December 2023

19 articles in this issue

### [Antioxidant activities in relation to the transport of heavy metals from the soil to different parts of \*Betula pendula\* \(Roth.\)](#)

Iwona Makuch-Pietraś, Dorota Grabek-Lejko ... Idalia Kasprzyk


Research | [Open Access](#) | Published: 06 March 2023

Article: 19




### [Customized biofilm device for antibiofilm and antibacterial screening of newly developed nanostructured silver and zinc coatings](#)

Daniele Ghezzi, Marco Boi ... Gabriela Graziani



[The ELF3 transcription factor is associated with an epithelial phenotype and represses epithelial-mesenchymal transition](#)


Ayalur Raghu Subbalakshmi, Sarthak Sahoo ... Mohit Kumar Jolly



[This is part of 2 collections](#)


[Exosome-mediated crosstalk between epithelial cells amplifies the cell injury cascade in CaOx stone formation](#)

Yuanyuan Yang, Senyuan Hong ... Yang Xun




[Dynamic interaction of injected liquid jet with skin layer interfaces revealed by microsecond imaging of optically cleared ex vivo skin tissue model](#)

Abdul Mohizin, Jakir Hossain Imran ... Jung Kyung Kim



[Synergistic improvement of cinnamylamine production by metabolic regulation](#)


Shan Yuan, Chao Xu ... Wei Liu






## [Therapeutic angiogenesis and tissue revascularization in ischemic vascular disease](#)

Xinyue Chen, Wenlu Yu ... Ao Shi




## [Assessing the capabilities of 2D fluorescence monitoring in microtiter plates with data-driven modeling for secondary substrate limitation experiments of \*Hansenula polymorpha\*](#)

Christoph Berg, Laura Herbst ... Jochen Büchs



## [Prospective use of amniotic mesenchymal stem cell metabolite products for tissue regeneration](#)

Andang Miatmoko, Berlian Sarasitha Hariawan ... Dewi Melani Hariyadi




## [LIFTOSCOPE: development of an automated AI-based module for time-effective and contactless analysis and isolation of cells in microtiter plates](#)

Florian Narrog, Richard Lensing ... Robert H. Schmitt

Research | [Open Access](#) | Published: 07 February 2023

Article: 10




## [CD146 expression profile in human skin and pre-vascularized dermo-epidermal skin substitutes in vivo](#)

Monica Nanni, Dominic Rüttsche ... Thomas Biedermann

Research | [Open Access](#) | Published: 31 January 2023

Article: 9



## [Research progress on black phosphorus hybrids hydrogel platforms for biomedical applications](#)

Hao-xuan Li, Kun-chi Zhao ... Qing-san Zhu

Review | [Open Access](#) | Published: 30 January 2023

Article: 8




## [RN-Autoencoder: Reduced Noise Autoencoder for classifying imbalanced cancer genomic data](#)


Ahmed Arafa, Nawal El-Fishawy ... Marwa Radad

Research | [Open Access](#) | Published: 30 January 2023

Article: 7




## [The synergistic and enhance effects of IL-6 and M-CSF to expand and differentiate functional dendritic cells from human monocytes under serum-free condition](#)




## [Tracking unlabeled cancer cells imaged with low resolution in wide migration chambers via U-NET class-1 probability \(pseudofluorescence\).](#)

Paola Antonello, Diego Morone ... Diego Ulisse Pizzagalli




## [T7Max transcription system](#)

Christopher Deich, Brock Cash ... Katarzyna P. Adamala




## [Investigation of two metabolic engineering approaches for \(R,R\)-2,3-butanediol production from glycerol in Bacillus subtilis](#)

Nunthaphan Vikromvarasiri, Shuhei Noda ... Akihiko Kondo



## [Mixed culture biocatalytic production of the high-value biochemical 7-methylxanthine](#)

Meredith B. Mock & Ryan M. Summers




## [Rapid remodeling observed at mid-term in-vivo study of a smart reinforced acellular vascular graft implanted on a rat model](#)

Francis O. Obiweluzor, Mukhammad Kayumov ... In-Seok Jeong

Research | [Open Access](#) | Published: 03 January 2023 |

Article: 1



## For authors

---

[Submission guidelines](#)

[Contact the journal](#)

Submit manuscript

## Explore

---

[Volumes and issues](#)

[Collections](#)

Not logged in - 180.247.145.54

East Java HE Consortium (3002712836) - 6763 SpringerLink Indonesia eJourn Consortium - Higher Education (3000122892) - Universitas Airlangga Faculty of Economics and Business (2000629929) - 11741 SpringerLink Indonesia eJourn Consortium (3000951794)

**SPRINGER NATURE**

© 2023 Springer Nature Switzerland AG. Part of [Springer Nature](#).



## Scientific Programming Papers


Open

Hindawi

## Journal of Biological Engineering

### COUNTRY

United Kingdom

 Universities and research institutions in United Kingdom

 Media Ranking in United Kingdom

### SUBJECT AREA AND CATEGORY

Biochemistry, Genetics and Molecular Biology  
Cell Biology  
Molecular Biology

Engineering  
Biomedical Engineering

Environmental Science  
Environmental Engineering

### PUBLISHER

BioMed Central Ltd.

### H-INDEX

**44**

### PUBLICATION TYPE

Journals

### ISSN

17541611

### COVERAGE


2007-2021

### INFORMATION

[Homepage](#)

[How to publish in this journal](#)





Hindawi

Open

---

## Scientific Programming Papers

Maximize The Impact, Reach & Visibility Of Your Next  
More On How To Submit

Submit

### SCOPE

Biological engineering is an emerging discipline that encompasses engineering theory and practice connected to and derived from the science of biology, just as mechanical engineering and electrical engineering are rooted in physics and chemical engineering in chemistry. Topical areas include, but are not limited to: Synthetic biology and cellular design Biomolecular, cellular and tissue engineering Bioproduction and metabolic engineering Biosensors Ecological and environmental engineering Biological engineering education and the biodesign process As the official journal of the Institute of Biological Engineering, Journal of Biological Engineering provides a home for the continuum from biological information science, molecules and cells, product formation, wastes and remediation, and educational advances in curriculum content and pedagogy at the undergraduate and graduate-levels. Manuscripts should explore commonalities with other fields of application by providing some discussion of the broader context of the work and how it connects to other areas within the field.

 Join the conversation about this journal



# Scientific Programming Papers


Maximize The Impact, Reach & Visibility Of Your Next Paper - How To Submit


Quartiles

11th Dec 2021


### FIND SIMILAR JOURNALS ?

1	2	3	4
<b>Frontiers in Bioengineering and Biotechnology</b> CHE	<b>Trends in Biotechnology</b> GBR	<b>Advanced Biosystems</b> DEU	<b>Advances in Experimental Medicine and Biology</b> USA
<b>79%</b> similarity	<b>71%</b> similarity	<b>71%</b> similarity	<b>68%</b> similarity






% International Collaboration





Citable documents

Non-citable documents



Cited documents

Uncited documents



**Journal of Biological Engineering**

Q1 Biomedical Engineering  
best quartile

SJR 2021  
1.05

powered by scimagojr.com


Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scimaç
```

**SCImago Graphica**

Explore, visually communicate and make sense of data with our **new data visualization tool**.



Metrics based on Scopus® data as of April 2022

**Leave a comment**

Name

Email

(will not be published)





REVIEW

Open Access



# Prospective use of amniotic mesenchymal stem cell metabolite products for tissue regeneration

Andang Miatmoko<sup>1,2\*</sup>, Berlian Sarasitha Hariawan<sup>3</sup>, Devy Maulidya Cahyani<sup>3</sup>, Syarifah Sutra Dewangga<sup>4</sup>, Kevin Ksatria Handoko<sup>4</sup>, Purwati<sup>2</sup>, Ram Kumar Sahu<sup>5</sup> and Dewi Melani Hariyadi<sup>1</sup>

## Abstract

Chronic disease can cause tissue and organ damage constituting the largest obstacle to therapy which, in turn, reduces patients' quality-adjusted life-year. Degenerative diseases such as osteoporosis, Alzheimer's disease, Parkinson's disease, and infectious conditions such as hepatitis, cause physical injury to organs. Moreover, damage resulting from chronic conditions such as diabetes can also culminate in the loss of organ function. In these cases, organ transplantation constitutes the therapy of choice, despite the associated problems of immunological rejection, potential disease transmission, and high morbidity rates. Tissue regeneration has the potential to heal or replace tissues and organs damaged by age, disease, or trauma, as well as to treat disabilities. Stem cell use represents an unprecedented strategy for these therapies. However, product availability and mass production remain challenges. A novel therapeutic alternative involving amniotic mesenchymal stem cell metabolite products (AMSC-MP) has been developed using metabolites from stem cells which contain cytokines and growth factors. Its potential role in regenerative therapy has recently been explored, enabling broad pharmacological applications including various gastrointestinal, lung, bladder and renal conditions, as well as the treatment of bone wounds, regeneration and skin aging due to its low immunogenicity and anti-inflammatory effects. The various kinds of growth factors present in AMSC-MP, namely bFGF, VEGF, TGF- $\beta$ , EGF and KGF, have their respective functions and activities. Each growth factor is formed by different proteins resulting in molecules with various physicochemical properties and levels of stability. This knowledge will assist in the manufacture and application of AMSC-MP as a therapeutic agent.

**Keywords** Quality-adjusted life-year, Amniotic mesenchymal stem cell metabolite products, Growth factors, Tissue injury, Stem cells, Molecular therapy

## \*Correspondence:

Andang Miatmoko  
andang-m@ff.unair.ac.id

<sup>1</sup> Department of Pharmaceutical Science, Faculty of Pharmacy, Universitas Airlangga, Nanizar Zaman Joenoes Building, Campus C UNAIR Mulyorejo, Mulyorejo, Surabaya, East Java 60115, Indonesia

<sup>2</sup> Stem Cell Research and Development Center, Universitas Airlangga, 2nd Floor Institute of Tropical Disease Building, Campus C UNAIR Mulyorejo, Surabaya 60115, Indonesia

<sup>3</sup> Master Program of Pharmaceutical Sciences, Faculty of Pharmacy, Universitas Airlangga, Campus C UNAIR Mulyorejo, Surabaya 60115, Indonesia

<sup>4</sup> Faculty of Pharmacy, Study Program of Pharmacy, Universitas Airlangga, Campus C UNAIR Mulyorejo, Surabaya 60115, Indonesia

<sup>5</sup> Department of Pharmaceutical Sciences, Hemvati Nandan Bahuguna Garhwal University (A Central University), Chauras Campus, Tehri Garhwal, Uttarakhand 249161, India



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

## Background

Degenerative diseases, physical injury to organs, and damage due to chronic conditions such as diabetes can cause organ function loss [1–3]. In the early stages of disease, pharmacological drug therapy is the first-line treatment choice, but it has some drawbacks. For example, there are only five types of drugs for reducing the symptoms of Alzheimer's disease [4], and pharmacotherapy to slow disease progression is not yet available. For this purpose, therapeutic agents should inhibit extracellular amyloid plaque deposition and intracellular neurofibrillary formation. Additionally, Alzheimer's therapies that have neuroprotective mechanisms and the use of anti-inflammatory stem cell therapies and the growth factor NDX-1017 are currently under investigation [5, 6].

Stem cells are undifferentiated cells that continuously divide, renew themselves, and differentiate into different types of cells. With the ability of self-renewal, pluripotency, and differentiation, stem cells have great potential for treating various diseases. Stem cells can be divided into two main groups based on their origin: embryonic stem cells and adult stem cells. However, stem cell therapy has two main problems that pose challenges for its therapeutic use. The first relates to availability and mass production. Cell therapy protocols using mesenchymal stem cells (MSCs) for tissue regeneration generally require hundreds of millions of MSCs per treatment. Therefore, expansion of the *in vitro* cell culture is necessary for longer periods and large bioreactors [7]. The need for cells to replace the disease-induced loss of hepatocytes, pancreatic cells, or cardiomyocytes involves approximately  $1$  to  $10 \times 10^9$  functional cells per patient. An even higher requirement has been calculated for the production of "in vitro blood," as approximately  $2.5 \times 10^{12}$  red blood cells are required per patient in transfusion treatments [8]. The main challenge in mass production lies in standardizing the process due to the complexity of pluripotent stem cells and processing cells on a large scale [9].

The use of stem cells can affect the recipient's immune system. The administered cells can directly induce an immune response or modulate the immune system. This is primarily in the case of cells that are not intended to be used for their essential function (nonhomologous use) or when administered to non-physiological sites, which may change the immunogenicity of the cells. Another risk is bacterial and viral infections. As a cell-based product, stem cell production does not allow for terminal sterilization, purification, virus removal, or inactivation processes. Thus, the risk of transmitting bacterial, viral, fungal, or prion pathogens from the donor to the recipient can lead to life-threatening and even fatal reactions [10].

Because of these limitations, new therapeutic strategies are needed. Stem cell-mediated tissue regeneration involves soluble factors secreted by these cells. Cytokines and growth factors, such as transforming growth factor beta (TGF- $\beta$ ), stromal cell-derived factor 1 (SDF-1), and vascular endothelial growth factor (VEGF), are secreted by stem cells and progenitor cells transplanted into the intestinal space or injected into blood vessels and stimulate many regenerative processes such as neovascularization, activation of tissue intrinsic progenitor cells, decreased apoptosis of endogenous cardiomyocytes, and registration of assistive cells for tissue repair [11, 12]. Additionally, mesenchymal stem cells (MSCs) secrete growth factors and cytokines, which promote wound repair. The combination of growth factors and cytokines successfully induces angiogenesis, reduces inflammation, and promotes fibroblast migration and collagen production [13].

## Amniotic membrane stem cell metabolite products (AMSC-MPs)

### Placental tissue is the primary source of AMSC-MPs

The amnion, chorion, amniotic fluid, and umbilical cord are of fetal origin. These components have been widely studied because of their potential use as cell sources for regenerative therapies. At delivery, the amniotic membrane is strong, protects the fetus from physical shocks, regulates the pH of the fluid membranes, and secretes various cell signals and bioactive molecules as antimicrobials and anti-inflammatory agents [14]. Amniotic membrane stem cells (AMSCs) are MSCs from the amniotic epithelium and the stroma of the amniotic membrane that are sources of epidermal growth factor (EGF) and keratinocyte growth factor (KGF). Furthermore, stem cells synthesize and secrete various extracellular matrix proteins, cytokines, growth factors, and other bioactive proteins that contribute to the healing and regenerative processes [15]. The collection of conditioned stem cell culture media rich in bioactive agents such as growth factor and cytokines secreted into the extracellular space is defined as a metabolite product. Since it is derived from amniotic mesenchymal stem cells, it is referred to as AMSC-MP [7].

The AMSC-MP collection procedures are conducted under sterile conditions. The first step is expanding cells to the desired passage. Then, upon trypsinization and counting of selected cells, the cell needs to be seeded again at a determined cell density, for example, 12,000 cells/cm<sup>2</sup>, and cultured in an appropriate growth medium. After 72 h, the growth medium is removed, and the cells are washed several times with phosphate-buffered saline, preferably without Ca<sup>2+</sup>/Mg<sup>2+</sup> content, at room temperature. After that, the cell culture medium

(pre-warmed to 37 °C) with only 1% antibiotic is added. The AMSC-MP was collected after being incubated for 48 h and centrifuged at 1500 rpm for 3 min two times, then filtered with a syringe filter 0.45 µm to remove debris and dead cells. The AMSC-MP then be dialyzed using a dialysis bag with a proper molecular-weight cut-off to replace the cell growth medium with the phosphate-buffered saline as the media [16, 17].


These molecules include basic fibroblast growth factor (bFGF), EGF, hyaluronic acid (HA), interleukins (IL-1 and IL-10), beta-defensins, TGF-β, elafin, human leukocyte antigen-G, matrix metalloproteinases, tissue inhibitors of metalloproteinases (TIMPs), and platelet-derived growth factor (PDGF) [15]. Additionally, amniotic tissue contains anti-inflammatory factors such as IL-1 and IL-10 receptor antagonists and regulators of catabolic enzymes such as TIMP1, TIMP2, TIMP3, and TIMP4. Furthermore, AMSC-MP is a potent down-regulator of TGF-β signaling, which stimulates the recruitment of fibroblasts and macrophages and upregulates collagen production [18].

#### Physicochemical properties and stability of AMSC-MP

AMSC-MP is a clear liquid containing various proteins, and some products form yellowish white lyophilized powder. A study by Kumala et al. (2020) compared the physicochemical stability of AMSC-MP in two forms, i.e., native AMSC-MP liquid and lyophilized AMSC-MP powder [19]. The results showed that AMSC-MP liquid began to change color after 7 days of storage at room temperature. In contrast, AMSC-MP liquid did not show color changes or an odor at cold temperatures. Its pH was 7–7.5 without significant changes during storage for 28 days [19].

AMSC-MP contains several proteins and the major component has a molecular weight of 75.33 kDa, as seen from the thick band in Fig. 1. A stability study measuring TGF-β with the enzyme-linked immunosorbent assay (ELISA) showed that the AMSC-MP liquid is less stable stored in room temperature (25°C) than when freeze-dried. Freeze-dried AMSC-MP has high crystallinity, which was supported by a scanning electron microscopy characterization showing a tetragonal crystal shape. The crystals have a melting point of 163.8 °C [19].

Specifically, stability data from each component of the AMSC-MP is still limited. It has been reported that the FGF1 and FGF2 lost most of their activity after 6 h at 37 °C [20]. Another study showed that the eye serum drops containing EGF, TGF-β1, and TGF-β2 remained stable in tube segments and vials made from polyvinyl chloride (PVC) with di-2-ethylhexyl phthalate (DEHP) and tris-2-ethylhexyl trimellitate (TOTM) as the plasticizers, respectively, for about 15 months by storage at



**Fig. 1** Qualitative determination of protein markers in fluid and freeze-dried (FD) AMSC-MP analysed using SDS-PAGE [19]

-30 °C. However, the stability was significantly decreased by storing samples at 4 °C and 22 °C, which could be reduced up to only 24 and 3 h for EGF and 48 and 9 h for TGF-β1 and TGF-β2, respectively. EGF was reportedly unstable at 37 °C [21]. Mizumachi & Ijima (2013) showed that the immobilized-VEGF prepared in a heparin-collagen substrate had higher stability than that of VEGF solution, resulting in about 43% of the initial VEGF retaining its biological activity, although after 16 days of pre-incubation at 37 °C [22]. Meanwhile, the research by Chen & Arakawa (1996) reported that the KGF is unstable in the aqueous media and it follows denaturation and aggregation when heated to 37 °C [23].

#### Biochemical components of AMSC-MP

##### bFGF

One of the growth factors in AMSC-MP is bFGF [7, 11, 18] (or FGF-2), which is a member of the FGF family that regulates various biological functions including proliferation, morphogenesis, and suppression of apoptosis

during development through a complex signal transduction system. bFGF is widely expressed in the nervous system, where it has multiple roles, and it supports the survival and growth of neuronal and neural stem cell cultures [24].

Members of the FGF family have a homologous core region of 120–130 amino acids arranged into 12 antiparallel strands ( $\beta 1$ – $\beta 12$ ) flanked by divergent amino and carboxyl functional groups. Generally, the sequence variation of the N- and C-terminal ends of specific FGF family members accounts for their differential ligand binding. The heparan sulfate glycosaminoglycan (HSGAG) binding site in the FGF core consists of a  $\beta 1$ – $\beta 2$  loop and parts of  $\beta 10$  and  $\beta 12$ . This section differs for each FGF family member and determines the endocrine properties of each. FGF binds to and activates the FGF receptor (FGFR) in the HSGAG-dependent tyrosine kinase receptor family. Upon ligand and HSGAG binding, FGFR dimerizes, allowing the cytoplasmic kinase domain to transphosphorylate and activate tyrosine loop A. Loop phosphorylation is followed by tyrosine phosphorylation in the C tail region, kinase insert, and juxtamembrane region. This process activates the Ras–mitogen-activated protein kinase (MAPK) and phosphoinositide 3-kinase–Akt signaling pathways [25].

In tissue regeneration therapy, FGFs from the paracrine group (FGF1–10, FGF16–18, FGF20, and FGF22) play a role. Paracrine FGFs have a high affinity for HSGAG, activating it and acting locally near the expression source.

## EGF

Another growth factor in AMSC-MP is EGF [11, 12, 17]. EGF is a small polypeptide mitogen present in many species that has been isolated and characterized in breast milk. EGF is a 6-kDa peptide derived from a 1207 amino acid precursor molecule that acts across a 170-kDa membrane glycoprotein receptor. EGF has intrinsic tyrosine kinase activity, like the IGF receptor, and functions in tyrosine kinase-mediated autophosphorylation. TGF- $\alpha$ , which has a 35% amino acid homology with murine EGF and 44% homology with human EGF, also acts through the EGF receptor [26].

Growth factors are naturally produced proteins that regulate cell proliferation, function, and differentiation through receptor signaling. EGF is one of the earliest known polypeptide growth factors and was the founder of the EGF-like family of proteins. EGF is an endogenous peptide that promotes cell growth, proliferation, and differentiation via ligand-receptor (EGFR) interactions [27, 28]. EGF was first isolated from the submaxillary glands of adult male rats by Cohen et al. in 1962. Currently, recombinant human EGF (rhEGF) can be mass produced from *Escherichia coli*, which has accelerated the

development of EGF formulations for treating skin conditions such as chronic wounds, burns, diabetic ulcers, and skin aging [28].

EGF has hydrophilic properties [29]. Structurally, EGF is a polypeptide chain with a molecular weight of 6045 Da consisting of 53 amino acids and 6 cysteine residues and has three intramolecular disulfide bonds [27]. EGF is characterized by the absence of three specific amino acid residues, lysine, alanine, and phenylalanine, and it lacks hexosamines and neutral sugars [30]. EGF has an optimal stability at pH 6.0–8.0 and an isoelectric point of about 4.6 [29]. EGF shows poor thermal stability because the protein structure begins to degenerate at 40 °C, and it has a transition midpoint at 55.5 °C. EGF can be completely denatured at temperatures above 76 °C [31].

EGF binds to EGFR; following ligand binding, EGFR (ErbB-1) dimerizes with itself or with a homolog, ErbB-2, ErbB-3, or ErbB-4, increasing intracellular tyrosine kinase activity. This process activates a signaling cascade that has multiple effects: cell proliferation, reduction of apoptosis, and angiogenesis [32].

## TGF- $\beta$

TGF- $\beta$  is an extracellular protein in AMSC-MP produced mainly by a subset of T cells [7, 11, 17]. TGF- $\beta$  belongs to a group of cytokines collectively referred to as the TGF- $\beta$  superfamily, whose members regulate epithelial cell growth, differentiation, motility, organization, apoptosis, and tumorigenesis [33]. The TGF- $\beta$  superfamily consists of a group of polypeptide morphogens. TGFs are divided into two subgroups: the TGF-like subgroup [TGF- $\beta$ s, activin, nodal, and multiple growth differentiation factors (GDFs)] and the BMP-like subgroup (BMP, GDF, and antimüllerian hormone). Members of the TGF-like subgroup exhibit functions in cell adhesion, growth, cytoskeletal organization, survival, proliferation migration, differentiation, chemotaxis, and immune cell activation in multicellular organism development [34].

Blood is the primary source of TGF- $\beta$ , which promotes healing and tissue regeneration during injury. Platelet aggregation and degranulation release high amounts of TGF- $\beta 1$  at wound healing sites. Additionally, recruited and activated leukocytes at wound sites secrete various cytokines, including TGF- $\beta 1$  to support the wound healing process [35]. TGF- $\beta$  is secreted in a latent or biologically inactive state. During cellular synthesis, the TGF- $\beta$  precursor undergoes intracellular proteolytic cleavage by furin endopeptidase, resulting in two proteins assembled into dimers via noncovalent associations.

TGF- $\beta$  signaling involves three parallel pathways, the bone morphogenic protein (BMP), TGF- $\beta$ , and activin

pathways, all of which are major regulators. TGF- $\beta$  signaling is transduced in cells by several SMAD protein modulators, which eventually enter the cell nucleus and influence the expression of target genes. Since all three pathways comprise ligands and receptors, the combination of different signals allows the regulation of many growth and developmental processes in highly specific ways [36].

### VEGF

VEGF regulates angiogenesis by inducing the proliferation, migration, and permeability of endothelial cells. VEGF is also found in AMSC-MP [7, 11, 17]. During the process of tissue regeneration, VEGF also plays an important role in cardiac repair by decreasing infarction size, reducing remodeling, decreasing endothelial cell apoptosis, supporting angiogenesis and neovascularization, increasing the number of mitotic cardiomyocytes in the border zone, and improving cardiac performance [37]. VEGF is produced by many cell types including tumor cells, macrophages, platelets, keratinocytes, and renal mesangial cells. VEGF activity is not confined to the vascular system; VEGF also plays a role in normal physiological functions such as bone formation, hematopoiesis, and wound healing [38].

The *VEGF* gene is located on chromosome 6p21.3 and is part of the *VEGF/PDGF* gene family, the cystine-knot superfamily of growth factors. Structurally, VEGF is a 40-kDa heterodimeric glycoprotein. VEGF contains a cystine-knot motif, characterized by disulfide bridges in the protein structure [39]. In humans, VEGF consists of several members: VEGF-A, which has several isoforms, VEGF-B, VEGF-C, VEGF-D, VEGF-E (VEGF virus), placenta growth factor (PlGF), and endothelial-derived VEGF [40].

There are three VEGF receptors: VEGFR-1, VEGFR-2, and VEGFR-3. Neuropilin-1 (NP-1) and neuropilin-2 (NP-2) coreceptors are non-tyrosine kinase receptors, and they selectively attach to certain VEGF subtypes or isoforms. The pro-angiogenic activity of VEGF occurs through the binding and activation of two receptor tyrosine kinases (TKs), which were initially identified as receptors for VEGF-A, namely VEGFR-1 and VEGFR-2. These receptors consist of seven extracellular Ig-like domains, a transmembrane domain, and an intracellular TK domain. Ligand binding induces receptor dimerization and phosphorylation. PlGF binds exclusively to VEGFR-1 with high affinity compared to VEGF-A and VEGF-B, and other family members also specifically bind VEGFR-1 [40].

PlGF was first identified in human placental tissue. It is involved in trophoblast growth and differentiation, trophoblast invasion, and blastocyst implantation. The

PlGF gene has four isoforms, PlGF-1 (PlGF131), PlGF-2 (PlGF152), PlGF-3 (PlGF203), and PlGF-4 (PlGF224), which differ in their molecular structure and biological properties. All isoforms have affinity for VEGFR-1. PlGF has no direct mitogenic effect and does not increase vascular permeability, but, under pathological conditions, binds to VEGFR-1, displaces VEGF-A from VEGFR-1, and allows binding of VEGF-A to VEGFR-2, indirectly enhancing the effect of VEGF-A. This increases vascular permeability, cell migration, and proliferation [40].

### KGF

Another growth factor reported to be present in AMSC-MP is KGF [7, 11, 17]. KGF is a monomeric polypeptide measuring 26–28 kDa and is an FGF family member [41]. KGF has been implicated in biological processes such as cell proliferation, development, and differentiation. KGF is encoded by the *FGF7* gene and is made up of 194 amino acids. The human KGF amino acid sequence in UniProtKB (P21781) indicates that this protein contains a signal peptide (residues 1–31) and a KGF chain (residues 32–194), and position 45 is glycosylated during posttranslational modification [41].

KGF has low stability in acidic and neutral pH conditions. Denatured KGF starts to aggregate at a moderate temperature. The calculated isoelectric point (pI) of KGF is 9.29. The pI is defined as the pH at which the molecule carries no electrical charge or is neutral on average. At a pH below the pI, proteins carry a positive charge, whereas above the pI, they carry a net negative charge [34, 35].

KGF has 19 negatively charged residues, such as Asp and Glu, and 29 positive charge residues, like Arg and Lys. KGF may be unstable under physiological conditions. The instability of the KGF protein is due to the repulsion between the positively charged residues. Therefore, the main cause KGF instability at a neutral pH is thought to be due to its high positive charge and repulsion forces, which lead to protein denaturing and irreversible aggregation [42].

Boroujeni et al. [43] investigated the stability of rhKGF, a truncated form of KGF, under different pH and temperature conditions using molecular dynamics simulation methods. That study showed that the stability of rhKGF increased with a decrease in the total charge at an alkaline pH and low temperature. KGF stability increased significantly at an alkaline pH of 8.5 and 9, presumably due to the presence of a high positive-charge residue. Acidic pH conditions caused instability and aggregation of KGF. In another study, KGF denatured at 400 °C. After a while, the protein began to aggregate, and irreversible particulate and precipitate formation occurred [44].

KGF is involved in various biological processes and has proliferation, antiapoptosis, cytoprotective, epithelial cell movement, and cytoskeletal reorganization effects. Additionally, its mitogenicity is useful for embryonic development, tissue patterning, cell growth, morphogenesis, wound healing, and tissue repair. As a growth factor, KGF acts through a variant of the receptor-2IIIb FGF (FGFR2b), which is expressed by epithelial cells [25]. The FGFR2b receptor is hereafter referred to as the KGF receptor (KGFR). KGFR is part of the FGFR family of receptor TKs, which are activated in the presence of heparin/ HSGAG. The binding of KGF to KGFR requires heparin/HS as a coreceptor. This process includes the dimerization of KGFR and activation of its kinase domain, inducing autophosphorylation of the receptor [45]. To bind to the receptor and its ligands, KGF has a positively charged site called the heparin binding site, and a neutral site that binds to KGFR.

The proliferative effect of KGF occurs due to activation of the RAS and Raf/MAPK/ERK pathways after KGFR dimerization and phosphorylation of the tyrosine kinase domain [46]. Its antiapoptotic effect occurs due to activation of p21-activated kinase 4 (PAK4), which then activates the antiapoptotic-Akt-dependent pathway by recruiting PI3K, which then arranges antiapoptotic genes [47]. Activation of the antiapoptotic pathway by the ERK1/2 pathway reduces the cellular inflammatory response by inducing the expression of the cytoprotective genes nuclear factor erythroid 2-related factor 2 (*NRF2*) and heme oxygenase-1 (*HO1*) in epithelial cells [48, 49].

KGF regulates epithelial cell motility and cytoskeletal reorganization through activation of Src-Cortactin, which phosphorylates paxillin and activates GTPases such as Rho, Rac, and Cdc42, resulting in lamellipodia extension, actin increase, and cell mobility and migration [48]. During mitogenicity, autophosphorylation of the tyrosine kinase domain of KGFR results in the activation of phosphatidylinositol hydrolysis. PLC $\gamma$  activation hydrolyzes phosphatidylinositol-4,5-diphosphate to inositol-1,4,5-triphosphate and diacylglycerol, which stimulate protein kinase C and increase intracellular Ca<sup>2+</sup> and subsequent mitogenic activity through upregulation of target genes [50]. Currently, palifermin, a recombinant preparation of human KGF, is a growth factor cocktail used for therapy. For example, a study by Spielberger et al. [51] showed that giving 60 g per kilogram of body weight of palifermin daily reduces the duration and severity of oral mucositis after intensive chemotherapy and radiotherapy for hematological cancer.

Based on the explanation above, it can be seen that various kinds of growth factors present in AMSC-MP, namely bFGF, VEGF, TGF- $\beta$ , EGF and KGF have their respective functions and activities. In addition, each growth factor is formed by different proteins resulting in molecules with different physicochemical properties and stability. This knowledge will assist in the manufacturing process and application of AMSC-MP as a therapeutic agent.

## **Biological activities of AMSC-MP for tissue regeneration**

### **Wound healing**

The wound repair process is divided into four main phases: hemostasis, inflammation, proliferation, and dermal remodeling [52]. Wound repair begins with the hemostasis phase, in which platelet formation blood loss and the initial fibrin matrix begins to form. Platelets are critical in the recruitment of immune cells to wound tissue, either by capturing immune cells directly or by releasing chemokine secretomes. Platelet secretomes also contain growth factors that stimulate resident skin cells, including fibroblasts and keratinocytes [52].

Furthermore, in the inflammation phase, necrotic cells and damaged tissue release damage-associated molecular patterns and resident immune cells, such as mast cells, Langerhans cells, T cells, and macrophages, respond by activating inflammatory pathways. In this phase, proinflammatory cytokine and chemokine release attracts leukocytes in the circulation to the injured tissue. Monocytes already in the wound tissue differentiate into macrophages. Macrophages engulf necrotic cell remnants and pathogenic material [52]. Activation of macrophages is influenced by proinflammatory stimuli, such as lipopolysaccharide and interferon-gamma (IFN- $\gamma$ ), and their activation enhances inflammation by releasing reactive oxygen species (ROS), inflammatory cytokines (e.g., IL-1, IL-6, and TNF-) and growth factors (e.g., VEGF and PDGF).

Macrophages control the degradation of extracellular connective tissue by enzyme secretion and phagocytosis and regulate wound matrix remodeling through the production of growth factors such as PDGF, TGF, ILs, and TNF [53]. These growth factors influence the regrowth process, epithelialization, fibroplasia, and angiogenesis [52, 54]. All three stages occur in the proliferation phase, starting when keratinocytes migrate to close the wound, followed by angiogenesis, and then fibroblasts replace the initial fibrin clot with granulation tissue [55]. Fibroplasia begins about 5 d after injury and continues for 2 weeks. Fibroblasts migrate into the wound and replicate in response to mediators released during inflammation. These mediators include C5a, fibronectin, PDGF,

FGF, and TGF $\beta$ 1. Remodeling of the extracellular matrix (ECM) spans the entire injury response, beginning with the initial deposition of a fibrin clot and ending several years later with the formation of a mature, type I collagen-rich scar.

Two types of FGF members play an important role in the wound healing process, among which the most important are KGF and bFGF. Both are present in AMSC-MP. Qu et al. (2018) showed that the combination of KGF and bFGF in a collagen delivery system increased cell migration in the wound healing process, accelerating wound closure [56]. bFGF improved wound healing in animal models and clinical studies. KGF is a cytokine that exerts a specific mitogenic effect in epithelial cells. This effect has been reported to be a key factor in wound healing, as it is weakly expressed in human skin but is strongly upregulated after skin injury [56].

The synergistic effect of bFGF and KGF can be observed during the wound healing process. Re-epithelialization begins within hours of injury, and bFGF and KGF promote cell proliferation. KGF stimulates keratinocyte migration, while bFGF promotes fibroblast migration and stimulates the production of collagenase, suggesting that bFGF and KGF have complementary roles in wound healing. Together, bFGF and KGF may also stimulate the accumulation of vascularization-related cells. However, other studies have shown that KGF affects ongoing inflammation and scar formation [57]. These negative effects can be minimized by bFGF as an antiscarring agent [58].

bFGF in wound healing was also investigated by Zhang et al. (2018), in which hydrogels combined with bFGF increased the wound healing process. Hydrogels derived from gum arabic, pectin, and divalent calcium ions help increase the stability of FGFs and provide a sustained release effect [59]. Fibroblast scratch assays showed that the hydrogel FGF formulation could close a wound within 12 h, while in controls, wound closure only started at the 12<sup>th</sup> hour. In an in vivo study conducted by creating a full-thickness skin incision on the back of mice, the hydrogel treatment group with bFGF showed the fastest wound closure compared to the negative control group [59].

Furthermore, the metabolite products of stem cells have a high potential for wound healing efficacy, based on preclinical and clinical studies on stem cells. The use of human AMSCs for wound healing has also been investigated in vivo in male mice in a heat-induced apoptosis model [60]. The administration of  $2 \times 10^6$  cells injected subcutaneously into the wounded skin showed accelerated re-epithelialization. Wound closure occurred on day 7, and cytokines, including PAI-1, C-GSE, periostin, and TIMP-1, have been reported to activate the PI3K/

AKT pathway, which plays a vital role in epithelial cell and dermal fibroblast migration and proliferation. In another study, the use of human adipose-derived MSCs and placenta-derived MSCs in amniotic membrane grafts accelerated wound healing in Wistar rats with an excisional wound splinting model at day 7 [61]. The processes of re-epithelialization, collagen remodeling, and neovascularization occur more quickly by embedding these cells on the amniotic membrane as wound dressing, and this membrane produces various growth factors such as TGF- $\alpha$ , TGF- $\beta$ , bFGF, EGF, and KGF, cytokines such as IL-4, IL-6, and IL-8, as well as matrix metalloproteinase inhibitors. A review by Huang et al. (2020), which explored the use of MSCs in preclinical and clinical studies for wound healing, showed that metabolite products play an active role in tissue repair and wound healing [53]. Tissue repair occurs through the stimulation of cell differentiation and paracrine action, involving growth factors such as bFGF, hepatic growth factor (HGF), EGF, KGF, VEGF, and TGF- $\beta$ , and also cytokines such as IL-10, to reduce inflammation and accelerate the angiogenesis, granulation, re-epithelialization, and wound closure processes.

### Potential uses of AMSC-MP in gastrointestinal injury therapy

In the gastric mucosa, TGF- $\alpha$  controls cell proliferation under normal conditions and after acute injury, while EGF controls cell proliferation during the healing of chronic ulcers. When the gastric mucosa is injured, growth factors predominantly restore the epithelial component, while bFGF and VEGF promote restoration of the connective tissue and angiogenesis in the injured mucosa. Granulated connective tissue, which grows under the stimulation of bFGF and VEGF, is the primary source for regenerating connective tissue lamina propria and microvessels within ulcer scars. Other growth factors such as insulin-like growth factor, KGF, hepatocyte growth factor, and trefoil peptides also act in gastrointestinal (gastric ulcers, colitis) regeneration following injury [62].

Research by Wei et al. (2022) showed that KGF in combination with polydopamine (PDA) and HA nanoparticles successfully prevented abdominal adhesions and promoted the repair of mesothelial cells in the injured peritoneum [63]. More importantly, PDA-KGF NPs combined with HA reduced collagen deposition and fibrosis and inhibited the inflammatory response [63].

KGF function is determined by phosphorylation of the protein tyrosine kinase SRC. When KGF binds to its receptor, SRC is phosphorylated by KGFR. In a study, we evaluated the phosphorylation level of rat Src in the injured peritoneum 7 d after surgery. The levels of phospho-Src protein in rat peritoneal tissue were higher in the

PDA-KGF NP treatment group compared to that treated with KGF alone. Thus, the *in vivo* positive effect of KGF is prolonged when KGF and PDA are administered as PDA-KGF nanogels [63].

KGF is also effective for treating ulcerative colitis. Ying-Zheng et al. (2019) reported that KGF encapsulated into neutrophil-like liposomes (KGF-Neus) effectively restored intestinal morphology and function in ulcerative colitis [64] because the neutrophil membrane vesicle (NEM) associated protein, KGF-Neus, is specifically internalized to the area of inflammation [64].

KGF and its receptors are present in the human fetal gastrointestinal tract, and *in vitro* stimulation of human fetal enterocytes with KGF results in cellular proliferation. KGF expression is increased in patients who undergo surgery for inflammatory bowel disease and is correlated with the degree of intestinal inflammation. In animal models of colitis, KGF administration reduces the degree of mucosal injury [65]. Recombinant KGF treatment has been studied for use in ulcerative colitis; in a clinical phase II study, recombinant KGF failed to induce remission in ulcerative colitis patients, but the maximal therapeutic dose used may have been too low [66].

Another AMSC-MP component, EGF, is associated with mucosal ulcer disease. Decreased EGF levels are associated with mucosal ulcer disease. Patients with duodenal ulcer disease also have decreased EGF levels. EGF supplementation promotes mucosal repair and regeneration in several conditions. In experiments in pigs, EGF significantly reduced esophageal ulceration, structural formation, and mucosal histological damage associated with sclerotherapy. In rats with gastric ulcers, orogastric EGF administered in combination with sucralate improved ulcer healing [67]. A small human study showed that treatment with intravenous EGF promoted better gastric ulcer healing compared with the antiulcer treatment cetraxate hydrochloride [67].

### Potential use of AMSC-MP in lung injury treatment

Growth factors are involved in all aspects of lung development. The spatial and temporal distribution of FGF10 in the lung determine the airway branching pattern. Some factors participate in more specific developmental programs, such as VEGF in blood vessel formation and FGF7 in type II alveolar cell differentiation [55]. KGF is a critical growth factor in lung development and is protective after lung injury. KGF is an important growth factor for local resident progenitor epithelial cell repair and for mobilization and enhanced engraftment of cytokeratin 5 circulating epithelial progenitor cells, which contributed to the repair of the proximal airway epithelium in a mouse model of syngeneic tracheal transplantation to the injured proximal airway epithelium [68].

KGF induces epithelial cell proliferation and protects against acute lung injury. Leblond et al. (2007) showed that 1 mg/kg of body weight of KGF given intravenously to rats injected with albumin as an asthma trigger reduced extravascular lung water levels. KGF treatment also reduced the number of inflammatory cells in the bronchoalveolar lavage fluid but not in the bronchial mucosa. KGF reduces allergen-induced changes in epithelial integrity and the expression of the intercellular junction proteins catenin and zonular occludens protein-1 [69].

Consistently, Wang et al. (2020) investigated the effect of KGF on the release of inflammatory-related cytokines by damaged bronchial epithelial cells. Compared with the healthy group, KGF and KGFR expression and apoptosis were significantly increased in asthmatic patients. An *in vitro* study showed that KGF treatment limited IFN- $\gamma$  and TNF- $\alpha$ -induced apoptosis by inhibiting apoptotic markers in the TNF signaling pathway. KGF limits the release of TSLP, IL-25, and IL-33 by damaging 16HBE 14o cells. In contrast, KGF promotes intercellular adhesion and wound closure of cultured 16HBE 14o cells through increased expression levels of the intercellular junction proteins ICAM-1,  $\beta$ -catenin, E-cad, and Dsc3. In summary, KGF and KGFR may aid bronchial epithelial cell repair in asthma by inhibiting epithelial cell apoptosis while promoting epithelial cell proliferation and migration [70].

TGF- $\beta$  is associated with acute lung injury. Research by Kan et al. 2014 showed an increase in the expression of TGF- $\beta$  in rat serum induced with paraquat, a compound that irritates the lungs [71]. *TGF- $\beta$ 1* mRNA expression in rat lungs was also significantly increased. Many inflammatory cells were observed infiltrating the alveoli of the injured lungs. The abnormal expression of *TGF- $\beta$ 1* was hypothesized to be important in the pathogenesis of chronic inflammatory and immune lung diseases, including asthma, chronic obstructive pulmonary disease, and pulmonary fibrosis [72]. In the future, cytokines and their inhibitors may provide new therapies for treating acute lung injury and pulmonary fibrosis.

Heparin-binding EGF-like growth factor (HB-EGF) reduces inflammation, maintains intestinal barrier function, and protects the lung from acute injury in several models of intestinal injury. Another study investigated the impact of HB-EGF by comparing burn-treated mice (25% of total body surface area) with burn-infected mice after two enteral doses of HB-EGF (1200 mg/kg/dose) [73]. The control mice had increased pulmonary myeloperoxidase levels, lung and spleen apoptosis, airway resistance and bronchial reactivity, and intestinal permeability.



These effects were significantly reduced in burn-injured mice treated with enteral HB-EGF [73].

FGF2 is closely involved in endothelial cell migration, proliferation, and injury repair. Recombinant FGF2 was injected peritoneally at a 0.1 mg/kg dose in septic mice induced by ligation and cecal puncture. FGF2 treatment reduced the inflammatory response, attenuated pulmonary capillary leakage, reduced lung injury, and increased survival in septic mice. Endothelial injury and macrophage inflammation induced by LPSs are inhibited by FGF2 administration via the AKT/P38/NF- $\kappa$ B signaling pathway [74].

### Potential use of AMSC-MP in bladder and renal injuries

Surgical and traumatic injuries to the bladder initiate a complex series of biological processes that result in wound healing. This involves cellular proliferation, migration, and differentiation; removal of damaged tissue; and production of extracellular matrix, all of which may be controlled by growth factors. KGF is induced in the skin following incisional injury. During the early phases of bladder wound healing, mRNA levels of *KGF* and *TGF- $\alpha$*  increased, and exogenous KGF directly affected urothelial proliferation [75].

Among the growth factors that affect the bladder are KGF and FGFs; KGF increases cyclophosphamide-induced bladder injury. Cyclophosphamide is often used to treat cancer and rheumatic and kidney diseases. Acrolein, its metabolite, is a toxic metabolite concentrated in urine that can cause acute hemorrhagic cystitis (7%–45% incidence rate) and urothelial cancer (4%–15% incidence rate, depending on the dose). In cyclophosphamide-induced urothelial injury, increased apoptosis of intermediate and basal cells was observed. KGF prevented apoptosis of deeper urothelial cells (UPK3 + intermediate and KRT5 + intermediate/basal cells), likely via activation of AKT [76].

Evidence suggests that KGF regulates bladder cell development and function and is directly responsible for urothelium proliferation. In a study by Narla et al. (2020), mice were given a 5 mg/kg injection of KGF dissolved in PBS 24 h before cyclophosphamide injection and showed increased urothelial regeneration compared to controls [76]. KGF pretreatment blocked cyclophosphamide-induced intermediate and basal cell apoptosis, possibly via phosphorylated AKT, and promoted ERK-mediated phosphorylated KRT5 + /KRT14 – cell proliferation, leading to urothelial regeneration. The effect of KGF on bladder injury was also found in a study by Jaal et al. (2007) [77], in which there was an increased positive response on day 2 in 50% of mice after a single injection of palifermin at a dose of 15 mg/kg [77].

In addition to KGF, bFGF also affects bladder injury. Chen et al. (2010) explored the ameliorative effect of collagen-based bFGF for bladder regeneration in a mouse model [78]. A bladder with a subtotal cystectomy was grafted with collagen membranes coupled with 0.56 nmol of CBD-bFGF. As a result, collagen/bFGF mice had faster collagen scaffold degradation and better bladder wall cell growth but no bladder stone formation [78].

In addition to bladder injury, FGF positively affects acute kidney disease. Zhou et al. investigated the effects of FGF2 in acute kidney disease using Sprague–Dawley and NRK-52E cells [79]. FGF2 significantly increased tissue apoptosis in acute kidney disease by inhibiting excessive ER stress. Moreover, FGF2 also reduced ER overstress and apoptosis in cultured NRK-52E cells injured with tert-butyl hydroperoxide [80].

### Potential uses of AMSC-MP for bone regeneration

The bone response to injury begins with an inflammatory phase. Bleeding from the fracture-surrounding soft tissue results in forming a fibrin clot and fracture hematoma. Subsequently, inflammatory cytokines are released, inducing angiogenesis and mesenchymal progenitor cell proliferation. These mesenchymal progenitors rapidly proliferate, forming an initial soft callus. The soft or primary callus response occurs within two weeks of injury. The degree of callus formation is proportional to the degree of motion at the fracture gap [81].

In the second stage of repair, the necrotic bone ends undergo resorption, and the mesenchymal progenitor cells proliferating at the injury site begin to differentiate into chondrocytes to form a cartilaginous callus and osteoblasts for intramembranous bone formation at the fracture margins. The mechanisms that control the influx, proliferation, and differentiation capacity of mesenchymal progenitor cells are critical components of the fracture healing process. For instance, BMP and Wnt signaling play roles in mesenchymal cell differentiation into osteoblasts and chondrocytes [81].

During the establishment and maturation of the soft callus, growth factors (e.g., TGF- $\beta$ , PDGF, GDF-5, FGF-1, and IGF-II) and hormones (e.g., PTHrP) are involved in the recruitment and proliferation of fibroblasts and MSCs. They also play an essential role in inducing MSC differentiation into osteoblasts or chondrocytes [82]. Once chondrocytes form, endochondral ossification occurs, and a hard callus is formed via woven bone in the third stage of healing. Whether this endochondral bone formation process is equivalent to that which occurs during bone growth is not completely understood [83]. The use of human AMSC in bone tissue regeneration has been reported in several studies, including in vitro and in vivo studies and clinical trials, as summarized

by Li et al. (2020) [84]. AMSC treatment in models of collagen-induced arthritis, intervertebral disc degeneration, rheumatoid arthritis, and osteoarthritis has shown anti-inflammatory, angiogenic, and immunomodulatory effects, all of which play important roles in tissue remodeling. Several growth factors and cytokines present in human AMSC play an important role in the regeneration process of bone defects. These include HGF, FGF7, BMP-2, VEGF, IL-6, and IL-8. Clinical trials in patients with bone defects of human AMSC administration either by subcutaneous injection or implantation into the hypodermis with a polymer or scaffold have shown increased proliferation, and osteoblastic differentiation of BMSCs increased osteogenesis and endogenous bone regeneration.

Itoh et al. (2007) confirmed that FGF/FGFR signaling plays a role in osteogenesis. FGF/FGFR signaling does not directly induce osteoblast differentiation but modulates it. FGF2 and FGF9 likely induce the proliferation of osteoblast cell lineages and the induction of angiogenesis, and FGF18 promotes osteoblast differentiation [85].

Wilkie et al. (2005) published an in vitro analysis of bone marrow-derived MSCs in which FGF18 enhanced osteoblast differentiation by activating FGFR1 or FGFR2 signaling [86]. Additionally, overexpression of FGF18 by lentiviral infection or direct addition of FGF18 to culture media induced the expression of osteoblast marker genes in C3H10T1/2 fibroblastic cells. Treatment with FGF18 in mouse-derived MSCs under differentiation-inducing conditions showed increased expression of osteoblast differentiation markers and mineralization [86].

Low-dose FGF18 treatment with osteogenic induction of bone morphogenetic protein 2 (BMP2)-dependent bone protein from MC3T3-E1 cells increased mineralization, whereas high-dose treatment inhibited the process. Additionally, FGF18-soaked heparin-coated acrylic beads accelerated osteoblast differentiation in mouse fetuses by regulating BMP2 expression in 90 osteoblast cell lineage cells [87]. FGF2 stimulates mitosis and cell proliferation, including of fibroblast and endothelial cells, which plays a vital role in maintaining these cells in tissue repair processes [88]. On the other hand, FGF18 stimulates cellular osteogenesis through the upregulation of bone morphogenetic proteins.

### Potential uses for skin rejuvenation in skin aging

Together with other growth factors and cytokines, EGF directly affects collagen, elastin, and ECM biosynthesis, but its binding and signaling diminish with age. Aged cells in the skin produce ROS, and the mitochondria of these cells disrupt tissue complexes by cleaving membrane-bound receptors, ECM proteins, growth factors, and other signaling ligands in the dermal

microenvironment [89]. Reduced EGF binding and signaling with age can cause collagen degradation in the skin [90, 91]. The rapid degradation of collagen in the skin leads to loss of elasticity and the appearance of skin wrinkling [92]. Decreased expression of EGFR also occurs in aging dermal fibroblasts in the ECM, is associated with reduced cell migration and proliferation, and ultimately leads to skin flexibility and elasticity loss. EGF helps reduce the effects of aging by supporting skin regeneration by stimulating cell renewal through the interaction of keratinocytes and fibroblasts. EGF plays an important role in forming fibroblasts in the dermis by stimulating collagen production via activation of EGFR [89]. The topical use of growth factors is a safe and effective medical treatment [93]. Applying EGF to aging skin can increase fibroblast proliferation [91]. Thus, EGF is a potential therapeutic antiaging agent for the skin.

A clinical trial of human umbilical cord-derived MSC-conditioned media administered by microneedle resulted in good efficacy as an antiaging product and provided an excellent potential for skin rejuvenation [94]. That study reported that the tested conditioned media contained growth factors, including EGF, VEGF-A, VEGF-D, HGF, FGF-2, and others. Furthermore, the administration of MSCs reduced the melanin index and brown spots on the skin. Additionally, wrinkles and skin pores were reduced, and there was an increase in skin elasticity, indicating an improvement in facial skin texture. Another study showed that administering amniotic fluid MSC-derived conditioned media with microneedles to the face improved the skin texture and reduced wrinkles [95]. Moreover, histologically, there was an increase in the number of dermal collagen bundles arranged more regularly, elastic fibers, and epidermis thickening.

AMSC-MP possess various growth factor and cytokines that enables them to become the agent therapeutics of many therapy.

### Future prospects for AMSCMP for tissue engineering

Many reports have demonstrated the potential use of AMSC-MP for tissue regeneration to improve the appearance of facial skin, in bone regeneration, and in tissue or organ repair. Preclinical reports, including in vitro and in vivo studies, have shown that growth factors and cytokines play an essential role in the tissue repair process, both through stimulation of cell differentiation and proliferation and via an indirect effect on regeneration, including antiinflammatory and angiogenic effects. Clinical trials using conditioned media and MSCs have also reported potent activity for tissue regeneration.

However, protein delivery systems of the cytokines and growth factors present in AMSC-MP still have many

shortcomings. Oral delivery is generally preferred, but oral delivery is not a viable method for proteins due to their poor absorption and degradation in the gastrointestinal tract and liver [96]. Oral administration of protein drugs leads to very low bioavailability [97]. Thus, the development of protein formulations is needed to overcome the low permeability of large molecules, the lack of lipophilicity, and rapid inactivation or enzymatic degradation in the gastrointestinal tract as well as protein physicochemical properties that are limiting [97]. For topical uses, growth factors have molecular weights higher than 500 Da, which makes the penetration of the stratum corneum difficult [91]. The ideal characteristics of a substance for a topical delivery system include a relatively low molecular weight (<500 Da), a low melting point (<200 °C), moderate lipophilicity (log P 1–3), and high water solubility (>1 mg/mL), as well as high pharmacological potential [98]. Therefore, a delivery system is needed to help these molecules penetrate the dermis.

The use of AMSC-MP is also limited due to its sensitivity to environmental factors such as temperature, pH, and reactivity during reconstitution. Because of this, the delivery system must optimize the growth factor dose, route of administration, and release kinetics for the safe and effective use of growth factors [99]. For example, KGF has poor *in vivo* bioactivity. KGF has a short biological half-life and poor stability, its biological activity is susceptible to environmental factors, and it cannot maintain bioactivity for a long time in the presence of other enzymes [45].

In addition to stability, using AMSC-MP for tissue regeneration therapy requires a carrier system capable of local delivery with the controlled release of growth factors. The uncontrolled release of growth factors can cause side effects. Using biomaterials as delivery systems is the most successful strategy for controlled delivery, and they have been developed into various commercially available systems [100]. Based on the explanation above, it can be seen that AMSC-MP has been widely studied for its benefits in various diseases. Thus AMSC-MP has potential as a tissue regeneration therapeutic agent.

### **Manufacturing of AMSC-MP for therapeutic products**

Presently, autologous cell therapy is primarily used for stem cell-based therapy. However, the small individual scale, insufficient reagents, and inefficient manufacturing process result in an expensive product. The proper indication for use and timing, adequate dosage, and appropriate route of administration still need to be determined for widespread use.

The possible scaling up of AMSC-MP would represent a more efficient means of manufacturing mass-scale

products to induce tissue regeneration of greater suitability to community needs rather than individual therapy. Moreover, it would reduce the possibility of immune system rejection of cell therapy. The process of manufacturing AMSC-MP from AMSCs is divided into at least three main stages: isolation of mesenchymal cells from placental tissue, cell culture and incubation, and the harvesting and purification of the metabolite products of cell cultures. Quality assurance involving validation needs to be carried out to identify a sustainable production process and guarantee the quality of AMSC-MP. The validation process includes cell culturing, cell stock storage, the harvesting of conditioned media containing cell metabolites, and their processing into AMSC-MP as bioactive materials. AMSC-MP standardization will then be required to ensure consistent and reliable product quality. Quality parameters such as physicochemical characteristics and growth factor content would constitute important specification parameters of bioactive materials. Through validation, mass-production is possible using tailor-made cell lines under controlled laboratory conditions which provide a high-quality source of bioactive factors necessary to produce mass products efficiently and safely.

Furthermore, large-scale manufacturing should be developed to minimize costs and provide affordable therapeutic products for the public; however, the standardization of AMSC-MP into biological products has substantial challenges. Process validation, quality control, and standardized protocols for isolating, culturing, and cultivating AMSC-MP are required, and these are major challenges for manufacturers. Moreover, regulations that support clinical use are also being developed, so AMSC-MP mass production remains under development. Despite these challenges, developing AMSC-MP-based therapeutic products is worthwhile for providing affordable advanced biological therapeutics for clinical practice.

### **Conclusions**

AMSC-MP has excellent potential for use in tissue regeneration therapy since it contains a variety of growth factors that provide better efficacy than single growth factors or cytokines alone. *In vitro* and *in vivo* preclinical studies have shown that AMSC-MP has biological activities related to wound healing, the repair of bone defects and other bone diseases, tissue repair of damaged organs, including the lungs, bladder, kidneys, and in the gastrointestinal tract, as well as skin rejuvenation related to antiaging effects, providing excellent efficacy and demonstrating a good safety profile. Some reports have shown satisfactory results in clinical trials in patients with certain diseases. There is a high clinical demand for

AMSC-MP as an alternative biological therapy, but further development is needed regarding its stability and the identification of a delivery system to provide maximum efficacy. Additionally, product development from the laboratory to a mass production scale requires further effort. Moreover, quality assurance is needed for biological product materials and complex manufacturing processes, which is the main challenge that must be addressed to optimize the use of AMSC-MP as a therapeutic agent.

#### Abbreviations

AMSC	Amniotic Membrane Stem Cell
AMSC-MP	Amniotic Membrane Stem Cell Metabolite Products
BAL	Bronchoalveolar lavage
BMP	Bone morphogenic protein
EGF	Epidermal growth factor
ELISA	Enzyme-linked immunosorbent assay
FD	Freeze-dried
GDF	Growth differentiation factors
HA	Hyaluronic acid
HGF	Hepatic growth factor
IGF	Insulin-like growth factor-I
ILD	Interstitial lung diseases
KGF	Keratinocyte growth factor
KGFR	KGF receptor
MAPK	Mitogen-activated protein kinase
MSC	Mesenchymal stem cells
PDGF	Platelet-derived growth factor
PIGF	Placenta growth factor
ROS	Reactive oxygen species
TGF	Transforming growth factor
TK	Tyrosine kinases
VEGF	Vascular endothelial growth factor

#### Acknowledgements

Not applicable.

#### Authors' contributions

Andang Miatmoko: 1) Conception and design of the work, data acquisition, data analysis, and interpretation; 2) drafting the article; 3) critically revising the article for important intellectual content; 4) final approval of the version to be published; 5) agrees to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of the work are appropriately investigated and resolved. Berlian Sarasitha Hariawan: 1) Conception and design of the work, data acquisition, data analysis, and interpretation; 2) drafting the article; 3) final approval of the version to be published; 4) agrees to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of the work are appropriately investigated and resolved. Devy Maulidya Cahyani: 1) Conception and design of the work, data acquisition, data analysis, and interpretation; 2) drafting the article; 3) final approval of the version to be published; 4) agrees to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of the work are appropriately investigated and resolved. Syarifah Sutra Dewangga: 1) Data analysis and interpretation; 2) drafting the article; 3) final approval of the version to be published; 4) agrees to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of the work are appropriately investigated and resolved. Kevin Ksatria Handoko: 1) Data analysis and interpretation; 2) drafting the article; 3) final approval of the version to be published; 4) agrees to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of the work are appropriately investigated and resolved. Purwati: 1) Data analysis and interpretation; 2) critically revising the article for important intellectual content; 3) final approval of the version to be published; 4) agrees to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of the work are appropriately investigated and resolved. Ram Kumar Sahu: 1) Data analysis and interpretation; 2) critically

revising the article for important intellectual content; 3) final approval of the version to be published; 4) agrees to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of the work are appropriately investigated and resolved. Dewi Melani Hariyadi: 1) Data analysis and interpretation; 2) critically revising the article for important intellectual content; 3) final approval of the version to be published; 4) agrees to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of the work are appropriately investigated and resolved. All authors reviewed the manuscript. The author(s) read and approved the final manuscript.

#### Authors' information

None.

#### Funding

This study was supported by Applied Research on Health (Penelitian Terapan Bidang Kesehatan) Grant No. 1554/UN3.15/PT/2021 provided by the Ministry of Education, Culture, Research and Technology of the Republic of Indonesia.

#### Availability of data and materials

Not applicable.

#### Declarations

#### Ethics approval and consent to participate

Not applicable.

#### Consent for publication

Not applicable.

#### Competing interests

The authors declare no competing interests.

Received: 9 September 2022 Accepted: 6 February 2023

Published online: 09 February 2023

#### References

1. Yardeni D, Ghany MG. Review article: hepatitis B-current and emerging therapies. *Aliment Pharmacol Ther.* 2022;55(7):805–19.
2. Checkoway H, Lundin JI, Kelada SN. Neurodegenerative diseases. *IARC Sci Publ.* 2011;163:407–19.
3. Tomé-Bermejo F, Piñera AR, Alvarez L. Osteoporosis and the management of spinal degenerative disease (II). *Arch Bone Jt Surg.* 2017;5(6):363–74.
4. Cummings JL, Tong G, Ballard C. Treatment combinations for Alzheimer's disease: current and future pharmacotherapy options. *J Alzheimer's Dis.* 2019;67(3):779–94.
5. Yiannopoulou KG, Papageorgiou SG. Current and future treatments in Alzheimer disease: An update. *J Cent Nerv Syst Dis.* 2020;12:117957352090739.
6. Cummings J, Lee G, Ritter A, Zhong K. Alzheimer's disease drug development pipeline: 2018. *Alzheimer's Dement Transl Res Clin Interv.* 2018;2018(4):195–214.
7. Ding M, Tegel H, Sivertsson Å, Hober S, Snijder A, Örmö M, et al. Secretome-based screening in target discovery. *SLAS Discov.* 2020;25(6):535–51.
8. Rahmadewi R, Retha R, Pitasari DA, Kusumastanto VA, Ardhaninggar AAA, Citrashanty I, et al. The efficacy of amniotic membrane stem cell (AMSC) metabolite product and vitamin E for wrinkles, spots, and pores in photoaging. *Dermatol Res Pract.* 2020;2020:1–5.
9. Kropp C, Massai D, Zweigerdt R. Progress and challenges in large-scale expansion of human pluripotent stem cells. *Process Biochem.* 2017;59:244–54.
10. Herberts CA, Kwa MSG, Hermesen HPH. Risk factors in the development of stem cell therapy. *J Transl Med.* 2011;9(1):29.
11. Plotnikov EY, Silachev DN, Popkov VA, Zorova LD, Pevzner IB, Zorov SD, et al. Intercellular signalling cross-talk: to kill, to heal and to rejuvenate. *Hear Lung Circ.* 2017;26(7):648–59.

12. Rikhtegar R, Pezeshkian M, Dolati S, Safaie N, Afrasiabi Rad A, Mahdipour M, et al. Stem cells as therapy for heart disease: iPSCs, ESCs, CSCs, and skeletal myoblasts. *Biomed Pharmacother*. 2018;2019(109):304–13.
13. Tamama K, Kerpedjjeva SS. Acceleration of wound healing by multiple growth factors and cytokines secreted from multipotential stromal cells/mesenchymal stem cells. *Adv Wound Care*. 2012;1(4):177–82.
14. Huddleston HP, Cohn MR, Haunschild ED, Wong SE, Farr J, Yanke AB. Amniotic product treatments: Clinical and basic science evidence. *Curr Rev Musculoskelet Med*. 2020;13(2):148–54.
15. Prakoeswa CRS, Pratiwi FD, Herwanto N, Citrashanty I, Indramaya DM, Murtiastutik D, et al. The effects of amniotic membrane stem cell-conditioned medium on photoaging. *J Dermatolog Treat*. 2019;30(5):478–82.
16. Damayanti RH, Rusdiana T, Wathoni N. Mesenchymal stem cell secretome for dermatology application: A review. *Clin Cosmet Investig Dermatol*. 2021;14:1401–12.
17. Teixeira FG, Serra SC, Salgado AJ. Tips on how to collect and administer the mesenchymal stem cell secretome for central nervous system applications. *Methods Mol Biol*. 2016;1416:457–65.
18. Bennett NT, Schultz GS. Growth factors and wound healing: Biochemical properties of growth factors and their receptors. *Am J Surg*. 1993;165(6):728–37.
19. Sari DIK, Erawati T, Miatmoko A, Prakoeswa CRS, Soeratri W. Characterization and stability study of amniotic membrane stem cell metabolite product (AMSC-MP). *Int J Pharma Res Heal Sci*. 2020;8(1):3126–30.
20. Chen G, Gulbranson DR, Yu P, Hou Z, Thomson JA. Thermal stability of fibroblast growth factor protein is a determinant factor in regulating self-renewal, differentiation, and reprogramming in human pluripotent stem cells. *Stem Cells*. 2012;30(4):623–30.
21. Tan JCG, Webb RG, Marks DC. Serum growth factor stability in different eye drop packaging systems during storage. *Transfus Apher Sci*. 2020;59(1):102608.
22. Mizumachi H, Iijima H. Measuring stability of vascular endothelial growth factor using an immobilization technique. *Adv Biomed Eng*. 2013;2:130–6.
23. Chen BLU, Arakawa T. Stabilization of recombinant human keratinocyte growth factor by osmolytes and salts. *J Pharm Sci*. 1996;85(4):419–22.
24. Zhang HY, Wang ZG, Wu FZ, Kong XX, Yang J, Lin BB, et al. Regulation of autophagy and ubiquitinated protein accumulation by bFGF promotes functional recovery and neural protection in a rat model of spinal cord injury. *Mol Neurobiol*. 2013;48(3):452–64.
25. Beenken A, Mohammadi M. The FGF family: biology, pathophysiology and therapy. *Nat Rev Drug Discov*. 2009;8(3):1–7.
26. Melmed S, Auchus RJ, Goldfine AB, Koenig RJ, Rosen CJ. *Williams Textbook of Endocrinology*. Elsevier. 2020;14:113–113.
27. Esquirol Caussa J, Herrero VE. Epidermal growth factor, innovation and safety. *Med Clínica*. 2015;145(7):305–12.
28. Surini S, Leonyza A, Suh CW. Formulation and in vitro penetration study of recombinant human epidermal growth factor-loaded transdermal emulgel. *Adv Pharm Bull*. 2020;10(4):586–94.
29. Santana H, González Y, Campana PT, Noda J, Amarantes O, Itri R, et al. Screening for stability and compatibility conditions of recombinant human epidermal growth factor for parenteral formulation: Effect of pH, buffers, and excipients. *Int J Pharm*. 2013;452(1–2):52–62.
30. Taylor JM, Mitchell WM, Cohen S. Epidermal growth factor. *J Biol Chem*. 1972;247(18):5928–34.
31. Yang CH, Wu PC, Huang YB, Tsai YH. A new approach for determining the stability of recombinant human epidermal growth factor by thermal Fourier transform infrared (FTIR) microspectroscopy. *J Biomol Struct Dyn*. 2004;22(1):101–10.
32. Laurent GJ, Shapiro SD. *Encyclopedia of Respiratory Medicine*. Elsevier Science. 2006. <https://books.google.co.id/books?id=OnZQAQAAIAAJ>.
33. Chin D, Boyle GM, Parsons PG, Coman WB. What is transforming growth factor-beta (TGF-β)? *Br J Plast Surg*. 2004;57(3):215–21.
34. Attisano L, Ayyaz A, Wrana JL. Recent advances in understanding contextual TGFβ signaling. *F1000Research*. 2017;6(May):1–9.
35. Ong CH, Tham CL, Harith HH, Firdaus N, Israf DA. TGF-β-induced fibrosis: A review on the underlying mechanism and potential therapeutic strategies. *Eur J Pharmacol*. 2021;911(September):174510.
36. Liang J, Zhou Y, Zhang N, Wang D, Cheng X, Li K, et al. The phosphorylation of the Smad2/3 linker region by nemo-like kinase regulates TGF-β signaling. *J Biol Chem*. 2021;296:100512.
37. Yaniz-Galende E, Hajjar RJ. Stem cell and gene therapy for cardiac regeneration. *Cardiac Regeneration and Repair*, vol. 1. Woodhead Publishing Limited. 2014. <https://doi.org/10.1533/9780857096708.4.347>.
38. Duffy AM, Bouchier-Hayes DJ, Harmey JH. Vascular endothelial growth factor (VEGF) and its role in non-endothelial cells: Autocrine signalling by VEGF. *VEGF Cancer*. 2004;1:133–44.
39. Melincovici CS, Boşca AB, Şuşman S, Mărginean M, Mihu C, Istrate M, et al. Vascular endothelial growth factor (VEGF) – key factor in normal and pathological angiogenesis. *Rom J Morphol Embryol*. 2018;59(2):455–67.
40. De Falco S. The discovery of placenta growth factor and its biological activity. *Exp Mol Med*. 2012;44(1):1–9.
41. Rubin JS, Bottaro DP, Chedid M, Miki T, Ron D, Cheon G, et al. Keratinocyte growth factor. *Cell Biol Int*. 1996;19(5):399–411.
42. Poorebrahim M, Sadeghi S, Ghorbani R, Asghari M, Abazari MF, Kalhor H, et al. In silico enhancement of the stability and activity of keratinocyte growth factor. *J Theor Biol*. 2017;418(January):111–21.
43. Borjian Boroujeni M, Shahbazi Dastjerdeh M, Shokrgozar MA, Rahimi H, Omidinia E. Computational driven molecular dynamics simulation of keratinocyte growth factor behavior at different pH conditions. *Informatics Med Unlocked*. 2021;23(October):100514.
44. Hsu E, Osslund T, Nybo R, Chen BL, Kenney WC, Morris CF, et al. Enhanced stability of recombinant keratinocyte growth factor by mutagenesis. *Protein Eng Des Sel*. 2006;19(4):147–53.
45. Sadeghi S, Kalhor H, Panahi M, Abolhasani H, Rahimi B, Kalhor R, et al. Keratinocyte growth factor in focus: A comprehensive review from structural and functional aspects to therapeutic applications of palifermin. *Int J Biol Macromol*. 2021;191(September):1175–90.
46. Hadari YR, Gotoh N, Kouhara H, Lax I, Schlessinger J. Critical role for the docking-protein FRS2α in FGF receptor-mediated signal transduction pathways. *Proc Natl Acad Sci U S A*. 2001;98(15):8578–83.
47. Chang HL, Sugimoto Y, Liu S, Wang LS, Huang YW, Ye W, et al. Keratinocyte growth factor (KGF) regulates estrogen receptor-α (ER-α) expression and cell apoptosis via phosphatidylinositol 3-kinase (PI3K)/Akt pathway in human breast cancer cells. *Anticancer Res*. 2009;29(8):3195–205.
48. Ceccarelli S, Cardinali G, Aspite N, Picardo M, Marchese C, Torrisi MR, et al. Cortactin involvement in the keratinocyte growth factor and fibroblast growth factor 10 promotion of migration and cortical actin assembly in human keratinocytes. *Exp Cell Res*. 2007;313(9):1758–77.
49. Braun S, Auf Dem Keller U, Steiling H, Werner S. Fibroblast growth factors in epithelial repair and cytoprotection. *Philos Trans R Soc B Biol Sci*. 2004;359(1445):753–7.
50. Lee JS, Liu JJ, Hong JW, Wilson SE. Differential expression analysis by gene array of cell cycle modulators in human corneal epithelial cells stimulated with epidermal growth factor (EGF), hepatocyte growth factor (HGF), or keratinocyte growth factor (KGF). *Curr Eye Res*. 2001;23(1):69–76.
51. Spielberger R, Stiff P, Bensinger W, Gentile T, Weisdorf D, Kewalramani T, et al. Palifermin for Oral Mucositis after Intensive Therapy for Hematologic Cancers. *N Engl J Med*. 2004;351(25):2590–8.
52. Wilkinson HN, Hardman MJ. Wound healing: cellular mechanisms and pathological outcomes: Cellular Mechanisms of Wound Repair. *Open Biol*. 2020;10(9):200223.
53. Huang Y, Gou M, Zhang W, Xie H. Mesenchymal Stem Cells for Chronic Wound Healing : Current Status of Preclinical and Clinical Studies. *Tissue Eng - Part B*. 2020;26(6):555–70.
54. Steed DL. The Role of Growth Factor in Wound Healing. *Surg Clin North Am*. 1997;77(3):13–5.
55. Desai TJ, Cardoso WV. Growth factors in lung development and disease: Friends and foe? *Respir Res*. 2001;3:1–4.
56. Qu Y, Cao C, Wu Q, Huang A, Song Y, Li H, et al. The dual delivery of KGF and bFGF by collagen membrane to promote skin wound healing. *J Tissue Eng Regen Med*. 2018;12(6):1508–18.
57. Braiman-Wikman L, Solomonik I, Spira R, Tennenbaum T. Novel Insights into Wound Healing Sequence of Events. *Toxicol Pathol*. 2007;35(6):767–79.
58. Akita S, Akino K, Hirano A. Basic Fibroblast Growth Factor in Scarless Wound Healing. *Adv Wound Care*. 2013;2(2):44–9.
59. Zhang X, Kang X, Jin L, Bai J, Liu W, Wang Z. Stimulation of wound healing using bioinspired hydrogels with basic fibroblast growth factor (bFGF). *Int J Nanomedicine*. 2018;13:3897–906.

60. Li J, Ren K, Zhang W, Xiao L, Wu H, Liu Q, et al. Human amniotic mesenchymal stem cells and their paracrine factors promote wound healing by inhibiting heat stress-induced skin cell apoptosis and enhancing their proliferation through activating PI3K / AKT signaling pathway. *Stem Cell Res Ther.* 2019;10(247):1–17.
61. Reza H, Motahareh A, Hosseini S, Gholami M, Mohamadi F. Mesenchymal stem cells' seeded amniotic membrane as a tissue - engineered dressing for wound healing. *Drug Deliv Transl Res.* 2022;12:538–49.
62. Jones MK, Tomikawa M, Mohajer B, Tarnawski AS. Gastrointestinal mucosal regeneration: role of growth factors. *Front Biosci.* 1999;4:D303–9.
63. Wei G, Wang Z, Liu R, Zhou C, Li E, Shen T, et al. A combination of hybrid polydopamine-human keratinocyte growth factor nanoparticles and sodium hyaluronate for the efficient prevention of postoperative abdominal adhesion formation. *Acta Biomater.* 2022;138:155–67.
64. Zhao YZ, ZhuGe DL, Tong MQ, Lin MT, Zheng YW, Jiang X, et al. Ulcerative colitis-specific delivery of keratinocyte growth factor by neutrophils-simulated liposomes facilitates the morphologic and functional recovery of the damaged colon through alleviating the inflammation. *J Control Release.* 2019;299:90–106.
65. Zeeh JM, Procaccino F, Hoffmann P, Aukerman SL, McRoberts JA, Soltani S, et al. Keratinocyte growth factor ameliorates mucosal injury in an experimental model of colitis in rats. *Gastroenterology.* 1996;110(4):1077–83.
66. Sandborn WJ, Sands BE, Wolf DC, Valentine JF, Safdi M, Katz S, et al. Repifermin (keratinocyte growth factor-2) for the treatment of active ulcerative colitis: a randomized, double-blind, placebo-controlled, dose-escalation trial. *Aliment Pharmacol Ther.* 2003;17(11):1355–64.
67. Itoh M, Matsuo Y. Gastric ulcer treatment with intravenous human epidermal growth factor: A double-blind controlled clinical study. *J Gastroenterol Hepatol.* 1994;9(1 S):S78–83.
68. Gomperts BN, Belperio JA, Fishbein MC, Keane MP, Burdick MD, Strieter RM. Keratinocyte growth factor improves repair in the injured tracheal epithelium. *Am J Respir Cell Mol Biol.* 2007;37(1):48–56.
69. Tillie-Leblond I, Gosset P, Le Berre R, Janin A, Prangère T, Tonnel AB, et al. Keratinocyte growth factor improves alterations of lung permeability and bronchial epithelium in allergic rats. *Eur Respir J.* 2007;30(1):31–9.
70. Wang J, Sun H, Liu Y. The proliferative and anti-apoptosis functions of KGF/KGFR contributes to bronchial epithelial repair in asthma. *Pulm Pharmacol Ther.* 2020;63:101931.
71. Kan B, Jian X, Zhou Q, Wang J, Yu G, Sun J, et al. Effect of transforming growth factor- $\beta$ 1 on acute lung injury caused by paraquat. *Mol Med Rep.* 2014;9(4):1232–6.
72. Kopyński P, Śladek K, Szczekliki J, Soja J, Szlubowski A, Balicka-Ślusarczyk B, et al. Expression of insulin-like growth factor-I (IGF-I) in alveolar macrophages and lymphocytes obtained by bronchoalveolar lavage (BAL) in interstitial lung diseases (ILD). Assessment of IGF-I as a potential local mitogen and antiapoptotic cytokine. *Folia Histochem Cytobiol.* 2006;44(4):249–58.
73. Lutmer J, Watkins D, Chen CL, Velten M, Besner G. Heparin-binding epidermal growth factor-like growth factor attenuates acute lung injury and multiorgan dysfunction after scald burn. *J Surg Res.* 2013;185(1):329–37.
74. Pan X, Xu S, Zhou Z, Wang F, Mao L, Li H, et al. Fibroblast growth factor-2 alleviates the capillary leakage and inflammation in sepsis. *Mol Med.* 2020;26(1):108.
75. Baskin LS, Sutherland RS, Thomson AA, Nguyen HT, Morgan DM, Hayward SW, et al. Growth factors in bladder wound healing. *J Urol.* 1997;157(6):2388–95.
76. Narla ST, Bushnell DS, Schaefer CM, Nouraie M, Bates CM. Keratinocyte growth factor reduces injury and leads to early recovery from cyclophosphamide bladder injury. *Am J Pathol.* 2020;190(1):108–24.
77. Jaal J, Dörr W. Effect of recombinant human keratinocyte growth factor (rHuKGF, Palifermin) on radiation-induced mouse urinary bladder dysfunction. *Int J Radiat Oncol Biol Phys.* 2007;69(2):528–33.
78. Chen W, Shi C, Yi S, Chen B, Zhang W, Fang Z, et al. Bladder regeneration by collagen scaffolds with collagen binding human basic fibroblast growth factor. *J Urol.* 2010;183(6):2432–9.
79. Chen G, Deng C, Li YP. TGF- $\beta$  and BMP signaling in osteoblast differentiation and bone formation. *Int J Biol Sci.* 2012;8(2):272–88.
80. Zou X, Chen K, Zou J, Han P, Hao J, Han Z. Single-cell RNA-seq data analysis on the receptor ACE2 expression reveals the potential risk of different human organs vulnerable to 2019-nCoV infection. *Front Med.* 2020;14(2):185–92.
81. Stewart S, Bryant SJ, Ahn J, Hankenson KD. Bone Regeneration. *Translational Regenerative Medicine.* Elsevier Inc. 2015. <https://doi.org/10.1016/B978-0-12-410396-2.00024-4>.
82. Zhu G, Zhang T, Chen M, Yao K, Huang X, Zhang B, et al. Bone physiological microenvironment and healing mechanism: Basis for future bone-tissue engineering scaffolds. *Bioact Mater.* 2021;6(11):4110–40.
83. Stewart S, Bryant SJ, Ahn J, Hankenson KD. Chapter 24 - Bone Regeneration. In: Atala A, Allickson JG, editors. *Translational Regenerative Medicine.* Boston: Academic Press; 2015. p. 313–33.
84. Li J, Zhou Z, Wen J, Jiang F, Xia Y. Human amniotic mesenchymal stem cells promote endogenous bone regeneration. *Front Endocrinol.* 2020;11(October):1–9.
85. Itoh N. The Fgf families in humans, mice, and zebrafish: Their evolutionary processes and roles in development, metabolism, and disease. *Biol Pharm Bull.* 2007;30(10):1819–25.
86. Wilkie AOM. Bad bones, absent smell, selfish testes: The pleiotropic consequences of human FGF receptor mutations. *Cytokine Growth Factor Rev.* 2005;16(2 SPEC. ISS.):187–203.
87. Nagayama T, Okuhara S, Ota MS, Tachikawa N, Kasugai S, Iseki S. FGF18 accelerates osteoblast differentiation by upregulating Bmp2 expression. *Congenit Anom.* 2013;53(2):83–8.
88. Morimoto N, Yoshimura K, Niimi M, Ito T, Aya R, Fujitaka J, et al. Novel collagen/gelatin scaffold with sustained release of basic fibroblast growth factor: Clinical trial for chronic skin ulcers. *Tissue Eng - Part A.* 2013;19(17–18):1931–40.
89. Kim D, Kim SY, Mun SK, Rhee S, Kim BJ. Epidermal growth factor improves the migration and contractility of aged fibroblasts cultured on 3D collagen matrices. *Int J Mol Med.* 2015;35(4):1017–25.
90. Ha JM, Lim CA, Han K, Ha JC, Lee HE, Lee Y, et al. The effect of microspicule containing epidermal growth factor on periorcular wrinkles. *Ann Dermatol.* 2017;29(2):187–93.
91. Eskens O, Amin S. Challenges and effective routes for formulating and delivery of epidermal growth factors in skin care. *Int J Cosmet Sci.* 2021;43(2):123–30.
92. Swift A, Liew S, Weinkle S, Garcia JK, Silberberg MB. The facial aging process from the "inside Out." *Aesthetic Surg J.* 2021;41(10):1107–19.
93. Fabi S, Sundaram H. The potential of topical and injectable growth factors and cytokines for skin rejuvenation. *Facial Plast Surg.* 2014;30(2):157–71.
94. Liang X, Li J, Yan Y, Xu Y, Wang X, Wu H. Efficacy of microneedling combined with local application of human umbilical cord-derived mesenchymal stem cells conditioned media in skin brightness and rejuvenation : A randomized controlled split-face study. *Front Med.* 2022;9:837332.
95. El-Domyati M, Mofteh NH, Nasif GA, Ameen SW, Ibrahim MR, Ragaie MH. Facial rejuvenation using stem cell conditioned media combined with skin needling: A split-face comparative study. *J Cosmet Dermatol.* 2020;19(9):2404–10.
96. Singh R, Singh S, Lillard JW. Past, present, and future technologies for oral delivery of therapeutic proteins. *J Pharm Sci.* 2008;97:2497–523.
97. Park K, Kwon IC, Park K. Oral protein delivery: Current status and future prospect. *React Funct Polym.* 2011;71(3):280–7.
98. Benson HAE, Namjoshi S. Proteins and peptides: Strategies for delivery to and across the skin. *J Pharm Sci.* 2008;97(9):3591–610.
99. Subbiah R, Guldberg RE. Materials science and design principles of growth factor delivery systems in tissue engineering and regenerative medicine. *Adv Healthc Mater.* 2019;8(1):1–24.
100. Atienza-Roca P, Cui X, Hooper GJ, Woodfield TBF, Lim KS. Growth factor delivery systems for tissue engineering and regenerative medicine. *Adv Exp Med Biol.* 2018;1078:245–69.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**WEB OF SCIENCE (WoS) JOURNAL INFO**

» [Journal of Biological Engineering](#)

<b>ABBREVIATION:</b>	J BIOL ENG
<b>ISSN:</b>	1754-1611
<b>EISSN:</b>	1754-1611
<b>CATEGORY/ QUARTILE:</b>	BIOTECHNOLOGY & APPLIED MICROBIOLOGY - SCIE(Q1) BIOCHEMICAL RESEARCH METHODS - SCIE(Q1)
<b>WoS CORE CITATION INDEXES:</b>	<a href="#">SCIE - Science Citation Index Expanded</a>
<b>IMPACT FACTOR (IF):</b>	6.248
<b>JOURNAL CITATION INDICATOR (JCI):</b>	0.91
<b>CITATIONS:</b>	2,081
<b>GOLD PERCENTAGE OPEN ACCESS:</b>	98.58 % <a href="#">Fully Open Access</a> . For more info, check it on <a href="#">DOAJ.ORG</a>
<b>COUNTRY:</b>	ENGLAND

[More Info from WoS »](#)

*\*In order to submit a manuscript to this journal, please read the guidelines for authors in the [journal homepage](#).*

REVIEW

Open Access



# Prospective use of amniotic mesenchymal stem cell metabolite products for tissue regeneration

Andang Miatmoko<sup>1,2\*</sup>, Berlian Sarasitha Hariawan<sup>3</sup>, Devy Maulidya Cahyani<sup>3</sup>, Syarifah Sutra Dewangga<sup>4</sup>, Kevin Ksatria Handoko<sup>4</sup>, Purwati<sup>2</sup>, Ram Kumar Sahu<sup>5</sup> and Dewi Melani Hariyadi<sup>1</sup>

## Abstract

Chronic disease can cause tissue and organ damage constituting the largest obstacle to therapy which, in turn, reduces patients' quality-adjusted life-year. Degenerative diseases such as osteoporosis, Alzheimer's disease, Parkinson's disease, and infectious conditions such as hepatitis, cause physical injury to organs. Moreover, damage resulting from chronic conditions such as diabetes can also culminate in the loss of organ function. In these cases, organ transplantation constitutes the therapy of choice, despite the associated problems of immunological rejection, potential disease transmission, and high morbidity rates. Tissue regeneration has the potential to heal or replace tissues and organs damaged by age, disease, or trauma, as well as to treat disabilities. Stem cell use represents an unprecedented strategy for these therapies. However, product availability and mass production remain challenges. A novel therapeutic alternative involving amniotic mesenchymal stem cell metabolite products (AMSC-MP) has been developed using metabolites from stem cells which contain cytokines and growth factors. Its potential role in regenerative therapy has recently been explored, enabling broad pharmacological applications including various gastrointestinal, lung, bladder and renal conditions, as well as the treatment of bone wounds, regeneration and skin aging due to its low immunogenicity and anti-inflammatory effects. The various kinds of growth factors present in AMSC-MP, namely bFGF, VEGF, TGF- $\beta$ , EGF and KGF, have their respective functions and activities. Each growth factor is formed by different proteins resulting in molecules with various physicochemical properties and levels of stability. This knowledge will assist in the manufacture and application of AMSC-MP as a therapeutic agent.

**Keywords** Quality-adjusted life-year, Amniotic mesenchymal stem cell metabolite products, Growth factors, Tissue injury, Stem cells, Molecular therapy

## \*Correspondence:

Andang Miatmoko  
andang-m@ff.unair.ac.id

<sup>1</sup> Department of Pharmaceutical Science, Faculty of Pharmacy, Universitas Airlangga, Nanizar Zaman Joenoes Building, Campus C UNAIR Mulyorejo, Mulyorejo, Surabaya, East Java 60115, Indonesia

<sup>2</sup> Stem Cell Research and Development Center, Universitas Airlangga, 2nd Floor Institute of Tropical Disease Building, Campus C UNAIR Mulyorejo, Surabaya 60115, Indonesia

<sup>3</sup> Master Program of Pharmaceutical Sciences, Faculty of Pharmacy, Universitas Airlangga, Campus C UNAIR Mulyorejo, Surabaya 60115, Indonesia

<sup>4</sup> Faculty of Pharmacy, Study Program of Pharmacy, Universitas Airlangga, Campus C UNAIR Mulyorejo, Surabaya 60115, Indonesia

<sup>5</sup> Department of Pharmaceutical Sciences, Hemvati Nandan Bahuguna Garhwal University (A Central University), Chauras Campus, Tehri Garhwal, Uttarakhand 249161, India



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.



## Background

Degenerative diseases, physical injury to organs, and damage due to chronic conditions such as diabetes can cause organ function loss [1–3]. In the early stages of disease, pharmacological drug therapy is the first-line treatment choice, but it has some drawbacks. For example, there are only five types of drugs for reducing the symptoms of Alzheimer's disease [4], and pharmacotherapy to slow disease progression is not yet available. For this purpose, therapeutic agents should inhibit extracellular amyloid plaque deposition and intracellular neurofibrillary formation. Additionally, Alzheimer's therapies that have neuroprotective mechanisms and the use of anti-inflammatory stem cell therapies and the growth factor NDX-1017 are currently under investigation [5, 6].

Stem cells are undifferentiated cells that continuously divide, renew themselves, and differentiate into different types of cells. With the ability of self-renewal, pluripotency, and differentiation, stem cells have great potential for treating various diseases. Stem cells can be divided into two main groups based on their origin: embryonic stem cells and adult stem cells. However, stem cell therapy has two main problems that pose challenges for its therapeutic use. The first relates to availability and mass production. Cell therapy protocols using mesenchymal stem cells (MSCs) for tissue regeneration generally require hundreds of millions of MSCs per treatment. Therefore, expansion of the *in vitro* cell culture is necessary for longer periods and large bioreactors [7]. The need for cells to replace the disease-induced loss of hepatocytes, pancreatic cells, or cardiomyocytes involves approximately  $1$  to  $10 \times 10^9$  functional cells per patient. An even higher requirement has been calculated for the production of "in vitro blood," as approximately  $2.5 \times 10^{12}$  red blood cells are required per patient in transfusion treatments [8]. The main challenge in mass production lies in standardizing the process due to the complexity of pluripotent stem cells and processing cells on a large scale [9].

The use of stem cells can affect the recipient's immune system. The administered cells can directly induce an immune response or modulate the immune system. This is primarily in the case of cells that are not intended to be used for their essential function (nonhomologous use) or when administered to non-physiological sites, which may change the immunogenicity of the cells. Another risk is bacterial and viral infections. As a cell-based product, stem cell production does not allow for terminal sterilization, purification, virus removal, or inactivation processes. Thus, the risk of transmitting bacterial, viral, fungal, or prion pathogens from the donor to the recipient can lead to life-threatening and even fatal reactions [10].

Because of these limitations, new therapeutic strategies are needed. Stem cell-mediated tissue regeneration involves soluble factors secreted by these cells. Cytokines and growth factors, such as transforming growth factor beta (TGF- $\beta$ ), stromal cell-derived factor 1 (SDF-1), and vascular endothelial growth factor (VEGF), are secreted by stem cells and progenitor cells transplanted into the intestinal space or injected into blood vessels and stimulate many regenerative processes such as neovascularization, activation of tissue intrinsic progenitor cells, decreased apoptosis of endogenous cardiomyocytes, and registration of assistive cells for tissue repair [11, 12]. Additionally, mesenchymal stem cells (MSCs) secrete growth factors and cytokines, which promote wound repair. The combination of growth factors and cytokines successfully induces angiogenesis, reduces inflammation, and promotes fibroblast migration and collagen production [13].

## Amniotic membrane stem cell metabolite products (AMSC-MPs)

### Placental tissue is the primary source of AMSC-MPs

The amnion, chorion, amniotic fluid, and umbilical cord are of fetal origin. These components have been widely studied because of their potential use as cell sources for regenerative therapies. At delivery, the amniotic membrane is strong, protects the fetus from physical shocks, regulates the pH of the fluid membranes, and secretes various cell signals and bioactive molecules as antimicrobials and anti-inflammatory agents [14]. Amniotic membrane stem cells (AMSCs) are MSCs from the amniotic epithelium and the stroma of the amniotic membrane that are sources of epidermal growth factor (EGF) and keratinocyte growth factor (KGF). Furthermore, stem cells synthesize and secrete various extracellular matrix proteins, cytokines, growth factors, and other bioactive proteins that contribute to the healing and regenerative processes [15]. The collection of conditioned stem cell culture media rich in bioactive agents such as growth factor and cytokines secreted into the extracellular space is defined as a metabolite product. Since it is derived from amniotic mesenchymal stem cells, it is referred to as AMSC-MP [7].

The AMSC-MP collection procedures are conducted under sterile conditions. The first step is expanding cells to the desired passage. Then, upon trypsinization and counting of selected cells, the cell needs to be seeded again at a determined cell density, for example, 12,000 cells/cm<sup>2</sup>, and cultured in an appropriate growth medium. After 72 h, the growth medium is removed, and the cells are washed several times with phosphate-buffered saline, preferably without Ca<sup>2+</sup>/Mg<sup>2+</sup> content, at room temperature. After that, the cell culture medium

(pre-warmed to 37 °C) with only 1% antibiotic is added. The AMSC-MP was collected after being incubated for 48 h and centrifuged at 1500 rpm for 3 min two times, then filtered with a syringe filter 0.45 µm to remove debris and dead cells. The AMSC-MP then be dialyzed using a dialysis bag with a proper molecular-weight cut-off to replace the cell growth medium with the phosphate-buffered saline as the media [16, 17].


These molecules include basic fibroblast growth factor (bFGF), EGF, hyaluronic acid (HA), interleukins (IL-1 and IL-10), beta-defensins, TGF-β, elafin, human leukocyte antigen-G, matrix metalloproteinases, tissue inhibitors of metalloproteinases (TIMPs), and platelet-derived growth factor (PDGF) [15]. Additionally, amniotic tissue contains anti-inflammatory factors such as IL-1 and IL-10 receptor antagonists and regulators of catabolic enzymes such as TIMP1, TIMP2, TIMP3, and TIMP4. Furthermore, AMSC-MP is a potent down-regulator of TGF-β signaling, which stimulates the recruitment of fibroblasts and macrophages and upregulates collagen production [18].

#### Physicochemical properties and stability of AMSC-MP

AMSC-MP is a clear liquid containing various proteins, and some products form yellowish white lyophilized powder. A study by Kumala et al. (2020) compared the physicochemical stability of AMSC-MP in two forms, i.e., native AMSC-MP liquid and lyophilized AMSC-MP powder [19]. The results showed that AMSC-MP liquid began to change color after 7 days of storage at room temperature. In contrast, AMSC-MP liquid did not show color changes or an odor at cold temperatures. Its pH was 7–7.5 without significant changes during storage for 28 days [19].

AMSC-MP contains several proteins and the major component has a molecular weight of 75.33 kDa, as seen from the thick band in Fig. 1. A stability study measuring TGF-β with the enzyme-linked immunosorbent assay (ELISA) showed that the AMSC-MP liquid is less stable stored in room temperature (25°C) than when freeze-dried. Freeze-dried AMSC-MP has high crystallinity, which was supported by a scanning electron microscopy characterization showing a tetragonal crystal shape. The crystals have a melting point of 163.8 °C [19].

Specifically, stability data from each component of the AMSC-MP is still limited. It has been reported that the FGF1 and FGF2 lost most of their activity after 6 h at 37 °C [20]. Another study showed that the eye serum drops containing EGF, TGF-β1, and TGF-β2 remained stable in tube segments and vials made from polyvinyl chloride (PVC) with di-2-ethylhexyl phthalate (DEHP) and tris-2-ethylhexyl trimellitate (TOTM) as the plasticizers, respectively, for about 15 months by storage at



**Fig. 1** Qualitative determination of protein markers in fluid and freeze-dried (FD) AMSC-MP analysed using SDS-PAGE [19]

-30 °C. However, the stability was significantly decreased by storing samples at 4 °C and 22 °C, which could be reduced up to only 24 and 3 h for EGF and 48 and 9 h for TGF-β1 and TGF-β2, respectively. EGF was reportedly unstable at 37 °C [21]. Mizumachi & Ijima (2013) showed that the immobilized-VEGF prepared in a heparin-collagen substrate had higher stability than that of VEGF solution, resulting in about 43% of the initial VEGF retaining its biological activity, although after 16 days of pre-incubation at 37 °C [22]. Meanwhile, the research by Chen & Arakawa (1996) reported that the KGF is unstable in the aqueous media and it follows denaturation and aggregation when heated to 37 °C [23].

#### Biochemical components of AMSC-MP

##### bFGF

One of the growth factors in AMSC-MP is bFGF [7, 11, 18] (or FGF-2), which is a member of the FGF family that regulates various biological functions including proliferation, morphogenesis, and suppression of apoptosis

during development through a complex signal transduction system. bFGF is widely expressed in the nervous system, where it has multiple roles, and it supports the survival and growth of neuronal and neural stem cell cultures [24].

Members of the FGF family have a homologous core region of 120–130 amino acids arranged into 12 antiparallel strands ( $\beta 1$ – $\beta 12$ ) flanked by divergent amino and carboxyl functional groups. Generally, the sequence variation of the N- and C-terminal ends of specific FGF family members accounts for their differential ligand binding. The heparan sulfate glycosaminoglycan (HSGAG) binding site in the FGF core consists of a  $\beta 1$ – $\beta 2$  loop and parts of  $\beta 10$  and  $\beta 12$ . This section differs for each FGF family member and determines the endocrine properties of each. FGF binds to and activates the FGF receptor (FGFR) in the HSGAG-dependent tyrosine kinase receptor family. Upon ligand and HSGAG binding, FGFR dimerizes, allowing the cytoplasmic kinase domain to transphosphorylate and activate tyrosine loop A. Loop phosphorylation is followed by tyrosine phosphorylation in the C tail region, kinase insert, and juxtamembrane region. This process activates the Ras–mitogen-activated protein kinase (MAPK) and phosphoinositide 3-kinase–Akt signaling pathways [25].

In tissue regeneration therapy, FGFs from the paracrine group (FGF1–10, FGF16–18, FGF20, and FGF22) play a role. Paracrine FGFs have a high affinity for HSGAG, activating it and acting locally near the expression source.

## EGF

Another growth factor in AMSC-MP is EGF [11, 12, 17]. EGF is a small polypeptide mitogen present in many species that has been isolated and characterized in breast milk. EGF is a 6-kDa peptide derived from a 1207 amino acid precursor molecule that acts across a 170-kDa membrane glycoprotein receptor. EGF has intrinsic tyrosine kinase activity, like the IGF receptor, and functions in tyrosine kinase-mediated autophosphorylation. TGF- $\alpha$ , which has a 35% amino acid homology with murine EGF and 44% homology with human EGF, also acts through the EGF receptor [26].

Growth factors are naturally produced proteins that regulate cell proliferation, function, and differentiation through receptor signaling. EGF is one of the earliest known polypeptide growth factors and was the founder of the EGF-like family of proteins. EGF is an endogenous peptide that promotes cell growth, proliferation, and differentiation via ligand-receptor (EGFR) interactions [27, 28]. EGF was first isolated from the submaxillary glands of adult male rats by Cohen et al. in 1962. Currently, recombinant human EGF (rhEGF) can be mass produced from *Escherichia coli*, which has accelerated the

development of EGF formulations for treating skin conditions such as chronic wounds, burns, diabetic ulcers, and skin aging [28].

EGF has hydrophilic properties [29]. Structurally, EGF is a polypeptide chain with a molecular weight of 6045 Da consisting of 53 amino acids and 6 cysteine residues and has three intramolecular disulfide bonds [27]. EGF is characterized by the absence of three specific amino acid residues, lysine, alanine, and phenylalanine, and it lacks hexosamines and neutral sugars [30]. EGF has an optimal stability at pH 6.0–8.0 and an isoelectric point of about 4.6 [29]. EGF shows poor thermal stability because the protein structure begins to degenerate at 40 °C, and it has a transition midpoint at 55.5 °C. EGF can be completely denatured at temperatures above 76 °C [31].

EGF binds to EGFR; following ligand binding, EGFR (ErbB-1) dimerizes with itself or with a homolog, ErbB-2, ErbB-3, or ErbB-4, increasing intracellular tyrosine kinase activity. This process activates a signaling cascade that has multiple effects: cell proliferation, reduction of apoptosis, and angiogenesis [32].

## TGF- $\beta$

TGF- $\beta$  is an extracellular protein in AMSC-MP produced mainly by a subset of T cells [7, 11, 17]. TGF- $\beta$  belongs to a group of cytokines collectively referred to as the TGF- $\beta$  superfamily, whose members regulate epithelial cell growth, differentiation, motility, organization, apoptosis, and tumorigenesis [33]. The TGF- $\beta$  superfamily consists of a group of polypeptide morphogens. TGFs are divided into two subgroups: the TGF-like subgroup [TGF- $\beta$ s, activin, nodal, and multiple growth differentiation factors (GDFs)] and the BMP-like subgroup (BMP, GDF, and antimüllerian hormone). Members of the TGF-like subgroup exhibit functions in cell adhesion, growth, cytoskeletal organization, survival, proliferation migration, differentiation, chemotaxis, and immune cell activation in multicellular organism development [34].

Blood is the primary source of TGF- $\beta$ , which promotes healing and tissue regeneration during injury. Platelet aggregation and degranulation release high amounts of TGF- $\beta 1$  at wound healing sites. Additionally, recruited and activated leukocytes at wound sites secrete various cytokines, including TGF- $\beta 1$  to support the wound healing process [35]. TGF- $\beta$  is secreted in a latent or biologically inactive state. During cellular synthesis, the TGF- $\beta$  precursor undergoes intracellular proteolytic cleavage by furin endopeptidase, resulting in two proteins assembled into dimers via noncovalent associations.

TGF- $\beta$  signaling involves three parallel pathways, the bone morphogenic protein (BMP), TGF- $\beta$ , and activin

pathways, all of which are major regulators. TGF- $\beta$  signaling is transduced in cells by several SMAD protein modulators, which eventually enter the cell nucleus and influence the expression of target genes. Since all three pathways comprise ligands and receptors, the combination of different signals allows the regulation of many growth and developmental processes in highly specific ways [36].

### VEGF

VEGF regulates angiogenesis by inducing the proliferation, migration, and permeability of endothelial cells. VEGF is also found in AMSC-MP [7, 11, 17]. During the process of tissue regeneration, VEGF also plays an important role in cardiac repair by decreasing infarction size, reducing remodeling, decreasing endothelial cell apoptosis, supporting angiogenesis and neovascularization, increasing the number of mitotic cardiomyocytes in the border zone, and improving cardiac performance [37]. VEGF is produced by many cell types including tumor cells, macrophages, platelets, keratinocytes, and renal mesangial cells. VEGF activity is not confined to the vascular system; VEGF also plays a role in normal physiological functions such as bone formation, hematopoiesis, and wound healing [38].

The *VEGF* gene is located on chromosome 6p21.3 and is part of the *VEGF/PDGF* gene family, the cystine-knot superfamily of growth factors. Structurally, VEGF is a 40-kDa heterodimeric glycoprotein. VEGF contains a cystine-knot motif, characterized by disulfide bridges in the protein structure [39]. In humans, VEGF consists of several members: VEGF-A, which has several isoforms, VEGF-B, VEGF-C, VEGF-D, VEGF-E (VEGF virus), placenta growth factor (PlGF), and endothelial-derived VEGF [40].

There are three VEGF receptors: VEGFR-1, VEGFR-2, and VEGFR-3. Neuropilin-1 (NP-1) and neuropilin-2 (NP-2) coreceptors are non-tyrosine kinase receptors, and they selectively attach to certain VEGF subtypes or isoforms. The pro-angiogenic activity of VEGF occurs through the binding and activation of two receptor tyrosine kinases (TKs), which were initially identified as receptors for VEGF-A, namely VEGFR-1 and VEGFR-2. These receptors consist of seven extracellular Ig-like domains, a transmembrane domain, and an intracellular TK domain. Ligand binding induces receptor dimerization and phosphorylation. PlGF binds exclusively to VEGFR-1 with high affinity compared to VEGF-A and VEGF-B, and other family members also specifically bind VEGFR-1 [40].

PlGF was first identified in human placental tissue. It is involved in trophoblast growth and differentiation, trophoblast invasion, and blastocyst implantation. The

PlGF gene has four isoforms, PlGF-1 (PlGF131), PlGF-2 (PlGF152), PlGF-3 (PlGF203), and PlGF-4 (PlGF224), which differ in their molecular structure and biological properties. All isoforms have affinity for VEGFR-1. PlGF has no direct mitogenic effect and does not increase vascular permeability, but, under pathological conditions, binds to VEGFR-1, displaces VEGF-A from VEGFR-1, and allows binding of VEGF-A to VEGFR-2, indirectly enhancing the effect of VEGF-A. This increases vascular permeability, cell migration, and proliferation [40].

### KGF

Another growth factor reported to be present in AMSC-MP is KGF [7, 11, 17]. KGF is a monomeric polypeptide measuring 26–28 kDa and is an FGF family member [41]. KGF has been implicated in biological processes such as cell proliferation, development, and differentiation. KGF is encoded by the *FGF7* gene and is made up of 194 amino acids. The human KGF amino acid sequence in UniProtKB (P21781) indicates that this protein contains a signal peptide (residues 1–31) and a KGF chain (residues 32–194), and position 45 is glycosylated during posttranslational modification [41].

KGF has low stability in acidic and neutral pH conditions. Denatured KGF starts to aggregate at a moderate temperature. The calculated isoelectric point (pI) of KGF is 9.29. The pI is defined as the pH at which the molecule carries no electrical charge or is neutral on average. At a pH below the pI, proteins carry a positive charge, whereas above the pI, they carry a net negative charge [34, 35].

KGF has 19 negatively charged residues, such as Asp and Glu, and 29 positive charge residues, like Arg and Lys. KGF may be unstable under physiological conditions. The instability of the KGF protein is due to the repulsion between the positively charged residues. Therefore, the main cause KGF instability at a neutral pH is thought to be due to its high positive charge and repulsion forces, which lead to protein denaturing and irreversible aggregation [42].

Boroujeni et al. [43] investigated the stability of rhKGF, a truncated form of KGF, under different pH and temperature conditions using molecular dynamics simulation methods. That study showed that the stability of rhKGF increased with a decrease in the total charge at an alkaline pH and low temperature. KGF stability increased significantly at an alkaline pH of 8.5 and 9, presumably due to the presence of a high positive-charge residue. Acidic pH conditions caused instability and aggregation of KGF. In another study, KGF denatured at 400 °C. After a while, the protein began to aggregate, and irreversible particulate and precipitate formation occurred [44].

KGF is involved in various biological processes and has proliferation, antiapoptosis, cytoprotective, epithelial cell movement, and cytoskeletal reorganization effects. Additionally, its mitogenicity is useful for embryonic development, tissue patterning, cell growth, morphogenesis, wound healing, and tissue repair. As a growth factor, KGF acts through a variant of the receptor-2IIIb FGF (FGFR2b), which is expressed by epithelial cells [25]. The FGFR2b receptor is hereafter referred to as the KGF receptor (KGFR). KGFR is part of the FGFR family of receptor TKs, which are activated in the presence of heparin/ HSGAG. The binding of KGF to KGFR requires heparin/HS as a coreceptor. This process includes the dimerization of KGFR and activation of its kinase domain, inducing autophosphorylation of the receptor [45]. To bind to the receptor and its ligands, KGF has a positively charged site called the heparin binding site, and a neutral site that binds to KGFR.

The proliferative effect of KGF occurs due to activation of the RAS and Raf/MAPK/ERK pathways after KGFR dimerization and phosphorylation of the tyrosine kinase domain [46]. Its antiapoptotic effect occurs due to activation of p21-activated kinase 4 (PAK4), which then activates the antiapoptotic-Akt-dependent pathway by recruiting PI3K, which then arranges antiapoptotic genes [47]. Activation of the antiapoptotic pathway by the ERK1/2 pathway reduces the cellular inflammatory response by inducing the expression of the cytoprotective genes nuclear factor erythroid 2-related factor 2 (*NRF2*) and heme oxygenase-1 (*HO1*) in epithelial cells [48, 49].

KGF regulates epithelial cell motility and cytoskeletal reorganization through activation of Src-Cortactin, which phosphorylates paxillin and activates GTPases such as Rho, Rac, and Cdc42, resulting in lamellipodia extension, actin increase, and cell mobility and migration [48]. During mitogenicity, autophosphorylation of the tyrosine kinase domain of KGFR results in the activation of phosphatidylinositol hydrolysis. PLC $\gamma$  activation hydrolyzes phosphatidylinositol-4,5-diphosphate to inositol-1,4,5-triphosphate and diacylglycerol, which stimulate protein kinase C and increase intracellular Ca<sup>2+</sup> and subsequent mitogenic activity through upregulation of target genes [50]. Currently, palifermin, a recombinant preparation of human KGF, is a growth factor cocktail used for therapy. For example, a study by Spielberger et al. [51] showed that giving 60 g per kilogram of body weight of palifermin daily reduces the duration and severity of oral mucositis after intensive chemotherapy and radiotherapy for hematological cancer.

Based on the explanation above, it can be seen that various kinds of growth factors present in AMSC-MP, namely bFGF, VEGF, TGF- $\beta$ , EGF and KGF have their respective functions and activities. In addition, each growth factor is formed by different proteins resulting in molecules with different physicochemical properties and stability. This knowledge will assist in the manufacturing process and application of AMSC-MP as a therapeutic agent.

## Biological activities of AMSC-MP for tissue regeneration

### Wound healing

The wound repair process is divided into four main phases: hemostasis, inflammation, proliferation, and dermal remodeling [52]. Wound repair begins with the hemostasis phase, in which platelet formation blood loss and the initial fibrin matrix begins to form. Platelets are critical in the recruitment of immune cells to wound tissue, either by capturing immune cells directly or by releasing chemokine secretomes. Platelet secretomes also contain growth factors that stimulate resident skin cells, including fibroblasts and keratinocytes [52].

Furthermore, in the inflammation phase, necrotic cells and damaged tissue release damage-associated molecular patterns and resident immune cells, such as mast cells, Langerhans cells, T cells, and macrophages, respond by activating inflammatory pathways. In this phase, proinflammatory cytokine and chemokine release attracts leukocytes in the circulation to the injured tissue. Monocytes already in the wound tissue differentiate into macrophages. Macrophages engulf necrotic cell remnants and pathogenic material [52]. Activation of macrophages is influenced by proinflammatory stimuli, such as lipopolysaccharide and interferon-gamma (IFN- $\gamma$ ), and their activation enhances inflammation by releasing reactive oxygen species (ROS), inflammatory cytokines (e.g., IL-1, IL-6, and TNF-) and growth factors (e.g., VEGF and PDGF).

Macrophages control the degradation of extracellular connective tissue by enzyme secretion and phagocytosis and regulate wound matrix remodeling through the production of growth factors such as PDGF, TGF, ILs, and TNF [53]. These growth factors influence the regrowth process, epithelialization, fibroplasia, and angiogenesis [52, 54]. All three stages occur in the proliferation phase, starting when keratinocytes migrate to close the wound, followed by angiogenesis, and then fibroblasts replace the initial fibrin clot with granulation tissue [55]. Fibroplasia begins about 5 d after injury and continues for 2 weeks. Fibroblasts migrate into the wound and replicate in response to mediators released during inflammation. These mediators include C5a, fibronectin, PDGF,

FGF, and TGF $\beta$ 1. Remodeling of the extracellular matrix (ECM) spans the entire injury response, beginning with the initial deposition of a fibrin clot and ending several years later with the formation of a mature, type I collagen-rich scar.

Two types of FGF members play an important role in the wound healing process, among which the most important are KGF and bFGF. Both are present in AMSC-MP. Qu et al. (2018) showed that the combination of KGF and bFGF in a collagen delivery system increased cell migration in the wound healing process, accelerating wound closure [56]. bFGF improved wound healing in animal models and clinical studies. KGF is a cytokine that exerts a specific mitogenic effect in epithelial cells. This effect has been reported to be a key factor in wound healing, as it is weakly expressed in human skin but is strongly upregulated after skin injury [56].

The synergistic effect of bFGF and KGF can be observed during the wound healing process. Re-epithelialization begins within hours of injury, and bFGF and KGF promote cell proliferation. KGF stimulates keratinocyte migration, while bFGF promotes fibroblast migration and stimulates the production of collagenase, suggesting that bFGF and KGF have complementary roles in wound healing. Together, bFGF and KGF may also stimulate the accumulation of vascularization-related cells. However, other studies have shown that KGF affects ongoing inflammation and scar formation [57]. These negative effects can be minimized by bFGF as an antiscarring agent [58].

bFGF in wound healing was also investigated by Zhang et al. (2018), in which hydrogels combined with bFGF increased the wound healing process. Hydrogels derived from gum arabic, pectin, and divalent calcium ions help increase the stability of FGFs and provide a sustained release effect [59]. Fibroblast scratch assays showed that the hydrogel FGF formulation could close a wound within 12 h, while in controls, wound closure only started at the 12<sup>th</sup> hour. In an in vivo study conducted by creating a full-thickness skin incision on the back of mice, the hydrogel treatment group with bFGF showed the fastest wound closure compared to the negative control group [59].

Furthermore, the metabolite products of stem cells have a high potential for wound healing efficacy, based on preclinical and clinical studies on stem cells. The use of human AMSCs for wound healing has also been investigated in vivo in male mice in a heat-induced apoptosis model [60]. The administration of  $2 \times 10^6$  cells injected subcutaneously into the wounded skin showed accelerated re-epithelialization. Wound closure occurred on day 7, and cytokines, including PAI-1, C-GSE, periostin, and TIMP-1, have been reported to activate the PI3K/

AKT pathway, which plays a vital role in epithelial cell and dermal fibroblast migration and proliferation. In another study, the use of human adipose-derived MSCs and placenta-derived MSCs in amniotic membrane grafts accelerated wound healing in Wistar rats with an excisional wound splinting model at day 7 [61]. The processes of re-epithelialization, collagen remodeling, and neovascularization occur more quickly by embedding these cells on the amniotic membrane as wound dressing, and this membrane produces various growth factors such as TGF- $\alpha$ , TGF- $\beta$ , bFGF, EGF, and KGF, cytokines such as IL-4, IL-6, and IL-8, as well as matrix metalloproteinase inhibitors. A review by Huang et al. (2020), which explored the use of MSCs in preclinical and clinical studies for wound healing, showed that metabolite products play an active role in tissue repair and wound healing [53]. Tissue repair occurs through the stimulation of cell differentiation and paracrine action, involving growth factors such as bFGF, hepatic growth factor (HGF), EGF, KGF, VEGF, and TGF- $\beta$ , and also cytokines such as IL-10, to reduce inflammation and accelerate the angiogenesis, granulation, re-epithelialization, and wound closure processes.

### Potential uses of AMSC-MP in gastrointestinal injury therapy

In the gastric mucosa, TGF- $\alpha$  controls cell proliferation under normal conditions and after acute injury, while EGF controls cell proliferation during the healing of chronic ulcers. When the gastric mucosa is injured, growth factors predominantly restore the epithelial component, while bFGF and VEGF promote restoration of the connective tissue and angiogenesis in the injured mucosa. Granulated connective tissue, which grows under the stimulation of bFGF and VEGF, is the primary source for regenerating connective tissue lamina propria and microvessels within ulcer scars. Other growth factors such as insulin-like growth factor, KGF, hepatocyte growth factor, and trefoil peptides also act in gastrointestinal (gastric ulcers, colitis) regeneration following injury [62].

Research by Wei et al. (2022) showed that KGF in combination with polydopamine (PDA) and HA nanoparticles successfully prevented abdominal adhesions and promoted the repair of mesothelial cells in the injured peritoneum [63]. More importantly, PDA-KGF NPs combined with HA reduced collagen deposition and fibrosis and inhibited the inflammatory response [63].

KGF function is determined by phosphorylation of the protein tyrosine kinase SRC. When KGF binds to its receptor, SRC is phosphorylated by KGFR. In a study, we evaluated the phosphorylation level of rat Src in the injured peritoneum 7 d after surgery. The levels of phospho-Src protein in rat peritoneal tissue were higher in the

PDA-KGF NP treatment group compared to that treated with KGF alone. Thus, the *in vivo* positive effect of KGF is prolonged when KGF and PDA are administered as PDA-KGF nanogels [63].

KGF is also effective for treating ulcerative colitis. Ying-Zheng et al. (2019) reported that KGF encapsulated into neutrophil-like liposomes (KGF-Neus) effectively restored intestinal morphology and function in ulcerative colitis [64] because the neutrophil membrane vesicle (NEM) associated protein, KGF-Neus, is specifically internalized to the area of inflammation [64].

KGF and its receptors are present in the human fetal gastrointestinal tract, and *in vitro* stimulation of human fetal enterocytes with KGF results in cellular proliferation. KGF expression is increased in patients who undergo surgery for inflammatory bowel disease and is correlated with the degree of intestinal inflammation. In animal models of colitis, KGF administration reduces the degree of mucosal injury [65]. Recombinant KGF treatment has been studied for use in ulcerative colitis; in a clinical phase II study, recombinant KGF failed to induce remission in ulcerative colitis patients, but the maximal therapeutic dose used may have been too low [66].

Another AMSC-MP component, EGF, is associated with mucosal ulcer disease. Decreased EGF levels are associated with mucosal ulcer disease. Patients with duodenal ulcer disease also have decreased EGF levels. EGF supplementation promotes mucosal repair and regeneration in several conditions. In experiments in pigs, EGF significantly reduced esophageal ulceration, structural formation, and mucosal histological damage associated with sclerotherapy. In rats with gastric ulcers, orogastric EGF administered in combination with sucralate improved ulcer healing [67]. A small human study showed that treatment with intravenous EGF promoted better gastric ulcer healing compared with the antiulcer treatment cetraxate hydrochloride [67].

### Potential use of AMSC-MP in lung injury treatment

Growth factors are involved in all aspects of lung development. The spatial and temporal distribution of FGF10 in the lung determine the airway branching pattern. Some factors participate in more specific developmental programs, such as VEGF in blood vessel formation and FGF7 in type II alveolar cell differentiation [55]. KGF is a critical growth factor in lung development and is protective after lung injury. KGF is an important growth factor for local resident progenitor epithelial cell repair and for mobilization and enhanced engraftment of cytokeratin 5 circulating epithelial progenitor cells, which contributed to the repair of the proximal airway epithelium in a mouse model of syngeneic tracheal transplantation to the injured proximal airway epithelium [68].

KGF induces epithelial cell proliferation and protects against acute lung injury. Leblond et al. (2007) showed that 1 mg/kg of body weight of KGF given intravenously to rats injected with albumin as an asthma trigger reduced extravascular lung water levels. KGF treatment also reduced the number of inflammatory cells in the bronchoalveolar lavage fluid but not in the bronchial mucosa. KGF reduces allergen-induced changes in epithelial integrity and the expression of the intercellular junction proteins catenin and zonular occludens protein-1 [69].

Consistently, Wang et al. (2020) investigated the effect of KGF on the release of inflammatory-related cytokines by damaged bronchial epithelial cells. Compared with the healthy group, KGF and KGFR expression and apoptosis were significantly increased in asthmatic patients. An *in vitro* study showed that KGF treatment limited IFN- $\gamma$  and TNF- $\alpha$ -induced apoptosis by inhibiting apoptotic markers in the TNF signaling pathway. KGF limits the release of TSLP, IL-25, and IL-33 by damaging 16HBE 14o cells. In contrast, KGF promotes intercellular adhesion and wound closure of cultured 16HBE 14o cells through increased expression levels of the intercellular junction proteins ICAM-1,  $\beta$ -catenin, E-cad, and Dsc3. In summary, KGF and KGFR may aid bronchial epithelial cell repair in asthma by inhibiting epithelial cell apoptosis while promoting epithelial cell proliferation and migration [70].

TGF- $\beta$  is associated with acute lung injury. Research by Kan et al. 2014 showed an increase in the expression of TGF- $\beta$  in rat serum induced with paraquat, a compound that irritates the lungs [71]. *TGF- $\beta$ 1* mRNA expression in rat lungs was also significantly increased. Many inflammatory cells were observed infiltrating the alveoli of the injured lungs. The abnormal expression of *TGF- $\beta$ 1* was hypothesized to be important in the pathogenesis of chronic inflammatory and immune lung diseases, including asthma, chronic obstructive pulmonary disease, and pulmonary fibrosis [72]. In the future, cytokines and their inhibitors may provide new therapies for treating acute lung injury and pulmonary fibrosis.

Heparin-binding EGF-like growth factor (HB-EGF) reduces inflammation, maintains intestinal barrier function, and protects the lung from acute injury in several models of intestinal injury. Another study investigated the impact of HB-EGF by comparing burn-treated mice (25% of total body surface area) with burn-infected mice after two enteral doses of HB-EGF (1200 mg/kg/dose) [73]. The control mice had increased pulmonary myeloperoxidase levels, lung and spleen apoptosis, airway resistance and bronchial reactivity, and intestinal permeability.

These effects were significantly reduced in burn-injured mice treated with enteral HB-EGF [73].

FGF2 is closely involved in endothelial cell migration, proliferation, and injury repair. Recombinant FGF2 was injected peritoneally at a 0.1 mg/kg dose in septic mice induced by ligation and cecal puncture. FGF2 treatment reduced the inflammatory response, attenuated pulmonary capillary leakage, reduced lung injury, and increased survival in septic mice. Endothelial injury and macrophage inflammation induced by LPSs are inhibited by FGF2 administration via the AKT/P38/NF- $\kappa$ B signaling pathway [74].

### Potential use of AMSC-MP in bladder and renal injuries

Surgical and traumatic injuries to the bladder initiate a complex series of biological processes that result in wound healing. This involves cellular proliferation, migration, and differentiation; removal of damaged tissue; and production of extracellular matrix, all of which may be controlled by growth factors. KGF is induced in the skin following incisional injury. During the early phases of bladder wound healing, mRNA levels of *KGF* and *TGF- $\alpha$*  increased, and exogenous KGF directly affected urothelial proliferation [75].

Among the growth factors that affect the bladder are KGF and FGFs; KGF increases cyclophosphamide-induced bladder injury. Cyclophosphamide is often used to treat cancer and rheumatic and kidney diseases. Acrolein, its metabolite, is a toxic metabolite concentrated in urine that can cause acute hemorrhagic cystitis (7%–45% incidence rate) and urothelial cancer (4%–15% incidence rate, depending on the dose). In cyclophosphamide-induced urothelial injury, increased apoptosis of intermediate and basal cells was observed. KGF prevented apoptosis of deeper urothelial cells (UPK3 + intermediate and KRT5 + intermediate/basal cells), likely via activation of AKT [76].

Evidence suggests that KGF regulates bladder cell development and function and is directly responsible for urothelium proliferation. In a study by Narla et al. (2020), mice were given a 5 mg/kg injection of KGF dissolved in PBS 24 h before cyclophosphamide injection and showed increased urothelial regeneration compared to controls [76]. KGF pretreatment blocked cyclophosphamide-induced intermediate and basal cell apoptosis, possibly via phosphorylated AKT, and promoted ERK-mediated phosphorylated KRT5 + /KRT14 – cell proliferation, leading to urothelial regeneration. The effect of KGF on bladder injury was also found in a study by Jaal et al. (2007) [77], in which there was an increased positive response on day 2 in 50% of mice after a single injection of palifermin at a dose of 15 mg/kg [77].

In addition to KGF, bFGF also affects bladder injury. Chen et al. (2010) explored the ameliorative effect of collagen-based bFGF for bladder regeneration in a mouse model [78]. A bladder with a subtotal cystectomy was grafted with collagen membranes coupled with 0.56 nmol of CBD-bFGF. As a result, collagen/bFGF mice had faster collagen scaffold degradation and better bladder wall cell growth but no bladder stone formation [78].

In addition to bladder injury, FGF positively affects acute kidney disease. Zhou et al. investigated the effects of FGF2 in acute kidney disease using Sprague–Dawley and NRK-52E cells [79]. FGF2 significantly increased tissue apoptosis in acute kidney disease by inhibiting excessive ER stress. Moreover, FGF2 also reduced ER overstress and apoptosis in cultured NRK-52E cells injured with tert-butyl hydroperoxide [80].

### Potential uses of AMSC-MP for bone regeneration

The bone response to injury begins with an inflammatory phase. Bleeding from the fracture-surrounding soft tissue results in forming a fibrin clot and fracture hematoma. Subsequently, inflammatory cytokines are released, inducing angiogenesis and mesenchymal progenitor cell proliferation. These mesenchymal progenitors rapidly proliferate, forming an initial soft callus. The soft or primary callus response occurs within two weeks of injury. The degree of callus formation is proportional to the degree of motion at the fracture gap [81].

In the second stage of repair, the necrotic bone ends undergo resorption, and the mesenchymal progenitor cells proliferating at the injury site begin to differentiate into chondrocytes to form a cartilaginous callus and osteoblasts for intramembranous bone formation at the fracture margins. The mechanisms that control the influx, proliferation, and differentiation capacity of mesenchymal progenitor cells are critical components of the fracture healing process. For instance, BMP and Wnt signaling play roles in mesenchymal cell differentiation into osteoblasts and chondrocytes [81].

During the establishment and maturation of the soft callus, growth factors (e.g., TGF- $\beta$ , PDGF, GDF-5, FGF-1, and IGF-II) and hormones (e.g., PTHrP) are involved in the recruitment and proliferation of fibroblasts and MSCs. They also play an essential role in inducing MSC differentiation into osteoblasts or chondrocytes [82]. Once chondrocytes form, endochondral ossification occurs, and a hard callus is formed via woven bone in the third stage of healing. Whether this endochondral bone formation process is equivalent to that which occurs during bone growth is not completely understood [83]. The use of human AMSC in bone tissue regeneration has been reported in several studies, including in vitro and in vivo studies and clinical trials, as summarized



by Li et al. (2020) [84]. AMSC treatment in models of collagen-induced arthritis, intervertebral disc degeneration, rheumatoid arthritis, and osteoarthritis has shown anti-inflammatory, angiogenic, and immunomodulatory effects, all of which play important roles in tissue remodeling. Several growth factors and cytokines present in human AMSC play an important role in the regeneration process of bone defects. These include HGF, FGF7, BMP-2, VEGF, IL-6, and IL-8. Clinical trials in patients with bone defects of human AMSC administration either by subcutaneous injection or implantation into the hypodermis with a polymer or scaffold have shown increased proliferation, and osteoblastic differentiation of BMSCs increased osteogenesis and endogenous bone regeneration.

Itoh et al. (2007) confirmed that FGF/FGFR signaling plays a role in osteogenesis. FGF/FGFR signaling does not directly induce osteoblast differentiation but modulates it. FGF2 and FGF9 likely induce the proliferation of osteoblast cell lineages and the induction of angiogenesis, and FGF18 promotes osteoblast differentiation [85].

Wilkie et al. (2005) published an in vitro analysis of bone marrow-derived MSCs in which FGF18 enhanced osteoblast differentiation by activating FGFR1 or FGFR2 signaling [86]. Additionally, overexpression of FGF18 by lentiviral infection or direct addition of FGF18 to culture media induced the expression of osteoblast marker genes in C3H10T1/2 fibroblastic cells. Treatment with FGF18 in mouse-derived MSCs under differentiation-inducing conditions showed increased expression of osteoblast differentiation markers and mineralization [86].

Low-dose FGF18 treatment with osteogenic induction of bone morphogenetic protein 2 (BMP2)-dependent bone protein from MC3T3-E1 cells increased mineralization, whereas high-dose treatment inhibited the process. Additionally, FGF18-soaked heparin-coated acrylic beads accelerated osteoblast differentiation in mouse fetuses by regulating BMP2 expression in 90 osteoblast cell lineage cells [87]. FGF2 stimulates mitosis and cell proliferation, including of fibroblast and endothelial cells, which plays a vital role in maintaining these cells in tissue repair processes [88]. On the other hand, FGF18 stimulates cellular osteogenesis through the upregulation of bone morphogenetic proteins.

### Potential uses for skin rejuvenation in skin aging

Together with other growth factors and cytokines, EGF directly affects collagen, elastin, and ECM biosynthesis, but its binding and signaling diminish with age. Aged cells in the skin produce ROS, and the mitochondria of these cells disrupt tissue complexes by cleaving membrane-bound receptors, ECM proteins, growth factors, and other signaling ligands in the dermal

microenvironment [89]. Reduced EGF binding and signaling with age can cause collagen degradation in the skin [90, 91]. The rapid degradation of collagen in the skin leads to loss of elasticity and the appearance of skin wrinkling [92]. Decreased expression of EGFR also occurs in aging dermal fibroblasts in the ECM, is associated with reduced cell migration and proliferation, and ultimately leads to skin flexibility and elasticity loss. EGF helps reduce the effects of aging by supporting skin regeneration by stimulating cell renewal through the interaction of keratinocytes and fibroblasts. EGF plays an important role in forming fibroblasts in the dermis by stimulating collagen production via activation of EGFR [89]. The topical use of growth factors is a safe and effective medical treatment [93]. Applying EGF to aging skin can increase fibroblast proliferation [91]. Thus, EGF is a potential therapeutic antiaging agent for the skin.

A clinical trial of human umbilical cord-derived MSC-conditioned media administered by microneedle resulted in good efficacy as an antiaging product and provided an excellent potential for skin rejuvenation [94]. That study reported that the tested conditioned media contained growth factors, including EGF, VEGF-A, VEGF-D, HGF, FGF-2, and others. Furthermore, the administration of MSCs reduced the melanin index and brown spots on the skin. Additionally, wrinkles and skin pores were reduced, and there was an increase in skin elasticity, indicating an improvement in facial skin texture. Another study showed that administering amniotic fluid MSC-derived conditioned media with microneedles to the face improved the skin texture and reduced wrinkles [95]. Moreover, histologically, there was an increase in the number of dermal collagen bundles arranged more regularly, elastic fibers, and epidermis thickening.

AMSC-MP possess various growth factor and cytokines that enables them to become the agent therapeutics of many therapy.

### Future prospects for AMSCMP for tissue engineering

Many reports have demonstrated the potential use of AMSC-MP for tissue regeneration to improve the appearance of facial skin, in bone regeneration, and in tissue or organ repair. Preclinical reports, including in vitro and in vivo studies, have shown that growth factors and cytokines play an essential role in the tissue repair process, both through stimulation of cell differentiation and proliferation and via an indirect effect on regeneration, including antiinflammatory and angiogenic effects. Clinical trials using conditioned media and MSCs have also reported potent activity for tissue regeneration.

However, protein delivery systems of the cytokines and growth factors present in AMSC-MP still have many

shortcomings. Oral delivery is generally preferred, but oral delivery is not a viable method for proteins due to their poor absorption and degradation in the gastrointestinal tract and liver [96]. Oral administration of protein drugs leads to very low bioavailability [97]. Thus, the development of protein formulations is needed to overcome the low permeability of large molecules, the lack of lipophilicity, and rapid inactivation or enzymatic degradation in the gastrointestinal tract as well as protein physicochemical properties that are limiting [97]. For topical uses, growth factors have molecular weights higher than 500 Da, which makes the penetration of the stratum corneum difficult [91]. The ideal characteristics of a substance for a topical delivery system include a relatively low molecular weight (<500 Da), a low melting point (<200 °C), moderate lipophilicity (log P 1–3), and high water solubility (>1 mg/mL), as well as high pharmacological potential [98]. Therefore, a delivery system is needed to help these molecules penetrate the dermis.

The use of AMSC-MP is also limited due to its sensitivity to environmental factors such as temperature, pH, and reactivity during reconstitution. Because of this, the delivery system must optimize the growth factor dose, route of administration, and release kinetics for the safe and effective use of growth factors [99]. For example, KGF has poor *in vivo* bioactivity. KGF has a short biological half-life and poor stability, its biological activity is susceptible to environmental factors, and it cannot maintain bioactivity for a long time in the presence of other enzymes [45].

In addition to stability, using AMSC-MP for tissue regeneration therapy requires a carrier system capable of local delivery with the controlled release of growth factors. The uncontrolled release of growth factors can cause side effects. Using biomaterials as delivery systems is the most successful strategy for controlled delivery, and they have been developed into various commercially available systems [100]. Based on the explanation above, it can be seen that AMSC-MP has been widely studied for its benefits in various diseases. Thus AMSC-MP has potential as a tissue regeneration therapeutic agent.

### **Manufacturing of AMSC-MP for therapeutic products**

Presently, autologous cell therapy is primarily used for stem cell-based therapy. However, the small individual scale, insufficient reagents, and inefficient manufacturing process result in an expensive product. The proper indication for use and timing, adequate dosage, and appropriate route of administration still need to be determined for widespread use.

The possible scaling up of AMSC-MP would represent a more efficient means of manufacturing mass-scale

products to induce tissue regeneration of greater suitability to community needs rather than individual therapy. Moreover, it would reduce the possibility of immune system rejection of cell therapy. The process of manufacturing AMSC-MP from AMSCs is divided into at least three main stages: isolation of mesenchymal cells from placental tissue, cell culture and incubation, and the harvesting and purification of the metabolite products of cell cultures. Quality assurance involving validation needs to be carried out to identify a sustainable production process and guarantee the quality of AMSC-MP. The validation process includes cell culturing, cell stock storage, the harvesting of conditioned media containing cell metabolites, and their processing into AMSC-MP as bioactive materials. AMSC-MP standardization will then be required to ensure consistent and reliable product quality. Quality parameters such as physicochemical characteristics and growth factor content would constitute important specification parameters of bioactive materials. Through validation, mass-production is possible using tailor-made cell lines under controlled laboratory conditions which provide a high-quality source of bioactive factors necessary to produce mass products efficiently and safely.

Furthermore, large-scale manufacturing should be developed to minimize costs and provide affordable therapeutic products for the public; however, the standardization of AMSC-MP into biological products has substantial challenges. Process validation, quality control, and standardized protocols for isolating, culturing, and cultivating AMSC-MP are required, and these are major challenges for manufacturers. Moreover, regulations that support clinical use are also being developed, so AMSC-MP mass production remains under development. Despite these challenges, developing AMSC-MP-based therapeutic products is worthwhile for providing affordable advanced biological therapeutics for clinical practice.

### **Conclusions**

AMSC-MP has excellent potential for use in tissue regeneration therapy since it contains a variety of growth factors that provide better efficacy than single growth factors or cytokines alone. *In vitro* and *in vivo* preclinical studies have shown that AMSC-MP has biological activities related to wound healing, the repair of bone defects and other bone diseases, tissue repair of damaged organs, including the lungs, bladder, kidneys, and in the gastrointestinal tract, as well as skin rejuvenation related to antiaging effects, providing excellent efficacy and demonstrating a good safety profile. Some reports have shown satisfactory results in clinical trials in patients with certain diseases. There is a high clinical demand for

AMSC-MP as an alternative biological therapy, but further development is needed regarding its stability and the identification of a delivery system to provide maximum efficacy. Additionally, product development from the laboratory to a mass production scale requires further effort. Moreover, quality assurance is needed for biological product materials and complex manufacturing processes, which is the main challenge that must be addressed to optimize the use of AMSC-MP as a therapeutic agent.

#### Abbreviations

AMSC	Amniotic Membrane Stem Cell
AMSC-MP	Amniotic Membrane Stem Cell Metabolite Products
BAL	Bronchoalveolar lavage
BMP	Bone morphogenic protein
EGF	Epidermal growth factor
ELISA	Enzyme-linked immunosorbent assay
FD	Freeze-dried
GDF	Growth differentiation factors
HA	Hyaluronic acid
HGF	Hepatic growth factor
IGF	Insulin-like growth factor-I
ILD	Interstitial lung diseases
KGF	Keratinocyte growth factor
KGFR	KGF receptor
MAPK	Mitogen-activated protein kinase
MSC	Mesenchymal stem cells
PDGF	Platelet-derived growth factor
PIGF	Placenta growth factor
ROS	Reactive oxygen species
TGF	Transforming growth factor
TK	Tyrosine kinases
VEGF	Vascular endothelial growth factor

#### Acknowledgements

Not applicable.

#### Authors' contributions

Andang Miatmoko: 1) Conception and design of the work, data acquisition, data analysis, and interpretation; 2) drafting the article; 3) critically revising the article for important intellectual content; 4) final approval of the version to be published; 5) agrees to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of the work are appropriately investigated and resolved. Berlian Sarasitha Hariawan: 1) Conception and design of the work, data acquisition, data analysis, and interpretation; 2) drafting the article; 3) final approval of the version to be published; 4) agrees to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of the work are appropriately investigated and resolved. Devy Maulidya Cahyani: 1) Conception and design of the work, data acquisition, data analysis, and interpretation; 2) drafting the article; 3) final approval of the version to be published; 4) agrees to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of the work are appropriately investigated and resolved. Syarifah Sutra Dewangga: 1) Data analysis and interpretation; 2) drafting the article; 3) final approval of the version to be published; 4) agrees to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of the work are appropriately investigated and resolved. Kevin Ksatria Handoko: 1) Data analysis and interpretation; 2) drafting the article; 3) final approval of the version to be published; 4) agrees to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of the work are appropriately investigated and resolved. Purwati: 1) Data analysis and interpretation; 2) critically revising the article for important intellectual content; 3) final approval of the version to be published; 4) agrees to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of the work are appropriately investigated and resolved. Ram Kumar Sahu: 1) Data analysis and interpretation; 2) critically

revising the article for important intellectual content; 3) final approval of the version to be published; 4) agrees to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of the work are appropriately investigated and resolved. Dewi Melani Hariyadi: 1) Data analysis and interpretation; 2) critically revising the article for important intellectual content; 3) final approval of the version to be published; 4) agrees to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of the work are appropriately investigated and resolved. All authors reviewed the manuscript. The author(s) read and approved the final manuscript.

#### Authors' information

None.

#### Funding

This study was supported by Applied Research on Health (Penelitian Terapan Bidang Kesehatan) Grant No. 1554/UN3.15/PT/2021 provided by the Ministry of Education, Culture, Research and Technology of the Republic of Indonesia.

#### Availability of data and materials

Not applicable.

#### Declarations

#### Ethics approval and consent to participate

Not applicable.

#### Consent for publication

Not applicable.

#### Competing interests

The authors declare no competing interests.

Received: 9 September 2022 Accepted: 6 February 2023

Published online: 09 February 2023

#### References

1. Yardeni D, Ghany MG. Review article: hepatitis B-current and emerging therapies. *Aliment Pharmacol Ther.* 2022;55(7):805–19.
2. Checkoway H, Lundin JI, Kelada SN. Neurodegenerative diseases. *IARC Sci Publ.* 2011;163:407–19.
3. Tomé-Bermejo F, Piñera AR, Alvarez L. Osteoporosis and the management of spinal degenerative disease (II). *Arch Bone Jt Surg.* 2017;5(6):363–74.
4. Cummings JL, Tong G, Ballard C. Treatment combinations for Alzheimer's disease: current and future pharmacotherapy options. *J Alzheimer's Dis.* 2019;67(3):779–94.
5. Yiannopoulou KG, Papageorgiou SG. Current and future treatments in Alzheimer disease: An update. *J Cent Nerv Syst Dis.* 2020;12:117957352090739.
6. Cummings J, Lee G, Ritter A, Zhong K. Alzheimer's disease drug development pipeline: 2018. *Alzheimer's Dement Transl Res Clin Interv.* 2018;2018(4):195–214.
7. Ding M, Tegel H, Sivertsson Å, Hober S, Snijder A, Örmö M, et al. Secretome-based screening in target discovery. *SLAS Discov.* 2020;25(6):535–51.
8. Rahmadewi R, Retha R, Pitasari DA, Kusumastanto VA, Ardhaninggar AAA, Citrashanty I, et al. The efficacy of amniotic membrane stem cell (AMSC) metabolite product and vitamin E for wrinkles, spots, and pores in photoaging. *Dermatol Res Pract.* 2020;2020:1–5.
9. Kropp C, Massai D, Zweigerdt R. Progress and challenges in large-scale expansion of human pluripotent stem cells. *Process Biochem.* 2017;59:244–54.
10. Herberts CA, Kwa MSG, Hermesen HPH. Risk factors in the development of stem cell therapy. *J Transl Med.* 2011;9(1):29.
11. Plotnikov EY, Silachev DN, Popkov VA, Zorova LD, Pevzner IB, Zorov SD, et al. Intercellular signalling cross-talk: to kill, to heal and to rejuvenate. *Hear Lung Circ.* 2017;26(7):648–59.

12. Rikhtegar R, Pezeshkian M, Dolati S, Safaie N, Afrasiabi Rad A, Mahdipour M, et al. Stem cells as therapy for heart disease: iPSCs, ESCs, CSCs, and skeletal myoblasts. *Biomed Pharmacother*. 2018;2019(109):304–13.
13. Tamama K, Kerpedjjeva SS. Acceleration of wound healing by multiple growth factors and cytokines secreted from multipotential stromal cells/mesenchymal stem cells. *Adv Wound Care*. 2012;1(4):177–82.
14. Huddleston HP, Cohn MR, Haunschild ED, Wong SE, Farr J, Yanke AB. Amniotic product treatments: Clinical and basic science evidence. *Curr Rev Musculoskelet Med*. 2020;13(2):148–54.
15. Prakoeswa CRS, Pratiwi FD, Herwanto N, Citrashanty I, Indramaya DM, Murtiastutik D, et al. The effects of amniotic membrane stem cell-conditioned medium on photoaging. *J Dermatolog Treat*. 2019;30(5):478–82.
16. Damayanti RH, Rusdiana T, Wathoni N. Mesenchymal stem cell secretome for dermatology application: A review. *Clin Cosmet Investig Dermatol*. 2021;14:1401–12.
17. Teixeira FG, Serra SC, Salgado AJ. Tips on how to collect and administer the mesenchymal stem cell secretome for central nervous system applications. *Methods Mol Biol*. 2016;1416:457–65.
18. Bennett NT, Schultz GS. Growth factors and wound healing: Biochemical properties of growth factors and their receptors. *Am J Surg*. 1993;165(6):728–37.
19. Sari DIK, Erawati T, Miatmoko A, Prakoeswa CRS, Soeratri W. Characterization and stability study of amniotic membrane stem cell metabolite product (AMSC-MP). *Int J Pharma Res Heal Sci*. 2020;8(1):3126–30.
20. Chen G, Gulbranson DR, Yu P, Hou Z, Thomson JA. Thermal stability of fibroblast growth factor protein is a determinant factor in regulating self-renewal, differentiation, and reprogramming in human pluripotent stem cells. *Stem Cells*. 2012;30(4):623–30.
21. Tan JCG, Webb RG, Marks DC. Serum growth factor stability in different eye drop packaging systems during storage. *Transfus Apher Sci*. 2020;59(1):102608.
22. Mizumachi H, Iijima H. Measuring stability of vascular endothelial growth factor using an immobilization technique. *Adv Biomed Eng*. 2013;2:130–6.
23. Chen BLU, Arakawa T. Stabilization of recombinant human keratinocyte growth factor by osmolytes and salts. *J Pharm Sci*. 1996;85(4):419–22.
24. Zhang HY, Wang ZG, Wu FZ, Kong XX, Yang J, Lin BB, et al. Regulation of autophagy and ubiquitinated protein accumulation by bFGF promotes functional recovery and neural protection in a rat model of spinal cord injury. *Mol Neurobiol*. 2013;48(3):452–64.
25. Beenken A, Mohammadi M. The FGF family: biology, pathophysiology and therapy. *Nat Rev Drug Discov*. 2009;8(3):1–7.
26. Melmed S, Auchus RJ, Goldfine AB, Koenig RJ, Rosen CJ. *Williams Textbook of Endocrinology*. Elsevier. 2020;14:113–113.
27. Esquirol Caussa J, Herrero VE. Epidermal growth factor, innovation and safety. *Med Clínica*. 2015;145(7):305–12.
28. Surini S, Leonyza A, Suh CW. Formulation and in vitro penetration study of recombinant human epidermal growth factor-loaded transdermal emulgel. *Adv Pharm Bull*. 2020;10(4):586–94.
29. Santana H, González Y, Campana PT, Noda J, Amarantes O, Itri R, et al. Screening for stability and compatibility conditions of recombinant human epidermal growth factor for parenteral formulation: Effect of pH, buffers, and excipients. *Int J Pharm*. 2013;452(1–2):52–62.
30. Taylor JM, Mitchell WM, Cohen S. Epidermal growth factor. *J Biol Chem*. 1972;247(18):5928–34.
31. Yang CH, Wu PC, Huang YB, Tsai YH. A new approach for determining the stability of recombinant human epidermal growth factor by thermal Fourier transform infrared (FTIR) microspectroscopy. *J Biomol Struct Dyn*. 2004;22(1):101–10.
32. Laurent GJ, Shapiro SD. *Encyclopedia of Respiratory Medicine*. Elsevier Science. 2006. <https://books.google.co.id/books?id=OnZQAQAAIAAJ>.
33. Chin D, Boyle GM, Parsons PG, Coman WB. What is transforming growth factor-beta (TGF-β)? *Br J Plast Surg*. 2004;57(3):215–21.
34. Attisano L, Ayyaz A, Wrana JL. Recent advances in understanding contextual TGFβ signaling. *F1000Research*. 2017;6(May):1–9.
35. Ong CH, Tham CL, Harith HH, Firdaus N, Israf DA. TGF-β-induced fibrosis: A review on the underlying mechanism and potential therapeutic strategies. *Eur J Pharmacol*. 2021;911(September):174510.
36. Liang J, Zhou Y, Zhang N, Wang D, Cheng X, Li K, et al. The phosphorylation of the Smad2/3 linker region by nemo-like kinase regulates TGF-β signaling. *J Biol Chem*. 2021;296:100512.
37. Yaniz-Galende E, Hajjar RJ. Stem cell and gene therapy for cardiac regeneration. *Cardiac Regeneration and Repair*, vol. 1. Woodhead Publishing Limited. 2014. <https://doi.org/10.1533/9780857096708.4.347>.
38. Duffy AM, Bouchier-Hayes DJ, Harmey JH. Vascular endothelial growth factor (VEGF) and its role in non-endothelial cells: Autocrine signalling by VEGF. *VEGF Cancer*. 2004;1:133–44.
39. Melincovici CS, Boşca AB, Şuşman S, Mărginean M, Mihu C, Istrate M, et al. Vascular endothelial growth factor (VEGF) – key factor in normal and pathological angiogenesis. *Rom J Morphol Embryol*. 2018;59(2):455–67.
40. De Falco S. The discovery of placenta growth factor and its biological activity. *Exp Mol Med*. 2012;44(1):1–9.
41. Rubin JS, Bottaro DP, Chedid M, Miki T, Ron D, Cheon G, et al. Keratinocyte growth factor. *Cell Biol Int*. 1996;19(5):399–411.
42. Poorebrahim M, Sadeghi S, Ghorbani R, Asghari M, Abazari MF, Kalhor H, et al. In silico enhancement of the stability and activity of keratinocyte growth factor. *J Theor Biol*. 2017;418(January):111–21.
43. Borjian Boroujeni M, Shahbazi Dastjerdeh M, Shokrgozar MA, Rahimi H, Omidinia E. Computational driven molecular dynamics simulation of keratinocyte growth factor behavior at different pH conditions. *Informatics Med Unlocked*. 2021;23(October):100514.
44. Hsu E, Osslund T, Nybo R, Chen BL, Kenney WC, Morris CF, et al. Enhanced stability of recombinant keratinocyte growth factor by mutagenesis. *Protein Eng Des Sel*. 2006;19(4):147–53.
45. Sadeghi S, Kalhor H, Panahi M, Abolhasani H, Rahimi B, Kalhor R, et al. Keratinocyte growth factor in focus: A comprehensive review from structural and functional aspects to therapeutic applications of palifermin. *Int J Biol Macromol*. 2021;191(September):1175–90.
46. Hadari YR, Gotoh N, Kouhara H, Lax I, Schlessinger J. Critical role for the docking-protein FRS2α in FGF receptor-mediated signal transduction pathways. *Proc Natl Acad Sci U S A*. 2001;98(15):8578–83.
47. Chang HL, Sugimoto Y, Liu S, Wang LS, Huang YW, Ye W, et al. Keratinocyte growth factor (KGF) regulates estrogen receptor-α (ER-α) expression and cell apoptosis via phosphatidylinositol 3-kinase (PI3K)/Akt pathway in human breast cancer cells. *Anticancer Res*. 2009;29(8):3195–205.
48. Ceccarelli S, Cardinali G, Aspite N, Picardo M, Marchese C, Torrisi MR, et al. Cortactin involvement in the keratinocyte growth factor and fibroblast growth factor 10 promotion of migration and cortical actin assembly in human keratinocytes. *Exp Cell Res*. 2007;313(9):1758–77.
49. Braun S, Auf Dem Keller U, Steiling H, Werner S. Fibroblast growth factors in epithelial repair and cytoprotection. *Philos Trans R Soc B Biol Sci*. 2004;359(1445):753–7.
50. Lee JS, Liu JJ, Hong JW, Wilson SE. Differential expression analysis by gene array of cell cycle modulators in human corneal epithelial cells stimulated with epidermal growth factor (EGF), hepatocyte growth factor (HGF), or keratinocyte growth factor (KGF). *Curr Eye Res*. 2001;23(1):69–76.
51. Spielberger R, Stiff P, Bensinger W, Gentile T, Weisdorf D, Kewalramani T, et al. Palifermin for Oral Mucositis after Intensive Therapy for Hematologic Cancers. *N Engl J Med*. 2004;351(25):2590–8.
52. Wilkinson HN, Hardman MJ. Wound healing: cellular mechanisms and pathological outcomes: Cellular Mechanisms of Wound Repair. *Open Biol*. 2020;10(9):200223.
53. Huang Y, Gou M, Zhang W, Xie H. Mesenchymal Stem Cells for Chronic Wound Healing : Current Status of Preclinical and Clinical Studies. *Tissue Eng - Part B*. 2020;26(6):555–70.
54. Steed DL. The Role of Growth Factor in Wound Healing. *Surg Clin North Am*. 1997;77(3):13–5.
55. Desai TJ, Cardoso WV. Growth factors in lung development and disease: Friends and foe? *Respir Res*. 2001;3:1–4.
56. Qu Y, Cao C, Wu Q, Huang A, Song Y, Li H, et al. The dual delivery of KGF and bFGF by collagen membrane to promote skin wound healing. *J Tissue Eng Regen Med*. 2018;12(6):1508–18.
57. Braiman-Wikman L, Solomonik I, Spira R, Tennenbaum T. Novel Insights into Wound Healing Sequence of Events. *Toxicol Pathol*. 2007;35(6):767–79.
58. Akita S, Akino K, Hirano A. Basic Fibroblast Growth Factor in Scarless Wound Healing. *Adv Wound Care*. 2013;2(2):44–9.
59. Zhang X, Kang X, Jin L, Bai J, Liu W, Wang Z. Stimulation of wound healing using bioinspired hydrogels with basic fibroblast growth factor (bFGF). *Int J Nanomedicine*. 2018;13:3897–906.

60. Li J, Ren K, Zhang W, Xiao L, Wu H, Liu Q, et al. Human amniotic mesenchymal stem cells and their paracrine factors promote wound healing by inhibiting heat stress-induced skin cell apoptosis and enhancing their proliferation through activating PI3K / AKT signaling pathway. *Stem Cell Res Ther.* 2019;10(247):1–17.
61. Reza H, Motahareh A, Hosseini S, Gholami M, Mohamadi F. Mesenchymal stem cells' seeded amniotic membrane as a tissue - engineered dressing for wound healing. *Drug Deliv Transl Res.* 2022;12:538–49.
62. Jones MK, Tomikawa M, Mohajer B, Tarnawski AS. Gastrointestinal mucosal regeneration: role of growth factors. *Front Biosci.* 1999;4:D303–9.
63. Wei G, Wang Z, Liu R, Zhou C, Li E, Shen T, et al. A combination of hybrid polydopamine-human keratinocyte growth factor nanoparticles and sodium hyaluronate for the efficient prevention of postoperative abdominal adhesion formation. *Acta Biomater.* 2022;138:155–67.
64. Zhao YZ, ZhuGe DL, Tong MQ, Lin MT, Zheng YW, Jiang X, et al. Ulcerative colitis-specific delivery of keratinocyte growth factor by neutrophils-simulated liposomes facilitates the morphologic and functional recovery of the damaged colon through alleviating the inflammation. *J Control Release.* 2019;299:90–106.
65. Zeeh JM, Procaccino F, Hoffmann P, Aukerman SL, McRoberts JA, Soltani S, et al. Keratinocyte growth factor ameliorates mucosal injury in an experimental model of colitis in rats. *Gastroenterology.* 1996;110(4):1077–83.
66. Sandborn WJ, Sands BE, Wolf DC, Valentine JF, Safdi M, Katz S, et al. Repifermin (keratinocyte growth factor-2) for the treatment of active ulcerative colitis: a randomized, double-blind, placebo-controlled, dose-escalation trial. *Aliment Pharmacol Ther.* 2003;17(11):1355–64.
67. Itoh M, Matsuo Y. Gastric ulcer treatment with intravenous human epidermal growth factor: A double-blind controlled clinical study. *J Gastroenterol Hepatol.* 1994;9(1 S):S78–83.
68. Gomperts BN, Belperio JA, Fishbein MC, Keane MP, Burdick MD, Strieter RM. Keratinocyte growth factor improves repair in the injured tracheal epithelium. *Am J Respir Cell Mol Biol.* 2007;37(1):48–56.
69. Tillie-Leblond I, Gosset P, Le Berre R, Janin A, Prangère T, Tonnel AB, et al. Keratinocyte growth factor improves alterations of lung permeability and bronchial epithelium in allergic rats. *Eur Respir J.* 2007;30(1):31–9.
70. Wang J, Sun H, Liu Y. The proliferative and anti-apoptosis functions of KGF/KGFR contributes to bronchial epithelial repair in asthma. *Pulm Pharmacol Ther.* 2020;63:101931.
71. Kan B, Jian X, Zhou Q, Wang J, Yu G, Sun J, et al. Effect of transforming growth factor- $\beta$ 1 on acute lung injury caused by paraquat. *Mol Med Rep.* 2014;9(4):1232–6.
72. Kopyński P, Śladek K, Szczekliki J, Soja J, Szlubowski A, Balicka-Ślusarczyk B, et al. Expression of insulin-like growth factor-I (IGF-I) in alveolar macrophages and lymphocytes obtained by bronchoalveolar lavage (BAL) in interstitial lung diseases (ILD). Assessment of IGF-I as a potential local mitogen and antiapoptotic cytokine. *Folia Histochem Cytobiol.* 2006;44(4):249–58.
73. Lutmer J, Watkins D, Chen CL, Velten M, Besner G. Heparin-binding epidermal growth factor-like growth factor attenuates acute lung injury and multiorgan dysfunction after scald burn. *J Surg Res.* 2013;185(1):329–37.
74. Pan X, Xu S, Zhou Z, Wang F, Mao L, Li H, et al. Fibroblast growth factor-2 alleviates the capillary leakage and inflammation in sepsis. *Mol Med.* 2020;26(1):108.
75. Baskin LS, Sutherland RS, Thomson AA, Nguyen HT, Morgan DM, Hayward SW, et al. Growth factors in bladder wound healing. *J Urol.* 1997;157(6):2388–95.
76. Narla ST, Bushnell DS, Schaefer CM, Nouraie M, Bates CM. Keratinocyte growth factor reduces injury and leads to early recovery from cyclophosphamide bladder injury. *Am J Pathol.* 2020;190(1):108–24.
77. Jaal J, Dörr W. Effect of recombinant human keratinocyte growth factor (rHuKGF, Palifermin) on radiation-induced mouse urinary bladder dysfunction. *Int J Radiat Oncol Biol Phys.* 2007;69(2):528–33.
78. Chen W, Shi C, Yi S, Chen B, Zhang W, Fang Z, et al. Bladder regeneration by collagen scaffolds with collagen binding human basic fibroblast growth factor. *J Urol.* 2010;183(6):2432–9.
79. Chen G, Deng C, Li YP. TGF- $\beta$  and BMP signaling in osteoblast differentiation and bone formation. *Int J Biol Sci.* 2012;8(2):272–88.
80. Zou X, Chen K, Zou J, Han P, Hao J, Han Z. Single-cell RNA-seq data analysis on the receptor ACE2 expression reveals the potential risk of different human organs vulnerable to 2019-nCoV infection. *Front Med.* 2020;14(2):185–92.
81. Stewart S, Bryant SJ, Ahn J, Hankenson KD. Bone Regeneration. *Translational Regenerative Medicine.* Elsevier Inc. 2015. <https://doi.org/10.1016/B978-0-12-410396-2.00024-4>.
82. Zhu G, Zhang T, Chen M, Yao K, Huang X, Zhang B, et al. Bone physiological microenvironment and healing mechanism: Basis for future bone-tissue engineering scaffolds. *Bioact Mater.* 2021;6(11):4110–40.
83. Stewart S, Bryant SJ, Ahn J, Hankenson KD. Chapter 24 - Bone Regeneration. In: Atala A, Allickson JG, editors. *Translational Regenerative Medicine.* Boston: Academic Press; 2015. p. 313–33.
84. Li J, Zhou Z, Wen J, Jiang F, Xia Y. Human amniotic mesenchymal stem cells promote endogenous bone regeneration. *Front Endocrinol.* 2020;11(October):1–9.
85. Itoh N. The Fgf families in humans, mice, and zebrafish: Their evolutionary processes and roles in development, metabolism, and disease. *Biol Pharm Bull.* 2007;30(10):1819–25.
86. Wilkie AOM. Bad bones, absent smell, selfish testes: The pleiotropic consequences of human FGF receptor mutations. *Cytokine Growth Factor Rev.* 2005;16(2 SPEC. ISS):187–203.
87. Nagayama T, Okuhara S, Ota MS, Tachikawa N, Kasugai S, Iseki S. FGF18 accelerates osteoblast differentiation by upregulating Bmp2 expression. *Congenit Anom.* 2013;53(2):83–8.
88. Morimoto N, Yoshimura K, Niimi M, Ito T, Aya R, Fujitaka J, et al. Novel collagen/gelatin scaffold with sustained release of basic fibroblast growth factor: Clinical trial for chronic skin ulcers. *Tissue Eng - Part A.* 2013;19(17–18):1931–40.
89. Kim D, Kim SY, Mun SK, Rhee S, Kim BJ. Epidermal growth factor improves the migration and contractility of aged fibroblasts cultured on 3D collagen matrices. *Int J Mol Med.* 2015;35(4):1017–25.
90. Ha JM, Lim CA, Han K, Ha JC, Lee HE, Lee Y, et al. The effect of microspicule containing epidermal growth factor on periorcular wrinkles. *Ann Dermatol.* 2017;29(2):187–93.
91. Eskens O, Amin S. Challenges and effective routes for formulating and delivery of epidermal growth factors in skin care. *Int J Cosmet Sci.* 2021;43(2):123–30.
92. Swift A, Liew S, Weinkle S, Garcia JK, Silberberg MB. The facial aging process from the "inside Out." *Aesthetic Surg J.* 2021;41(10):1107–19.
93. Fabi S, Sundaram H. The potential of topical and injectable growth factors and cytokines for skin rejuvenation. *Facial Plast Surg.* 2014;30(2):157–71.
94. Liang X, Li J, Yan Y, Xu Y, Wang X, Wu H. Efficacy of microneedling combined with local application of human umbilical cord-derived mesenchymal stem cells conditioned media in skin brightness and rejuvenation : A randomized controlled split-face study. *Front Med.* 2022;9:837332.
95. El-Domyati M, Mofteh NH, Nasif GA, Ameen SW, Ibrahim MR, Ragaie MH. Facial rejuvenation using stem cell conditioned media combined with skin needling: A split-face comparative study. *J Cosmet Dermatol.* 2020;19(9):2404–10.
96. Singh R, Singh S, Lillard JW. Past, present, and future technologies for oral delivery of therapeutic proteins. *J Pharm Sci.* 2008;97:2497–523.
97. Park K, Kwon IC, Park K. Oral protein delivery: Current status and future prospect. *React Funct Polym.* 2011;71(3):280–7.
98. Benson HAE, Namjoshi S. Proteins and peptides: Strategies for delivery to and across the skin. *J Pharm Sci.* 2008;97(9):3591–610.
99. Subbiah R, Guldberg RE. Materials science and design principles of growth factor delivery systems in tissue engineering and regenerative medicine. *Adv Healthc Mater.* 2019;8(1):1–24.
100. Atienza-Roca P, Cui X, Hooper GJ, Woodfield TBF, Lim KS. Growth factor delivery systems for tissue engineering and regenerative medicine. *Adv Exp Med Biol.* 2018;1078:245–69.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



## Journal of Biological Engineering

### COUNTRY

United Kingdom



Universities and research institutions in United Kingdom



Media Ranking in United Kingdom

### SUBJECT AREA AND CATEGORY

Biochemistry, Genetics and Molecular Biology

Cell Biology

Molecular Biology

Engineering

Biomedical Engineering

Environmental Science

Environmental Engineering

### PUBLISHER

BioMed

Central

Ltd.

### H-INDEX

44

### PUBLICATION TYPE

Journals

### ISSN

17541611

### COVERAGE

2007-  
2021

### INFORMATION

[Homepage](#)

[How to  
publish in  
this journal](#)

### SCOPE

Biological engineering is an emerging discipline that encompasses engineering theory and practice connected to and derived from the science of biology, just as mechanical engineering and electrical engineering are rooted in physics and chemical engineering in chemistry. Topical areas include, but are not limited to: Synthetic biology and cellular design Biomolecular, cellular and tissue


engineering Bioproduction and metabolic engineering Biosensors Ecological and environmental engineering Biological engineering education and the biodesign process As the official journal of the Institute of Biological Engineering, Journal of Biological Engineering provides a home for the continuum from biological information science, molecules and cells, product formation, wastes and remediation, and educational advances in curriculum content and pedagogy at the undergraduate and graduate-levels. Manuscripts should explore commonalities with other fields of application by providing some discussion of the broader context of the work and how it connects to other areas within the field.







Join the conversation about this journal

Quartiles

FIND SIMILAR JOURNALS ?

<p>1 <b>Frontiers in Bioengineering and Biotechnology</b> CHE</p> <p><b>79%</b> similarity</p>	<p>2 <b>Trends in Biotechnology</b> GBR</p> <p><b>71%</b> similarity</p>	<p>3 <b>Advanced Biosystems</b> DEU</p> <p><b>71%</b> similarity</p>	<p>4 <b>Advances in Experimental Medicine and Biology</b> USA</p> <p><b>68%</b> similarity</p>
--	--	--	--





**Journal of Biological Engineering**

Q1 Biomedical Engineering  
best quartile

SJR 2021 1.05

powered by scimagojr.com


← Show this widget in your own website

Just copy the code below and paste within your html code:

`<a href="https://www.scimaç`

**SCImago Graphica**

Explore, visually communicate and make sense of data with our **new data visualization tool.**





**Leave a comment**

**Name**

**Email**

(will not be published)

I'm not a robot reCAPTCHA  
Privacy - Terms

Submit

The users of Scimago Journal & Country Rank have the possibility to dialogue through comments linked to a specific journal. The purpose is to have a forum in which general doubts about the processes of publication in the journal, experiences and other issues derived from the publication of papers are resolved. For topics on particular articles, maintain the dialogue through the usual channels with your editor.

---

Developed by:



Powered by:



Follow us on @ScimagoJR

Scimago Lab, Copyright 2007-2022. Data Source: Scopus®

EST MODUS IN REBUS

Horatio (Satire 1, 1, 106)

[Cookie settings](#)

[Cookie policy](#)

---

# Source details

## Journal of Biological Engineering

Open Access ⓘ

Scopus coverage years: from 2007 to Present

Publisher: Springer Nature

ISSN: 1754-1611

Subject area: Environmental Science: Environmental Engineering Engineering: Biomedical Engineering  
Biochemistry, Genetics and Molecular Biology: Molecular Biology  
Biochemistry, Genetics and Molecular Biology: Cell Biology

CiteScore 2021 **9.0** ⓘ

SJR 2021 **1.053** ⓘ

SNIP 2021 **1.183** ⓘ

Source type: Journal

[View all documents >](#) [Set document alert](#) [Save to source list](#) [Source Homepage](#)


[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)

[Export content for category](#)

CiteScore rank ⓘ 2021 In category: Environmental Engineering

☆	Rank	Source title	CiteScore 2021	Percentile
☆	#18	Journal of Biological Engineering	9.0	89th percentile
	173			
☆	#1	Critical Reviews in Environmental Science and Technology	18.2	99th percentile
☆	#2	Water Research	18.0	99th percentile
☆	#3	Bioresource Technology	17.4	98th percentile
☆	#4	Reviews in Environmental Science and Biotechnology	16.0	97th percentile
☆	#5	Journal of Hazardous Materials	14.7	97th percentile
☆	#6	Engineering	14.5	96th percentile
☆	#7	Science of the Total Environment	14.1	96th percentile
☆	#8	Biofuel Research Journal	13.0	95th percentile
☆	#9	Chemosphere	11.7	95th percentile
☆	#10	Journal of Environmental Management	11.4	94th percentile

### CiteScore trend



☆	Rank	Source title	CiteScore 2021	Percentile
☆	#11	Agronomy for Sustainable Development	11.1	93rd percentile
☆	#12	Journal of Environmental Sciences	11.1	93rd percentile
☆	#13	Building and Environment	10.7	92nd percentile
☆	#14	Sustainable Environment Research	10.2	92nd percentile
☆	#15	Process Safety and Environmental Protection	9.9	91st percentile
☆	#16	Environmental Modelling and Software	9.5	91st percentile
☆	#17	Journal of King Saud University, Engineering Sciences	9.2	90th percentile
☆	#18	Journal of Biological Engineering	9.0	89th percentile
☆	#19	Coastal Engineering	8.6	89th percentile
☆	#20	Indoor Air	8.4	88th percentile
☆	#21	Journal of Flood Risk Management	8.2	88th percentile
☆	#22	Sustainable Production and Consumption	8.1	87th percentile
☆	#23	Groundwater for Sustainable Development	7.9	86th percentile
☆	#24	Ecological Engineering	7.7	86th percentile
☆	#25	Environmental Science: Water Research and Technology	7.6	85th percentile
☆	#26	International Journal of Sustainable Transportation	7.6	85th percentile
☆	#27	Elementa	7.5	84th percentile
☆	#28	Journal of Aerosol Science	7.4	84th percentile
☆	#29	AIChE Journal	7.1	83rd percentile
☆	#30	Ecological Engineering: X	7.1	82nd percentile
☆	#31	Environmental Geochemistry and Health	6.9	82nd percentile
☆	#32	Journal of Polymers and the Environment	6.8	81st percentile
☆	#33	Biochemical Engineering Journal	6.7	81st percentile
☆	#34	Ocean Engineering	6.5	80th percentile
☆	#35	Journal of Sustainable Mining	6.4	80th percentile

☆	Rank	Source title	CiteScore 2021	Percentile
☆	#36	Stochastic Environmental Research and Risk Assessment	6.4	79th percentile
☆	#37	Bioresource Technology Reports	6.3	78th percentile
☆	#38	Clean Technologies and Environmental Policy	6.2	78th percentile
☆	#39	Biodegradation	6.0	77th percentile
☆	#40	Global Journal of Environmental Science and Management	6.0	77th percentile
☆	#41	Waste Management and Research	5.9	76th percentile
☆	#42	Journal of Environmental Quality	5.8	76th percentile
☆	#43	International Journal of Environmental Science and Technology	5.6	75th percentile
☆	#44	Energy and Environment	5.5	74th percentile
☆	#45	Environmental Engineering Research	5.5	74th percentile
☆	#46	Journal of Ocean Engineering and Science	5.4	73rd percentile
☆	#47	Engineering in Life Sciences	5.4	73rd percentile
☆	#48	Chinese Journal of Chemical Engineering	5.3	72nd percentile
☆	#49	Environmental Technology Reviews	5.3	71st percentile
☆	#50	Waste and Biomass Valorization	5.1	71st percentile
☆	#51	Journal of Hydro-Environment Research	5.1	70th percentile
☆	#52	Remediation	5.0	70th percentile
☆	#53	Environmental Science and Ecotechnology	4.9	69th percentile
☆	#54	International Journal of Energy and Environmental Engineering	4.8	69th percentile
☆	#55	Journal of Water Supply: Research and Technology - AQUA	4.7	68th percentile
☆	#56	Environmental Progress and Sustainable Energy	4.6	67th percentile
☆	#57	Case Studies in Chemical and Environmental Engineering	4.5	67th percentile
☆	#58	Ozone: Science and Engineering	4.4	66th percentile
☆	#59	Environmental Geotechnics	4.4	66th percentile
☆	#60	Mediterranean Marine Science	4.3	65th percentile

☆	Rank	Source title	CiteScore 2021	Percentile
☆	#61	IET Energy Systems Integration	4.2	65th percentile
☆	#62	European Journal of Environmental and Civil Engineering	4.2	64th percentile
☆	#62	International Journal of Geotechnical Engineering	4.2	64th percentile
☆	#64	Clean Energy	4.1	63rd percentile
☆	#65	Environmental Processes	4.0	62nd percentile
☆	#66	ISH Journal of Hydraulic Engineering	4.0	62nd percentile
☆	#67	Water Cycle	3.9	61st percentile
☆	#68	Water, Air, and Soil Pollution	3.8	60th percentile
☆	#69	Greenhouse Gases: Science and Technology	3.8	60th percentile
☆	#70	Journal of Environmental Science and Health - Part A Toxic/Hazardous Substances and Environmental Engineering	3.7	59th percentile
☆	#71	Nanotechnology for Environmental Engineering	3.7	59th percentile
☆	#72	Civil Engineering Journal (Iran)	3.6	58th percentile
☆	#73	Science and Technology for the Built Environment	3.5	58th percentile
☆	#74	Water Science and Technology	3.4	57th percentile
☆	#75	International Journal on Energy Conversion	3.4	56th percentile
☆	#76	BioResources	3.4	56th percentile
☆	#77	Journal of Environmental Engineering, ASCE	3.3	55th percentile
☆	#78	Paddy and Water Environment	3.2	55th percentile
☆	#79	Theoretical and Applied Mechanics Letters	3.1	54th percentile
☆	#80	Ecological Chemistry and Engineering S	3.1	54th percentile
☆	#81	Journal of Environmental Health Science and Engineering	3.0	53rd percentile
☆	#82	Environment	2.9	52nd percentile
☆	#83	Water and Environment Journal	2.9	52nd percentile
☆	#84	Journal of Environmental and Engineering Geophysics	2.8	51st percentile

☆	Rank	Source title	CiteScore 2021	Percentile
☆	#85	International Journal of Renewable Energy Development	2.7	51st percentile
☆	#86	Journal of Hazardous, Toxic, and Radioactive Waste	2.7	50th percentile
☆	#87	Innovative Infrastructure Solutions	2.6	50th percentile
☆	#88	Energy, Environment, and Sustainability	2.6	49th percentile
☆	#89	Geosystem Engineering	2.4	48th percentile
☆	#90	Transportation Infrastructure Geotechnology	2.3	48th percentile
☆	#91	Journal of Engineering Thermophysics	2.3	47th percentile
☆	#92	Detritus	2.2	47th percentile
☆	#93	Open Engineering	2.2	46th percentile
☆	#94	Revista Brasileira de Engenharia Agricola e Ambiental	2.1	45th percentile
☆	#95	Motivation Science	2.1	45th percentile
☆	#96	Journal of Environmental Engineering and Landscape Management	2.0	44th percentile
☆	#97	Resources, Environment and Sustainability	2.0	44th percentile
☆	#98	Applied Environmental Biotechnology	2.0	43rd percentile
☆	#99	Journal of Green Building	1.9	43rd percentile
☆	#100	Journal of Water and Land Development	1.8	42nd percentile
☆	#101	Journal of Water and Environment Technology	1.8	41st percentile
☆	#102	International Journal of Social Ecology and Sustainable Development	1.8	41st percentile
☆	#103	Environmental Engineering and Management Journal	1.8	40th percentile
☆	#104	Environmental Research, Engineering and Management	1.7	40th percentile
☆	#105	Journal of Integrated Coastal Zone Management	1.7	39th percentile
☆	#106	International Journal of GEOMATE	1.7	39th percentile
☆	#107	International Journal of Hydrology Science and Technology	1.6	38th percentile
☆	#108	Acta Hydrotechnica	1.5	37th percentile

☆	Rank	Source title	CiteScore 2021	Percentile
☆	#108	Greening of Industry Networks Studies	1.5	37th percentile
☆	#110	Acta Innovations	1.5	36th percentile
☆	#111	Environmental and Engineering Geoscience	1.4	36th percentile
☆	#112	Journal of Environmental Biology	1.3	35th percentile
☆	#113	Transactions of the Royal Institution of Naval Architects Part A: International Journal of Maritime Engineering	1.3	34th percentile
☆	#114	International Journal of Environmental Technology and Management	1.3	34th percentile
☆	#115	WSEAS Transactions on Environment and Development	1.2	33rd percentile
☆	#116	Huanjing Kexue Xuebao / Acta Scientiae Circumstantiae	1.2	33rd percentile
☆	#117	Cleaner Environmental Systems	1.2	32nd percentile
☆	#118	International Review of Applied Sciences and Engineering	1.2	31st percentile
☆	#118	Journal of Korean Society for Atmospheric Environment	1.2	31st percentile
☆	#120	Journal of Urban and Environmental Engineering	1.2	30th percentile
☆	#121	Mathematical and Computational Forestry and Natural-Resource Sciences	1.1	30th percentile
☆	#122	Geomatics and Environmental Engineering	1.1	29th percentile
☆	#123	Civil and Environmental Engineering	1.1	29th percentile
☆	#124	Journal of Resources and Ecology	1.0	28th percentile
☆	#125	Advances in Technology Innovation	1.0	28th percentile
☆	#126	Journal of Marine Environmental Engineering	0.9	27th percentile
☆	#127	Cleaner Engineering and Technology	0.9	26th percentile
☆	#128	Environmental Protection Engineering	0.9	26th percentile
☆	#129	Italian Journal of Engineering Geology and Environment	0.9	25th percentile
☆	#130	International Journal of Environment and Waste Management	0.9	25th percentile
☆	#131	Nafta - Gaz	0.9	24th percentile



☆	Rank	Source title	CiteScore 2021	Percentile
☆	#132	Environmental Challenges	0.9	23rd percentile
☆	#133	Acque Sotterranee - Italian Journal of Groundwater	0.9	23rd percentile
☆	#134	Water Resources Development and Management	0.9	22nd percentile
☆	#135	Sustainable Production, Life Cycle Engineering and Management	0.8	22nd percentile
☆	#136	Bauphysik	0.8	21st percentile
☆	#137	Cleaner and Responsible Consumption	0.8	21st percentile
☆	#138	Progress in Agricultural Engineering Sciences	0.8	20th percentile
☆	#139	Geingegneria Ambientale e Mineraria	0.8	19th percentile
☆	#140	Sustainable Civil Infrastructures	0.7	19th percentile
☆	#141	Current Applied Science and Technology	0.7	18th percentile
☆	#142	Chemistry, Didactics, Ecology, Metrology	0.7	18th percentile
☆	#143	Clinical Archives of Communication Disorders	0.7	17th percentile
☆	#144	Ecological Chemistry and Engineering A	0.6	17th percentile
☆	#145	Informes de la Construccion	0.6	16th percentile
☆	#146	Journal of Irrigation and Drainage	0.6	15th percentile
☆	#147	Journal of Environmental Engineering (Japan)	0.6	15th percentile
☆	#148	Journal of Environmental Health and Sustainable Development	0.6	14th percentile
☆	#149	Journal of Japan Society of Civil Engineers	0.5	14th percentile
☆	#150	Ecological Engineering and Environmental Technology	0.5	13th percentile
☆	#151	International Journal of Sustainable Construction Engineering and Technology	0.5	13th percentile
☆	#152	Agricultural Engineering	0.5	12th percentile
☆	#152	Chinese Journal of Environmental Engineering	0.5	12th percentile
☆	#154	Waste Forum	0.5	11th percentile
☆	#155	ASEAN Engineering Journal	0.5	10th percentile
☆	#156	Aqua Water Infrastructure, Ecosystems and Society	0.4	10th percentile

☆	Rank	Source title	CiteScore 2021	Percentile
☆	#157	Pakistan Journal of Engineering and Applied Sciences	0.4	9th percentile
☆	#158	Environmental Science and Engineering	0.3	8th percentile
☆	#158	Paliva	0.3	8th percentile
☆	#160	Journal of Renewable Energy and Environment	0.3	7th percentile
☆	#161	Advances in Environmental Technology	0.3	7th percentile
☆	#162	Sciences in Cold and Arid Regions	0.2	6th percentile
☆	#163	Taiwan Water Conservancy	0.2	6th percentile
☆	#164	Journal of Engineering Geology	0.2	5th percentile
☆	#165	Clean Air and Containment Review	0.2	4th percentile
☆	#166	Vytapeni, Vetrani, Instalace	0.2	4th percentile
☆	#167	Environmental Engineering(Lithuania)	0.2	3rd percentile
☆	#168	Journal of the IEST	0.1	3rd percentile
☆	#169	Advances in Geophysical and Environmental Mechanics and Mathematics	0.1	2nd percentile
☆	#170	Revista Politecnica	0.0	2nd percentile
☆	#171	Avicenna Journal of Environmental Health Engineering	0.0	0th percentile
☆	#171	Energy Studies Review	0.0	0th percentile
☆	#171	Exploration Geophysics, Remote Sensing and Environment	0.0	0th percentile

---

## About Scopus

[What is Scopus](#)

[Content coverage](#)

[Scopus blog](#)

[Scopus API](#)

[Privacy matters](#)

## Language

[日本語版を表示する](#)

[查看简体中文版本](#)

[查看繁體中文版本](#)

[Просмотр версии на русском языке](#)

## Customer Service

[Help](#)

[Tutorials](#)

[Contact us](#)

---

## ELSEVIER

[Terms and conditions](#) ↗ [Privacy policy](#) ↗

Copyright © Elsevier B.V. ↗. All rights reserved. Scopus® is a registered trademark of Elsevier B.V.

We use cookies to help provide and enhance our service and tailor content. By continuing, you agree to the use of cookies ↗.

