



# Frontiers in Pharmacology





Search in this journal

## 11,257 editors

[Role](#)   [Most viewed](#)   [Most publications](#)[Role ▾](#)[Sections ▾](#)

### Field Chief Editor

**Heike  
Wulff**University of  
California...Davis ,  
United States[Field Chief...  
Frontiers in  
Pharmacology](#)

### Specialty Chief Editors

**Giuseppe  
Remuzzi**  
Istituto di  
Ricerche...  
Bergamo , Italy[Specialty Chief...  
Renal  
Pharmacology](#)**Dianwen Ju**  
School of  
Pharmacy, Fuda...  
Shanghai , China[Specialty Chief Editor  
Pharmacology of  
Anti-Cancer Drugs](#)**Eliot Ohlstein**  
Drexel University  
Philadelphia , United  
States[Specialty Chief Editor  
Cardiovascular and  
Smooth Muscle...](#)**Paolo  
Montuschi**  
Catholic  
University of th...  
Milano , Italy[Specialty Chief...  
Respiratory  
Pharmacology](#)



**Dieter  
Steinhilber**  
Goethe  
University...  
Frankfurt ,  
Germany



**Jaime Kapitulnik**  
Hebrew University  
of Jerusalem  
Jerusalem , Israel



**Michael Heinrich**  
School of Pharmacy,  
Faculty of Life...  
London , United  
Kingdom



**Dov  
Greenbaum**  
Yale University  
New Haven ,  
United States

Specialty Chief...  
Inflammation  
Pharmacology

Specialty Chief Editor  
Drug Metabolism and  
Transport

Specialty Chief Editor  
Ethnopharmacology

Specialty Chief...  
ELSI in Science and  
Genetics



**José A G  
Agúndez**  
Institute of  
Molecular...  
Cáceres , Spain



**Ursula Gundert-  
Remy**  
Charité  
Universitätsmediz...  
Berlin , Germany



**Philippe Lory**  
Institut de  
Génomique...  
Montpellier , France



**Nicholas M  
Barnes**  
University of  
Birmingham  
Birmingham ,  
United Kingdom

Specialty Chief...  
Pharmacogenetics  
and...

Specialty Chief Editor  
Predictive Toxicology

Specialty Chief Editor  
Pharmacology of Ion  
Channels and...

Specialty Chief...  
Neuropharmacology



**Francesco  
Rossi**  
University of  
Campania Lui...  
Caserta , Italy



**Angelo A Izzo**  
University of  
Naples Federico II  
Naples , Italy



**Anick Bérard**  
Université de  
Montréal  
Montreal , Canada



**George Joseph  
Christ**  
University of  
Virginia  
Charlottesville ,  
United States

Specialty Chief...  
Cardiovascular and  
Smooth Muscle...

Specialty Chief Editor  
Gastrointestinal and  
Hepatic...

Specialty Chief Editor  
Pharmacoepidemiology

Specialty Chief...  
Integrative and  
Regenerative...



**Alex F. Chen**  
Xiangya Hospital,  
Central South...  
Changsha .



**Jun-Yan Liu**  
Chongqing  
Medical University  
Chonqaina . China



**Olivier Feron**  
Pôle de  
Pharmacologie et ...  
Brussels . Belaium



**Jeffrey Scott  
Barrett**  
Critical Path  
Institute

**Catherine M T  
Sherwin**Wright State  
UniversityDayton , United  
StatesSpecialty Chief...  
Obstetric and  
Pediatric...**Hendrik W. Van  
Veen**University of  
CambridgeCambridge , United  
KingdomSpecialty Chief Editor  
Pharmacology of  
Infectious Diseases**Jean-Marie  
Boeynaems**Université libre de  
Bruxelles

Brussels , Belgium

Specialty Chief Editor  
Pharmaceutical  
Medicine and...**Paola  
Patrignani**University of  
Studies G....

Chieti , Italy

Specialty Chief...  
Inflammation  
Pharmacology**Alastair  
George...**The University of  
Melbourne  
Parkville ,  
AustraliaSpecialty Chief...  
Translational  
Pharmacology**Diana Conte  
Camerino**University of Bari  
Aldo Moro  
Bari , Italy**Brian Godman**Strathclyde Institute  
of Pharmacy and...  
Glasgow , United  
KingdomSpecialty Chief Editor  
Drugs Outcomes  
Research and Policies**Assistant Specialty Chief Editor****Javier  
Echeverria**University of  
Santiago  
Santiago , ChileAssistant Specialt...  
Chief Editor  
Ethnopharmacology



Pavel  
Anzenbacher

Palacký  
University,...  
  
Olomouc ,  
Czechia

Associate Editor  
Drug Metabolism  
and Transport



Ted W. Simon

TedSimon LLC  
  
Winston , United  
States

Associate Editor  
Predictive Toxicology



Howard  
Mcleod

University of  
North Carolina ...  
  
Chapel Hill ,  
United States

Associate Editor  
Pharmacogenetics  
and...



Stelios Loukides

National and Kapodistria  
University of Athens  
  
Athens , Greece

Associate Editor  
Respiratory Pharmacolog...



Neeraj Vij

Precision  
Theranostics Inc  
  
Baltimore , United  
States

Associate Editor  
Respiratory  
Pharmacology



Noel Edward  
Cranswick

The University of  
Melbourne  
  
Parkville , Australia



Yohana James  
Mashalla

University of  
Botswana  
  
Gaborone ,  
Botswana



Sitta Sittampalam

University of Kansas  
Medical Center  
  
Kansas City , United Stat...

Associate Editor  
Integrative and Regenerativ...



Vijay Suppiah

University of  
South Australia  
  
Adelaide ,  
Australia

Associate Editor  
Pharmacogenetics  
and...



Giuseppe Testa

European Institute  
of Oncology (IEO)  
  
Milan , Italy



Michael R.  
Ruggieri

Lewis Katz School  
of Medicine,...  
  
Philadelphia ,  
United States



Mahmoud Iravani

University of Hertfordshi  
Hatfield , United Kingdo...

Associate Editor  
Neuropharmacology



Modena , Italy

United States

Associate Editor  
Drugs Outcomes  
Research and...

Associate Editor  
Translational  
Pharmacology

Associate Editor  
Neuropharmacology

Associate Editor  
Neuropharmacology



**Mary Vore**  
University of Kentucky  
Lexington , United States



**Trinity Jude Bivalacqua**  
School of Medicine, Johns Hopkins...  
Baltimore , United States



**Sebastien Fuchs**  
Western University of...  
Pomona , United States



**Travis J Obrien**  
George Washington University  
Washington, D.C. , United States

Associate Editor  
Drug Metabolism and Transport

Associate Editor  
Integrative and Regenerative...

Associate Editor  
Neuropharmacology

Associate Editor  
Pharmacogenetics and Pharmacogenomics



**Minoli A Perera**  
Northwestern University  
Evanston , United States



**David S Miller**  
National Institute of Environmental...  
Durham , United States



**Muzamil Ahmad**  
Indian Institute of Integrative...  
Jammu , India



**Owen Mcmanus**  
Johns Hopkins University  
Baltimore , United States

Associate Editor  
Pharmacogenetics and...

Associate Editor  
Drug Metabolism and Transport

Associate Editor  
Neuropharmacology

Associate Editor  
Pharmacology of Ion Channels and Channelopathies



**Jeff Schmitt**  
Wake Forest School of...  
Winston-Salem , United States



**Natalia Prevarskaya**  
Lille University of Science and...  
Villeneuve-d'Ascq , France



**Gunter Peter Eckert**  
University of Giessen  
Giessen , Germany



**Luigi Milella**  
University of Basilicata  
Potenza , Italy

Associate Editor  
Integrative and Regenerative...

Associate Editor  
Pharmacology of Ion Channels and...

Associate Editor  
Ethnopharmacology

Associate Editor  
Integrative and Regenerative Pharmacology

Department of  
Earth Sciences...  
Rabat , Morocco

Umrao...  
Central Institute of  
Medicinal and...  
Lucknow , India

Dyke  
McGill University  
Montreal ,  
Canada

China Medical Universit  
(Taiwan)  
Taichung , Taiwan

Associate Editor  
Ethnopharmacology

Associate Editor  
Ethnopharmacology

Associate Editor  
ELSI in Science and  
Genetics

Associate Editor  
Pharmacology of Anti-Canc  
Drugs



**Dennis Smith**  
Pfizer (United  
States)  
New York , United  
States



**Mamdouh M. El-  
Shishtawy**  
Mansoura University  
Mansoura , Egypt



**Ryan M Pelis**  
Novartis Institutes  
for BioMedical...  
Cambridge ,  
United States



**Mara H Hutz**  
Federal University of Rio  
Grande do Sul  
Porto Alegre , Brazil

Associate Editor  
Predictive  
Toxicology

Associate Editor  
Translational  
Pharmacology

Associate Editor  
Drug Metabolism  
and Transport

Associate Editor  
Pharmacogenetics and  
Pharmacogenomics



**Antony Vinh**  
Department of  
Physiology,...  
Melbourne ,  
Australia



**Rezzan Gülgan  
(Aker)**  
School of Medicine,  
Marmara University  
Maltepe , Turkey



**Ari Haukkala**  
University of  
Helsinki  
Helsinki , Finland



**Nicholas Allen Moor**  
RenaSci (United Kingdor  
Nottingham , United  
Kingdom

Associate Editor  
Cardiovascular and  
Smooth Muscle...

Associate Editor  
Neuropharmacology

Associate Editor  
ELSI in Science and  
Genetics

Associate Editor  
Neuropharmacology



**Chi-Feng Hung**  
College of  
Medicine, Fu Je...  
New Taipei ,  
Taiwan



**Olivier Cuvillier**  
UPR8241  
Laboratoire de...  
Toulouse , France



**Alberto Bardelli**  
Institute for  
Cancer Researc...  
Candiolo , Italy



**Satoshi Murakami**  
Tokyo Institute of  
Technology  
Meguro , Japan

Associate Editor  
Ethnopharmacology

Associate Editor  
Pharmacology of Anti-  
Cancer Drugs

Associate Editor  
Pharmacology of  
Anti-Cancer Drugs

Associate Editor  
Pharmacology of Infectiou  
Diseases

Christophe Duranton  
UMR7370  
Laboratoire de...  
Nice , France

Associate Editor  
Pharmacology of  
Ion Channels an...

Yoh Takuwa  
Kanazawa  
University  
Kanazawa , Japan

Associate Editor  
Integrative and  
Regenerative...

Renato  
Bernardini  
University of  
Catania  
Catania , Italy

Associate Editor  
Experimental  
Pharmacology an...

Raghunatha Yammani  
Wake Forest School  
of Medicine  
Winston-Salem ,  
United States

Associate Editor  
Neuropharmacology

Sunita A Ramesh  
Flinders University  
Adelaide , Australia

Associate Editor  
Pharmacology of Ion  
Channels and...

Seong-Gyu Ko  
Kyung Hee  
University  
Seoul , South Korea

Associate Editor  
Pharmacology of Anti-  
Cancer Drugs

Tomohiro Nabekura  
School of  
Pharmacy, Aich...  
Nagoya , Japan

Associate Editor  
Drug Metabolism  
and Transport

Wenji Li  
Yangzhou  
University  
Yangzhou , China

Associate Editor  
Drugs Outcomes  
Research and...

José Ramón López-López  
University of  
Valladolid  
Valladolid , Spain

Associate Editor  
Cardiovascular and  
Smooth Muscle...

Chong Li  
Southwest  
University  
Chongqing ,  
China

Li Li  
The University of  
Queensland  
Brisbane , Australia

Pedro D'orléans-...  
Université de  
Sherbrooke  
Sherbrooke ,  
Canada



Benedict Green

Agricultural Research  
Service, United States.  
Washington D.C. , Unite  
States

Associate Editor  
Ethnopharmacology



Val Miranda

Oregon State University  
Corvallis , United State:

Associate Editor  
Experimental Pharmacolog  
and Drug Discovery



Sulayman D Dib-Haj  
Yale University  
New Haven , United Stat

Associate Editor  
Pharmacology of Ion Chann  
and Channelopathies



Chong Li  
Southwest  
University  
Chongqing ,  
China

Li Li  
The University of  
Queensland  
Brisbane , Australia

Pedro D'orléans-...  
Université de  
Sherbrooke  
Sherbrooke ,  
Canada

Philippe H Beaune  
Université Paris Cité  
Paris , France

**Tingjun Hou**

Zhejiang University

Hangzhou , China

**Ajay Sharma**

Chapman University  
Orange , United States

**Maohua Shi**

First People's Hospital of...  
Foshan , China

**Wolff Mayer Kirsch**

Loma Linda University Medical Center (LLUMC)  
Loma Linda , United Stat

---

Associate Editor  
Experimental Pharmacology an...

Associate Editor  
Experimental Pharmacology and...

Associate Editor  
Inflammation Pharmacology

Associate Editor  
Experimental Pharmacology and Drug Discovery

**Kiran Sharma**

Baylor College of Medicine  
Houston , United States

---

Associate Editor  
Experimental Pharmacology an...

**Alessandro Varella**

University of Salerno  
Fisciano , Italy

**Johannes Van Staden**

University of KwaZulu-Natal  
Durban , South Africa

**Valentina Vellecco**

University of Naples Federico II  
Naples , Italy

Associate Editor  
Experimental Pharmacology and Drug Discovery

---

Associate Editor  
Experimental Pharmacology and...

Associate Editor  
Ethnopharmacology

**Choel Kim**

Baylor College of Medicine  
Houston , United States

---

Associate Editor  
Experimental Pharmacology an...

**Robert Gensure**

Children's Hospital at Dartmouth...  
Lebanon , United States

**Supriya Dinkar Mahajan**

University at Buffalo  
Buffalo , United States

**Haladhar Dev Sarmah**

Bhabha Atomic Research Centre (BARC)  
Mumbai , India

Associate Editor  
Experimental Pharmacology and Drug Discovery

---

Associate Editor  
Experimental Pharmacology and...

Associate Editor  
Neuropharmacology

**Brian Sansbury**

School of Medicine,...  
Louisville , United

**Edward Jonathan Okello**

Newcastle University  
Newcastle upon

**Giovanni Zifarelli**

National Research Coun...

**Nikolaos Pitsikas**

University of Thessaly  
Volos , Greece

---

School of Medicine,...  
Louisville , United

Submit your research



**Stefano  
Giovagnoli**  
University of  
Perugia  
Perugia , Italy

Associate Editor  
Pharmaceutical  
Medicine and...



**Zhongzhen Zhao**  
Hong Kong Baptist  
University  
Kowloon , Hong  
Kong, SAR China

Associate Editor  
Ethnopharmacology



**Ricard Farre**  
Translational  
Research Cent...  
Leuven , Belgium



**Stella E. Tsirka**  
Stony Brook University  
Stony Brook , United Stat...

Associate Editor  
Neuropharmacology



**Jinyao Li**  
Xinjiang  
University  
Ürümqi , China

Associate Editor  
Drug Metabolism  
and Transport



**Filipa Alves Da  
Costa**  
Faculty of Pharmacy,  
University of Lisbon  
Lisbon , Portugal

Associate Editor  
Drugs Outcomes  
Research and Policies



**Tais Freire  
Galvao**  
State University of  
Campinas  
Campinas , Brazil



**Ricky Yuet-Kin Leun**  
University of Arkansas fo...  
Medical Sciences  
Little Rock , United State...

Associate Editor  
Pharmacology of Anti-Canc...  
Drugs



**Andrea Lapucci**  
University of  
Florence  
Florence , Italy

Associate Editor  
Pharmacology of  
Anti-Cancer Drugs



**Pinarosa Avato**  
University of Bari  
Aldo Moro  
Bari , Italy

Associate Editor  
Ethnopharmacology



**Pasquale  
Pagliano**  
University of  
Salerno  
Fisciano , Italy



**Lukasz Szczerbinski**  
Broad Institute  
Cambridge , United Stat...

Associate Editor  
Pharmacogenetics and  
Pharmacogenomics



**Ravinder  
Koundal**



**Paul Alexander  
Foster**



**Ginetta Collo**



**Pier Maria Fornasari**

Associate Editor  
Neuropharmacology

Associate Editor  
Experimental  
Pharmacology and...

Associate Editor  
Neuropharmacology

Associate Editor  
Inflammation Pharmacolo...



**Zahid Hussain**  
Faculty of  
Pharmacy,...  
Sharjah , United  
Arab Emirates



**Shih-Min Wang**  
National Cheng  
Kung University  
Tainan , Taiwan



**Elisabetta  
Vegeto**  
University of  
Milan  
Milan , Italy



**Eva Brandl**  
Charité Universitätsmedi:  
Berlin  
Berlin , Germany

Associate Editor  
Experimental  
Pharmacology an...

Associate Editor  
Inflammation  
Pharmacology

Associate Editor  
Pharmacology of  
Anti-Cancer Drugs

Associate Editor  
Pharmacogenetics and  
Pharmacogenomics



**Akshay Sood**  
University of New  
Mexico School ...  
Albuquerque ,  
United States



**Tushar Suhas  
Khare**  
Department of  
Environmental...  
Pune , India



**Narasimham L  
Parinandi**  
Division of  
Pulmonary,...  
Columbus ,  
United States



**Nazir M Khan**  
School of Medicine, Emc  
University  
Atlanta , United States

Associate Editor  
Respiratory  
Pharmacology

Associate Editor  
Experimental  
Pharmacology and...

Associate Editor  
Translational  
Pharmacology

Associate Editor  
Inflammation Pharmacolo...



**Claudia Maria  
Hattinger**  
Rizzoli  
Orthopedic...  
Bologna , Italy



**Rick Kittles**  
City of Hope  
National Medical...  
Duarte , United  
States



**Marije  
Koenders**  
Radboud  
University...  
Nijmegen ,  
Netherlands



**Sheikh Raisuddin**  
Department of Medica  
Elementology and...  
New Delhi , India

Associate Editor  
Pharmacogenetics  
and...

Associate Editor  
Pharmacogenetics and  
Pharmacogenomics

Associate Editor  
Inflammation  
Pharmacology

Associate Editor  
Gastrointestinal and Hepat  
Pharmacology

University of  
Catania  
Catania , Italy

College of Korean  
Medicine, Kyung...  
Seoul , South Korea

Prostran  
University of  
Belgrade  
Belgrade , Serbia

Sam Higginbottom  
University of Agriculture  
Allahabad , India

Associate Editor  
Experimental  
Pharmacology an...

Associate Editor  
Neuropharmacology

Associate Editor  
Neuropharmacology

Associate Editor  
Experimental Pharmacology  
and Drug Discovery



Tomas Drabek  
University of  
Pittsburgh  
Pittsburgh ,  
United States



Jacques Joubert  
University of the  
Western Cape  
Bellville , South  
Africa



Kylie Michelle  
Wagstaff  
Monash  
University  
Melbourne ,  
Australia



Damiana Leo  
University of Mons  
Mons , Belgium

Associate Editor  
Neuropharmacology

Associate Editor  
Neuropharmacology

Associate Editor  
Experimental  
Pharmacology an...

Associate Editor  
Neuropharmacology



Robyn Mary  
Brown  
Florey Institute of  
Neuroscience...  
Parkville ,  
Australia



Rita Citraro  
Department of  
Health Sciences,...  
Catanzaro , Italy



Beatriz Suarez-  
Alvarez  
Central University  
Hospital of...  
Oviedo , Spain



Bart Ellenbroek  
Victoria University of  
Wellington  
Wellington , New Zealand

Associate Editor  
Neuropharmacology

Associate Editor  
Neuropharmacology

Associate Editor  
Renal Pharmacology

Associate Editor  
Neuropharmacology



Donato  
Cappetta  
University of  
Campania Luigi...  
Caserta , Italy



Daniel Prieto-  
Alhambra  
University of Oxford  
Oxford , United  
Kingdom



Ignacio Segarra  
Catholic  
University San...  
Guadalupe , Spain



Rodrigo Pacheco  
Fundación Ciencia y  
Vida  
Santiago , Chile

Associate Editor  
Cardiovascular and  
Smooth Muscle...

Associate Editor  
Pharmacoepidemiology

Associate Editor  
ELSI in Science and  
Genetics

Associate Editor  
Neuropharmacology

Sonia Maria  
Oliani  
São Paulo State  
University  
São Paulo , Brazil

Associate Editor  
Translational  
Pharmacology



Jesse J. Swen  
Leiden University  
Medical Cente...  
Leiden ,  
Netherlands

Associate Editor  
Pharmacogenetics  
and...



Nehad M  
Ayoub  
Jordan University  
of Science and...  
Irbid , Jordan

Associate Editor  
Pharmacology of  
Anti-Cancer Drugs



Jianxiang  
Zhang  
Army Medical  
University  
Chongqing ,

Tzer-Bin Lin  
Taipei Medical  
University  
Taipei , Taiwan

Associate Editor  
Neuropharmacology



Rosa Serio  
University of  
Palermo  
Palermo , Italy

Associate Editor  
Gastrointestinal and  
Hepatic Pharmacology



Raquel  
Montenegro  
Federal University of  
Ceara  
Fortaleza , Brazil

Associate Editor  
Pharmacology of Anti-  
Cancer Drugs



Francesc J  
Moreso  
Vall d'Hebron  
University Hospital  
Barcelona , Spain

Jing Li  
Guangzhou  
University of...  
Guangzhou ,  
China

Associate Editor  
Pharmacology of  
Ion Channels an...



Junshan Liu  
Southern Medical  
University  
Guangzhou ,  
China

Associate Editor  
Inflammation  
Pharmacology



Jennifer R. Fishman  
McGill University  
Montreal , Canada

Associate Editor  
ELSI in Science and Genetic



Martin Schulz  
Freie Universität Berlin  
Berlin , Germany

Associate Editor  
Drugs Outcomes Research a  
Policies



Domenico Trombett  
Department of Chemical  
Biological, Pharmaceutical  
and Environmental Sciences  
Messina , Italy



Masakazu Ichinose  
Wakayama Medical  
University  
Wakayama , Japan



**Carolina  
Pellegrini**  
University of Pisa  
Pisa , Italy



**Maria Javier  
Ramirez**  
University of Navarra  
Pamplona , Spain



**Deborah  
Carmen Mash**  
University of  
Miami  
Coral Gables ,  
United States



**Ioanna Chinou**  
National and Kapodistrian  
University of Athens  
Athens , Greece

---

Associate Editor  
Inflammation  
Pharmacology

---

Associate Editor  
Neuropharmacology

---

Associate Editor  
Neuropharmacology

---

Associate Editor  
Ethnopharmacology



**Tong-Chuan  
He**  
University of  
Chicago...  
Chicago , United  
States



**Veljko Dubljevic**  
North Carolina State  
University  
Raleigh , United  
States



**Maurizio  
Popoli**  
University of  
Milan  
Milan , Italy



**Alex McKeown**  
University of Oxford  
Oxford , United Kingdom

---

Associate Editor  
Pharmacology of  
Anti-Cancer Drugs

---

Associate Editor  
ELSI in Science and  
Genetics

---

Associate Editor  
Neuropharmacology

---

Associate Editor  
ELSI in Science and Genetic



**Valeria Bruno**  
Department of  
Physiology an...  
Rome , Italy



**George Qian Li**  
Australian Eureka  
Bee Products  
DENISTONE ,  
Australia



**Victor Kuete**  
University of  
Dschang  
Dschang ,  
Cameroon



**Dirk Steinritz**  
Ludwig Maximilian  
University of Munich  
Munich , Germany

---

Associate Editor  
Experimental  
Pharmacology an...

---

Associate Editor  
Ethnopharmacology

---

Associate Editor  
Ethnopharmacology

---

Associate Editor  
Predictive Toxicology



**Carlos Alberto**



**Hany Omar**



**Massimo Grilli**



**Anindya Bhattacharya**

Associate Editor  
Pharmacology of  
Anti-Cancer Drugs

Associate Editor  
Pharmacology of Anti-  
Cancer Drugs

Associate Editor  
Neuropharmacology

Associate Editor  
Neuropharmacology



**Sheng-Teng  
Huang**  
China Medical  
University...  
Taichung , Taiwan



**Babu L Tekwani**  
University of  
Mississippi  
Oxford , United  
States



**George Priya  
Doss C**  
VIT University  
Vellore , India



**Federica Barbieri**  
University of Genoa  
Genoa , Italy

Associate Editor  
Ethnopharmacology

Associate Editor  
Pharmacogenetics and  
Pharmacogenomics

Associate Editor  
Experimental  
Pharmacology an...

Associate Editor  
Pharmacology of Anti-Canc  
Drugs



**Sevtap Arikan-  
Akdagli**  
Hacettepe  
University  
Ankara , Turkey



**Rong-Rong He**  
Jinan University  
Guangzhou , China



**Christian  
Waeber**  
University  
College Cork  
Cork , Ireland



**Kenneth Kc Lee**  
School of Pharmacy,  
Taylor's University  
Selangor , Malaysia

Associate Editor  
Pharmacology of  
Infectious Diseases

Associate Editor  
Ethnopharmacology

Associate Editor  
Experimental  
Pharmacology an...

Associate Editor  
Drugs Outcomes Research a  
Policies



**Ye Shen**  
University of  
Georgia  
Athens , United  
States



**Thomas  
Brzozowski**  
Jagiellonian  
University Medica...  
Kraków , Poland



**Brad Randal  
Scott...**  
Monash  
University  
Melbourne ,  
Australia

Associate Editor  
Drugs Outcomes  
Research and...

Associate Editor  
Gastrointestinal and  
Hepatic Pharmacology

Associate Editor  
Cardiovascular and  
Smooth Muscle...

- Guidelines ▾
- Explore ▾
- Outreach ▾
- Connect ▾

Follow us





# The Inaugural Monash International Health Science and Technology Conference: Pharmacology Perspectives

**967**

Total Downloads

**9,172**

Views

Overview

**Articles** 5

Authors 33

Impact

**5 Articles**

Sort by:

Views ▾

REVIEW

Published on 24 Dec 2021

## Nanoparticles use for Delivering Ursolic Acid in Cancer Therapy: A Scoping Review

Andang Miatmoko · Ester Adelia Mianing · Retno Sari · Esti Hendradi

Parameter	Lipo E	Nano A	Nano G	Poli B
Administration route	intragastric	intravenous	intravenous	intravenous
UA dose (mg/kg)	80	10	10	11
T <sub>1/2</sub> (hour)	8.6	8.3 and 10	8.7	4.9 and 5.2
AUC (µg.h/ml)	134.061	NA	NA	NA



# Nanoparticles use for Delivering Ursolic Acid in Cancer Therapy: A Scoping Review

Andang Miatmoko<sup>1,2\*</sup>, Ester Adelia Mianing<sup>3</sup>, Retno Sari<sup>1</sup> and Esti Hendradi<sup>1</sup>

<sup>1</sup>Department of Pharmaceutical Sciences, Faculty of Pharmacy, Universitas Airlangga, Surabaya, Indonesia, <sup>2</sup>Stem Cell Research and Development Center, Universitas Airlangga, Surabaya, Indonesia, <sup>3</sup>Study Program of Pharmacy, Faculty of Pharmacy, Universitas Airlangga, Surabaya, Indonesia

## OPEN ACCESS

### Edited by:

Michael Whittaker,  
Monash University, Australia

### Reviewed by:

Khalid Saad Alharbi,  
Jouf University, Saudi Arabia  
Yong Sze Ong,  
Monash University Malaysia, Malaysia

### \*Correspondence:

Andang Miatmoko  
andang-m@ff.unair.ac.id

### Specialty section:

This article was submitted to  
Pharmacology of Anti-Cancer Drugs,  
a section of the journal  
*Frontiers in Pharmacology*

**Received:** 30 September 2021

**Accepted:** 29 November 2021

**Published:** 24 December 2021

### Citation:

Miatmoko A, Mianing EA, Sari R and Hendradi E (2021) Nanoparticles use for Delivering Ursolic Acid in Cancer Therapy: A Scoping Review. *Front. Pharmacol.* 12:787226.  
doi: 10.3389/fphar.2021.787226

Ursolic acid is a natural pentacyclic triterpenoid that exerts a potent anticancer effect. Furthermore, it is classified as a BCS class IV compound possessing low permeability and water solubility, consequently demonstrating limited bioavailability in addition to low therapeutic effectiveness. Nanoparticles are developed to modify the physical characteristics of drug and can often be produced in the range of 30–200 nm, providing highly effective cancer therapy due to the Enhanced Permeation and Retention (EPR) Effect. This study aims to provide a review of the efficacy and safety of various types of Ursolic Acid-loading nanoparticles within the setting of preclinical and clinical anticancer studies. This literature study used scoping review method, where the extracted data must comply with the journal inclusion criteria of within years of 2010–2020. The identification stage produced 237 suitable articles. Duplicate screening was then conducted followed by the initial selection of 18 articles that had been reviewed and extracted for data analysis. Based on this review, the use of nanoparticles can be seen to increase the anticancer efficacy of Ursolic Acid in terms of several parameters including pharmacokinetic data, survival rates and inhibition rates, as well as the absence of serious toxicity in preclinical and clinical trials in terms of several parameters including body weight, blood clinical chemistry, and organ histopathology. Based on this review, the use of nanoparticles has been able to increase the anticancer efficacy of Ursolic Acid, as well as show the absence of serious toxicity in preclinical and clinical trials. Evenmore, the liposome carrier provides development data that has reached the clinical trial phase I. The use of nanoparticle provides high potential for Ursolic Acid delivery in cancer therapy.

**Keywords:** ursolic acid, cancer, nanoparticle, liposome, nanosphere, polymeric micelle, efficacy, toxicity

## INTRODUCTION

Cancer is a disease that can occur in almost any organ or tissue of the body when abnormal cells grow uncontrollably beyond their usual limits to invade adjacent parts of the body and/or spread to other organs (Kamińska et al., 2015). Data from the Global Burden of Cancer released by the World Health Organization (WHO) reported that the number of cases and deaths from cancer in 2018 totaled 18.1 million and 9.6 million respectively. Cancer-related deaths are predicted to increase to more than 13.1 million by 2030 (Internatioanl Agency for Research on Cancer, 2018). The most common types affecting men include lung, prostate, colorectal, and liver cancer, while in women, they comprise breast, colorectal, lung, cervical and thyroid cancers. In 2018, the incidence rate of liver cancer in

Indonesia ranked eighth highest in Southeast Asia, while in Asia as a whole it occupied 23rd position with 348,809 cases (Globocan. Indonesia - Global Cancer Observatory WHO, 2018).

The first-line options for cancer treatment include surgery, radiotherapy and the administering of biological and chemical drugs (chemotherapy, hormones, and biological therapy). However, such forms of treatment fail to control metastatic tumors that have spread to other organs (Kumari et al., 2016). Chemotherapeutic agents are predominantly toxic compounds that primarily inhibit the rapid proliferation of cancer cells, while also potentially restricting the growth of hair follicle, bone marrow and gastrointestinal cells culminating in severe undesired side effects (Pérez-Herrero and Fernández-Medarde, 2015). Consequently, the effect of certain natural compounds have been widely explored and it has been scientifically proven that Ursolic Acid (UA), an active agent, inhibits the proliferation of cancer cells (Kashyap et al., 2016).

UA is a natural pentacyclic triterpene of the cyclosqualenoid family present in many plants which can modulate cellular transcription factors, growth factor receptors, inflammatory cytokines, and numerous other molecular targets that regulate cell proliferation, metastasis, apoptosis, angiogenesis, and autophagy. The anticancer effects of UA have been reported for many types of cancer, one of which is liver cancer (Khwaza et al., 2020). The mechanisms of UA which produce such effects include nuclear-kappa B (NF- $\kappa$ B) factors and apoptosis signaling in cancer cells (Seo et al., 2018). The protease activity involving urokinase and cathepsin B, both of which are known to be associated with tumor invasion and metastasis, is also significantly inhibited by UA, interleukin-1  $\beta$  (IL-1 $\beta$ ), Tumor necrotic factor- $\alpha$  (TNF- $\alpha$ ), and the expression of MMP-2 and MMP-9 (Mitochondria-Mediated Pathway) (Kashyap et al., 2016; Ali et al., 2019). Prolonged administration of excessive doses of UA, with an LD50 value of 9.26 g/kg in acute toxicity tests on mice (Sun et al., 2020), has the potential to cause liver cytotoxicity which is not classified as genetic toxicity. Within the Biopharmaceutical Classification System (BCS), UA is categorized as a BCS class IV compound demonstrating low permeability and solubility (Sun et al., 2020) which, consequently, requires a nanotechnology-based drug delivery system to reach the desired target (Biopharmaceutical, 2011). In particular, the development of drug delivery systems entails the use of nanoparticles targeted at cancer cells which significantly improve therapeutic and diagnostic efficacy as well as reducing unwanted side effects.

Nanotechnology represents a new therapeutic platform that employs nanoparticles (NPs) in the diagnosis and treatment of cancer. NPs are used in cancer therapy due to their relatively small size, 1–1,000 nm, the fact that they frequently fall within the range of 10–200 nm, and the presence of the EPR effect (Rajesh, 2000; Reddy, 2005; Fang et al., 2011). Nano-drug delivery systems have been lauded for their excellent biocompatibility properties, low toxicity, increased solubility in water, in addition to their ability to deliver targeted drugs to tissues which limits their accumulation in the kidneys, liver, spleen, and other non-targeted organs, while improving therapeutic efficacy (Kumari et al., 2016; Li et al., 2017). The delivery of nanoparticles to tumor

tissues through systemic circulation can be executed through two targeting strategies, including passive targeting when nanoparticles entering circulation will accumulate at the tumor site due to enhanced permeation and retention (EPR). In contrast, active targeting generally employs ligand molecules such as antibodies and peptides to recognize specific antigens expressed in tumor cells or the microenvironment (Kumari et al., 2016).

Many types of nanoparticles exist, including polymeric therapy (polymer-protein and polymer-drug conjugates) in which drugs are covalently bound or conjugated to polymer structures and nanoparticulate drugs. The drugs are physically trapped in assembled particles composed of different materials such as polymers (polymer micelles, dendrimers and polymer nanoparticles), lipids (liposomes), or organometallic compounds (carbon nanotubes). The first generation of anticancer nanoparticles approved by the FDA include liposomal drugs and polymer conjugates (Barenholz, 2012; Pérez-Herrero and Fernández-Medarde, 2015). However, certain products can be subjected to *in vivo* and clinical trials, while others remain limited to *in vitro* studies. Therefore, the effectiveness and safety of the nanoparticle drug constitute important assessed parameters.

As for the development of drugs with nanoparticle carriers, one example is Doxil<sup>®</sup>, the first nano-drug using liposomes approved by the FDA, which demonstrates the clinical performance advantages of doxorubicin in a variety of neoplastic conditions due to pharmacokinetics and the unique EPR-related biodistribution of liposomal doxorubicin (Barenholz, 2012; Wei et al., 2016). Doxil<sup>®</sup> can reduce side effects, especially that of cardiac toxicity, and improve patients' adherence and quality of life (Barenholz, 2012). Cisplatin is an anticancer drug prepared with a polymeric micelle through the formation of a metal-polymer complex between cisplatin and poly-(ethylene glycol)-poly(glutamic acid) block copolymers (Nishiyama et al., 2003; Uchino et al., 2005). These micelles are 28 nm in size with a very narrow distribution, demonstrate extremely extended blood circulation, and accumulate effectively in solid tumor models of Lewis lung carcinoma cells. However, because they undergo chemical synthesis, toxicity and scale-up production become major issues (Mizumura et al., 2001; Plummer et al., 2011; Mitchell et al., 2021). In addition, the development of Abraxane<sup>®</sup>, a paclitaxel-albumin-bound nanoparticle ~130 nm in size, was approved by the FDA in 2005 for the treatment of metastatic breast cancer (Miele et al., 2009; Ma and Mumper, 2013). This formulation has been shown to have several advantages in terms of toxicity reduction compared to Taxol<sup>®</sup>. Moreover, the total dose can be given within 30 min without pre-treatment. However, the manner in which Abraxane<sup>®</sup> can improve survival rates and overcome P-GP-mediated drug resistance remains unclear (Ma and Mumper, 2013).

Certain nanoparticles have been used in the delivery of UA as a cancer therapy including liposomes, polymeric nanoparticles, and polymeric micelles (Zaimy et al., 2017). However, at the time of writing, in contrast to other chemotherapy drugs such as Doxorubicin (Doxil<sup>®</sup>), Cisplatin, Paclitaxel (Taxol<sup>®</sup>), or Amphotericin B (Ambisome<sup>®</sup>), no review of the effectiveness

**TABLE 1 |** The inclusion and exclusion criteria for article screening and selection.

Test parameters	Inclusion criteria	Exclusion criteria
Type of research	a) Randomized or non-randomized phases 1, 2, or 3 clinical trials  b) <i>In vivo</i> studies in animals	a) <i>In vitro</i> study b) <i>Ex vivo</i> study c) Review article
Intervention	a) Native UA as an active ingredient b) Nanoparticles (lipids, polymers, hybrid nanoparticles as carriers)	a) Extracts containing UA and UA derivates b) Microparticles or other carrier systems more than 1,000 nm in size c) Administration routes other than those meeting the inclusion criteria (topical, transdermal)
Comparison	c) Administration routes comprise oral route in addition to intravenous, intraperitoneal, and intratumoral injection d) Healthy patients and those suffering from all types of cancer (both individuals who have undergone surgery and those who have not)	-
Outcome	a) No comparison with other drugs, only negative controls b) Comparison with other drugs	-
Types of Publications	a) Primary efficacy outcomes (improved lifespan, enhanced survival rate, tumor growth inhibition) b) Secondary efficacy outcomes (e.g., blood parameters, no complaints); improvement in physical condition (body weight, tissue histopathology); clinical and non-clinical improvements c) Toxicity (body weight, blood parameters, clinical parameters, non-clinical parameters, adverse events, organ histopathology)	The article is not written in English
	a) Articles are written in English b) Not included as predatory journals	

and safety of several types of nanoparticles for the delivery of UA for cancer therapy has been conducted (Zaimy et al., 2017). This study aims to provide a literature review related to the anticancer effectiveness and safety of UA delivered with various types of nanoparticles based on pre-clinical and clinical trial results from existing research published between 2011 and 2021.

## METHODS

### Article Selection Criteria

This study uses the scoping review method involving literature accessible through the PubMed, Sciencedirect, Scopus, and Google Scholar databases consisting of online research publications dating from 2011 to 2021. Clinical trial articles were sourced using the keywords "Clinical trial", "Ursolic Acid", "Cancer", and "OR Nanoparticle Liposome". As for the search for articles relating to *in vivo* studies, this employed the keywords "Pre-Clinical OR *in vivo* OR Animal", Ursolic Acid", "Cancer", "Nanoparticle". Within this study, several inclusion and exclusion criteria were applied to select and screen articles for review as shown in Table 1.

### Evaluation of Physical Characteristics of UA Nanoparticles

Data analysis involved comparing the physical characteristics of different types of nanoparticles identified in the selected articles.

### Analysis of Particle Size

Particle size and particle size distribution produce significant impacts on drug loading variation, drug release, bioavailability, and efficacy (Kharia et al., 2012). In addition, particle sizes of 150 nm to 4.5  $\mu$ m will be easily recognized by macrophages and

dendritic cells during phagocytosis (Koppolu and Zaharoff, 2013). Instruments used in nanoparticle size and morphology determination include Dynamic Light Scattering (DLS), Scanning Electron Microscopy (SEM), Transmission Electron Microscopy (TEM), and Atomic Force Microscopy (AFM).

### Analysis of $\zeta$ -Potential

Zeta potential is used to predict the stability of colloidal dispersion systems during storage. Generally,  $\zeta$ -potential values above  $\pm 30$  mV resulted in more stable particles because the repulsive force between particles can prevent aggregation. The  $\zeta$ -potential is affected by surfactants, polymers, the surface active agent component of nanoparticles, the presence of absorbing compounds, dispersed phase media, ionic strength, and pH (Kharia et al., 2012). The  $\zeta$ -potential can be analyzed using the Electrophoretic Light Scattering (ELS) method (Miatmoko et al., 2017; Wang et al., 2017).

### Analysis of Encapsulation Efficiency

Entrapment Efficiency (EE%), or encapsulation efficiency, is defined as the portion of drugs encapsulated in nanoparticles. Free drugs that are not encapsulated in the drug delivery carriers or nanoparticles can be separated by means of centrifugation, dialysis, or gel chromatography. The concentration of entrapped and un-entrapped drugs can be determined through the use of instruments such as spectrophotometers or high-performance liquid chromatography (HPLC). The encapsulation efficiency percentage is calculated using the following equation:

$$EE(\%) = \frac{W_T - W_U}{W_T} \times 100\%$$

where,  $W_T$  is the total weight of AU and  $W_U$  is an un-entrapped UA weight (Poudel et al., 2020).

## Pharmacokinetic Evaluation of UA Nanoparticles

Plasma concentration versus time data was analyzed using non-compartmental methods. Peak plasma concentration ( $C_{max}$ ) and time-to-peak plasma concentrations ( $T_{max}$ ) were obtained through experimental observation. In elimination half-life ( $t_{1/2}$ ) calculated as  $0.693/\lambda z$ ,  $\lambda z$  is the slope of the terminal phase. In areas under the curve (AUC<sub>0-t</sub>) of plasma concentration versus time from zero to infinity, AUC<sub>0-∞</sub> is equivalent to the total area from time = 0 to the last measurable concentration time. The value is calculated using the linear trapezoidal method up to  $C_{max}$ , log-trapezoidal method (until the last measured concentration), and extrapolated areas (Zhu et al., 2013). In this study, the analysis was conducted by comparing pharmacokinetic profiles from various studies contained in the articles reviewed.

## Evaluation of the Effectiveness of Ursolic Acid Nanoparticles

The analysis was conducted by comparing the results of efficacy studies including survival rate, tumor growth inhibition, tumor weight, and tumor volume, as well as tumor tissue histopathology extracted from reviewed articles.

### Cancerous Tissue Histopathology

The histological section of the liver was stained with haematoxylin and eosin (H&E) staining and subsequently compared to the histopathological appearance of each organ in the control and treatment groups (Melo et al., 2020). The microstructure and morphology of tissues were observed using a light microscope (Wang et al., 2017). Hematoxylin is a base dye that has an affinity for the acidic components of cells, primarily the nucleic acids contained in the nucleus, while eosin is an acidic dye that binds to the cell cytoplasm. As a result, H&E stained the core blue and cytoplasm orange-red (Slaoui and Fiette, 2011).

### Analysis of Relative Tumor Volume

In the articles, the size of the tumor is determined by means of a calliper, while its volume is calculated using the following equation:

$$V = 0.5xy^2$$

where x is the longest and y the shortest diameter (Hattori et al., 2014; Yang et al., 2014; Wang et al., 2016; Miatmoko et al., 2017; Liu et al., 2021). In this study, the relative tumor volume is calculated using the following formulas:

$$\text{relative tumor volume} = \frac{V_T \text{ NC}}{V_T \text{ AUNP}}$$

or

$$\text{relative tumor volume} = \frac{V_T \text{ AU}}{V_T \text{ AUNP}}$$

where  $V_T \text{ NC}$  is the volume of the negative control group tumor,  $V_T \text{ AU}$  is the volume of the native UA treatment group tumor,

and  $V_T \text{ AUNP}$  is the volume of the UA nanoparticle treatment group tumor.

## Analysis of Relative Tumor Weight and Growth Inhibition Rate

The antitumor activity of nanoparticles is assessed through the tumor growth inhibition rate (IR) or tumor growth inhibition (TGI) at the experimental endpoint, calculated using the following equations:

$$\text{IR (\%)} \text{ or } \text{TGI (\%)} = \frac{W_T \text{ of negative control group} - W_T \text{ of treatment group}}{W_T \text{ of negative control group}}$$

where  $W_T$  is the weight of the tumor (Wang et al., 2017; Shen et al., 2018; Fu et al., 2021). In this study, the relative tumor weight and relative inhibition rate were each calculated using the following formulas:

$$\text{Relative tumor weight} = \frac{W_T \text{ NC}}{W_T \text{ AUNP}}$$

or

$$\text{relative tumor weight} = \frac{W_T \text{ AU}}{W_T \text{ AUNP}}$$

where  $W_T \text{ NC}$  is the weight of the negative control group tumor,  $W_T \text{ AU}$  is the tumor weight of the native UA treatment group, and  $W_T \text{ AUNP}$  is the volume of the UA nanoparticle treatment group tumor.

$$\text{Relative inhibition rate} = \frac{I_R \text{ AUNP}}{I_R \text{ AU}}$$

where  $I_R \text{ AUNP}$  is the tumor inhibition rate of UA nanoparticle treatment group, and  $I_R \text{ AU}$  represents the tumor inhibition rate of the native UA treatment group.

## Analysis of Relative Survival Rate

Survival rates can be used as a standard assessment of effective therapy. The survival period is usually calculated from the date of diagnosis or commencement of the treatment period. The survival curve of each group was estimated using the Kaplan-Meier method with the average survival time difference being assessed by means of a log-rank test (Yang et al., 2014). This method involves non-parametric estimation of the survival function commonly used to describe the survival of a single population or compare the survival of two populations. The Kaplan-Meier estimate is one of the most effective statistical methods of measuring the probability of a patient's survival observed during a post-treatment period (Etikan, 2017). In this study, the relative survival rate was calculated using the following formula:

$$\text{Relative survival rate} = \frac{S_R \text{ AUNP}}{S_R \text{ NC}}$$

where  $S_R \text{ AUNP}$  is the survival rate following the administration of UA nanoparticles and  $S_R \text{ NC}$  is the survival rate of the negative controls.

## Evaluation of Toxicity of UA Nanoparticles in Pre-Clinical Trials

### Weight Analysis

Weight loss represents a significant parameter of biological safety analysis or drug safety. The weight of the mice subjects is measured on the day of tumor inoculation and continues daily, or at least several times per week, during treatment. Each treatment group mouse is quantitatively weighed with the result being compared to that of a normal mouse in order to identify any significant difference between the two groups (Valcourt et al., 2020). In this study, the weight of the mice was calculated using the following formulas:

$$\text{Relative body weights} = \frac{W_B \text{ NC}}{W_B \text{ AUNP}}$$

or

$$\text{Relative body weights} = \frac{W_B \text{ AU}}{W_B \text{ AUNP}}$$

Where  $W_B \text{ NC}$  is the mice body weight in the administration of negative control,  $W_B \text{ AU}$  is the mice body weight in the administration of native UA, and  $W_B \text{ AUNP}$  represents the mice body weight in the administration of UA nanoparticles.

### Other Toxicity Analysis

Other toxicity data on the *in vivo* studies was derived by data recapitulation that included: tissue histopathology, increased levels of ALT and AST, and the amount of WBC as an indicator of hematological toxicity (Zhao et al., 2018; Li et al., 2019; Liu et al., 2021).

## Evaluation of AU-NP Toxicity in Clinical Trials

### Analysis of Clinical Chemistry Data

Toxicity was evaluated in all subjects treated with at least one cycle of UA Liposome therapy. Hematological parameters (red blood cells, WBC, hemoglobin, ANC, and platelets), urinary routines (urine protein, glucose, erythrocytes, leukocytes, and urine bilirubin), and stool routines (stool erythrocytes and stool leukocytes) were evaluated. Blood biochemistry including ALT, AST, ALP, gamma-glutamyl transpeptidase (GGT) were further analyzed (Qian et al., 2015). In this study, the analysis was conducted by comparing clinical laboratory data (ALT, AST, GGT, DBIL, and TBIL) extracted from reviewed articles.

### Analysis of Adverse Events

Adverse events are used to assess unintended events (AE) in healthy adult volunteers and patients with advanced solid tumor disease. In addition, the toxicity can be seen from the value of the maximum tolerated dose (MTD) used to determine the highest dose of the drug that can be administered without adverse events. The adverse events documented during the study were classified as mild, moderate, or severe based on the dosage (Valcourt et al., 2020). In this study, the analysis was conducted by comparing adverse events or side effects occurring in subjects who had received the treatment mentioned in reviewed articles.

## RESULTS AND DISCUSSION

This study provides a literature review focusing on the anticancer effectiveness and safety of UA delivered with various types of nanoparticles to increase its anticancer effects as confirmed by both pre-clinical and clinical trials. Literature searches of all four databases using pre-determined keywords identified 237 articles in the prescreening stage as can be seen in **Figure 1**. Of the total literature reviewed, duplication screening was conducted using the Mendeley application to produce a final body of 226 articles. Application of exclusion criteria resulting in a body of 196 selected articles which were subsequently subjected to inclusion criteria to produce a final total of 30. The initial selection process identified 24 articles which were subsequently reviewed, culminating in 18 which satisfied the inclusion criteria. The summary of reviewed articles is presented in **Table 2**.


The data extraction of the literature used can be seen in **Table 2**.

### Nanoparticle Characterization Results

From the review of the 18 articles, it was clear that three types of drug represent the most frequent delivery carriers of UA as an anticancer agent, namely; Liposome (50%), Nanosphere (39%) and Polymeric Misel (11%), (see **Figure 2A**).

According to the review results, several characterization parameters of liposomes, nanospheres, and polymeric micelles exist, including particle size,  $\zeta$ -potential, and encapsulation efficiency (EE). From the data analysis of the 18 articles, the size of liposome particles was found to range from 70 to 200 nm (67%); nanosphere particle size to be between 70 and 200 nm (100%); and micelle polymeric particle size to be between 30 and 70 nm (50%). The  $\zeta$ -potential of liposomes ranged from (-)30 to 0 mV (11%) and 0 to (+)30 mV (33%), while the nanosphere  $\zeta$ -potentials were between (-)30 to 0 mV (57%), 0 to (+)30 mV (43%); and  $\zeta$  liposomes of (-)30 to 0 mV (100%). For the EE of liposomes  $\geq 90\%$  (11%) and 30–90% (22%); EE nanospheres  $\geq 90\%$  (14%) and 30–90% (14%); as for EE polymeric micelles, these are not mentioned in the article, as presented in **Figure 2B**.

Characterization of liposome particle size is important because it affects the interaction of liposomes with target cells as well as the elimination, penetration and retention of drugs in the target sites (Monteiro et al., 2014). Phospholipids represent the main constituents of liposomal membranes and the use of lipid types and ratios within different preparation methods can affect the size of liposomes (Monteiro et al., 2014; Shaker et al., 2017). From **Figure 2A** it can be seen that liposomes prepared with ethanol injection and thin-film hydration methods have particle sizes ranging from 70 to 200 nm. This finding is in accordance with that of previous research arguing that, with the ethanol injection method, liposome could be generated as SUVs with diameters of 30–110 nm (Akbarzadeh et al., 2013; Monteiro et al., 2014), while with the adoption of thin film hydration methods, continued use of sonication or extrusion processes can produce liposomes as 25 nm to 1  $\mu\text{m}$ -sized ULVs (Monteiro et al., 2014). Liposome size depends on that of the phospholipid molecule assembly whose average dimensions depend on their lipid composition, while it is supposed that the size of liposome particles increases slightly with



a reduction in the molar ratio of HSPC/SPC in the range of 119–143 nm (Chen et al., 2006). On the other hand, liposomes made from DMPC, DSPC, and HSPC (at a weight ratio of 2:1 to cholesterol) experienced different increases in particle size, e.g., DMPC:Chol liposomes increased in size from 149 to 190 nm, DSPC:Chol expanded from 83 to 104 nm, and HSPC:Chol liposomes from 88 to 122 nm (Webb et al., 2019).

For nanospheres, particle size, which is greatly affected by lipid type, ranges from 70 to 200 nm. This is in accordance with a previous report stating that nanospheres have a diameter of 10–200 nm (Baldim et al., 2020). With regard to polymeric micelles, studies show that particle sizes ranging from 30–70 nm are affected by polymer types based on the characteristics of hydrophilic and hydrophobic block copolymers. This finding is in keeping with that of earlier research which reported that the size of polymeric micelle particles is determined by the ratio of hydrophobic and hydrophilic block chains and can produce particle sizes of  $\leq 50$  nm (Cabral et al., 2011). Increased targeting of drugs to cancer cells within the tumor tissues with the use of long-circulating polymeric micelles depends on the size of the micelle and the vascular permeability of the tumor tissues. In hypervasculature tumors with highly permeable vascular structures, sub-100 nm polymeric micelles show no limits for drug extravasation and tumor penetration. In contrast, only micelles smaller than 50 nm can penetrate hypovascular tumors whose vascular permeability is poor (Cabral et al., 2011).

The zeta potentials which reflect the liposome surface charges were evaluated (Alves et al., 2017). **Figure 2B** shows that

liposomes and nanospheres had zeta potentials ranging from (-30) to 0 mV and 0 to (+)30 mV, while those of polymeric micelles varied from (-30) to 0 mV. If the system has a strong negative or positive zeta potential the particles will tend to repulse each other and no aggregation occurs. Therefore, if the system has zeta potential  $>+30$  mV or  $<-30$  mV, then it can be considered stable (Laouini et al., 2012; Lunardi et al., 2021). The positive or negative charges measured in nanoparticles are highly dependent on lipid components. Analysis of the composition and intracellular delivery mechanisms confirmed that conventional liposomes had a relative neutral charge due to their neutral phospholipid composition such as HSPC and became negatively charged when added to cholesterol. pH sensitive liposomes contained a DOPE-like phospholipid component with CHEMS causing their negative charges; cationic liposomes had a cationic lipid composition such as DDAB, DOGS, DOTAP, DOTMA, DMRIE, DORIE with DOPE; Long-circulating liposomes (LCL) had a high  $T_c$  neutral lipid composition, cholesterol, added to approximately 5–10 mol % of PEG-DSPE rendering these liposomes stable when under protein opsonization (Sharma and Sharma, 1997; Gabizon et al., 2012).

The tendency of a drug to interact with polar or non-polar bonds and/or electrostatic interactions with lipid bilayer will determine whether it will be encapsulated into inner aqueous compartments or the lipid bilayer membrane, or whether it will be closely related to the polar head group of the bilayer membrane through electrostatic interactions. It will correlate to encapsulation efficiency (EE) or loading capacity, which is usually defined as a fraction of the percentage of the total

**TABLE 2 |** The summary of literature reviews for UA-loaded nanoparticles.

No	Code	Carrier type	Formula-tion	Type of research	Information research	Administra-tion route	Reference
1	Lipo A	Liposomes	Not Available	Phase I Clinical Trials	Safety Evaluation of Double Dose and Antitumor Activity of Ursolic Acid (UAL) Liposomes in Subjects with Advanced Solid Tumors including: Non-Hodgkin Lymphoma (24%), Hodgkin Lymphoma (24%), Renal Carcinoma (5%), Hepatocellular Carcinoma (5%), Breast Cancer (9%), Lung Cancer (9%), Other Cancers (19%)	Intravenous 4 h infusion at doses equivalent to 54, 74, and 96 mg UA/m <sup>2</sup> for 14 consecutive days	Qian et al. (2015)
2	Lipo B	Liposomes	Not Available	Phase I Clinical Trials	Toxicity evaluation of a single dose of intravenous ursolic acid liposomes (UAL) in healthy adult volunteers and patients with advanced solid tumors including Non-Hodgkin Lymphoma, Hodgkin Lymphoma, Renal Carcinoma, and Hepatocellular Carcinoma	Intravenous (IV) route at doses equivalent to 11, 22, 37, 56, 74, and 96, and 130 mg UA/m <sup>2</sup> administered as a 4 h infusion	Wang et al. (2013)
3	Lipo C	Liposomes	Not Available	Phase I Clinical Trials	Toxicity evaluation of Ursolic Acid Nanoliposome (UANL) in healthy volunteers and patients with advanced solid tumors including: Non-Hodgkin Lymphoma (50%), Hodgkin Lymphoma (12.5%), Gut Cancer (12.5%), Hepatocellular Carcinoma (25%)	Intravenous (IV) route at doses equivalent to 74 mg/m <sup>2</sup> as a single dose, 98 mg/m <sup>2</sup> , and 74 mg/m <sup>2</sup> as double doses daily for 14 days via a 4 h infusion	Zhu et al. (2013)
4	Lipo D	Liposomes	Hydrophobic components (PC, Chl, and UA) at a weight ratio of 2:1:0.5; ethanol injection method	Preclinical or <i>in vivo</i> study	Tumor inhibition activity and toxicity studies of UA-PLL-HA in SCC-7 tumor-induced mice	Intravenous (IV) at a dose of equivalent to 20 mg UA/Kg mouse for 5 times every 4 days	Poudel et al. (2020)
5	Lipo E	Liposomes	PEGylated UA Liposomes composed of SPC, CHOL, and UA at a weight ratio of 50:8:5, respectively; ethanol injection method	Preclinical or <i>in vivo</i> study	Tumor growth inhibition study and cytotoxicity of UA PEGylated liposomes in mice with U14 cervical carcinoma cells	Intragastric route at a dose of equivalent to 80 mg UA/kg mouse twice a day for a total of 14 days	Wang et al. (2016)
6	Lipo F	Liposomes	Liposomes composed of hydrophobic components (SPC, CHOL and UA) at a weight ratio of 0:6:5; ethanol injection method	Preclinical or <i>in vivo</i> study	Tumor growth inhibition and toxicity studies of CS-UA-L in mice with U14 cervical carcinoma cells	Intragastric route at a dose of equivalent to 80 mg UA/Kg mouse once a day for 14 days	Wang et al. (2017)
7	Lipo G	Liposomes	Lipids-UA (HSPC/Kolesterol/DSPE-PEG2000/UA = 90/0/5/5 and 90/0/5/10, (molar ratio); thin film hydration method	Preclinical or <i>in vivo</i> study	Tumor and growth inhibition study of UA-L in mice with 4T1 tumors (breast cancer)	Intravenous (IV) route at a dose of equivalent to 10 mg UA/kg mouse for 5 times every other day	Zhang et al. (2020)
8	Lipo H	Liposomes	Lipid components of FA-UA-L: DOTAP/CHOL/MPEG-DSPE2000/FA-PEG-CHEMS at a molar ratio of 40:55:4, 5:0, 5 (equal to weight ratio of 28; 21,3; 12,6, dan 2, 1 mg). The ratio of UA to lipid is 1:20 (w/w); thin film hydration method	Preclinical or <i>in vivo</i> study	Tumor growth inhibition and toxicity studies of FA-UA/siRNA-L in mice with human KB cells tumor	Intravenous (IV) injection with the dose of 4.5 mg/kg for UA and 170 µg/kg for siRNA for 5 times every other day	Li et al. (2019)
9	Lipo I	Liposomes	Lipid composition: HSPC/CHOL/mPEG-DSPE2000/FA-PEG-CHEMS at molar ratio 63:32:4.5:0.5 (equal to weights amount of 48, 12, 13.4, and 5 mg), respectively. The ratio of UA to lipids is 1:20 (w/w); thin film hydration method	Preclinical or <i>in vivo</i> study	Efficacy study of FTL-UA for tumor inhibition in mice with human KB tumor cells	Intravenous (IV) at a dose of equivalent to 4.5 mg UA/kg mouse for 5 times every other day, which is similar to 23 mg/kg or 98 mg/m <sup>2</sup> drug administration	Yang et al. (2014)
10	Nano A	Nanospheres	Not available	Preclinical or <i>in vivo</i> study	Tumor growth inhibition and toxicity studies of HCPT @F-Pt-PU NPs in mice with H22 subcutaneous tumors (liver cancer)	Intravenous (IV) injection at a dose of equivalent to 10 mg UA/kg mouse for 5 times every 2 days	Liu et al. (2021)

(Continued on following page)

**TABLE 2 | (Continued)** The summary of literature reviews for UA-loaded nanoparticles.

No	Code	Carrier type	Formula-tion	Type of research	Information research	Administra-tion route	Reference
11	Nano B	Nanospheres	NP composed of 32 mg chitosan, 10 mg UA, 30 mg EDC, and 8 mg NHS. The ratio of UA to lipids is 1:10 (w/w); overnight magnetic stirring method	Preclinical or <i>in vivo</i> study	Tumor inhibition study of CH-UA-NPs in mice with H22 subcutaneous tumors (liver cancer)	Oral administration at a dose of equivalent to 11 mg UA/Kg mouse once every 2 days for a total of 8 times	Jin et al. (2016a)
12	Nano C	Nanospheres	NP composed of 32 mg chitosan, 10 mg UA, 30 mg EDC, and 8 mg NHS. The ratio of UA to lipids is 1:10 (w/w); overnight magnetic stirring method	Preclinical or <i>in vivo</i> study	FA-CS-UA-NPs tumor inhibiting activity study in MCF-7 xenograft bearing models (breast cancer)	Intraperitoneal (IP) injection at a dose of equivalent to 12.5 mg UA/kg mouse once a day for 9 times	Jin et al. (2016b)
13	Nano D	Nanospheres	Not available	Preclinical or <i>in vivo</i> study	Tumor growth inhibition efficacy and toxicity studies of UA-LA-ICG NPs in tumor bearing mice by murine H22-hepatocarcinoma cells induced tumor xenograft models	Intravenous (IV) injection at a dose of 10 mg/kg of UA and 2.5 mg/kg of ICG with 5 min irradiation at 24 h post injection	Zhao et al. (2018)
14	Nano E	Nanospheres	Prepared by making 3 mg UA solution in ethanol (1 ml, 6,569 mM) in 10 ml of water. The ratio of UA and NPs was 1:10, respectively; solvent exchange preparation method	Preclinical or <i>in vivo</i> study	Tumor inhibition efficacy and toxicity studies of UA NPs in mice bearing A549 xenograft models (lung cancer)	Intravenous (IV) injection at a dose of 8 mg/kg of UA for 21 days	Fan et al. (2018)
15	Nano F	Nanospheres	Not available	Preclinical or <i>in vivo</i> study	Tumor inhibiting activity and toxicity studies of UA NPs in H22-induced mice (Hepatocellular carcinoma)	Intraperitoneal (IP) injection at a daily dose of 50 mg/kg of UA for 10 days	Zhang et al. (2015)
16	Nano G	Nanospheres	Self-assembly method of polymer deposition	Preclinical or <i>in vivo</i> study	Antitumor activity and toxicity studies of Pec-8PUH NPs in mice with 4T1 tumors (breast cancer)	Intravenous (IV) injection at a dose of 10 mg/kg of UA once every 2 days for 5 times	Liu et al. (2018)
17	Poli A	Polymeric Micelles	PM composed of UA (4 mg) and mPEG2000-PLA2000 (40 mg) at a weight ratio of 1:10; thin film dispersion method	Preclinical or <i>in vivo</i> study	Antitumor activity and toxicity studies of UA-PMs in H22-induced mice (Hepatocellular carcinoma)	Intraperitoneal (IP) injection at a dose of 50 mg/kg of UA every 2 days for 6 times	Zhou et al. (2019)
18	Poli B	Polymeric Micelles	Solvent evaporation method	Preclinical or <i>in vivo</i> study	Antitumor activity and toxicity studies of U-SS-M in tumor bearing MG-63/Osteosarcoma (OS)	Intravenous (IV) injection at a dose of 11 mg/kg of UA every 3 days for 5 times	Fu et al. (2021)


Notes: UAL, Ursolic Acid Liposome; UANL, Ursolic Acid Nanoliposome; UA-PLL-HA, Ursolic Acid-Poly-L-Lysine-Hyaluronic Acid; UA-PEGylated, Ursolic Acid-Polietilenglikolisis; CH-UA-NPs, Chitosan-Ursolic Acid-Nanoparticles; CS-UA-L, Chitosan-Ursolic Acid-Liposome; CHOL/Chl, Cholesterol; DSPE-PEG2000, 1,2-distearoyl-sn-glycero-3-phosphoethanolamine-N [methoxy (poly-ethylene glycol) -2000]; DOTAP, 1,2-dioleoyl-3-trimethylammonium-propane; EDC, Ethyl-(3-dimethylaminopropyl) carbodimide hydrochloride; FA-CS-UA-NPs, Folate-Chitosan-Ursolic Acid-Nanoparticles; FA-PEG-CHEMS, Folate Polyethylene Glycol Cholestryl hemisuccinate; FA-UA/siRNA-L, Folate- Ursolic Acid/Small Interfering RNA-Liposome; F-Pt-PU, Folia Acid-Pectin-Eight-Arm PEG-UA conjugate; FTL-UA, Folate Receptor Targeted Liposome-Ursolic Acid; HCPT @F-Pt-PU NPs, Hydroxycamptothecin @folic acid-pectin-eight-arm PEG-UA nanoparticle; HSPC, Hydrogenated Soybean Phosphatidyl Choline; mPEG2000-PLA2000, Monomethoxy Polyethylene Glycol 2000 Poly Lactic Acid 2000; MPEG-DSPE2000, Monomethoxy Polyethylene Glycol 2000-Distearoyl Phosphatidylethanolamine; NHS, N-Hydroxy-Succinimide; PC, Phosphatidylcholine; Pec-8PUH NPs, pectin-eight-arm polyethylene glycol-ursolic acid/hydroxycamptothecin nanoparticle; SPC, Soybean Phosphatidyl Choline; UA-NPs, Ursolic Acid-Nanoparticles; UA-LA-ICG NPs, Ursolic Acid- Lactobionic Acid -Indocyanine Green; UA-PMs, Ursolic Acid-Polymer Micelles; U-SS-M, Micelles assembled by PEG-SS-UA (polyethylene glycol using a disulfide bond)

encapsulated drug, in the bilayer membrane or aqueous intravesicular compartments or the matrix of nanoparticles (Kulkarni et al., 1995). UA has poor permeability and low water (Sun et al., 2020), thus causing possibly encapsulated within membrane bilayer of lipid vesicles. As can be seen in **Figure 2B**, the EE of liposomes and nanospheres ranges from 30% to ≥90%, while the EE of polymeric micelles is not mentioned. This suggests that drugs are successfully encapsulated in nanoparticles in order to increase the amount of drugs delivered to the target sites.

## Pharmacokinetic Data in Clinical Trials

From the review of pharmacokinetic data relating to clinical trials, it was found that in Lipo A the average  $t_{1/2}$  of UAL was 4.00–4.58 h, a

low value of  $t_{1/2}$  resulting in its rapid elimination from the blood circulation as shown by the contents of **Table 3**. This suggests that UAL does not accumulate in the body but must be infused repeatedly to ensure the steady plasma concentration of UA and further enhance its antitumor effect (Qian et al., 2015). In Lipo B, a linear relationship exists between  $C_{max}$  or  $AUC_{0 \rightarrow 24h}$  or  $AUC_{0 \rightarrow \infty}$  and increased doses of UAL, signifying that UAL has a linear pharmacokinetic profile (Wang et al., 2013). In Lipo C, after administration of a single IV dose, the total concentration of UA in all subjects experienced a two-fold decrease. On completion of IV infusion, the plasma concentration of UA rapidly decreases to one approximately ten times lower than the peak concentration after 2 hours. The pharmacokinetics profiles of UAL are linear and dosage



**FIGURE 2 | (A)** Types of drug carrier extracted from the article review regarding the preclinical and clinical studies of nanoparticle use for UA delivery within cancer therapy, **(B)** the physical characteristics of UA-loaded nanoparticles including particle size, zeta potential, and efficiency of encapsulation.

**TABLE 3 |** Pharmacokinetic data from clinical trials of UA-loading nanoparticles.

Parameter	Lipo A		Lipo B		Lipo C		
	Administration Route	Intravenous	Intravenous		Intravenous		
Dose (mg/m <sup>2</sup> )	74 (double dose)	37	74	98	37	74 (single dose)	98
t <sub>1/2</sub> T/2(hours)	4.58 ± 2.04	4.59 ± 2.44	4.46 ± 1.41	3.90 ± 2.08	4.59 ± 2.44	4.46 ± 1.41	3.90 ± 2.08
V <sub>d</sub> (L/m <sup>2</sup> )	NA	NA	NA	NA	58.7 ± 33.0	64.3 ± 17.9	55.4 ± 28.1
CL (L/h/m <sup>2</sup> )	NA	8.65 ± 1.09	10.2 ± 1.46	9.94 ± 1.13	8.67 ± 1.07	10.20 ± 1.46	9.94 ± 1.13
AUC <sub>0-t</sub> (ng·h/mL)<	5,172 ± 1,136	4,213 ± 606	7,175 ± 999	9,696 ± 1,134	4,203 ± 588	7,175 ± 999	9,696 ± 1,134
AUC <sub>0-∞</sub> (ng·h/mL)	5,498 ± 1,525	4,339 ± 574	7,418 ± 1,057	9,971 ± 1,144	4,329 ± 556	7,418 ± 1,057	9,971 ± 1,144
MRT <sub>0-t</sub> (hour)	NA	3.69 ± 0.36	3.93 ± 0.37	3.84 ± 0.34	3.69 ± 0.36	3.93 ± 0.37	3.84 ± 0.34
MRT <sub>0-∞</sub> (hour)	NA	4.28 ± 0.91	4.56 ± 0.88	4.41 ± 0.95	4.29 ± 0.90	4.56 ± 0.88	4.41 ± 0.95
C <sub>max</sub> (ng/ml)	1,589 ± 635	1835 ± 438	2,865 ± 868	3,457 ± 856	1835 ± 438	2,865 ± 868	3,457 ± 856
T <sub>max</sub> (hour)	NA	4.03 ± 0.04	4.02 ± 0.04	4.0 ± 0.00	4.03 ± 0.04	4.02 ± 0.04	4.00 ± 0.00

Notes: t<sub>1/2</sub>, half-life time; V<sub>d</sub>, distribution volume; CL, clearance; AUC, area under curve of concentration vs time; MRT, mean retention time; C<sub>max</sub>, maximum plasma concentration; T<sub>max</sub>, time required to reach maximum plasma concentration.

proportional at a range of 37 mg/m<sup>2</sup> to 98 mg/m<sup>2</sup>. No accumulation of UA was observed following repeated doses of UAL in eight patients after receiving continuous IV infusion 74 mg/m<sup>2</sup> over a 14-day period (Zhu et al., 2013).

## Pharmacokinetic Data Review on *In Vivo* Studies

The pharmacokinetic data on *in vivo* studies of Lipo E shows that the highest plasma UA concentration in the PEGylated liposome treatment group was 19.87 ug/mL, which exceeded that of both the Ursolic Acid Liposomes (UAL) and Ursolic Acid (UA)

solution groups. In addition, as seen from **Table 4**, PEG-modified liposomes have the longest t<sub>1/2</sub>, while C<sub>max</sub> and AUC in the bloodstream have similar trends. This suggests that PEGylated UA liposomes may extend the time required for the drug to circulate in the circulatory system and produce a slow release effect (Wang et al., 2016).

In Nano A, after administration of a hydrophobic drug, i.e., hydroxycamptothecin (HCPT), conjugated to folic acid-pectin-eight-arm PEG-UA (F-Pt-PU) the concentration of UA and HCPT in plasma decreases slowly, resulting in the longer circulation period of native UA, which may be due to the breaking of ester bonds between 8 arm-PEG and UA. The concentration of

**TABLE 4 |** Pharmacokinetic data summary of preclinical studies of nanoparticles containing UA.

Parameter	Lipo E	Nano A	Nano G	Poli B
Administration route	intragastric	intravenous	intravenous	intravenous
UA dose (mg/kg)	80	10	10	11
T <sub>1/2</sub> (hour)	8.6	8.3 and 10	8.7	4.9 and 5.2
AUC (μg.h/ml)	134.061	NA	NA	NA
C <sub>max</sub> (μg/ml)	19.87	NA	NA	NA
T <sub>max</sub> (hour)	NA	80	80	48

Notes: T<sub>1/2</sub>, half-life time; AUC, area under curve of concentration versus time; C<sub>max</sub>, maximum plasma concentration; T<sub>max</sub>, time required to reach maximum plasma concentration.

NP HCPT@F-Pt-PU in the bloodstream, still detectable at 80 h, is higher than that of native UA (7 h) and HCPT (8 h). The concentration of NP HCPT @F-Pt-PU in plasma is higher than that of np F-Pt-PU, possibly because the conjugation of HCPT into polymers increases the strength of the hydrophobic bonds in the particle cores, thereby reducing the hydrolysis rate of nanoparticles (Liu et al., 2021).

In Nano G, UA blood circulation in pectin-eight-arm polyethylene glycol-ursolic acid/hydrooxycamptothecin nanoparticles (NP Pec-8PUH) at 80 h can be maintained at a higher concentration in plasma, while native UA and HCPT rapidly disappear from plasma. The half-life of UA blood circulation in NP Pec-8PUH is 8.7 h which is 7.3 times longer than in native UA (Liu et al., 2018).

In Poly B, polymeric drug conjugates are synthesized by conjugating UA into polyethylene glycol using disulfide bonds (U-SS-M), while UA is eliminated relatively slowly and maintained at high concentrations in plasma for up to 48 h after administration. U-SS-M exhibits a similar pattern of biodistribution and accumulates mainly in the liver and kidneys before being subsequently eliminated by these organs. In tumor tissue, the concentration of UA decreases over time, although the amount delivered by the polymer-drug conjugate gradually increases. The concentration of U-SS-M in tumor tissue is significantly higher than that of native UA at both 6 and 12 h after administration (Fu et al., 2021).

## Recapitulation of Pre-Clinical and Clinical Research Relating to UA Nanoparticles

The results indicate that the available articles which discuss pre-clinical/*in vivo* trials amounted of 83%, including the use of

nanoparticle carrier types of nanospheres (47%), liposomes (40%), and polymeric micelles (13%). As for those that discussed clinical trials (17%), as seen in **Figure 3A–C**, these featured only the use of liposomes (100%). Clinical trials are still being conducted in phase 1, indicating that they remain at the stage of evaluating dose levels, acute toxicity, and drug distribution in humans (Valcourt et al., 2020).

## In vivo Anti-Cancer Efficacy of Nanoparticles Containing UA

### Analysis of Tumor Tissue Histopathology


The anticancer effectiveness of nanoparticles containing UA compared to negative control and free UA are shown to have a significant effect on tumor growth inhibition as shown in **Table 5**.


### Comparative Analysis of Tumor Growth Inhibition

The results indicate that the normal tumor volume when compared to administration of UA-loaded liposomes (Lipo D, E, F, G, H, I), nanospheres (Nano A, B, D, E, F, G) and polymeric micelles (Poly A) decreased in relative tumor volume by approximately 2.0–21.2 times. The tumor volume of native UA compared to the administration of UA liposomes (Lipo D, E, F, G, H, I), nanospheres (Nano A, D, E, F, G) and polymeric micelle (Poly A) showed a relative reduction in tumor volume of about 1.6–15.9 times lower than that of the native UA group, as presented in **Figure 5A**. This suggests that nanoparticles can improve UA effectiveness in inhibiting expansions in tumor volume.

The relative tumor weight analysis results relating to groups treated with UA liposomes (Lipo D, E, F), nanospheres (Nano C, E) and polymeric micelles (Poly A,B) indicated a relative reduction in tumor weight approximately 1.9–5.3 times that of the negative control group. Tumor weight in the native UA group compared to that of groups administered with UA liposomes (Lipo D, E, F), nanospheres (Nano C,E) and polymeric micelles (Poly A,B) showed a relative reduction of about 1.6–3.2x, as shown in **Figure 5B**. This suggests that nanoparticles may increase the effectiveness of UA in inhibiting tumor growth resulted in reduction of tumor weight.

The relative inhibition rate analysis results indicate that the administration of UA liposomes (Lipo E, F), nanospheres (Nano A) and polymeric micelles (Poly B) produces an increase in the

**FIGURE 3 |** Research recapitulation of **(A)** clinical and preclinical studies, **(B)** types of nanoparticle use in clinical trials, **(C)** and pre-clinical trials.



**TABLE 5 |** Tissue histopathology of liver cancer after administration of negative control, native UA and nanoparticles containing UA.

Code	Tissue histopathology		
	Negative control	Free UA	Nanoparticles containing UA
Lipo E	It features no hemorrhagic or necrosis phenomena and the cell is round or polygonal	Tumor cells and angiogenesis occur in native UA solution and conventional UA liposomes treatment groups, which become rare with slight necrosis	The tumor cells of the UA liposome with polyethylene glycols (PEGylated UA Liposome) group undergo severe necrosis, the nucleus/pulp ratio is significantly reduced, and apoptosis occurs due to a large number of scattered single tumor cells
Lipo F	The nucleus size and tumor cell shape are irregular. The tumor cells have clear cellular morphology and chromatin indicating that the tumor cells are growing quickly	A limited shrinkage and fragmentation of the nucleus indicates a low rate of tumor cell necrosis	Most tumor tissue cells in the group treated with Chitosan-Ursolic Acid-Liposomes (CS-UA-L) undergo apoptosis or necrosis, indicating good potential for killing cancer cells
Nano B	There are numerous sinusoids and small blood vessels filled with blood (indicated by the arrow) spreading through the hepatocellular carcinoma trabeculae	Not available	Several sinusoid liver or blood vessels can be observed in Chitosan-Ursolic Acid-Nanoparticle (CH-UA-NP) group with the exception of liver sinusoid dysplasia. Massive necrotic tissue can still be observed in hepatocellular carcinoma
Poli A	Tumor necrosis is undefined in the saline treatment group	Tumor cells and angiogenesis become rare with little necrosis	Most cancer cells in the high-dose Ursolic Acid-Polymer Micelles (UA-PMs) group at 100 mg/kg showed a high degree of H22 cell necrosis


relative tumor inhibition rate of approximately 1.9-3.4x compared to the native UA group. Of the three types of drug carriers, liposomes (Lipo F) experienced the highest relative inhibition rate increase of 3.4x the native UA group, as seen in **Figure 4A**. This suggests that nanoparticles may increase the effectiveness of UA in inhibiting tumor growth.

#### Analysis of Survival Rate

Based on the results, the administration of liposomes (Lipo E, F), nanospheres (Nano A) and polymeric micelles (Poly B) produced

an increase in the relative survival rate about 1.3-2.2x higher when compared to that of the negative control group, as seen in **Figure 4B**. This suggests that nanoparticles may increase the effectiveness of UA associated with improved survival rate.

The increased anti-tumor activity observed from the volume and weight of the tumor was associated with necrosis in the tumor tissues caused by large dose exposures of UA reaching cancer cells due to the increased permeability of small nanoparticles with high drug loading due to the EPR effect. Furthermore, the drug will be released into the extracellular and/or intracellular matrix. In the



**FIGURE 5 | (A)** Relative tumor volume in animal models treated with UA-loaded nanoparticles compared to negative control (black bars) and UA-free treatment groups (grey bars), **(B)** relative tumor tissue weight of animal models treated with UA-loaded nanoparticles compared to negative control (black bars) and native UA-treatment groups (grey bars).

extracellular fluids, the drug will leak from nanoparticles and subsequently diffuse into cancer cells, while in intracellular fluids nanoparticles will experience endocytosis and the matrix will be destroyed in the endosome and release free drugs which then diffuse into the cytoplasm and nucleus subsequently causing cell necrosis. These results show that the use of nanoparticles as carriers within anticancer drug delivery can increase the *in vivo* survival rate.

Other studies have suggested that when nanoparticles such as liposomes interact with cells, drug delivery and diffusion into target cells can occur in several ways. Liposomes can penetrate the tumor tissue matrix resulting in degradation of carrier lipids by enzymes, such as lipase, or by mechanical strain inducing release of active substances into the extracellular fluid. This process induces drug diffusion into cell membranes culminating in cytoplasm and nucleus delivery. However, the latter process cannot easily be achieved by the use of hydrophilic drugs. Secondly, liposome membranes will fuse with those of the target cell leading to the release of liposomes directly into the cytoplasm. The third and most frequent method is that of receptor-mediated endocytosis. This process involves only vesicles with a maximum diameter of 150 nm and active substances demonstrating significant stability in such an acidic lysosome environment where liposomes are metabolized enzymatically. Phagocytosis may also ensue but involving large size nanoparticles affected by specialized immune system cells, such as macrophages, monocytes, and Kupffer cells. This process may eliminate the nanoparticles from the circulatory system (Bozzuto and Molinari, 2015).

The survival rate of liposomes is higher than that of other nanoparticles indicating the stability of the system in the blood circulation which ensures that the trapped drug is carried by the nanoparticles for further release into the cancer cells. In addition, because of the biomimetic property of liposome components that resemble phospholipid cell membranes it is easier for them to be absorbed by the cell.

## In vivo Toxicity Studies of Nanoparticles Containing UA

### Pre-Clinical Toxicity Based on the Analysis of Relative Body Weight

From the results of relative body weight calculations contained in Table 6, no significant differences existed in the weight of the mice, proving that nanoparticle administration neither caused side effects nor produced symptoms of toxicity (Poudel et al., 2020; Liu et al., 2021; Fu et al., 2021). This result is also supported by other toxicity data presented in Table 7, which shows that there was no clear cell damage and no morphological changes in the major organs i.e. heart, liver, spleen, lungs, and kidneys. However, ALT, AST and WBC levels all decreased after administration of UA nanoparticles when compared to native UA (Zhang et al., 2015; Zhao et al., 2018). This suggested that UA nanoparticles do not cause serious toxicity, indeed, do not even produce toxicity. Rather, the effect is mild and of short duration (Li et al., 2019).

**TABLE 6 |** The relative body weight of animal models treated with UA-loaded nanoparticles compared to negative control and native UA-treatment groups.

Code	Toxicity
Lipo H	ALT and AST levels were significantly higher following an injection of FA-UA/siRNA-L compared to that of saline solution. The AST/ALT ratio of the FA-UA/siRNA-L group was significantly lower than that of the saline group. These results suggest that liver toxicity caused by liposomes produces mild, temporary liver toxicity
Nano A	The number of rat WBCs in the NP HCPT@F-Pt-PU treatment group increased more rapidly than in the native UA group which suggests that folate-targeted pectin delivery systems may prevent serious hematological toxicity
Nano D	There was no obvious cell damage or morphological changes in the major organs i.e., heart, liver, spleen, lungs, and kidneys in the NP UA-LA-ICG treatment group members compared to those of the negative control group
Nano E	ALT levels in mice treated with UA-NP were significantly lower than in the CCl4 group members, but there were no changes in the native UA- treatment group. In addition, AST levels in the UA-NP treatment group were also significantly lower compared to the CCl4 group and the native UA-treatment groups
Nano F	The native UA group experienced necropsy in the central section of the tumor tissue. These results partly suggest that native UA causes more toxicity than UA-NP. Meanwhile, H&E staining indicated that there were no obvious abnormalities or inflammatory lesions in any of the five organs, i.e., heart, liver, spleen, lungs, kidneys for the UA-NP treatment group when compared to their negative control and native UA counterparts
Nano G	Rats treated with the Pec-8PUH-NPs group did not experience any significant reduction in WBC counts as an indicator of hematotoxicity suggesting that the use of nanoparticles might prevent hematological toxicity

**TABLE 7 |** Recapitulation of other preclinical toxicities.

Relative body weight	Results		
	Code	NC/AU-NP	AU/AU-NP
Lipo D	Decreased by 1.2x normal value (Not significant)		Increased by 1.0x AU value (Not significant)
Lipo E	Decreased by 1.1x normal value (Not significant)		Increased by 1.0x AU value (Not significant)
Lipo F	Decreased by 1.2x normal value (Not significant)		Decreased by 1.0x AU value (Not significant)
Lipo G	There is no obvious difference		Not available
Lipo H	Decreased by 1.0x normal value (Not significant)		There is no obvious difference
Nano A	There is no obvious difference		There is no obvious difference
Nano D	There is no obvious difference		There is no obvious difference
Nano E	There is no obvious difference		There is no obvious difference
Nano F	Decreased by 1.0x normal value (Not significant)		Increased by 0.9x AU value (Not significant)
Nano G	There is no obvious difference		There is no obvious difference
Poli A	Decreased by 1.1x normal value (Not significant)		Decreased by 1.0x AU value (Not significant)
Poli B	There is no obvious difference		There is no obvious difference

## Safety Aspects of the Use of Nanoparticles Containing UA Based on the Clinical Trials Toxicity Based on Clinical Laboratory Parameters


The content of the graphs contained in **Figure 6A**, confirm an increase in levels of AST, ALT, GGT, TG, DBIL, and TBIL after UA liposome (Lipo A, B, C) administration occurred. It can also be seen that Dose Limiting Toxicity (DLT) related to hepatotoxicity, which was monitored for substantial side effect parameters, was at a moderate level (Valcourt et al., 2020).

In Lipo A, an increase in levels of AST (5%), ALT (5%), GGT (14%), TG (5%) was observed in 21 subjects who received doses of 56, 74, and 98 mg/m<sup>2</sup>. In Lipo B, elevated levels of AST (13%), ALT (13%), GGT (15%), TG (8%), DBIL (5%) and TBIL (8%) were recorded in 39 subjects receiving doses of 11, 22, 37, 56, 74, 98, and 130 mg/m<sup>2</sup>. In Lipo C, there was an increase in AST (4%), ALT (4%), GGT (2%), TG (9%), and TBIL (4%) levels observed in 24 subjects who received


doses of 74 mg/m<sup>2</sup> as a single dose and 98 and 74 mg/m<sup>2</sup> as multiple doses.

## Clinical Toxicity Based on the Occurrences of Adverse Events

According to the data contained in **Figure 6B**, three subjects (14%) treated with a dose of 56 mg/m<sup>2</sup> of Lipo A experienced a mild fever but recovered after 2 hours without receiving treatment. Moreover, three subjects (14%) treated with sequential doses of 56, 74, and 98 mg/m<sup>2</sup> of Lipo A experienced an increase in GGT, two subjects (10%) administered with doses of 56 and 74 mg/m<sup>2</sup> of Lipo A experienced abdominal distension, and one patient (5%) experienced a rise in ALT levels. Other mild symptoms included increased AST and TG, pruritus, arthralgia, and hypokalemia. The most common adverse conditions included fever, increased GGT, and flatulence. These results indicated that a 4-h intravenous administration of Lipo A was tolerable and safe if a timetable of three doses per day for 14 consecutive days



**FIGURE 6 | (A)** Total subjects with increasing blood marker levels on clinical trials of Lipo A, Lipo B, and Lipo C, **(B)** Adverse events of Lipo A during phase I clinical trials. Notes: AST, Alanine Aminotransferase/SGPT (serum glutamic pyruvic transaminase); ALT, Aspartate Aminotransferase/SGOT (serum glutamic oxaloacetic transaminase); GGT, Gamma Glutamyl Transpeptidase; TG, Triglycerides; DBIL, Direct Bilirubin; TBIL, Total Bilirubin.



**FIGURE 7 |** Total subjects with adverse events of **(A)** Lipo B, and **(B)** Lipo C in phase I clinical trials. AST, Alanine Aminotransferase/SGPT (serum glutamic pyruvic transaminase); ALT, Aspartate Aminotransferase/SGOT (serum glutamic oxaloacetic transaminase); GGT, Gamma Glutamyl Transpeptidase; DBIL, Direct Bilirubin; TBIL, Total Bilirubin; TG, Triglycerides

followed by a break lasting 7 days within each 21-days cycle was adhered to. Therefore, a  $98 \text{ mg/m}^2$  dose of Lipo A is the recommended dose for phase II trials (Qian et al., 2015).

In addition, from the contents of **Figure 7A**, it can be seen that one patient treated with a  $11 \text{ mg/m}^2$  dose experienced a first degree skin rash which healed untreated after 3 days. In addition, two patients who had been administered with a  $98 \text{ mg/m}^2$  dose experienced vascular stimulation. First degree microscopic hematuria was observed in three subjects (7.7%) suffering from hepatoma malignancy who had received 11 doses of 11, 74, and  $130 \text{ mg/m}^2$ , respectively. However, these side effects disappeared after 3 days without any treatment being administered. Elevated levels of AST, ALT, GGT, DBIL, and TBIL were observed in several subjects receiving doses of 74, 98, and  $130 \text{ mg/m}^2$ . Dose Limiting Toxicity (DLT) resulted in hepatotoxicity: two subjects (5.1%) experienced an increase in AST, four subjects (10.3%) an increase in ALT, one subject (2.6%) an increase in GGT, and one subject (2.6%) an increase in DBIL. Diarrhea (2.6%) constitutes another DLT. Other drug-related side effects included nausea reported by one subject (2.6%), abdominal distension observed in another (2.6%), vascular stimulation occurred in two subjects (5.1%), while elevated TG was reported in three subjects (7.7%). Other reported adverse events included one subject (2.6%) suffering a skin rash and another (2.6%) experiencing higher serum sodium levels. At a dose of  $74 \text{ mg/m}^2$ , one of six subjects experienced DLT, which is a form of non-hematological toxicity, including increased AST/ALT and diarrhea. At a dose of  $98 \text{ mg/m}^2$ , one of the eleven subjects experienced DLT, i.e., non-hematological toxicity including increased ALT/GGT). At a dose of  $130 \text{ mg/m}^2$ , two thirds of the subjects experienced DLT (increased ALT, AST, and DBIL). Therefore, the increased dosage was suspended and MTD was confirmed to be  $98 \text{ mg/m}^2$ . Double administration of trial doses of UAL at recommended levels of 56, 74, and  $98 \text{ mg/m}^2$  was completed (Wang et al., 2013).

From the graph in **Figure 7B**, it is clear that all subjects in the study tolerated the Lipo C treatment. Most adverse events varying from mild to moderate related to Lipo C, which is Ursolic Acid Nanoliposome (UANL), were non-dose dependent. The most commonly observed adverse events included abdominal distension, nausea, and diarrhea. The adverse events after a 14-days continuous infusion of UANL comprised skin pruritus, arthrisgia, and increased triglycerides levels. UANL has minimal toxic effects. The limiting toxicity of UANL dose is hepatotoxicity. In this study, intravenous UANL infusions were well tolerated both by healthy volunteers and patients with advanced tumors (Zhu et al., 2013).

Based on the review analysis, only three articles which focused on liposomes as the drug carrier discussed clinical trials of UA. Although UA is classified as a BCS class IV drug, its permeability and solubility can be enhanced with the use of liposomes. It is related to the natural phase properties of the liposomal membrane that significantly affect permeability, aggregation, protein binding and liposome fusion. Membrane permeability largely depends on lipid components. Lipids that contain

saturated chains or do not have carbon double bonds are more stable because they demonstrate greater resistance to oxidation. Lipid bilayer and liposome membranes possess a good lipid-packing order or gel phase below the lipid phase transition temperature ( $T_c$ ), where the temperature is in a balanced proportion in the two phases. The fluidity of the lipid bilayer can be controlled by the selection and combined use of lipids, as the various  $T_c$ s depend on the length and origin sources (saturated or unsaturated) of fatty acid chains. For example, the incorporation of cholesterol at low concentrations into the lipid bilayer leads to increased transmembrane permeability, where the incorporation of large amounts ( $>30 \text{ mol\%}$ ) of cholesterol can reduce the transition phase and decrease membrane permeability at higher  $T_c$  temperatures (Corvera et al., 1992). Liposome permeability is related to the rate of solute diffusion through the lipid bilayer. The liposome membrane will achieve the highest permeability in the transition temperature phase, while its permeability is lower in gel form than in the fluid phase. The temperature of the bilayer phase transition is determined by the composition of the liposome. In the transition temperature phase, the permeability of liposomes to molecules such as protons and water increases (Koda et al., 2008; Yoshimoto et al., 2013; Hansen et al., 2015). In addition, the *in vivo* biodistribution and disposition of liposomes varies depending on the composition of the lipids, particle size, potential charge and degree of steric surface/hydration. In addition, the administration route may affect the *in vivo* disposition of liposomes. During intravenous administration, liposomes usually interact with serum proteins and are absorbed by RES cells, thus accumulating in the liver or spleen (Sercombe et al., 2015).

The development of nanoparticles for drug delivery, one of which is Doxil<sup>®</sup> (Doxorubicin HCl liposome injection), the first nanoliposomal drug approved by FDA in 1995, was based on three principles: 1) prolonging drug circulation time and RES avoidance due to the PEGylation of nanoliposomes; 2) higher stable loading of doxorubicin driven by the transmembrane ammonium sulfate gradient which also allows the re-release of the drug in tumors; and 3) having lipid bilayer liposomes in a "liquid ordered" phase consisting of phosphatidylcholine with a high melting temperature ( $T_m = 53^\circ\text{C}$ ), and which largely use cholesterol as a membrane stabilizer (Barenholz, 2012). In addition, various drug formulas in liposomes have received approval to be marketed and are widely used in clinical settings including Myocet<sup>®</sup> (Elan Pharmaceuticals Inc., Princeton, NJ, United States of America). This is an encapsulation of doxorubicin in liposomes (Swenson et al., 2001; Chastagner et al., 2015); Daunoxome<sup>®</sup> (Gilead Sciences), daunorubicin formulated into liposomes (Forssen, 1997; Dawidczyk et al., 2014); Marqibo<sup>®</sup> non-PEGylated liposomal vincristine developed in 2012 as a therapy for various cancers including lymphoma, brain, leukemia, or melanoma (Silverman and Deitcher, 2012); Onivyde<sup>®</sup> MM-398, which is a PEGylated liposomal irinotecan developed in 2015 as a drug to treat metastatic pancreatic cancer (Zhang, 2016), and many other forms of cancer (Ventola, 2017; Anselmo and Mitragotri, 2019). Various developments of the liposome delivery system

indicated that liposomes possess non-toxic, flexible, biocompatible, and biodegradable properties that can enhance the therapeutic effects, safety, and efficacy of various anticancer drugs (Akbarzadeh et al., 2013).

As for the development of cisplatin therapy, which incorporates the use of an anticancer drug, this involves a polymeric micelle delivery system. Polymeric micelles were prepared through the formation of a metal-polymer complex between cisplatin (CDDP) and poly-(ethylene glycol)-poly(glutamic acid) block copolymers. Cisplatin polymeric micelles (CDDP/m) are 28 nm in size with the ability to distribute themselves through narrow spaces such as blood vessels in pancreatic tissue. These micelles undergo lengthy blood circulation and accumulate effectively in solid tumors of Lewis lung carcinoma cells. However, because they are produced synthetically, the toxicity and safety aspects as well as manufacturing production scale constitute extremely important issues (Plummer et al., 2011).

Abraxane<sup>®</sup>, a paclitaxel albumin-bound nanoparticle with a particle size of ~130 nm which received FDA approval in 2005 for the treatment of metastatic breast cancer successfully reduces toxicities in comparison to Taxol<sup>®</sup>. Moreover, it enables a complete dose to be administered within 30 min without the need for any pre-treatment. Nevertheless, the mechanism of Abraxane<sup>®</sup> in improving survival rate and overcoming P-GP-mediated drug resistance remains unclear (Ma and Mumper, 2013).

The findings of this scoping review suggest that liposomes provide more comprehensive data than other forms of nanoparticles. This is demonstrated by the existence of *in vivo* studies of anticancer effectiveness assessed using several parameters such as increasing relative survival rate; more robust tumor growth inhibition (increasing relative inhibition rate, decreasing relative tumor weight, and reducing tumor volume); and improvements in tumor tissue histopathology. In addition, *in vivo* studies related to safety were also evaluated employing several parameters, i.e., weight loss, and other toxicity (lowering AST, ALT, and WBC), and well-tolerated toxicity by healthy volunteers and patients with advanced tumors.

There needs multi-faceted views of the use of nanoparticles for reviewing drug delivery. The components of the nanoparticle formulation would greatly affect the characteristics of the nanoparticles including particle size, potential charges, stealth and biomimetic properties, and others, which are closely related to drug delivery to cancer tissue, due to the Enhanced Permeation and Retention (EPR) effects. In addition, *in vivo* analysis of different types of cancer, where each type of cancer cell has different biological properties, also requires an in-depth study to provide data on supporting the effectiveness of drug delivery to target cancerous tissues. Moreover, the route of administration, dose, and frequency of drug administration related to the physicochemical properties and pharmacokinetic profile of the drug also greatly affect the systemic bioavailability and effective drug amount capable of reaching cancer tissue as the target of drug delivery. All these aspects provide important

views for comprehensive study of the drug delivery system in cancer therapy.

## CONCLUSION

Based on the scoping review of the relevant literature, it can be concluded that UA loaded into nanoparticles is effective as a form of anticancer therapy. Pre-clinical trials confirm that it increases the relative survival rate; tumor resistance (increasing the relative inhibition rate, lowering the relative tumor weight, and decreasing tumor volume); and improves tumor tissue histopathology. In addition, UA-loaded nanoparticles have been proven safe for anticancer therapy based on the evaluation of weight loss and other toxicity (decreased AST/ALT). The results from the last 10-years analysis have indicated that, compared to nanospheres and polymeric micelles, liposomes have been assessed as more effective and safer during more comprehensive pre-clinical and clinical trials. This finding highlights the potential for liposomes to be further developed as a means of delivering UA as an anticancer therapy.

## AUTHOR CONTRIBUTIONS

AM: 1) conception and design of the work, data acquisition, data analysis and interpretation; 2) critically revising the article for important intellectual content; 3) Final approval of the version to be published; 4) Agreement to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of the work are appropriately investigated and resolved. EM: 1) data acquisition; 2) Drafting the article; 3) Final approval of the version to be published; 4) Agreement to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of the work are appropriately investigated and resolved. RS: 1) data analysis and interpretation; 2) Final approval of the version to be published; 3) Agreement to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of the work are appropriately investigated and resolved. EH: 1) data analysis and interpretation; 2) Final approval of the version to be published; 3) Agreement to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of the work are appropriately investigated.

## FUNDING

This study was supported by a Preliminary Research on Excellence in Higher Education Institutions (Penelitian Dasar Unggulan Perguruan Tinggi, PDUPT) Grant No. 8/E1/KPT/2021, No. 4/E1/KP.PTNBH/2021, and No. 385/UN3.15PT/2021 provided by the Ministry of Research and Technology-National Research and Innovation Agency of Republic of Indonesia.

## REFERENCES

- Akbarzadeh, A., Rezaei-Sadabady, R., Davaran, S., Joo, S. W., Zarghami, N., Hanifehpour, Y., et al. (2013). Liposome: Classification, Preparation, and Applications. *Nanoscale Res. Lett.* 8 (1), 102. doi:10.1186/1556-276X-8-102
- Ali, S. A., Ibrahim, N. A., Mohammed, M. M. D., El-Hawary, S., and Refaat, E. A. (2019). The Potential Chemo Preventive Effect of Ursolic Acid Isolated from Paulownia Toomentosa, against N-Diethylnitrosamine: Initiated and Promoted Hepatocarcinogenesis. *Heliyon* 5 (5), e01769. doi:10.1016/j.heliyon.2019.e01769
- Alves, R. F., Favaro, M. T. P., Balbino, T. A., de Toledo, M. A. S., de la Torre, L. G., and Azzoni, A. R. (2017). Recombinant Protein-Based Nanocarriers and Their Association with Cationic Liposomes: Characterization and *In Vitro* Evaluation. *Colloids Surf. A: Physicochemical Eng. Aspects* 513, 1–10. doi:10.1016/j.colsurfa.2016.11.019
- Anselmo, A. C., and Mitragotri, S. (2019). Nanoparticles in the Clinic: An Update. *Bioeng. Transl. Med.* 4 (3), e10143–16. doi:10.1002/btm2.10143
- Baldim, I., Oliveira, W. P., Kadian, V., Rao, R., Yadav, N., Mahant, S., et al. (2020). Natural Ergot Alkaloids in Ocular Pharmacotherapy: Known Molecules for Novel Nanoparticle-Based Delivery Systems. *Biomolecules* 10 (7), 1–21. doi:10.3390/biom10070980
- Barenholz, Y. (2012). Doxil®—the First FDA-Approved Nano-Drug: Lessons Learned. *J. Control. Release* 160 (2), 117–134. doi:10.1016/j.jconrel.2012.03.020
- Biopharmaceutical, B. (2011). Classification System and Formulation Development. *Part. Sci.* 9, 1–4.
- Bozzuto, G., and Molinari, A. (2015). Liposomes as Nanomedical Devices. *Int. J. Nanomedicine* 10, 975–999. doi:10.2147/IJN.S68861
- Cabral, H., Matsumoto, Y., Mizuno, K., Chen, Q., Murakami, M., Kimura, M., et al. (2011). Accumulation of Sub-100 Nm Polymeric Micelles in Poorly Permeable Tumours Depends on Size. *Nat. Nanotechnol* 6 (12), 815–823. doi:10.1038/nnano.2011.166
- Chastagner, P., Devictor, B., Geoerger, B., Aerts, I., Leblond, P., Frappaz, D., et al. (2015). Phase I Study of Non-pegylated Liposomal Doxorubicin in Children with Recurrent/refractory High-Grade Glioma. *Cancer Chemother. Pharmacol.* 76 (2), 425–432. doi:10.1007/s00280-015-2781-0
- Chen, J., Ping, Q. N., Guo, J. X., Chu, X. Z., and Song, M. M. (2006). Effect of Phospholipid Composition on Characterization of Liposomes Containing 9-nitrocampothecin. *Drug Dev. Ind. Pharm.* 32 (6), 719–726. Jul. doi:10.1080/03639040500529077
- Corvera, E., Mouritsen, O. G., Singer, M. A., and Zuckermann, M. J. (1992). The Permeability and the Effect of Acyl-Chain Length for Phospholipid Bilayers Containing Cholesterol: Theory and experiment. *Biochim. Biophys. Acta* 1107 (2), 261–270. doi:10.1016/0005-2736(92)90413-g
- Dawidczyk, C. M., Kim, C., Park, J. H., Russell, L. M., Lee, K. H., Pomper, M. G., et al. (2014). State-of-the-art in Design Rules for Drug Delivery Platforms: Lessons Learned from FDA-Approved Nanomedicines. *J. Control. Release* 187, 133–144. doi:10.1016/j.jconrel.2014.05.036
- Etikan, I. (2017). The Kaplan Meier Estimate in Survival Analysis. *Biometrics Biostat Int. J.* 5 (2), 55–59. doi:10.15406/bbij.2017.05.00128
- Fan, L., Zhang, B., Xu, A., Shen, Z., Guo, Y., Zhao, R., et al. (2018). Carrier-Free, Pure Nanodrug Formed by the Self-Assembly of an Anticancer Drug for Cancer Immune Therapy. *Mol. Pharm.* 15 (6), 2466–2478. Jun. doi:10.1021/acs.molpharmaceut.8b00444
- Fang, J., Nakamura, H., and Maeda, H. (2011). The EPR Effect: Unique Features of Tumor Blood Vessels for Drug Delivery, Factors Involved, and Limitations and Augmentation of the Effect. *Adv. Drug Deliv. Rev.* 63 (3), 136–151. Available from: doi:10.1016/j.addr.2010.04.009
- Forsen, E. A. (1997). The Design and Development of DaunoXome® for Solid Tumor Targeting *In Vivo*. *Adv. Drug Deliv. Rev.* 24 (2), 133–150. doi:10.1016/s0169-409x(96)00453-x
- Fu, D., Ni, Z., Wu, K., Cheng, P., Ji, X., Li, G., et al. (2021). A Novel Redox-Responsive Ursolic Acid Polymeric Prodrug Delivery System for Osteosarcoma Therapy. *Drug Deliv.* 28 (1), 195–205. doi:10.1080/10717544.2020.1870583
- Gabizon, A., Shmeeda, H., and Grenader, T. (2012). Pharmacological Basis of Pegylated Liposomal Doxorubicin: Impact on Cancer Therapy. *Eur. J. Pharm. Sci.* 45 (4), 388–398. doi:10.1016/j.ejps.2011.09.006
- Globocan. Indonesia - Global Cancer Observatory WHO. International Agency for Research on Cancer, 2018. Available at: <Https://GcoIarcFr/Today/Data/Factsheets/Populations/360-Indonesia-Fact-SheetsPdf>. 256. ( Accessed 2020 1 2).
- Hansen, A. E., Petersen, A. L., Henriksen, J. R., Boerresen, B., Rasmussen, P., Elema, D. R., et al. (2015). Positron Emission Tomography Based Elucidation of the Enhanced Permeability and Retention Effect in Dogs with Cancer Using Copper-64 Liposomes. *ACS Nano* 9 (7), 6985–6995. doi:10.1021/acsnano.5b01324
- Hattori, Y., Yamashita, J., Sakaida, C., Kawano, K., and Yonemochi, E. (2014). Evaluation of Antitumor Effect of Zoledronic Acid Entrapped in Folate-Linked Liposome for Targeting to Tumor-Associated Macrophages. *J. Liposome Res.* 25 (2), 131–140. doi:10.3109/08982104.2014.954128
- Internatioanl Agency for Research on Cancer (2018). *Latest Global Cancer Data : Cancer burden Rises to 18.1 Million New Cases and 9 . 6 Million Cancer Deaths in 2018*.
- Jin, H., Pi, J., Yang, F., Jiang, J., Wang, X., Bai, H., et al. (2016). Folate-Chitosan Nanoparticles Loaded with Ursolic Acid Confer Anti-breast Cancer Activities *In Vitro* and *In Vivo*. *Sci. Rep.* 6, 30782, 2016. July 2016. doi:10.1038/srep30782
- Jin, H., Pi, J., Yang, F., Wu, C., Cheng, X., Bai, H., et al. (2016). Ursolic Acid-Loaded Chitosan Nanoparticles Induce Potent Anti-angiogenesis in Tumor. *Appl. Microbiol. Biotechnol.* 100, 6643–6652. doi:10.1007/s00253-016-7360-8
- Kamińska, K., Szczylak, C., Bielecka, Z. F., Bartnik, E., Porta, C., Lian, F., et al. (2015). The Role of the Cell-Cell Interactions in Cancer Progression. *J. Cel Mol Med* 19 (2), 283–296. doi:10.1111/jcmm.12408
- Kashyap, D., Tuli, H. S., and Sharma, A. K. (2016). Ursolic Acid (UA): A Metabolite with Promising Therapeutic Potential. *Life Sci.* 146, 201–213. doi:10.1016/j.lfs.2016.01.017
- Kharia, A. A., Singhai, A. K., and Verma, R. (2012). Formulation and Evaluation of Polymeric Nanoparticles of an Antiviral Drug for Gastoretention. *PCI-Accredited-IJPSN* 4 (4), 1557–1562. doi:10.37285/ijpsn.2011.4.4.6
- Khwaza, V., Oyedele, O. O., and Aderibigbe, B. A. (2020). *Ursolic Acid-Based Derivatives as Potential Anti-cancer Agents : An Update*.
- Koda, Y., Liang, M. T., Blanchfield, J. T., and Toth, I. (2008). *In Vitro* stability and Permeability Studies of Liposomal Delivery Systems for a Novel Lipophilic Endomorphin 1 Analogue. *Int. J. Pharm.* 356 (1–2), 37–43. doi:10.1016/j.ijpharm.2007.12.036
- Koppolu, B., and Zaharoff, D. A. (2013). The Effect of Antigen Encapsulation in Chitosan Particles on Uptake, Activation and Presentation by Antigen Presenting Cells. *Biomaterials* 34 (9), 2359–2369. Mar. doi:10.1016/j.biomaterials.2012.11.066
- Kulkarni, S. B., Betageri, G. V., and Singh, M. (1995). Factors Affecting Microencapsulation of Drugs in Liposomes. *J. Microencapsul* 12 (3), 229–246. doi:10.3109/02652049509010292
- Kumari, P., Ghosh, B., and Biswas, S. (2016). Nanocarriers for Cancer-Targeted Drug Delivery. *J. Drug Target.* 24 (3), 179–191. doi:10.3109/1061186X.2015.1051049
- Laouini, A., Jaafar-Maalej, C., Limayem-Bloua, I., Sfar, S., Charcosset, C., Fessi, H., et al. (2012). Preparation, Characterization and Applications of Liposomes: State of the Art. *J Coll. Sci. Biotechnol.* 1, 147–168. doi:10.1166/jcsb.2012.1020
- Li, W., Yan, R., Liu, Y., He, C., Zhang, X., Lu, Y., et al. (2019). Co-delivery of Bmi1 Small Interfering RNA with Ursolic Acid by Folate Receptor-Targeted Cationic Liposomes Enhances Anti-tumor Activity of Ursolic Acid *In Vitro* and *In Vivo*. *Drug Deliv.* 26 (1), 794–802. doi:10.1080/10717544.2019.1645244
- Li, Z., Tan, S., Li, S., Shen, Q., and Wang, K. (2017). Cancer Drug Delivery in the Nano Era: An Overview and Perspectives (Review). *Oncol. Rep.* 38 (2), 611–624. doi:10.3892/or.2017.5718
- Liu, Y., Kong, T., Yang, Z., Zhang, Y., Lei, J., and Zhao, P. (2021). Self-Assembled Folic Acid-Targeted Pectin-Multi-Arm Polyethylene Glycol Nanoparticles for Tumor Intracellular Chemotherapy. *ACS omega* 6 (2), 1223–1234. Jan. doi:10.1021/acsomega.0c04350
- Liu, Y., Liu, K., Li, X., Xiao, S., Zheng, D., Zhu, P., et al. (2018). A Novel Self-Assembled Nanoparticle Platform Based on Pectin-Eight-Arm Polyethylene Glycol-Drug Conjugates for Co-delivery of Anticancer Drugs. *Mater. Sci. Eng. C Mater. Biol. Appl.* 86, 28–41. doi:10.1016/j.msec.2017.12.018
- Lunardi, C. N., Gomes, A. J., Rocha, F. S., De Tommaso, J., and Patience, G. S. (2021). Experimental Methods in Chemical Engineering: Zeta Potential. *Can. J. Chem. Eng.* 99 (3), 627–639. Mar 1. doi:10.1002/cjce.23914

- Ma, P., and Mumper, R. J. (2013). Paclitaxel Nano-Delivery Systems: A Comprehensive Review. *J. Nanomed. Nanotechnol.* 4 (2), 1000164. doi:10.4172/2157-7439.1000164
- Melo, R. C. N., Raas, M. W. D., Palazzi, C., Neves, V. H., Malta, K. K., and Silva, T. P. (2020). Whole Slide Imaging and its Applications to Histopathological Studies of Liver Disorders. *Front. Med. (Lausanne)* 6 (January), 310–313. doi:10.3389/fmed.2019.00310
- Miatmoko, A., Kawano, K., Yoda, H., Yonemochi, E., and Hattori, Y. (2017). Tumor Delivery of Liposomal Doxorubicin Prepared with Poly-L-Glutamic Acid as a Drug-Trapping Agent. *J. Liposome Res.* 27 (2), 99–107. doi:10.3109/08982104.2016.1166511
- Miele, E., Spinelli, G. P., Miele, E., Tomao, F., and Tomao, S. (2009). Albumin-bound Formulation of Paclitaxel (Abraxane ABI-007) in the Treatment of Breast Cancer. *Int. J. Nanomedicine* 4, 99–105. 2009/04/20. doi:10.2147/ijn.s3061
- Mitchell, M. J., Billingsley, M. M., Haley, R. M., Wechsler, M. E., Peppas, N. A., and Langer, R. (2021). Engineering Precision Nanoparticles for Drug Delivery. *Nat. Rev. Drug Discov.* 20, 101–124. doi:10.1038/s41573-020-0090-8
- Mizumura, Y., Matsumura, Y., Hamaguchi, T., Nishiyama, N., Kataoka, K., Kawaguchi, T., et al. (2001). Cisplatin-incorporated Polymeric Micelles Eliminate Nephrotoxicity, while Maintaining Antitumor Activity. *Jpn. J. Cancer Res.* 92 (March), 328–336. doi:10.1111/j.1349-7006.2001.tb01099.x
- Monteiro, N., Martins, A., Reis, R. L., and Neves, N. M. (2014). Liposomes in Tissue Engineering and Regenerative Medicine. *J. R. Soc. Interf.* 11 (101), 20140459. doi:10.1098/rsif.2014.0459
- Nishiyama, N., Okazaki, S., Cabral, H., Miyamoto, M., Kato, Y., Sugiyama, Y., et al. (2003). Novel Cisplatin-Incorporated Polymeric Micelles Can Eradicate Solid Tumors in Mice. *Cancer Res.* 63 (24), 8977–8983.
- Pérez-Herrero, E., and Fernández-Medarde, A. (2015). Advanced Targeted Therapies in Cancer: Drug Nanocarriers, the Future of Chemotherapy. *Eur. J. Pharm. Biopharm.* 93, 52–79.
- Plummer, R., Wilson, R. H., Calvert, H., Boddy, A. V., Griffin, M., Sludden, J., et al. (2011). A Phase I Clinical Study of Cisplatin-Incorporated Polymeric Micelles (NC-6004) in Patients with Solid Tumours. *Br. J. Cancer* 104 (4), 593–598. doi:10.1038/bjc.2011.6
- Poudel, K., Gautam, M., Maharjan, S., Jeong, J. H., Choi, H. G., Khan, G. M., et al. (2020). Dual Stimuli-Responsive Ursolic Acid-Embedded Nanophytoliposome for Targeted Antitumor Therapy. *Int. J. Pharm.* 582 (December 2019), 119330. doi:10.1016/j.ijpharm.2020.119330
- Qian, Z., Wang, X., Song, Z., Zhang, H., Zhou, S., Zhao, J., et al. (2015). A Phase I Trial to Evaluate the Multiple-Dose Safety and Antitumor Activity of Ursolic Acid Liposomes in Subjects with Advanced Solid Tumors. *Biomed. Res. Int.* 2015, 809714–809717. doi:10.1155/2015/809714
- Rajesh, S. J. W. L., Jr. (2000). Nanoparticle-based Targeted Drug Delivery. *Exp. Mol. Pathol.* 86 (3), 215–223.
- Reddy, L. H. (2005). Drug Delivery to Tumours: Recent Strategies, Internet. *J. Pharm. Pharmacol.* 57 (10), 1231–1242. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/16259751>. doi:10.1211/jpp.57.10.0001
- Seo, D. Y., Lee, S. R., Heo, J. W., No, M. H., Rhee, B. D., Ko, K. S., et al. (2018). Ursolic Acid in Health and Disease. *Korean J. Physiol. Pharmacol.* 22 (3), 235–248. doi:10.4196/kjpp.2018.22.3.235
- Sercombe, L., Veerati, T., Moheimani, F., Wu, S. Y., Sood, A. K., and Hua, S. (2015). Advances and Challenges of Liposome Assisted Drug Delivery. *Front. Pharmacol.* 6 (DEC), 286. doi:10.3389/fphar.2015.00286
- Shaker, S., Gardouh, A. R., and Ghorab, M. M. (2017). Factors Affecting Liposomes Particle Size Prepared by Ethanol Injection Method. *Res. Pharm. Sci.* 12 (5), 346–352. doi:10.4103/1735-5362.213979
- Sharma, A., and Sharma, U. S. (1997). Liposomes in Drug Delivery: Progress and Limitations. *Int. J. Pharmaceutics* 154 (2), 123–140. doi:10.1016/s0378-5173(97)00135-x
- Shen, Z., Li, B., Liu, Y., Zheng, G., Guo, Y., Zhao, R., et al. (2018). A Self-Assembly Nanodrug Delivery System Based on Amphiphilic Low Generations of PAMAM Dendrimers-Ursolic Acid Conjugate Modified by Lactobionic Acid for HCC Targeting Therapy. *Nanomedicine* 14 (2), 227–236. doi:10.1016/j.nano.2017.10.007
- Silverman, J. A., and Deitcher, S. R. (2012). Marqibo® (Vincristine Sulfate Liposome Injection) Improves the Pharmacokinetics and Pharmacodynamics of Vincristine. *Cancer Chemother. Pharmacol.* 71 (3), 555–564. 2013 Mar. doi:10.1007/s00280-012-2042-4
- Slaoui, M., and Fiette, L. (2011). Histopathology Procedures: from Tissue Sampling to Histopathological Evaluation. *Methods Mol. Biol.* 691 (December 2014), 69–82. doi:10.1007/978-1-60761-849-2\_4
- Sun, Q., He, M., Zhang, M., Zeng, S., Chen, L., Zhou, L., et al. (2020). Ursolic Acid: A Systematic Review of its Pharmacology, Toxicity and Rethink on its Pharmacokinetics Based on PK-PD Model. *Fitoterapia* 147, 104735. doi:10.1016/j.fitote.2020.104735
- Swenson, C. E., Perkins, W. R., Roberts, P., and Janoff, A. S. (2001). Liposome Technology and the Development of Myocet (Liposomal Doxorubicin Citrate). *The Breast* 10, 1–7. doi:10.1016/s0960-9776(01)80001-1
- Uchino, H., Matsumura, Y., Negishi, T., Koizumi, F., Hayashi, T., Honda, T., et al. (2005). Cisplatin-incorporating Polymeric Micelles (NC-6004) Can Reduce Nephrotoxicity and Neurotoxicity of Cisplatin in Rats. *Br. J. Cancer* 93 (6), 678–687. doi:10.1038/sj.bjc.6602772
- Valcourt, D. M., Kapadia, C. H., Scully, M. A., Dang, M. N., and Day, E. S. (2020). Best Practices for Preclinical *In Vivo* Testing of Cancer Nanomedicines. *Adv. Healthc. Mater.* 9 (12), e2000110–15. doi:10.1002/adhm.202000110
- Ventola, C. L. (2017). Progress in Nanomedicine: Approved and Investigational Nanodrugs. *P. T.* 42 (12), 742–755.
- Wang, M., Zhao, T., Liu, Y., Wang, Q., Xing, S., Li, L., et al. (2017). Ursolic Acid Liposomes with Chitosan Modification: Promising Antitumor Drug Delivery and Efficacy. *Mater. Sci. Eng. C Mater. Biol. Appl.* 71, 1231–1240. doi:10.1016/j.msec.2016.11.014
- Wang, Q., Zhao, T., Liu, Y., Xing, S., Li, L., and Gao, D. (2016). An Evaluation of Anti-tumor Effect and Toxicity of PEGylated Ursolic Acid Liposomes. *J. Nanoparticle Res.* 18 (2), 1–13. doi:10.1007/s11051-016-3339-8
- Wang, X. H., Zhou, S. Y., Qian, Z. Z., Zhang, H. L., Qiu, L. H., Song, Z., et al. (2013). Evaluation of Toxicity and Single-Dose Pharmacokinetics of Intravenous Ursolic Acid Liposomes in Healthy Adult Volunteers and Patients with Advanced Solid Tumors. *Expert Opin. Drug Metab. Toxicol.* 9 (2), 117–125. doi:10.1517/17425255.2013.738667
- Webb, C., Khadke, S., Schmidt, S. T., Roces, C. B., Forbes, N., Berrie, G., et al. (2019). The Impact of Solvent Selection: Strategies to Guide the Manufacturing of Liposomes Using Microfluidics. *Pharmaceutics* 11 (12), 653. doi:10.3390/pharmaceutics11120653
- Wei, X., Cohen, R., and Barenholz, Y. (2016). Insights into Composition/structure/function Relationships of Doxil® Gained from "High-Sensitivity" Differential Scanning Calorimetry, Internet. *Eur. J. Pharm. Biopharm.* 104, 260–270. Available from: <http://linkinghub.elsevier.com/retrieve/pii/S0939641116301357>. doi:10.1016/j.ejpb.2016.04.011
- Yang, G., Yang, T., Zhang, W., Lu, M., Ma, X., and Xiang, G. (2014). Vitro and *In Vivo* Antitumor E Ff Ects of Folate-Targeted Ursolic AcidIn Vitro and *In Vivo* Antitumor Effects of Folate-Targeted Ursolic Acid Stealth Liposome. *J. Agric. Food Chem.* 62 (10), 2207–2215. doi:10.1021/jf405675g
- Yoshimoto, M., Tamura, R., and Natsume, T. (2013). Liposome Clusters with Shear Stress-Induced Membrane Permeability. *Chem. Phys. Lipids* 174, 8–16. doi:10.1016/j.chemphyslip.2013.06.001
- Zaimy, M. A., Saffarzadeh, N., Mohammadi, A., Pourghadamayari, H., Izadi, P., Sarli, A., et al. (2017). New Methods in the Diagnosis of Cancer and Gene Therapy of Cancer Based on Nanoparticles. *Cancer Gene Ther.* 24 (6), 233–243. doi:10.1038/cgt.2017.16
- Zhang, H. (2016). Onivyde for the Therapy of Multiple Solid Tumors. *Oncotargets Ther.* 9, 3001–3007. doi:10.2147/OTT.S105587
- Zhang, H., Zheng, D., Ding, J., Xu, H., Li, X., and Sun, W. (2015). Efficient Delivery of Ursolic Acid by poly(N-Vinylpyrrolidone)-Block-Poly( $\epsilon$ -Caprolactone) Nanoparticles for Inhibiting the Growth of Hepatocellular Carcinoma *In Vitro* and *In Vivo*. *Int. J. Nanomedicine* 10, 1909–1920. doi:10.2147/IJN.S77125
- Zhang, N., Liu, S., Shi, S., Chen, Y., Xu, F., Wei, X., et al. (2020). Solubilization and Delivery of Ursolic-Acid for Modulating Tumor Microenvironment and Regulatory T Cell Activities in Cancer Immunotherapy. *J. Control. Release* 320 (January), 168–178. doi:10.1016/j.jconrel.2020.01.015
- Zhao, R., Zheng, G., Fan, L., Shen, Z., Jiang, K., Guo, Y., et al. (2018). Carrier-free Nanodrug by Co-assembly of Chemotherapeutic Agent and Photosensitizer for

- Cancer Imaging and Chemo-Photo Combination Therapy. *Acta Biomater.* 70, 197–210. doi:10.1016/j.actbio.2018.01.028
- Zhou, M., Yi, Y., Liu, L., Lin, Y., Li, J., Ruan, J., et al. (2019). Polymeric Micelles Loading with Ursolic Acid Enhancing Anti-tumor Effect on Hepatocellular Carcinoma. *J. Cancer* 10 (23), 5820–5831. doi:10.7150/jca.30865
- Zhu, Z., Qian, Z., Yan, Z., Zhao, C., Wang, H., and Ying, G. (2013). A Phase I Pharmacokinetic Study of Ursolic Acid Nanoliposomes in Healthy Volunteers and Patients with Advanced Solid Tumors. *Int. J. Nanomedicine* 8, 129–136. doi:10.2147/IJN.S38271

**Conflict of Interest:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

**Publisher's Note:** All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2021 Miatmoko, Mianing, Sari and Hendradi. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

←

Ads by Google

[Stop seeing this ad](#) [Why this ad? ⓘ](#)

## Frontiers in Pharmacology

**COUNTRY**[Switzerland](#) Universities and research institutions in Switzerland**SUBJECT AREA AND CATEGORY**

[Medicine](#)  
[Pharmacology \(medical\)](#)  
[Pharmacology, Toxicology and Pharmaceutics](#)  
[Pharmacology](#)

**PUBLISHER**[Frontiers Media S.A.](#)**H-INDEX****104**

←

Ads by Google

[Stop seeing this ad](#)[Why this ad? ⓘ](#)**PUBLICATION TYPE**[Journals](#)**ISSN**

16639812

**COVERAGE**

2010-2021

**INFORMATION**[Homepage](#)[How to publish in this journal](#)

Ads by Google

[Stop seeing this ad](#) [Why this ad? ⓘ](#)

← Ads by Google

[Stop seeing this ad](#)

[Why this ad? ⓘ](#)

## SCOPE

Frontiers in Pharmacology is a leading journal in its field, publishing rigorously peer-reviewed research across disciplines, including basic and clinical pharmacology, medicinal chemistry, pharmacy and toxicology. Field Chief Editor Heike Wulff at UC Davis is supported by an outstanding Editorial Board of international researchers. This multidisciplinary open-access journal is at the forefront of disseminating and communicating scientific knowledge and impactful discoveries to researchers, academics, clinicians and the public worldwide.

Join the conversation about this journal

←

Ads by Google

[Stop seeing this ad](#)

[Why this ad? ⓘ](#)

Quartiles  
I

## FIND SIMILAR JOURNALS ⓘ

←

Ads by Google

[Stop seeing this ad](#)

[Why this ad? ⓘ](#)

1  
**Pharmacological Research**

USA

**87%**

similarity

2  
**Life Sciences**

USA

**83%**

similarity

3  
**Current Drug Targets**

ARE

**82%**

similarity

4  
**Acta Pharmacologica Sinica**

GBR


**79%**

similarity

Ads by Google

[Stop seeing this ad](#)

[Why this ad? ⓘ](#)



## SCImago Graphica

Explore, visually communicate and make sense of data with our [new data visualization tool](#).



← Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scimagojr.com/widget.php?...>
```



16 Types

[Learn more](#)

Project Management

[Learn more](#)

CSS

[Learn more](#)

Metrics based on Scopus® data as of April 2022

Ads by Google

[Stop seeing this ad](#) [Why this ad?](#)



# Source details

## Frontiers in Pharmacology

Open Access i

Scopus coverage years: from 2010 to Present

Publisher: Frontiers Media S.A.

ISSN: 1663-9812

CiteScore 2021

**6.6**i

SJR 2021

**1.143**i

Subject area: Medicine: Pharmacology (medical) Pharmacology, Toxicology and Pharmaceutics: Pharmacology

Source type: Journal


SNIP 2021

**1.388**i[View all documents >](#)[Set document alert](#) [Save to source list](#) [Source Homepage](#)[CiteScore](#)[CiteScore rank & trend](#)[Scopus content coverage](#) [Export content for category](#)

### CiteScore rank i 2021

In category: Pharmacology (medical)

### CiteScore trend



#47  
255

Frontiers in Pharmacology

6.6

81st percentile

Rank	Source title	CiteScore 2021	Percentile
#1	Drug Resistance Updates	30.8	99th percentile
#2	Pharmacology and Therapeutics	22.2	99th percentile
#3	OpenNano	21.9	99th percentile
#4	Drugs	16.3	98th percentile
#5	International Journal of Antimicrobial Agents	16.1	98th percentile
#6	Journal of Biomedical Science	15.7	97th percentile
#7	Alimentary Pharmacology and Therapeutics	15.2	97th percentile
#8	American Journal of Transplantation	13.1	97th percentile
#9	Current Neuropharmacology	11.7	96th percentile
#10	Acta Pharmacologica Sinica	11.3	96th percentile
#11	Clinical Pharmacology and Therapeutics	10.7	95th percentile
#12	BioDrugs	10.5	95th percentile

Rank	Source title	CiteScore 2021	Percentile
★ #13	Clinical Pharmacokinetics	10.3	95th percentile
★ #14	Neurotherapeutics	10.0	94th percentile
★ #15	CNS Drugs	9.8	94th percentile
★ #16	npj Vaccines	9.4	93rd percentile
★ #17	Journal of Antimicrobial Chemotherapy	9.4	93rd percentile
★ #18	Antimicrobial Agents and Chemotherapy	9.3	93rd percentile
★ #19	International Journal of Neuropsychopharmacology	9.1	92nd percentile
★ #20	CNS Neuroscience and Therapeutics	8.9	92nd percentile
★ #21	Drug Metabolism Reviews	8.8	91st percentile
★ #22	Expert Opinion on Investigational Drugs	8.6	91st percentile
★ #23	Pharmacopsychiatry	8.5	91st percentile
★ #24	Rheumatology	8.5	90th percentile
★ #25	Antimicrobial Resistance and Infection Control	8.3	90th percentile
★ #26	Drug Safety	8.3	90th percentile
★ #27	Pharmacotherapy	8.2	89th percentile
★ #28	European Archives of Psychiatry and Clinical Neuroscience	7.9	89th percentile
★ #29	European Heart Journal - Cardiovascular Pharmacotherapy	7.8	88th percentile
★ #30	Cochrane Database of Systematic Reviews	7.6	88th percentile
★ #31	Advanced Therapeutics	7.5	88th percentile
★ #32	Inflammopharmacology	7.4	87th percentile
★ #33	Pharmaceutical Research	7.4	87th percentile
★ #34	Cannabis and Cannabinoid Research	7.3	86th percentile
★ #35	Targeted Oncology	7.2	86th percentile
★ #36	Current Treatment Options in Oncology	7.0	86th percentile
★ #37	OncoTargets and Therapy	7.0	85th percentile
★ #38	European Neuropsychopharmacology	6.9	85th percentile

Rank	Source title	CiteScore 2021	Percentile
☆ #39	International Journal for Parasitology: Drugs and Drug Resistance	6.9	84th percentile
☆ #40	Therapeutic Advances in Respiratory Disease	6.8	84th percentile
☆ #41	Expert Review of Clinical Pharmacology	6.8	84th percentile
☆ #42	Journal of Psychopharmacology	6.8	83rd percentile
☆ #43	Drugs and Aging	6.7	83rd percentile
☆ #44	Molecular Therapy - Oncolytics	6.7	82nd percentile
☆ #45	Expert Review of Anticancer Therapy	6.7	82nd percentile
☆ #46	Journal of Psychiatry and Neuroscience	6.6	82nd percentile
☆ #47	Frontiers in Pharmacology	6.6	81st percentile
☆ #48	Therapeutic Advances in Vaccines and Immunotherapy	6.5	81st percentile
☆ #49	npj Breast Cancer	6.5	80th percentile
☆ #50	Expert Review of Neurotherapeutics	6.5	80th percentile
☆ #51	British Journal of Clinical Pharmacology	6.3	80th percentile
☆ #52	Therapeutic Advances in Drug Safety	6.3	79th percentile
☆ #53	Investigational New Drugs	6.2	79th percentile
☆ #54	Expert Opinion on Drug Safety	6.1	79th percentile
☆ #55	Drug and Alcohol Dependence	6.1	78th percentile
☆ #56	European Review for Medical and Pharmacological Sciences	6.0	78th percentile
☆ #57	Recent Patents on Anti-Cancer Drug Discovery	5.9	77th percentile
☆ #58	World Journal of Men's Health	5.9	77th percentile
☆ #59	Breast Cancer	5.9	77th percentile
☆ #60	Cancer Chemotherapy and Pharmacology	5.9	76th percentile
☆ #61	American Journal of Cardiovascular Drugs	5.9	76th percentile
☆ #62	Medical Cannabis and Cannabinoids	5.8	75th percentile
☆ #63	Pulmonary Pharmacology and Therapeutics	5.8	75th percentile
☆ #64	Biologics: Targets and Therapy	5.7	75th percentile

Rank	Source title	CiteScore 2021	Percentile
☆ #65	Cardiovascular Drugs and Therapy	5.6	74th percentile
☆ #66	Expert Opinion on Emerging Drugs	5.6	74th percentile
☆ #67	Infection and Drug Resistance	5.6	73rd percentile
☆ #68	Expert Opinion on Pharmacotherapy	5.5	73rd percentile
☆ #69	Advances in Pharmacological and Pharmaceutical Sciences	5.5	73rd percentile
☆ #70	Paediatric Drugs	5.4	72nd percentile
☆ #71	Journal of acquired immune deficiency syndromes (1999)	5.3	72nd percentile
☆ #72	Clinical Therapeutics	5.2	71st percentile
☆ #73	Clinical Drug Investigation	5.2	71st percentile
☆ #73	Orphanet Journal of Rare Diseases	5.2	71st percentile
☆ #75	ACS Pharmacology and Translational Science	5.1	70th percentile
☆ #76	Journal of Aerosol Medicine and Pulmonary Drug Delivery	5.1	70th percentile
☆ #77	Antiviral Therapy	5.1	70th percentile
☆ #78	CPT: Pharmacometrics and Systems Pharmacology	5.0	69th percentile
☆ #79	Journal of Clinical Pharmacology	5.0	69th percentile
☆ #80	Advances in Therapy	5.0	68th percentile
☆ #81	Annals of Pharmacotherapy	5.0	68th percentile
☆ #82	Journal of Dietary Supplements	5.0	68th percentile
☆ #83	Therapeutic Advances in Cardiovascular Disease	4.9	67th percentile
☆ #84	Photodiagnosis and Photodynamic Therapy	4.9	67th percentile
☆ #85	PPAR Research	4.9	66th percentile
☆ #86	European Journal of Clinical Pharmacology	4.8	66th percentile
☆ #87	Vascular Health and Risk Management	4.8	66th percentile
☆ #88	Fundamental and Clinical Pharmacology	4.7	65th percentile
☆ #89	Journal of Addiction Medicine	4.7	65th percentile

Rank	Source title	CiteScore 2021	Percentile
☆ #90	Experimental and Clinical Psychopharmacology	4.7	64th percentile
☆ #91	Cardiovascular Therapeutics	4.6	64th percentile
☆ #92	Journal of Ocular Pharmacology and Therapeutics	4.6	64th percentile
☆ #93	Current Reviews in Clinical and Experimental Pharmacology	4.6	63rd percentile
☆ #94	Chemotherapy	4.5	63rd percentile
☆ #95	Vaccines	4.5	62nd percentile
☆ #96	Therapeutics and Clinical Risk Management	4.5	62nd percentile
☆ #97	HIV Medicine	4.5	62nd percentile
☆ #98	Drug Metabolism and Pharmacokinetics	4.5	61st percentile
☆ #99	Primary Care - Clinics in Office Practice	4.4	61st percentile
☆ #100	European Journal of Drug Metabolism and Pharmacokinetics	4.3	60th percentile
☆ #101	Drug Target Insights	4.3	60th percentile
☆ #102	BMC pharmacology & toxicology	4.2	60th percentile
☆ #103	Journal of Cardiovascular Pharmacology and Therapeutics	4.2	59th percentile
☆ #104	Journal of Child and Adolescent Psychopharmacology	4.1	59th percentile
☆ #105	Clinical Psychopharmacology and Neuroscience	4.1	59th percentile
☆ #106	Chinese Journal of Integrative Medicine	4.0	58th percentile
☆ #107	Pharmacoepidemiology and Drug Safety	4.0	58th percentile
☆ #108	Antibiotics	3.9	57th percentile
☆ #109	PharmaNutrition	3.9	57th percentile
☆ #110	Infection and Chemotherapy	3.9	57th percentile
☆ #111	Australian Prescriber	3.8	56th percentile
☆ #112	Cancer Drug Resistance	3.7	56th percentile
☆ #113	International Clinical Psychopharmacology	3.7	55th percentile
☆ #114	Journal of Cardiovascular Development and Disease	3.6	55th percentile

Rank	Source title	CiteScore 2021	Percentile
☆ #115	Clinical Pharmacology in Drug Development	3.6	55th percentile
☆ #116	Clinical Pharmacology: Advances and Applications	3.6	54th percentile
☆ #117	Iranian Journal of Pharmaceutical Research	3.5	54th percentile
☆ #118	Journal of Infection and Chemotherapy	3.5	53rd percentile
☆ #119	Journal of Clinical Psychopharmacology	3.4	53rd percentile
☆ #120	Trials	3.4	53rd percentile
☆ #121	Expert Review of Pharmacoeconomics and Outcomes Research	3.4	52nd percentile
☆ #122	Therapies	3.3	52nd percentile
☆ #123	International Journal of Clinical Pharmacy	3.3	51st percentile
☆ #124	ADMET and DMPK	3.2	51st percentile
☆ #125	Contemporary Clinical Trials	3.2	51st percentile
☆ #126	Biopharmaceutics and Drug Disposition	3.1	50th percentile
☆ #127	Journal of Chemotherapy	3.1	50th percentile
☆ #128	Journal of Clinical Pharmacy and Therapeutics	3.1	50th percentile
☆ #129	Advances in Clinical and Experimental Medicine	3.0	49th percentile
☆ #130	Journal of Pharmacy Practice	3.0	49th percentile
☆ #131	International Journal of Breast Cancer	3.0	48th percentile
☆ #132	Pharmaceutical Medicine	3.0	48th percentile
☆ #133	Synergy	3.0	48th percentile
☆ #134	PharmacoEconomics - Open	2.9	47th percentile
☆ #135	Current Drug Safety	2.9	47th percentile
☆ #136	Drug Metabolism and Personalized Therapy	2.9	46th percentile
☆ #137	Recent Advances in Anti-Infective Drug Discovery	2.8	46th percentile
☆ #138	Journal of Experimental Pharmacology	2.8	46th percentile
☆ #139	Human Psychopharmacology	2.8	45th percentile
☆ #140	HIV Research and Clinical Practice	2.7	45th percentile

Rank	Source title	CiteScore 2021	Percentile
☆ #141	Drug Metabolism Letters	2.7	44th percentile
☆ #142	Drugs - Real World Outcomes	2.7	44th percentile
☆ #143	Mental Health Clinician	2.7	44th percentile
☆ #144	Medicine in Drug Discovery	2.6	43rd percentile
☆ #145	American Journal of Therapeutics	2.6	43rd percentile
☆ #146	Cardiology	2.6	42nd percentile
☆ #147	European Journal of Contraception and Reproductive Health Care	2.6	42nd percentile
☆ #148	Neuropsychopharmacology Reports	2.5	42nd percentile
☆ #149	Current Therapeutic Research	2.5	41st percentile
☆ #150	AIDS Research and Therapy	2.5	41st percentile
☆ #151	Journal of Oncology Pharmacy Practice	2.4	40th percentile
☆ #152	Journal of Applied Pharmaceutical Science	2.3	40th percentile
☆ #153	International Journal of STD and AIDS	2.3	40th percentile
☆ #154	Journal of Pediatric Pharmacology and Therapeutics	2.3	39th percentile
☆ #155	Nanomedicine Research Journal	2.2	39th percentile
☆ #156	Therapeutic Innovation and Regulatory Science	2.2	39th percentile
☆ #157	EJNMMI Radiopharmacy and Chemistry	2.2	38th percentile
☆ #158	Pharmacia	2.1	38th percentile
☆ #159	Clinica e Investigacion en Arteriosclerosis	2.1	37th percentile
☆ #160	Pharmaceutical Statistics	2.0	37th percentile
☆ #161	Journal of Anaesthesiology Clinical Pharmacology	2.0	37th percentile
☆ #162	Research Results in Pharmacology	2.0	36th percentile
☆ #163	Translational and Clinical Pharmacology	2.0	36th percentile
☆ #164	Current Signal Transduction Therapy	1.9	35th percentile
☆ #165	Reading Teacher	1.9	35th percentile
☆ #166	Journal of Population Therapeutics and Clinical Pharmacology	1.8	35th percentile

Rank	Source title	CiteScore 2021	Percentile
☆ #167	Journal of Biopharmaceutical Statistics	1.8	34th percentile
☆ #168	Expert Opinion on Orphan Drugs	1.7	34th percentile
☆ #169	Journal of Nephropharmacology	1.5	33rd percentile
☆ #170	Journal of Cellular and Molecular Anesthesia	1.5	33rd percentile
☆ #171	Methods in Pharmacology and Toxicology	1.5	33rd percentile
☆ #172	Molecular and Integrative Toxicology	1.5	32nd percentile
☆ #173	Therapeutic Advances in Infectious Disease	1.5	32nd percentile
☆ #174	JACCP Journal of the American College of Clinical Pharmacy	1.4	31st percentile
☆ #175	Canadian Journal of Hospital Pharmacy	1.4	31st percentile
☆ #176	Journal of Pharmacy Practice and Research	1.3	31st percentile
☆ #177	Psychiatry and Clinical Psychopharmacology	1.7	32nd percentile
☆ #177	Psychopharmacology Bulletin	1.3	30th percentile
☆ #178	Journal of Pain and Palliative Care Pharmacotherapy	1.3	30th percentile
☆ #179	Research Results in Biomedicine	1.3	30th percentile
☆ #180	Research Journal of Pharmacy and Technology	1.3	29th percentile
☆ #181	Journal of Research in Pharmacy	1.3	29th percentile
☆ #182	Hospital Pharmacy	1.2	28th percentile
☆ #183	Infectio	1.2	28th percentile
☆ #184	Current Cardiovascular Risk Reports	1.2	28th percentile
☆ #185	Open Access Journal of Clinical Trials	1.1	27th percentile
☆ #186	International Journal of Cancer Management	1.0	27th percentile
☆ #187	Current Drug Therapy	0.9	26th percentile
☆ #188	Journal of Reports in Pharmaceutical Sciences	0.9	26th percentile
☆ #189	Tropical Journal of Pharmaceutical Research	0.9	26th percentile
☆ #190	Journal of Psychedelic Studies	0.9	25th percentile
☆ #191	Rational Pharmacotherapy in Cardiology	0.9	25th percentile

Rank	Source title	CiteScore 2021	Percentile
☆ #192	Zhongguo Zhongyao Zazhi	0.9	24th percentile
☆ #193	Archivos Venezolanos de Farmacología y Terapéutica	0.9	24th percentile
☆ #194	Drugs and Therapy Perspectives	0.9	24th percentile
☆ #195	Frontiers in Clinical Drug Research - Anti Infectives	0.8	23rd percentile
☆ #196	Medical Letter on Drugs and Therapeutics	0.8	23rd percentile
☆ #197	Indonesian Journal of Pharmacy	0.7	22nd percentile
☆ #198	Farmatsiya i Farmakologiya	0.7	22nd percentile
☆ #199	International Journal of Nutrition, Pharmacology, Neurological Diseases	0.7	22nd percentile
☆ #200	Journal of Pharmaceutical Health Care and Sciences	0.7	21st percentile
☆ #201	Chinese Herbal Medicines	0.7	21st percentile
☆ #202	Side Effects of Drugs Annual	0.6	20th percentile
☆ #203	Journal of Pharmacology and Pharmacotherapeutics	0.6	20th percentile
☆ #204	Iraqi Journal of Pharmaceutical Sciences	0.6	20th percentile
☆ #205	Drug and Therapeutics Bulletin	0.6	19th percentile
☆ #206	Pharmacien Hospitalier et Clinicien	0.6	19th percentile
☆ #207	Chinese Traditional and Herbal Drugs	0.5	19th percentile
☆ #208	Pharmaceutical Sciences Asia	0.5	18th percentile
☆ #209	Chinese Journal of Infection and Chemotherapy	0.5	18th percentile
☆ #210	Precision Cancer Medicine	0.5	17th percentile
☆ #211	Prescrire International	0.4	17th percentile
☆ #212	Current Topics in Pharmacology	0.4	17th percentile
☆ #213	Geneesmiddelenbulletin	0.4	16th percentile
☆ #214	Japanese Pharmacology and Therapeutics	0.4	16th percentile
☆ #215	Epitheorese Klinikes Farmakologias kai Farmakokinetikes	0.4	15th percentile

Rank	Source title	CiteScore 2021	Percentile
☆ #216	Review of Clinical Pharmacology and Pharmacokinetics, International Edition	0.4	15th percentile
☆ #216	Vojnosanitetski Pregled	0.4	15th percentile
☆ #218	Adverse Drug Reaction Bulletin	0.3	14th percentile
☆ #218	Biomedical and Biopharmaceutical Research	0.3	14th percentile
☆ #220	Annals of Cancer Research and Therapy	0.3	13th percentile
☆ #221	Cancer and Chemotherapy Reviews	0.3	13th percentile
☆ #222	Psychopharmakotherapie	0.3	13th percentile
☆ #223	Giornale Italiano di Farmacia Clinica	0.3	12th percentile
☆ #224	Drugs of the Future	0.2	12th percentile
☆ #225	Sovremennaya Revmatologiya	0.2	11th percentile
☆ #226	Krankenhauspharmazie	0.2	11th percentile
☆ #227	WHO Drug Information	0.2	11th percentile
☆ #228	Frontiers in Clinical Drug Research - Anti Allergy Agents	0.2	10th percentile
☆ #229	Chinese Journal of New Drugs	0.2	10th percentile
☆ #230	Actualites Pharmaceutiques	0.1	9th percentile
☆ #230	Opuholi Golovy i Sei	0.1	9th percentile
☆ #232	Anti-Obesity Drug Discovery and Development	0.1	9th percentile
☆ #233	Opuholi Zenskoj Reproduktivnoj Sistemy	0.1	8th percentile
☆ #234	Japanese Journal of Chemotherapy	0.1	8th percentile
☆ #235	Oral Therapeutics and Pharmacology	0.1	8th percentile
☆ #236	Adverse Drug Reactions Journal	0.1	7th percentile
☆ #237	Japanese Journal of Clinical Pharmacology and Therapeutics	0.1	7th percentile
☆ #238	Journal de Pharmacie Clinique	0.1	6th percentile
☆ #239	Medizinische Monatsschrift fur Pharmazeuten	0.1	6th percentile
☆ #240	Arzneimitteltherapie	0.1	6th percentile
☆ #241	European Journal of Clinical Pharmacy	0.1	5th percentile

Rank	Source title	CiteScore 2021	Percentile
☆ #242	Trends in Immunotherapy	0.1	5th percentile
☆ #243	Journal of Arrhythmology	0.1	4th percentile
☆ #244	International Journal of Pharmaceutical Sciences and Nanotechnology	0.0	4th percentile
☆ #245	Journal fur Pharmakologie und Therapie	0.0	4th percentile
☆ #246	Drugs and Clinic	0.0	3rd percentile
☆ #247	Circular Farmaceutica	0.0	3rd percentile
☆ #248	Deutsche Apotheker Zeitung	0.0	2nd percentile
☆ #249	Pharmaceutisch Weekblad	0.0	2nd percentile
☆ #250	Pharmazeutische Zeitung	0.0	2nd percentile
☆ #251	Frontiers in Clinical Drug Research - CNS and Neurological Disorders	0.0	0th percentile
☆ #251	Frontiers in Clinical Drug Research - HIV	0.0	0th percentile
☆ #251	Health Biotechnology and Biopharma	0.0	0th percentile
☆ #251	Kliniceskaa Mikrobiologa i Antimikrobnaa Himioterapia	0.0	0th percentile
☆ #251	Revista Colombiana de Ciencias Quimico-Farmaceuticas(Colombia)	0.0	0th percentile

## About Scopus

[What is Scopus](#)

[Content coverage](#)

[Scopus blog](#)

[Scopus API](#)

[Privacy matters](#)

## Language

[日本語版を表示する](#)

[查看简体中文版本](#)

[查看繁體中文版本](#)

[Просмотр версии на русском языке](#)

## Customer Service

[Help](#)

[Tutorials](#)

[Contact us](#)

---

## ELSEVIER

[Terms and conditions](#) ↗ [Privacy policy](#) ↗

Copyright © Elsevier B.V. ↗. All rights reserved. Scopus® is a registered trademark of Elsevier B.V.

We use cookies to help provide and enhance our service and tailor content. By continuing, you agree to the use of cookies ↗.

