

E-ISSN: 1309-0001

www.namaraiphamj.com

Volume 22 + Issue 3 + July 2018

JRP

Journal of
Research in Pharmacy

AN INTERNATIONAL OPEN ACCESS JOURNAL OF
PHARMACY AND PHARMACEUTICAL SCIENCES

Vol. 22 (3)
2018

Editor in Chief

Dray Kizilgizel

Associate Editor

Ewa Talar

Section Editors

Ali Deniz Sözer

Bülent Kaymakçalan

Orhan Tamer

Hüseyin Karaman

Özge Karataş

Turgut Şişir

Technical Editor

Ayhan Özdemir

Ekin Başer

Period

Quarterly

Publisher

Namara University Press

Formerly published as Namara Pharmaceutical Journal

Journal of Research in Pharmacy

An international open-access journal of pharmacy and pharmaceutical sciences
Formerly published as Marmara Pharmaceutical Journal

(<https://jrespharm.com/>)

JRP

ISSN : 1309-0801



[Current Issue \(https://jrespharm.com/content.php?id=78\)](https://jrespharm.com/content.php?id=78)

[Articles in Press \(https://jrespharm.com/content.php?id=46\)](https://jrespharm.com/content.php?id=46)

[Archive \(https://jrespharm.com/archive.php\)](https://jrespharm.com/archive.php)

[Search \(https://jrespharm.com/search.php\)](https://jrespharm.com/search.php)

[Author Benefits \(https://jrespharm.com/static.php?id=11\)](https://jrespharm.com/static.php?id=11)

Editor-in-Chief

Hatice Kübra Elçioğlu

Vice Editors

Levent Kabasakal

Esra Tatar

Online ISSN

2630-6344

Publisher

Marmara University

Frequency

Bimonthly (Six issues / year)

Abbreviation

J.Res.Pharm.

Former Name

Marmara Pharmaceutical Journal

Editorial Board

Editor-in-Chief

Hatice Kübra ELÇİOĞLU

Department of Pharmacology, Faculty of Pharmacy, Marmara University, Istanbul, Turkey
kubra.elcioglu@marmara.edu.tr (mailto:kubra.elcioglu@marmara.edu.tr)

Vice Editors

Levent KABASAKAL

Department of Pharmacology, Faculty of Pharmacy, Marmara University, Istanbul, Turkey
lkabasakal@marmara.edu.tr (mailto:lkabasakal@marmara.edu.tr)

Esra TATAR

Department of Pharmaceutical Chemistry, Faculty of Pharmacy, Marmara University, Istanbul, Turkey
etatar@marmara.edu.tr (mailto:etatar@marmara.edu.tr)

Guest Editors for the Special Issue : The International Conference on Pharmaceutical Research and Practice (ICPRP) 2022

Arba Pramundita RAMADANI

Department of Pharmacy, Universitas Islam Indonesia, Yogyakarta, Indonesia
arba.pramundita@uii.ac.id (mailto:arba.pramundita@uii.ac.id)

Asih TRIASTUTI

Department of Pharmacy, Universitas Islam Indonesia, Yogyakarta, Indonesia
asih.triastuti@uii.ac.id (mailto:asih.triastuti@uii.ac.id)

Analytical Chemistry & Therapeutic Drug Monitoring

Anisa ELHAMILI

Department of Medicinal & Pharmaceutical Chemistry, Faculty of Pharmacy, University of Tripoli, Tripoli, Libya
aaelhamili2000@gmail.com (mailto:aaelhamili2000@gmail.com)

Lorena MEMUSHAJ

Department of Pharmacy, Faculty of Medical Sciences, Aldent University, Tirana, Albania
lorena.memushaj@ual.edu.al (mailto:lorena.memushaj@ual.edu.al)

Mohd Younis RATHER

Multidisciplinary Research Unit, Government Medical College Srinagar, Srinagar, India
younis.rather78@gmail.com (mailto:younis.rather78@gmail.com)

Pablo MIRALLES IBARRA

1. Foundation for the Promotion of Health and Biomedical Research in the Valencian Region (FISABIO-Public Health), Valencia, Spain
2. Department of Analytical Chemistry, Faculty of Chemistry, University of Valencia, Burjassot, Spain miralles_pabiba@gva.es (mailto:miralles_pabiba@gva.es)

Pınar TALAY PINAR

Department of Analytical Chemistry, Faculty of Pharmacy, Yüzüncü Yıl University, Van, Turkey
ptalay@gmail.com (mailto:ptalay@gmail.com)

Biochemistry & Cancer Research

Derya ÖZSAVCI

Department of Biochemistry, Faculty of Pharmacy, Marmara University, Istanbul, Turkey
derya.ozsavci@marmara.edu.tr (mailto:derya.ozsavci@marmara.edu.tr)

Gülberk UÇAR

Department of Biochemistry, Faculty of Pharmacy, Hacettepe University, Ankara, Turkey
gulberk@hacettepe.edu.tr (mailto:gulberk@hacettepe.edu.tr)

Lokman AYAZ

Department of Biochemistry, Faculty of Pharmacy, Trakya University, Edirne, Turkey
lokmanayaz@yahoo.com (mailto:lokmanayaz@yahoo.com)

Biotechnology

Ali Demir SEZER

Department of Pharmaceutical Biotechnology, Faculty of Pharmacy, Marmara University, Istanbul, Turkey
adsezer@marmara.edu.tr (mailto:adsezer@marmara.edu.tr)

Ammad Ahmad FAROOQI

Department of Molecular Oncology, Institute of Biomedical and Genetic Engineering (IBGE), Islamabad, Pakistan
farooqiamdahmad@gmail.com (mailto:farooqiamdahmad@gmail.com)

Murat DOĞAN

Department of Pharmaceutical Biotechnology, Faculty of Pharmacy, Cumhuriyet University, Sivas, Turkey
mdogan@cumhuriyet.edu.tr (mailto:mdogan@cumhuriyet.edu.tr)

Uğur KARAGÖZ

Department of Pharmaceutical Biotechnology, Faculty of Pharmacy, Trakya University, Edirne, Turkey
ugurkaragoz@trakya.edu.tr (mailto:ugurkaragoz@trakya.edu.tr)

Clinical Pharmacy & Social Pharmacy

Abdikarim Mohammed ABDI

Department of Clinical Pharmacy, Faculty of Pharmacy, Yeditepe University, Istanbul, Turkey
abdikarim.abdi@yeditepe.edu.tr (mailto:abdikarim.abdi@yeditepe.edu.tr)

Betül OKUYAN

Department of Clinical Pharmacy, Faculty of Pharmacy, Marmara University, Istanbul, Turkey
betul.okuyan@marmara.edu.tr (mailto:betul.okuyan@marmara.edu.tr)

Maja ORTNER HADŽIABDIĆ

Centre for Applied Pharmacy, Faculty of Pharmacy and Biochemistry, University of Zagreb, Zagreb, Croatia
mortner@pharma.hr (mailto:mortner@pharma.hr)

Mesut SANCAR

Department of Pharmacology, Faculty of Pharmacy, Marmara University, Istanbul, Turkey
mesut.sancar@marmara.edu.tr (mailto:mesut.sancar@marmara.edu.tr)

Tarik CATIĆ

Department of Pharmacy, Sarajevo School of Science and Technology, Sarajevo, Bosnia and Herzegovina
tarik.catic@ssst.edu.ba (mailto:tarik.catic@ssst.edu.ba)

In Silico Studies

Gizem TATAR YILMAZ

Department of Biostatistics and Medical Informatics, Faculty of Medicine, Karadeniz Technical University, Trabzon, Turkey
gizemtatar@gmail.com (mailto:gizemtatar@gmail.com)

Medicinal Chemistry

Bahadır BÜLBÜL

Department of Pharmaceutical Chemistry, Faculty of Pharmacy, Düzce University, Düzce, Turkey
bahadir.bulbul@yahoo.com.tr (mailto:bahadir.bulbul@yahoo.com.tr)

Entela HALOCI

Faculty of Pharmacy, University of Medicine, Tirana, Albania
entela.haloci@umed.edu.al (mailto:entela.haloci@umed.edu.al)

Hasan Erdinç SELLİTEPE

Department of Pharmaceutical Chemistry, Faculty of Pharmacy, Karadeniz Technical University, Trabzon, Turkey
esellitepe@ktu.edu.tr (mailto:esellitepe@ktu.edu.tr)

Somaieh SOLTANI

Faculty of Pharmacy, Tabriz University of Medical Sciences, Tabriz, Iran
soltanis@tbzmed.ac.ir (mailto:soltanis@tbzmed.ac.ir)

*Microbiology & Immunology***Erkan RAYAMAN**

Department of Pharmaceutical Microbiology, Faculty of Pharmacy, Marmara University, Istanbul, Turkey
erayaman@marmara.edu.tr (mailto:erayaman@marmara.edu.tr)

Gülgün TINAZ

Department of Basic Pharmaceutical Sciences, Faculty of Pharmacy, Marmara University, Istanbul, Turkey
gulgun.tinaz@marmara.edu.tr (mailto:gulgun.tinaz@marmara.edu.tr)

Zahraa AMER HASHIM

Department of Microbiology and Immunology, College of Pharmacy, Mosul University, Mosul, Iraq
hashimz@uomosul.edu.iq (mailto:hashimz@uomosul.edu.iq)

*Pharmaceutical Botany & Pharmacognosy & Chemistry of Natural Products***Ahmet EMİR**

Department of Pharmacognosy, Faculty of Pharmacy, Ege University, Izmir, Turkey
ahmet.emir@ege.edu.tr (mailto:ahmet.emir@ege.edu.tr)

Annalisa CHIAVAROLI

Department of Pharmacology, Faculty of Pharmacy, G. d'Annunzio University of Chieti-Pescara, Chieti, Italy
annalisa.chiavaroli@unich.it (mailto:annalisa.chiavaroli@unich.it)

Antoaneta TRENDAFILOVA

Institute of Organic Chemistry with Centre of Phytochemistry, Bulgarian Academy of Sciences, Sofia, Bulgaria
antoaneta.trendafilova@orgchm.bas.bg (mailto:antoaneta.trendafilova@orgchm.bas.bg)

Ayşe Esra KARADAĞ

Department of Pharmacognosy, Faculty of Pharmacy, Istanbul Medipol University, Istanbul, Turkey
ayseesraguler@gmail.com (mailto:ayseesraguler@gmail.com)

Ceren EMİR

Department of Pharmacognosy, Faculty of Pharmacy, Ege University, Izmir, Turkey
ceren.acir@ege.edu.tr (mailto:ceren.acir@ege.edu.tr)

Claudio FERRANTE

Department of Pharmacology, Faculty of Pharmacy, G. d'Annunzio University of Chieti-Pescara, Chieti, Italy
claudio.ferrante@unich.it (mailto:claudio.ferrante@unich.it)

İ. İrem TATLI ÇANKAYA

Department of Pharmaceutical Botany, Faculty of Pharmacy, Hacettepe University, Ankara, Turkey
iremcanakaya@gmail.com (mailto:iremcanakaya@gmail.com)

Lejla KLEPO

Department of Chemistry, Faculty of Science, University of Sarajevo, Sarajevo, Bosnia and Herzegovina
klepolejla@gmail.com (mailto:klepolejla@gmail.com)

Nurettin YAYLI

Department of Pharmacognosy, Faculty of Pharmacy, Karadeniz Technical University, Trabzon, Turkey
yayli@ktu.edu.tr (mailto:yayli@ktu.edu.tr)

*Pharmacognosy***Randolph ARROO**

Leicester School of Pharmacy, De Montfort University, Leicester, UK
rrjarroo@dmu.ac.uk (mailto:rrjarroo@dmu.ac.uk)

Turgut TAŞKIN

Department of Pharmacognosy, Faculty of Pharmacy, Marmara University, Istanbul, Turkey
turguttaskin@marmara.edu.tr (mailto:turguttaskin@marmara.edu.tr)

Vildan ÇELİKSOY

School of Optometry and Vision Sciences, Cardiff University, Cardiff, UK
celiksoyv@cardiff.ac.uk (mailto:celiksoyv@cardiff.ac.uk)

Zoran ZEKOVIĆ

Faculty of Technology, University of Novi Sad, Novi Sad, Serbia
zzekovic@tf.uns.ac.rs (mailto:zzekovic@tf.uns.ac.rs)

*Pharmaceutics***Afife Büşra UĞUR KAPLAN**

Department of Pharmaceutical Technology, Faculty of Pharmacy, Atatürk University, Erzurum, Turkey
afife.busra.ugur@gmail.com (mailto:afife.busra.ugur@gmail.com)

Dinesh KUMAR

Department of Pharmaceutical Engineering & Technology, Indian Institute of Technology (BHU), Varanasi, India
dinesh.phe@itbhu.ac.in (mailto:dinesh.phe@itbhu.ac.in)

Ebru ALTUNTAŞ

Department of Pharmaceutical Technology, Faculty of Pharmacy, Istanbul University, Istanbul, Turkey
ebru.altuntas@istanbul.edu.tr (mailto:ebru.altuntas@istanbul.edu.tr)

Enkelejda GOCI

Pharmacotherapeutic Research Center, Aldent University, Tirana, Albania
enkelejda.goci@ual.edu.al (mailto:enkelejda.goci@ual.edu.al)

Gülşah GEDİK

Department of Pharmaceutical Technology, Faculty of Pharmacy, Trakya University, Edirne, Turkey
gulsahgedik@trakya.edu.tr (mailto:gulsahgedik@trakya.edu.tr)

Ongun Mehmet SAKA

Department of Pharmaceutical Technology and Biotechnology, Faculty of Pharmacy, Ankara University, Ankara, Turkey
omsaka@gmail.com (mailto:omsaka@gmail.com)

Oya KERİMOĞLU

Department of Pharmaceutical Technology, Faculty of Pharmacy, Marmara University, Istanbul, Turkey
osipahigil@marmara.edu.tr (mailto:osipahigil@marmara.edu.tr)

Rezarta SHKRELI

Department of Pharmacy, Faculty of Medical Sciences, Aldent University, Tirana, Albania
rezarta.shkreli@ual.edu.al (mailto:rezarta.shkreli@ual.edu.al)

Saeideh SOLTANI

Novel Drug Research Center, School of Pharmacy and Pharmaceutical Sciences, Isfahan University of Medical Sciences, Isfahan, Iran
soltanisa@pharm.mui.ac.ir (mailto:soltanisa@pharm.mui.ac.ir)

*Pharmacology & Toxicology***Ana V. PEJČIĆ**

Department of Pharmacology and Toxicology, Faculty of Medical Sciences, University of Kragujevac, Kragujevac, Serbia
anapejic201502@yahoo.com (mailto:anapejic201502@yahoo.com)

Ayfer BECEREN

Department of Pharmaceutical Toxicology, Faculty of Pharmacy, Marmara University, Istanbul, Turkey
ayfer.tozan@marmara.edu.tr (mailto:ayfer.tozan@marmara.edu.tr)

Büşra ERTAŞ

Department of Pharmacology, Faculty of Pharmacy, Marmara University, Istanbul, Turkey
busra.ertas@marmara.edu.tr (mailto:ayfer.tozan@marmara.edu.tr)

Klodiola DHAMO

Faculty of Technical Medical Sciences, Aldent University, Tirana, Albania
klodiola.dhamo@ual.edu.al (mailto:klodiola.dhamo@ual.edu.al)

Nasrin MALEKI DIZAJI

Faculty of Pharmacy, Tabriz University of Medical Sciences, Tabriz, Iran
malekins2014@gmail.com (mailto:malekins2014@gmail.com)

Rümeysa KELEŞ KAYA

Department of Medical Pharmacology, Faculty of Medicine, Sakarya University, Sakarya, Turkey
rumeysakeles@sakarya.edu.tr (mailto:rumeysakeles@sakarya.edu.tr)

Ünzile YAMAN

Department of Pharmaceutical Toxicology, Faculty of Pharmacy, Katip Çelebi University, İzmir, Turkey
unzileyaman@gmail.com (mailto:unzileyaman@gmail.com)

Zeina ALTHANOON

Department of Pharmacology and Toxicology, College of Pharmacy, Mosul University, Mosul, Iraq
dr.zeina@uomosul.edu.iq (mailto:dr.zeina@uomosul.edu.iq)

Rümeysa Keleş Kaya

Department of Medical Pharmacology, Faculty of Medicine, Sakarya University, Sakarya, Turkey
rumeysakeles@sakarya.edu.tr (mailto:rumeysakeles@sakarya.edu.tr)

*Copy Editor***Ayşe Nur HAZAR YAVUZ**

Department of Pharmacology, Faculty of Pharmacy, Marmara University, Istanbul, Turkey
ayse.hazar@marmara.edu.tr (mailto:ayse.hazar@marmara.edu.tr)

Büşra ERGEN

Department of Clinical Pharmacy, Faculty of Pharmacy, Marmara University, Istanbul, Turkey
busra.ergen@marmara.edu.tr (mailto:busra.ergen@marmara.edu.tr)

Fatih Taha ÇİFTÇİ

Department of Clinical Pharmacy, Faculty of Pharmacy, Marmara University, Istanbul, Turkey
fatih.ciftci@marmara.edu.tr (mailto:fatih.ciftci@marmara.edu.tr)

Müzeyyen AKSOY

Department of Clinical Pharmacy, Faculty of Pharmacy, Marmara University, Istanbul, Turkey
muzeyyen.aksoy@marmara.edu.tr (mailto:muzeyyen.aksoy@marmara.edu.tr)

Ömer Faruk ÖZKANLI

Department of Clinical Pharmacy, Faculty of Pharmacy, Marmara University, Istanbul, Turkey
oozkanli@marmara.edu.tr (mailto:oozkanli@marmara.edu.tr)

Şeyma GÖZELİZMİR

Department of Clinical Pharmacy, Faculty of Pharmacy, Marmara University, Istanbul, Turkey
seyma.gozelizmir@marmara.edu.tr (mailto:seyma.gozelizmir@marmara.edu.tr)

Language Editor

Khadija ALJESRI

Department of Pharmacology, Institute of Health Sciences, Marmara University, Istanbul, Turkey

Biostatistics Editor

Gülnaz NURAL BEKİROĞLU

Department of Biostatistics, Faculty of Medicine, Marmara University, Istanbul, Turkey
nural@marmara.edu.tr (mailto:nural@marmara.edu.tr)

Marmara University

[Home \(https://jrespharm.com/\)](https://jrespharm.com/)

[Aims and Scope \(static.php?id=1\)](#)

[Editorial Board \(static.php?id=2\)](#)

[Advisory Board \(content.php?id=78\)](#)

[Table of Contents \(static.php?id=4\)](#)

[Author Guidelines \(static.php?id=7\)](#)

[Abstracting / Indexing \(static.php?id=9\)](#)

[Contact \(static.php?id=8\)](#)

Journal of Research in Pharmacy

An international open-access journal of pharmacy and pharmaceutical sciences
Formerly published as Marmara Pharmaceutical Journal

(<https://jrespharm.com/>)

JRP

ISSN : 2630-6344



[Current Issue \(https://jrespharm.com/content.php?id=78\)](https://jrespharm.com/content.php?id=78)

[Articles in Press \(https://jrespharm.com/content.php?id=46\)](https://jrespharm.com/content.php?id=46)

[Archive \(https://jrespharm.com/archive.php\)](https://jrespharm.com/archive.php)

[Search \(https://jrespharm.com/search.php\)](https://jrespharm.com/search.php)

[Author Benefits \(https://jrespharm.com/static.php?id=11\)](https://jrespharm.com/static.php?id=11)

Editor-in-Chief

Hatice Kübra Elçioğlu

Vice Editors

Levent Kabasakal

Esra Tatar

Online ISSN

2630-6344

Publisher

Marmara University

Frequency

Bimonthly (Six issues / year)

Abbreviation

J.Res.Pharm.

Former Name

Marmara Pharmaceutical Journal

Other Issues :

[◀ Previous \(content.php?id=76\)](https://jrespharm.com/content.php?id=76)

[Next ▶ \(content.php?id=78\)](https://jrespharm.com/content.php?id=78)

[Issue XML \(issue-xml.php?id=77\)](https://jrespharm.com/issue-xml.php?id=77)

Journal of Research in Pharmacy

2022 , Vol 26 , Issue 4

➔ Review Article (1)

➔ Research Article (32)

Review Article

[Clear](#) [View Selected Abstracts](#)

Reporting quality of animal studies published in journals listed in ULAKBIM TR index: a systematic review on compliance to the ARRIVE guidelines (abstract.php?id=1055) *Review Article*

Pages 704-713

Selda EMRE AYDINGÖZ,Oğuzhan Ekin EFE,Gökçe ÇALIŞKAN

DOI : 10.29228/jrp.168

[Abstract \(abstract.php?lang=en&id=1055\)](#)

[Full Text PDF \(pdf.php?&id=1055\)](#)

[Similar Articles \(similar.php?&id=1055\)](#)

[E-mail to Author \(mailto:ecz.gokcecaliskan@gmail.com\)](mailto:ecz.gokcecaliskan@gmail.com)

[How to Cite \(ajax_handler.php?id=1055\)](#)

Research Article

[Clear](#) [View Selected Abstracts](#)

Forty Micromole Hydroxychloroquine Enhanced Cytotoxic Effect of Doxorubicin Against Laryngeal Cancer Cell Line HEp-2 (abstract.php?id=1056) *Research Article*

Pages 714-721

Hasanain Kamil OWADH,Rana A. GHALEB ,Fatima Adnan ALZUBAIDI

DOI : 10.29228/jrp.169

[Abstract \(abstract.php?lang=en&id=1056\)](#)

[Full Text PDF \(pdf.php?&id=1056\)](#)

[Similar Articles \(similar.php?&id=1056\)](#)

[E-mail to Author \(mailto:hasanain@mustaqbal-college.edu.iq\)](mailto:hasanain@mustaqbal-college.edu.iq)

[How to Cite \(ajax_handler.php?id=1056\)](#)

Investigation of the pharmacological potential of myricetin on alcohol addiction in mice (abstract.php?id=1057)

Research Article

Pages 722-733

Oruç YUNUSOĞLU, Andleeb SHAHZADI, Canan AKÜNAL TÜREL, Muhammed Hamdi DEMIRKOL, Mehmet BERKÖZ, Ahmet Gökhan AKKAN

DOI : 10.29228/jrp.170

[Abstract \(abstract.php?lang=en&id=1057\)](#)

[Full Text PDF \(pdf.php?&id=1057\)](#)

[Similar Articles \(similar.php?&id=1057\)](#)

[E-mail to Author \(mailto:orucfarm@gmail.com\)](mailto:orucfarm@gmail.com)

[How to Cite \(ajax_handler.php?id=1057\)](#)

Evaluation of Anti-Inflammatory and Gastroprotective Activity of Aqueous Peel Extract of Allium Cepa (abstract.php?id=1058) *Research Article*

Pages 734-741

Rashidat AYANNIYI, Hidayah OLUMOH-ABDUL, Fatimoh OJUJADE, Juliana AKINTOLA

DOI : 10.29228/jrp.171

[Abstract \(abstract.php?lang=en&id=1058\)](#)

[Full Text PDF \(pdf.php?&id=1058\)](#)

[Similar Articles \(similar.php?&id=1058\)](#)

[E-mail to Author \(mailto:ayanniyi.ro@unilorin.edu.ng\)](mailto:ayanniyi.ro@unilorin.edu.ng)

[How to Cite \(ajax_handler.php?id=1058\)](#)

Combined ethanol extract of Hypselodelphys poggeana and Spermacoce radiata leaves ameliorate benign prostatic hyperplasia in rats via modulation of serum sex hormonal levels (abstract.php?id=1059) *Research Article*

Pages 742-752

ROBERT IKECHUKWU UROKO, SOLOMON NNAH IJIOMA, ELISHA UKO OGWO, FAVOUR MATTHEW AWAH

DOI : 10.29228/jrp.201

[Abstract \(abstract.php?lang=en&id=1059\)](#)

[Full Text PDF \(pdf.php?&id=1059\)](#)

[Similar Articles \(similar.php?&id=1059\)](#)

[E-mail to Author \(mailto:ir.uoko@mouau.edu.ng\)](mailto:ir.uoko@mouau.edu.ng)

[How to Cite \(ajax_handler.php?id=1059\)](#)

Euryale ferox Salisb ameliorates arthritis in experimental animals (abstract.php?id=1060) *Research Article*

Pages 753-769

Shradhanjali SINGH,Varsha CHAUDHARY,Shanti Bhushan MISHRA,Anil Kumar SINGH

DOI : 10.29228/jrp.173

[Abstract \(abstract.php?lang=en&id=1060\)](#)

[Full Text PDF \(pdf.php?&id=1060\)](#)

[Similar Articles \(similar.php?&id=1060\)](#)

[E-mail to Author \(mailto:singhshradhanjali@gmail.com\)](mailto:singhshradhanjali@gmail.com)

[How to Cite \(ajax_handler.php?id=1060\)](#)

Methanolic extract of Eryngium creticum Lam leaves, flowers and roots: Quantification and qualification of phenolic contents, antitumor effect and antioxidant capacity (abstract.php?id=1061) *Research Article*

Pages 770-780

Shreen NUSAIR,Mohammad AHMAD,Tamam EL-ELIMAT,Khansaa AL-ESSA,Luay ABU-QATOUSEH,Rawand KHASAWNEH

DOI : 10.29228/jrp.174

[Abstract \(abstract.php?lang=en&id=1061\)](#)

[Full Text PDF \(pdf.php?&id=1061\)](#)

[Similar Articles \(similar.php?&id=1061\)](#)

[E-mail to Author \(mailto:sdnusair@just.edu.jo\)](mailto:sdnusair@just.edu.jo)

[How to Cite \(ajax_handler.php?id=1061\)](#)

Chemical composition and in vitro mutagenic, antioxidant and anti-inflammatory activities of Lavandula angustifolia Mill. essential oil from Turkey (abstract.php?id=1062) *Research Article*

Pages 781-789

Pınar Sinem OMURTAG ÖZGEN, Sevde Nur BİLTEKİN, Tuğba İDUĞ, Çağlar MACİT, İrem AYRAN, Sadiye Ayşe ÇELİK, Şükran KÜLTÜR, Yüksel KAN, Gülden Zehra OMURTAG

DOI : 10.29228/jrp.175

[Abstract \(abstract.php?lang=en&id=1062\)](#)

[Full Text PDF \(pdf.php?&id=1062\)](#)

[Similar Articles \(similar.php?&id=1062\)](#)

[E-mail to Author \(mailto:tugbaidug@gmail.com\)](mailto:tugbaidug@gmail.com)

[How to Cite \(ajax_handler.php?id=1062\)](#)

Antioxidant, Antityrosinase Activities and Composition of Essential Oils Obtained from Roots and Fruits of *Ferulago longistylis* Boiss (abstract.php?id=1063) Research Article

Pages 790-798

Mine KÜRKÇÜOĞLU, Hale Gamze AĞALAR, Burak TEMİZ, Ahmet DURAN, Mehmet Yavuz PAKSOY, Kemal Hüsnü Can BAŞER

DOI : 10.29228/jrp.176

[Abstract \(abstract.php?lang=en&id=1063\)](#)

[Full Text PDF \(pdf.php?&id=1063\)](#)

[Similar Articles \(similar.php?&id=1063\)](#)

[E-mail to Author \(mailto:mkurkcuo@anadolu.edu.tr\)](mailto:mkurkcuo@anadolu.edu.tr)

[How to Cite \(ajax_handler.php?id=1063\)](#)

Phenolic profile, antioxidant and anticandidal activities of *Inula oculus-christi* L. from Turkey (abstract.php?id=1064) Research Article

Pages 799-808

Nagehan SALTAN, Yavuz Bülent KÖSE, Fatih GÖGER, Gökalp İŞCAN

DOI : 10.29228/jrp.177

[Abstract \(abstract.php?lang=en&id=1064\)](#)

[Full Text PDF \(pdf.php?&id=1064\)](#)

[Similar Articles \(similar.php?&id=1064\)](#)

E-mail to Author (<mailto:ndagdeviren@anadolu.edu.tr>)

How to Cite ([ajax_handler.php?id=1064](#))

Comparison of Commercial *Calendula officinalis* L. Samples with Pharmacopeial Drug: Antiradical Activities and Chemical Profiles ([abstract.php?id=1065](#)) *Research Article*

Pages 809-819

Reza Chamansara,Romina Rashidfarokhi,Ezgi Ak Sakallı,Nesrin Öztinen,Müberra Koşar

DOI : 10.29228/jrp.178

Abstract ([abstract.php?lang=en&id=1065](#))

Full Text PDF ([pdf.php?&id=1065](#))

Similar Articles ([similar.php?&id=1065](#))

E-mail to Author (<mailto:muberra.kosar@emu.edu.tr>)

How to Cite ([ajax_handler.php?id=1065](#))

***Ammi visnaga* L. and Nanocarrier Approaches in the Treatment of Skin Diseases ([abstract.php?id=1066](#))**

Research Article

Pages 820-827

İmren ESENTÜRK-GÜZEL,Sehernaz TOPUZOĞLU,Lüceyn ABDO,Eda SÖNMEZ GÜRER,Evren ALGIN YAPAR

DOI : 10.29228/jrp.181

Abstract ([abstract.php?lang=en&id=1066](#))

Full Text PDF ([pdf.php?&id=1066](#))

Similar Articles ([similar.php?&id=1066](#))

E-mail to Author (<mailto:imrenesenturk@gmail.com>)

How to Cite ([ajax_handler.php?id=1066](#))

Formulation and evaluation of herbal antioxidant face cream using extract of the marine seaweed *Sargassum* spp. (Phaeophyceae) ([abstract.php?id=1067](#)) *Research Article*

Pages 828-833

DOI : 10.29228/jrp.180

Abstract (abstract.php?lang=en&id=1067)
Full Text PDF (pdf.php?&id=1067)
Similar Articles (similar.php?&id=1067)
E-mail to Author (mailto:z.arabeyyat@ju.edu.jo)
How to Cite (ajax_handler.php?id=1067)

Preliminary research on ibuprofen self-emulsifying formulation (abstract.php?id=1068) *Research Article*

Pages 834-841

Zwanden Sule YAHAYA,Fatima Shuaibu KURFI,Danjuma MALLAM,Rukayyat Bukola OLOYEDE,Olutayo Ademola ADELEYE,Gideon Owoicho OKPANACHI

DOI : 10.29228/jrp.182

Abstract (abstract.php?lang=en&id=1068)
Full Text PDF (pdf.php?&id=1068)
Similar Articles (similar.php?&id=1068)
E-mail to Author (mailto:pharmzwanzu@gmail.com)
How to Cite (ajax_handler.php?id=1068)

Development of bergamot flavors according to their enantiomeric state (abstract.php?id=1069) *Research Article*

Pages 842-848

Gulsah GEDIK,Nagihan GULHAN,Ralf ELHADEF,Yasemin KUTUK,Gozde GOKDENIZ,M. Onur YAKAK,Hilal EMIRLE,M. Fatih ALBAYRAK

DOI : 10.29228/jrp.183

Abstract (abstract.php?lang=en&id=1069)
Full Text PDF (pdf.php?&id=1069)
Similar Articles (similar.php?&id=1069)
E-mail to Author (mailto:gulsahgedik@gmail.com)

[How to Cite \(ajax_handler.php?id=1069\)](#)

Intravaginal delivery of reverse micellar epigallocatechin loaded in κ -carrageenan and HPMC K100M-based gel (abstract.php?id=1070) *Research Article*

Pages 849-858

Noorma ROSITA, Dewi Melani HARIYADI, Cordellia Calista AMELIA, Alyajilan Madani MAMURRIE, Andang MIATMOKO

DOI : 10.29228/jrp.184

[Abstract \(abstract.php?lang=en&id=1070\)](#)

[Full Text PDF \(pdf.php?&id=1070\)](#)

[Similar Articles \(similar.php?&id=1070\)](#)

[E-mail to Author \(mailto:noorma-r@ff.unair.ac.id\)](mailto:noorma-r@ff.unair.ac.id)

[How to Cite \(ajax_handler.php?id=1070\)](#)

Development of novel formulation technology for oral delivery of Sterculia Gum (abstract.php?id=1071) *Research Article*

Pages 859-872

Harshad KAPARE, Sohan CHITLANGE, Kaveri AHER, Shivaji AHER

DOI : 10.29228/jrp.185

[Abstract \(abstract.php?lang=en&id=1071\)](#)

[Full Text PDF \(pdf.php?&id=1071\)](#)

[Similar Articles \(similar.php?&id=1071\)](#)

[E-mail to Author \(mailto:hskapare@yahoo.in\)](mailto:hskapare@yahoo.in)

[How to Cite \(ajax_handler.php?id=1071\)](#)

In Situ Gels Triggered by Temperature for Ocular Delivery of Dexamethasone and Dexamethasone/ SBE- β -CD Complex (abstract.php?id=1072) *Research Article*

Pages 873-883

H. Kerem POLAT

DOI : 10.29228/jrp.186

Abstract (abstract.php?lang=en&id=1072)
Full Text PDF (pdf.php?&id=1072)
Similar Articles (similar.php?&id=1072)
E-mail to Author (mailto:kerem_hybt@hotmail.com)
How to Cite (ajax_handler.php?id=1072)

Lisinopril dihydrate loaded nano-spanlastic bio-adhesive gel for intranasal delivery: 23 factorial optimization, fabrication and ex-vivo studies for enhanced mucosal permeation (abstract.php?id=1073) *Research Article*

Pages 884-899

Suresh PRIYANKA,Radhakrishnan NITHYA

DOI : 10.29228/jrp.187

Abstract (abstract.php?lang=en&id=1073)
Full Text PDF (pdf.php?&id=1073)
Similar Articles (similar.php?&id=1073)
E-mail to Author (mailto:82nithikrish@gmail.com)
How to Cite (ajax_handler.php?id=1073)
Supporting Info (supporting.php?&id=1073)

Pectin/poly(vinylpyrrolidone) composite sponges as wound dressings for delivery of silver sulfadiazine (abstract.php?id=1074) *Research Article*

Pages 900-910

Emine ALARÇIN

DOI : 10.29228/jrp.188

Abstract (abstract.php?lang=en&id=1074)
Full Text PDF (pdf.php?&id=1074)
Similar Articles (similar.php?&id=1074)
E-mail to Author (mailto:emine.alarcin@marmara.redu.tr)

[How to Cite \(ajax_handler.php?id=1074\)](#)

Preparation of resveratrol containing poly (lactic-coglycolic acid) nanoparticles and investigation of its cytotoxic effect in C6 glioma cells (abstract.php?id=1075) *Research Article*

Pages 911-919

MURAT DOĞAN,FATMA ZİŞAN GÜRGAH

DOI : /10.29228/jrp.189

[Abstract \(abstract.php?lang=en&id=1075\)](#)

[Full Text PDF \(pdf.php?&id=1075\)](#)

[Similar Articles \(similar.php?&id=1075\)](#)

[E-mail to Author \(mailto:mdogan@cumhuriyet.edu.tr\)](mailto:mdogan@cumhuriyet.edu.tr)

[How to Cite \(ajax_handler.php?id=1075\)](#)

Chemical constituents and cockroach repellent activity of Sudanese Cyperus rotundus rhizomes essential oil (abstract.php?id=1076) *Research Article*

Pages 920-930

Azhari H. NOUR,Abeer A. IDRIS,Omer A. ISHAG,Abdurahman H. NOUR,Oluwaseun R. ALARA

DOI : 10.29228/jrp.190

[Abstract \(abstract.php?lang=en&id=1076\)](#)

[Full Text PDF \(pdf.php?&id=1076\)](#)

[Similar Articles \(similar.php?&id=1076\)](#)

[E-mail to Author \(mailto:abrahman@ump.edu.my\)](mailto:abrahman@ump.edu.my)

[How to Cite \(ajax_handler.php?id=1076\)](#)

Synthesis and potential antitumor activities of mandelic acid linked 2-aryl-1,3-thiazolidin-4-ones (abstract.php?id=1077) *Research Article*

Pages 931-940

Kübra DEMİR-YAZICI,Özlen GÜZEL-AKDEMİR

DOI : 10.29228/jrp.191

[Abstract \(abstract.php?lang=en&id=1077\)](#)

[Full Text PDF \(pdf.php?&id=1077\)](#)

[Similar Articles \(similar.php?&id=1077\)](#)

[E-mail to Author \(mailto:oguzel@istanbul.edu.tr\)](mailto:oguzel@istanbul.edu.tr)

[How to Cite \(ajax_handler.php?id=1077\)](#)

Synthesis, structure elucidation and cytotoxic activities of 2,5-disubstituted-1,3,4-thiadiazole and 1,2,4-triazole-3-thione derivatives (abstract.php?id=1078) *Research Article*

Pages 941-953

Levent KANDEMİR,Sevgi KARAKUŞ,Suna ÖZBAŞ,Sevim ROLLAS,Julide AKBUĞA

DOI : 10.29228/jrp.192

[Abstract \(abstract.php?lang=en&id=1078\)](#)

[Full Text PDF \(pdf.php?&id=1078\)](#)

[Similar Articles \(similar.php?&id=1078\)](#)

[E-mail to Author \(mailto:skarakus@marmara.edu.tr\)](mailto:skarakus@marmara.edu.tr)

[How to Cite \(ajax_handler.php?id=1078\)](#)

Synthesis and antioxidant activity of benzo[a]pyrano[2,3- c] phenazine derivative compound via one-pot multicomponent reaction (abstract.php?id=1079) *Research Article*

Pages 954-961

Agus Rimus Liandi,Antonius Herry Cahyana

DOI : 10.29228/jrp.193

[Abstract \(abstract.php?lang=en&id=1079\)](#)

[Full Text PDF \(pdf.php?&id=1079\)](#)

[Similar Articles \(similar.php?&id=1079\)](#)

[E-mail to Author \(mailto:agusrimus@uinjkt.ac.id\)](mailto:agusrimus@uinjkt.ac.id)

[How to Cite \(ajax_handler.php?id=1079\)](#)

Glypican-3 as a potential hepatocellular carcinoma peptide vaccine candidate for the Indonesian population (abstract.php?id=1080) Research Article

Pages 962-979

Angelika, Arli Aditya PARIKESIT, Michael JONATHAN, Nadya Marcelina JULIANTO

DOI : 10.29228/jrp.194

[Abstract \(abstract.php?lang=en&id=1080\)](#)

[Full Text PDF \(pdf.php?&id=1080\)](#)

[Similar Articles \(similar.php?&id=1080\)](#)

[E-mail to Author \(mailto:arli.parikesit@i3l.ac.id\)](mailto:arli.parikesit@i3l.ac.id)

[How to Cite \(ajax_handler.php?id=1080\)](#)

Investigation of the effects of bicalutamide and TRAIL signaling mechanism on docetaxel resistant prostate cancer cells (abstract.php?id=1081) Research Article

Pages 980-991

Hilal ACIDERELI, Ozge CEVIK

DOI : 10.29228/jrp.195

[Abstract \(abstract.php?lang=en&id=1081\)](#)

[Full Text PDF \(pdf.php?&id=1081\)](#)

[Similar Articles \(similar.php?&id=1081\)](#)

[E-mail to Author \(mailto:drozgecevik@gmail.com\)](mailto:drozgecevik@gmail.com)

[How to Cite \(ajax_handler.php?id=1081\)](#)

The Cytotoxicity and the Antimicrobial Activities of Arum italicum Miller (abstract.php?id=1082) Research Article

Pages 992-996

Melek ULUSOYLU, Burçak GÜRBÜZ

DOI : 10.29228/jrp.196

[Abstract \(abstract.php?lang=en&id=1082\)](#)

Full Text PDF (pdf.php?&id=1082)
Similar Articles (similar.php?&id=1082)
E-mail to Author (mailto:ulusoylumelek@hotmail.com)
How to Cite (ajax_handler.php?id=1082)

Fullerene modified single-use electrodes as a convenient biosensor platform for electrochemical monitoring of drug- DNA interaction (abstract.php?id=1083) *Research Article*

Pages 997-1006

Ece EKSIN,Arzum ERDEM

DOI : 10.29228/jrp.197

Abstract (abstract.php?lang=en&id=1083)
Full Text PDF (pdf.php?&id=1083)
Similar Articles (similar.php?&id=1083)
E-mail to Author (mailto:arzum.erdem@ede.edu.tr)
How to Cite (ajax_handler.php?id=1083)
Supporting Info (supporting.php?&id=1083)

Assessment of drug-related problems in pediatric inpatients by clinical pharmacist-led medication review: An observational study (abstract.php?id=1084) *Research Article*

Pages 1007-1015

Yeliz ŞAHİN,Çağatay NUHOĞLU,Betul OKUYAN,Mesut SANCAR

DOI : 10.29228/jrp.198

Abstract (abstract.php?lang=en&id=1084)
Full Text PDF (pdf.php?&id=1084)
Similar Articles (similar.php?&id=1084)
E-mail to Author (mailto:yelizsahin89@gmail.com)
How to Cite (ajax_handler.php?id=1084)

Determining levels of unemployment anxiety among pharmacy students at Ege University (abstract.php?id=1085) *Research Article*

Pages 1016-1024

Bülent KIRAN,Elif Gizem KARACA,Büşra BALI

DOI : 10.29228/jrp.199

[Abstract \(abstract.php?lang=en&id=1085\)](#)

[Full Text PDF \(pdf.php?&id=1085\)](#)

[Similar Articles \(similar.php?&id=1085\)](#)

[E-mail to Author \(mailto:bulent.kiran@ege.edu.tr\)](mailto:bulent.kiran@ege.edu.tr)

[How to Cite \(ajax_handler.php?id=1085\)](#)

Health Literacy Among University Students and Associated Factors: A Cross-Sectional Survey Among Pharmacy Students (abstract.php?id=1086) *Research Article*

Pages 1025-1036

Bülent KIRAN,Elif Gizem KARACA,Hür HASSOY

DOI : 10.29228/jrp.200

[Abstract \(abstract.php?lang=en&id=1086\)](#)

[Full Text PDF \(pdf.php?&id=1086\)](#)

[Similar Articles \(similar.php?&id=1086\)](#)

[E-mail to Author \(mailto:bulent.kiran@ege.edu.tr\)](mailto:bulent.kiran@ege.edu.tr)

[How to Cite \(ajax_handler.php?id=1086\)](#)

Formulation and in-vitro characterization of herbal licorice gel for periodontal therapy (abstract.php?id=1088) *Research Article*

Pages 1037-1051

Safiya Fatima KHAN,Bhavya SHETTY,Asim Mustafa KHAN,Ibrahim FAZAL

DOI : 10.29228/jrp.202

[Abstract \(abstract.php?lang=en&id=1088\)](#)

[Full Text PDF \(pdf.php?&id=1088\)](#)

[Similar Articles \(similar.php?&id=1088\)](#)

[E-mail to Author \(mailto:safisupernova@gmail.com\)](mailto:safisupernova@gmail.com)

[How to Cite \(ajax_handler.php?id=1088\)](#)

Marmara University

[Home \(https://jrespharm.com/\)](https://jrespharm.com/)

[Aims and Scope \(static.php?id=1\)](#)

[Editorial Board \(static.php?id=2\)](#)

[Advisory Board \(content.php?id=78\)](#)

[Table of Contents \(static.php?id=4\)](#)

[Author Guidelines \(static.php?id=7\)](#)

[Abstracting / Indexing \(static.php?id=9\)](#)

[Contact \(static.php?id=8\)](#)

Intravaginal delivery of reverse micellar epigallocatechin loaded in κ -carrageenan and HPMC K100M-based gel

Noorma ROSITA^{1,*} , Dewi Melani HARIYADI¹ , Cordellia Calista AMELIA² , Alyajilan Madani MAMURRIE² , Andang MIATMOKO^{1,3} 

¹ Department of Pharmaceutical Sciences, Faculty of Pharmacy, Universitas Airlangga, Nanizar Zaman Joenoes Building, Campus C UNAIR Mulyorejo, Surabaya, 60115, Indonesia

² Bachelor Program of Pharmacy, Department of Pharmaceutical Sciences, Faculty of Pharmacy, Universitas Airlangga, Nanizar Zaman Joenoes Building, Campus C UNAIR Mulyorejo, Surabaya, 60115, Indonesia

³ Stem Cell Research and Development Center, Universitas Airlangga, Institute of Tropical Disease Building 2nd Floor, Campus C UNAIR Mulyorejo, Surabaya, 60115, Indonesia

* Corresponding Author. E-mail: noorma-r@ff.unair.ac.id(N.R.); Tel. +6231-5935249.

Received: 14 January 2022 / Revised: 22 April 2022 / Accepted: 22 April 2022

ABSTRACT: A gel with mucoadhesive properties and a controlled release profile is a suitable dosage form for reverse micellar EGCG delivery. In this study, κ -Carrageenan and HPMC K100M were used as the gel components at a weight ratio of 1:1.5, respectively, for loading native and reverse micellar EGCG. The characteristics of the gel were determined based on pH, swelling index, disintegration time, hardness, and entrapment efficiency. The *in vitro* EGCG release rate was further determined for EGCG levels. Moreover, *in vivo* cervical penetration studies of rhodamine-labeled EGCG gels in mice at two and six hours after intravaginal administrations were conducted. The results showed that the pH and hardness characteristics of the gels for each formula did not differ significantly, while the gel-loaded reverse micellar EGCG had a higher swelling index than that of native EGCG gels. In addition, the rate of release and cervical penetration of rhodamine-labeled reverse micellar EGCG loaded in gels was higher than those of rhodamine-labeled native EGCG gels. Therefore, it can be concluded that loading reverse micelles EGCG into gels prepared with κ -Carrageenan and HPMC K100M successfully controlled the release rate and improved cervical penetration, thereby enabling its potential use in cervical cancer treatment.

KEYWORDS: Epigallocatechin gallate; cancer; hydroxypropyl methylcellulose; κ -carrageenan; reverse micelle; cervical penetration

1. INTRODUCTION

Cervical cancer has a high prevalence in women. World Health Organization states that cervical cancer is the fourth most common cancer in women. In 2018, 570,000 women were diagnosed with cervical cancer worldwide, and about 311,000 women died from the disease(1).It is reported that the persistent infection with carcinogenic human papillomavirus (HPV) types is the leading cause in triggering the development of cervical cancer (2). HPV16 and HPV18 were the most prevalent HPV subtypes among cervical cancer patients, followed by HPV58, HPV53, and HPV33 HPV16 and HPV18 were the most pervasive HPV subtypes among cervical cancer patients(3). Risk factors of developing cervical cancer are low socioeconomic status, smoking, marrying before age 18 years, young age at the first coitus, multiple sexual partners, multiple sexual partners of a spouse, and multiple childbirths (4).Management of cervical cancer is primarily by surgery, radiation therapy, or chemotherapy(5). However, chemotherapy can cause several serious side effects(6), encouraging researchers to look for alternative therapies, including natural products. One natural active ingredient that has been widely reported for cancer treatment is Epigallocatechin gallate (EGCG).

EGCG represents the most effective anticancer polyphenol among the ten contained in green tea (7) and has been shown to have anti-proliferative, anti-metastatic, and pro-apoptotic properties (8).It has been reported to be capable of inhibiting HeLa cell viability, inducing HeLa cell apoptosis, and reducing matrix metalloproteinase-9 (MMP-9) in HeLa cells (9). However, EGCG has low lipophilicity, is easily degraded at high temperatures, and demonstrates low tissue permeability (10).Moreover, it lacks high bioavailability for

Rosita N, Hariyadi DM, Amelia CC, Mamurrie AM, Miatmoko A. Intravaginal delivery of reverse micellar EGCG loaded in κ -Carrageenan and HPMC-based gel. J Res Pharm.2022; 26 (4): 849-858

oral preparations because it can be degraded by the metabolism of digestive bacteria (11). It has previously been reported that the concentration of EGCG decreases at 50°C during storage periods (12). However, although degraded at pH 6.8, it remains stable at pH 4.5 (13). Intravaginal administration at a pH level of 3.5-4.5 with cervical tissue as the target renders gels appropriate as delivery carriers of EGCG in the treatment of cervical cancer.

Gels have certain advantages in the delivery of drugs that protect against first-pass metabolism and enzymatic degradation in the digestive tracts (14). Moreover, the use of mucoadhesive polymers in intravaginal gels can increase the contact time of therapeutic drugs with the tissue mucosa, thus prolonging the therapeutic onset time (15). The most commonly used mucoadhesive polymers include Hydroxy Propyl Methyl Cellulose (HPMC) and κ -Carrageenan (16). HPMC is non-toxic, stable within a wide pH range, and a non-irritating polymer (17). In the previous studies of vaginal tablets prepared with HPMC as the polymer matrix, it has been shown that almost 100% of drugs can be released within 40–150 hours (18,19).

Meanwhile, κ -Carrageenan is a family of gel-forming polysaccharides extracted from certain red seaweed species (20). Gel made from κ -Carrageenan possesses strong and rigid physical properties. Potassium ions have been identified as the optimum crosslinking agents for κ -Carrageenan gels. κ -Carrageenan, as a mucoadhesive polymer of vaginal tablets, successfully releases 100% of Acyclovir within 48 hours (21).

It is known that drug release from the carrier matrix constitutes the initial stage before its penetration into tissues. This process is influenced by several factors, namely, the amount of drug, polymer types, other pharmaceutical excipients, physical or chemical interactions between components, and manufacturing methods (22). The drug release profile can be defined by the mechanism, kinetics, and rate of release. The drug release mechanisms in polymer-based delivery carriers include controlled diffusion, osmotic-controlled, and swelling drug release (22). After the dissolution of the drug, tissue permeability, which is represented by the Log P value, will become an important factor determining concentration in target organs or tissues (23). EGCG has a log P value of 1.1 at pH 4.0 (24). However, optimal lipophilicity for topical drug penetration is indicated by a log P value of between 2 and 3 (25). Modification of EGCG into reverse micelle with the addition of Tween 80 and Span 80 as a surfactant combined with a Hydrophilic-Lipophilic Balance (HLB) value of 6 at pH 5.0 \pm 0.5 was proven to increase the log P value of EGCG to 2.1 (24). This increased lipophilicity of EGCG has also been shown to increase the penetration of EGCG into the skin and also increase its cellular uptake, as indicated by its high cytotoxicity against HeLa cells (24).

The presence of surfactants as components of EGCG reverse micelles at very low concentrations, combined with the polymer, allows aggregate formation (13,26). The aggregates will be formed spontaneously and are reversible at levels above the Critical Aggregation Concentration (CAC) (27). The systems containing polyelectrolytes and non-ionic surfactants have lower CAC. However, interactions will still occur compared to their Critical Micelle Concentration (CMC), albeit in very weak states (13). Consequently, possible physical interactions result from the combination of anionic polymers, including κ -Carrageenan and non-ionic polymers such as HPMC with non-ionic surfactants, i.e., Tween 80 and Span 80 (26).

In this study, the effect of reverse micellar EGCG formation on the characteristics of gels prepared with κ -Carrageenan and HPMC K100M, as well as the release profiles of EGCG and cervical penetration, were investigated. These evaluations proved beneficial for determining the potential use of gels for cervical delivery of reverse micellar EGCG.

2. RESULTS

2.1. Physical characteristics of gel loading reverse micellar EGCG

The physical characteristics of the gels are shown in Table 1. It can be seen that the pH of gel loading native EGCG (EGCG-Gel) was not significantly different from the gel loading reverse micellar EGCG (RM-EGCG-Gel) ($P > 0.05$), being 4.24 and 4.23, respectively. Moreover, the gel hardness of both formulas was approximately 49.75 N, which was higher than the hardness value of a suppository ranging between 17.7–24.5 N (28,29). However, these gels were produced in good firmness resulting in easy intravaginal insertion. The swelling index evaluation revealed a significant difference between the two formulas in that EGCG-Gel had a larger swelling index than that of RM-EGCG-Gel. The use of surfactants for RM-EGCG-Gel can probably form aggregates with polymers from the gel matrix (13), thereby reducing the wettability, which prevents water from causing the polymer to swell. Crucially, such polymer swelling may affect stability and the drug release profile.

Table 1. The physical characteristic of gel loading EGCG prepared with κ -Carrageenan and HPMC K100M. The data represent the mean \pm SD (* $P < 0.05$).

Parameter	EGCG-Gel	RM-EGCG-Gel
pH (n=3)	4.24 \pm 0.01	4.23 \pm 0.06
Hardness (N, n=10)	49.76 \pm 0.11	49.74 \pm 0.04
Swelling Index (n=3)	2.95 \pm 0.54	1.46 \pm 0.19 [*]

2.2. Profiles of EGCG released from gels

The release study showed that approximately 35% of EGCG was released from the gels during the first two hours, as shown in Figure 1. This burst of drug release was probably due to the presence of EGCG on the gel surface and was beneficial in inducing an immediate therapeutic effect. Moreover, the release occurred continuously and was maintained at a slower rate.




Figure 1. Release Profiles of EGCG from EGCG-Gel and RM-EGCG-Gel prepared with κ -Carrageenan and HPMC K100M. The data represent the mean \pm SD (n=3).

This data was subjected to a further calculation for release kinetics modeling. As can be seen from Table 2, the release kinetics of EGCG from the EGCG-Gel fitted the Higuchi model. In contrast, the EGCG released from RM-EGCG-Gel is based on Korsmeyer-Peppas release kinetics. The Higuchi mechanism requires water absorption for drugs to dissolve and diffuse throughout the media. This is in accordance with the increasing index value of EGCG-Gel, which is greater than that of RM-EGCG-Gel.

Table 2. Release kinetic models and their coefficient correlation to match EGCG release profiles from gels

Release Kinetic Model	Coefficient correlation(R ²)	
	EGCG-Gel	RM-EGCG-Gel
Zero Order	0.684	0.701
First Order	0.449	0.582
Korsmeyer-Peppas	0.758 (n=0.54)	0.898 (n=0.38)
Higuchi	0.824	0.849
Hixson-Crowell	0.495	0.617

There was a significant difference in the rate of release or EGCG flux between these two formulas, while the release rate of EGCG gels was obtained from the slope value of each corresponding release kinetics regression. The use of surfactants in reverse micellar EGCG gels (RM-EGCG-Gels) increased the rate of EGCG release from gels prepared with a combination of HPMC K100M polymer and κ -Carrageenan, as shown in Figure 2.

The Korsmeyer-Peppas mechanism determines that drug release does not involve the water absorption process, thus producing a higher RM-EGCG-Gel release rate than EGCG-gels.




Figure 2. The flux of EGCG release from gels prepared with κ -Carrageenan and HPMC K100M. The data represent the mean \pm SD (n=3, *P<0.05).

2.3. In vivo penetration study of gels loading EGCG

The fluorescence intensities of Rhodamine-Labelled EGCG gels at two and six hours of intravaginal administration are shown in Figure 3. There was no Rhodamine or low intensities of the dye observed in the surfactant-blank control group, while the use of surfactants improved its cervical penetration, especially after six hours of administration.




Figure 3. The penetration of Rhodamine-Labelled gel loading native and reverse micellar EGCG in the cervical tissue observed using a fluorescence microscope at two and six hours after intravaginal administration.

Figure 4 shows that the penetration of Rhodamine-labelled EGCG gels after two hours was to a depth of $161.92 \pm 12.30 \mu\text{m}$, while after six hours, it reached a depth of $184.67 \pm 52.98 \mu\text{m}$ tissue penetration, indicating that the longer the contact time of the gels, the deeper the penetration of EGCG. However, both results are based on weak-moderate fluorescence intensity. On the other hand, after two hours, the gel loading reverse micellar EGCG reached a tissue depth of $173.38 \pm 30.79 \mu\text{m}$ with moderate intensity. Then, six hours after administration, the penetration depth was $210.29 \pm 24.11 \mu\text{m}$ with a moderate fluorescence intensity stronger than that of the native EGCG gels.




Figure 4. The depth of EGCG penetration was observed at two and six hours after intravaginal administration of gels. The data represent the mean \pm SD ($n=4$, $*P<0.05$ compared to blank surfactant gel).

3. DISCUSSION

The presence of surfactants in reverse micellar EGCG produces a lower swelling index of the gel prepared with κ -Carrageenan and HPMC K100M (1:1.5) polymers than that of native EGCG. The interaction between polymer-surfactants probably affects the swelling characteristics of polymer-based gels (28). HPMC is a non-ionic polymer, while κ -Carrageenan is an anionic polyelectrolyte polymer (29–31). On the other hand, Tween 80 and Span 80 are non-ionic surfactants (26). It has been reported that interactions may still occur in the systems containing polyelectrolytes, anionic and/or non-ionic polymers with non-ionic surfactants even though their intensity is very low (13,26). These interactions may form aggregates, thereby reducing water wettability and causing the polymer to expand. However, there was no difference between these two formulas in terms of gel hardness.

According to the drug release profiles, during the first two hours, there was an initial burst of EGCG to a level of 35%. The rapid release at this incipient stage may have been due to the intrinsic characteristics of κ -Carrageenan and HPMCK100M related to the water uptake and degree of polymer swelling (32). Nevertheless, the presence of surfactants in reverse micellar EGCG affected the release rate. The existence of this burst produced an immediate therapeutic effect. Furthermore, the release occurred continuously with sustained effects. It can be seen from the contents of Table 2 that the kinetics of the EGCG release from EGCG-Gel matched those of Higuchi release, while the RM-EGCG-gel drug release kinetics fitted the Korsmeyer-Peppas model. Within the Higuchi kinetic model, the release of the drug is based on a diffusion process following Fick's law (33). This mechanism usually occurs in a tablet matrix containing water-soluble drugs with drug release following Higuchi's model, which requires water uptake in the early stages (33–35). It correlates well with the higher swelling index value of EGCG-Gel rather than that of RM-EGCG-Gel.

The release rate analysis indicated that RM-EGCG-Gel has an EGCG release rate greater than that of EGCG-Gel. It appears that the formation of aggregates between surfactants present in the RM-EGCG-Gel had no effect on the EGCG release process. In this case, the release process was more influenced by the polymer-dependent-release mechanism. The rate of drug release of a nanocarrier was also affected by other factors, such as polymer and other excipients, physical or chemical interactions between components, and manufacturing methods(22). The type and number of monomer units of polymers remarkably determined the release of the drug from the dosage form (36).

The penetration study showed that application time significantly affects the depth of EGCG penetration. The rhodamine-labeled EGCG observed during the sixth hour after application produced higher intensity in cervical tissue, which experienced deeper EGCG penetration than those tissues observed two hours after topical administration. In addition, EGCG penetration of the RM-EGCG-Gel indicated greater penetration than that of the EGCG-Gel. The presence of surfactants in reverse micellar EGCG produced an enhancing effect on EGCG penetration. This could also have been caused by an increased EGCG partition coefficient. Physical modification of EGCG with the addition of surfactant Tween 80 and Span 80 produced more lipophilic EGCG by increasing the coefficient partition of EGCG from 1.89 to 2.1 (24). It has been established that the optimal Log P value for tissue penetration is between 2 and 3 (23,24). The addition of Tween 80 and Span 80 surfactants to reverse micellar EGCG systems can also increase membrane fluidity which is affected by the fatty acid chain of surfactants (37). Unsaturated fatty acids of Tween 80 and Span 80 have a double bond that can cause the membrane structure to stretch and render, making it easier for EGCG to penetrate the membrane (37).

In this research, the intensity of rhodamine indicated that EGCG could penetrate as far as the deep layer of cervical tissue. However, the concentration of EGCG required in the target tissue to support anticancer therapy should be further analyzed.

4. CONCLUSION

The characteristics of the gel-loaded reverse micellar EGCG were not significantly different from those in the original EGCG. However, the use of surfactants in these gels produced a higher swelling index than that of native EGCG. Moreover, surfactant Tween 80 and Span 80 in reverse micellar EGCG increased the release flux and cervical penetration of EGCG from κ -Carrageenan and HPMC K100M-based gels, indicating the future application of EGCG for therapeutical purposes.

5. MATERIALS AND METHODS

5.1. Materials

EGCG with a purity grade of 98% was obtained from Xi'an Rhongsheng Biotechnology Co., Ltd. (Shaanxi, China). HPMC K100M is a product of Aldrich (USA). κ -Carrageenan was purchased from Danisco-Cultor (Copenhagen, Denmark). Tween 80 is a product of KOA Corporation (Tokyo, Japan). Span 80 was acquired from Nanhang Industrial Co. Ltd. (Zhejiang, China). Other chemicals and reagents used represented available non-technical grades.

5.2. Preparation of EGCG Gels

Gels were prepared using the components in Table 3. κ -Carrageenan and HPMC K100M were initially dispersed in citrate buffer pH 4.2 and mixed after complete swelling had occurred. The EGCG solution in citrate buffer pH 4.2 \pm 0.3 was subsequently added to the polymer dispersion, followed by 0.025 M KCl solution as the crosslinking agent and stirred until homogeneous. At that point, the gels were put into molds and stored at 4°C for 24 hours.

Table 3. Formulation of gels loading native and reverse micellar EGCG

Components	Function	EGCG-Gel		RM-EGCG-Gel	
		%	mg	%	mg
Native EGCG	Active substance	3	90	-	-
EGCG reverse micelle	Active substance	-	-	Equivalent to 3% EGCG	
Tween 80	Surfactant	-	-	0.01	0.3
Span 80	Surfactant	-	-	0.05	1.5
κ -Carrageenan	Polymer	7.5	225	7.5	225
HPMC K100M	Polymer	5	150	5	150
0.025 M KCl solution	Crosslinking agent		1 mL		1 mL
Citrate Buffer pH 4.2	Solvent		Up to 3 grams		Up to 3 grams

In preparing gel loading reverse micellar EGCG, the modified reverse micelle EGCG was first prepared in the manner previously described (24) using Tween 80 and Span 80 HLB 6 in citrate buffer at pH 4.2. The gels were then prepared to employ the same method as gel loading native EGCG.

5.3. Characterization of gel loading EGCG

For preparing the gel, the gel mass was shaped into a mold and stored at a cool temperature for 24 hours, and then the hard gel was left at room temperature for a while and measured for further evaluation. The pH was initially evaluated by dissolving the gels in warm water until all matrix components had melted and dispersed. At that point, the pH was measured at 37 \pm 0.5 °C using a pH meter (38). Moreover, a gel hardness evaluation was carried out on ten gels using an Erweka® hardness tester at a temperature of 25 \pm 0.5 °C (38–40). The swelling index was further evaluated by weighing gels individually (W_0). Firstly, the beaker glass, funnel, and filter paper were weighed before approximately 50 mL citrate buffer pH 4.2 was added and the weight of the buffer solution was calculated. A gel was then introduced into the expanding medium. After six hours, it was transferred to another container, and the final buffer solution was weighed. The weight of the buffer solution absorbed by the gel was obtained, giving the final gel weight (W_t). The swelling index was subsequently calculated using the following formula (41)

$$\text{Swelling Index} = \frac{W_t - W_0}{W_0}$$

where W_t : weight of gel at time t , and W_0 : initial weight of the gel

5.4. Evaluation of EGCG released from gels

The EGCG released from gels was examined for gels equivalent to 270 mg of EGCG using 100 mL citrate buffer pH 4.2 as the release media at 37°C with continuous stirring at 100 rpm. Each sample of approximately 5 mL was extracted with a syringe at several determined points up to 720 minutes prior to filtering through paper with a pore size of 0.45 μm . The same volume of citrate buffer pH 4.2 was then added to the release medium to replace the extracted volume of samples. The EGCG level in the sample was determined using a UV-Vis Spectrophotometer at a maximum wavelength of 271 nm prior to the EGCG release profile being analyzed. To determine the release kinetics of native EGCG and reverse micellar EGCG from HPMC and κ -Carrageenan-based gels, the release profile was further plotted and analyzed for zero-order, first-order, Korsmeyer-Peppas, Higuchi, and Hixson-Crowell kinetics models.

5.5. *In vivo* cervical penetration study of EGCG gels

The protocol of this experimental study was approved by the Ethical Committee of the Faculty of Veterinary Medicine, Universitas Airlangga, Indonesia, through ethical clearance letter No. 2.KE.095.06.22021. Twenty-four female mice were divided into six groups of four subjects that had been acclimatized for a week prior to initiation of the study.

The samples were prepared for the *in vivo* penetration study by adding about 2.56 mg Rhodamine-B to the mixtures of surfactants and gel bases as presented in Table 2 for preparing Rhodamine-labelled Blank Surfactant, EGCG- and RM-EGCG Gels.

The subjects were anesthetized with ketamine at a dose of 70 mg/kg body weight and administered intraperitoneally. The sample was subsequently applied at an amount equivalent to 0.234 mg EGCG intravaginally using a syringe connected to a catheter. After two and six hours of application, the subjects were sacrificed using cervical dislocation. Cervical tissue sections were then sliced to a thickness of 5 μm at -20°C using a cryo-microtome (Leica Cryocut, Germany) before being observed using a fluorescence microscope (FSX100 Olympus, Japan).

Data analysis was performed using a fluorescence intensity scoring system based on sample penetration after two and six hours of intravaginal administration. The fluorescence intensity scores were (-) for no fluorescence intensity, (+) for weak fluorescence intensity, (++) for moderate fluorescence intensity, and (+++) for strong fluorescence intensity.

5.6. Statistical analysis

In this study, the data represent the mean \pm standard deviation (SD). The data was analyzed statistically using Independent Samples t-Test with IBM SPSS Statistics v.22 software. The P -value less than 0.05 expressed the significant difference between samples.

Acknowledgments: This study was financially supported by a Preliminary Research on Excellence in Higher Education Institutions (Penelitian Dasar Unggulan Perguruan Tinggi, PDUPT) No. 992/UN3.14/PT/2020 provided by the Ministry of Research and Technology-National Research and Innovation Agency of the Republic of Indonesia.

Author contributions: Concept - N.R., D.M.H., A.M.; Design - N.R., D.M.H., A.M.; Supervision - N.R., D.M.H., A.M.; Resources - N.R., D.M.H.; Materials - N.R., D.M.H., A.M.; Data Collection and/or Processing - C.C.A., A.M.M., N.R.; Analysis and/or Interpretation - C.C.A., A.M.M., N.R., D.M.H., A.M.; Literature Search - C.C.A., A.M.M.; Writing - C.C.A., A.M.M., N.R.; Critical Reviews - N.R., D.M.H., A.M.

Conflict of interest statement: The authors declared no conflict of interest.

REFERENCES

1. World Health Organization (WHO). Cervical Cancer. <https://www.who.int/cancer/prevention/diagnosis-screening/cervical-cancer/en/> (accessed on 14 January 2017)
2. Arbyn M, Weiderpass E, Bruni L, de Sanjosé S, Saraiya M, Ferlay J, et al. Estimates of incidence and mortality of cervical cancer in 2018: a worldwide analysis. *Lancet Glob Heal*. 2020;8(2):e191-203. [CrossRef]
3. Li K, Yin R, Wang D, Li Q. Human papillomavirus subtypes distribution among 2309 cervical cancer patients in West China. *Oncotarget*. 2017;8(17):28502-9. [CrossRef]

4. Kashyap N, Krishnan N, Kaur S, Ghai S. Risk Factors of Cervical Cancer: A Case-Control Study. *Asia-Pacific J Oncol Nurs.* 2019;6(3):308–14. [\[CrossRef\]](#)
5. Bhatla N, Aoki D, Sharma DN, Sankaranarayanan R. Cancer of the cervix uteri. *Int J Gynecol Obstet.* 2018;143:22–36. [\[CrossRef\]](#)
6. Nurgali K, Jagoe RT, Abalo R. Editorial: Adverse effects of cancer chemotherapy: Anything new to improve tolerance and reduce sequelae? *Front Pharmacol.* 2018;9(MAR):1–3. [\[CrossRef\]](#)
7. Du GJ, Wang CZ, Qi LW, Zhang ZY, Calway T, He TC, et al. The synergistic apoptotic interaction of panaxadiol and epigallocatechin gallate in human colorectal cancer cells. *Phyther Res.* 2013;27(2):272–7. [\[CrossRef\]](#)
8. Almatroodi SA, Almatroodi A, Khan AA, Alhumaydhi FA, Alsahli MA, Rahmani AH. Potential therapeutic targets of epigallocatechin gallate (egcg), the most abundant catechin in green tea, and its role in the therapy of various types of cancer. *Molecules.* 2020 Jul 9;25(14):3146. [\[CrossRef\]](#)
9. Sharma C, Nusri QEA, Begum S, Javed E, Rizvi TA, Hussain A. (-)-Epigallocatechin-3-gallate induces apoptosis and inhibits invasion and migration of human cervical cancer cells. *Asian Pacific J Cancer Prev.* 2012;13(9):4815–22. [\[CrossRef\]](#)
10. Yoshino S, Mitoma T, Tsuruta K, Todo H, Sugibayashi K. Effect of emulsification on the skin permeation and UV protection of catechin. *Pharm Dev Technol.* 2014;19(4):395–400. [\[CrossRef\]](#)
11. Gan RY, Li H Bin, Sui ZQ, Corke H. Absorption, metabolism, anticancer effect and molecular targets of epigallocatechin gallate (EGCG): An updated review. *Crit Rev Food Sci Nutr.* 2018;58(6):924–41. [\[CrossRef\]](#)
12. Rosita N, Nailufa Y, Hariyadi DM. Characteristics, stability and activity of epigallocatechin gallate (EGCG)-chitosan microspheres: Effect of polymer concentration. *Res J Pharm Technol.* 2020;13(5):2303–9. [\[CrossRef\]](#)
13. Diamant H, Andelman D. Onset of self-assembly in polymer-surfactant systems. *Europhys Lett.* 1999;48(2):170–6. [\[CrossRef\]](#)
14. Loyd V. Allen J, Ansel HC. *Ansel's Pharmaceutical Dosage Forms and Drug Delivery Systems.* 10th ed. Lippincott Williams & Wilkins, Philadelphia, PA 2014.
15. De Araújo Pereira RR, Bruschi ML. Vaginal mucoadhesive drug delivery systems. *Drug Dev Ind Pharm.* 2012;38(6):643–52. [\[CrossRef\]](#)
16. Acarturk F. Mucoadhesive Vaginal Drug Delivery Systems. *Recent Pat Drug Deliv Formul.* 2009;3(3):193–205. [\[CrossRef\]](#)
17. Rowe RC, Sheskey PJ, Quinn ME. *Handbook of Pharmaceutical Excipients.* 6th ed. USA: RPS Publishing; 2009.
18. Pacheco-Quito EM, Ruiz-Caro R, Rubio J, Tamayo A, Veiga MD. Carrageenan-based acyclovir mucoadhesive vaginal tablets for prevention of genital herpes. *Mar Drugs.* 2020;18(5). [\[CrossRef\]](#)
19. Perioli L, Ambrogi V, Pagano C, Massetti E, Rossi C. New solid mucoadhesive systems for benzydamine vaginal administration. *Colloids Surfaces B Biointerfaces.* 2011;84(2):413–20. [\[CrossRef\]](#)
20. Necas J, Bartosikova L. Carrageenan: a review. *Vet Med (Praha).* 2013;58(4):187–205. [\[CrossRef\]](#).
21. Sánchez-Sánchez M. P., Martín-Illana A., Ruiz-Caro R., Bermejo P., Abad M. J., Carro R., Bedoya L. M., Tamayo A., Rubio J., Fernández-Ferreiro A., Otero-Espinar F., & Veiga M. D. Chitosan and kappa-carrageenan vaginal acyclovir formulations for prevention of genital herpes. *in vitro and ex vivo evaluation.* *Mar Drugs.* 2015;13(9):5976–92. [\[CrossRef\]](#)
22. Son GH, Lee BJ, Cho CW. Mechanisms of drug release from advanced drug formulations such as polymeric-based drug-delivery systems and lipid nanoparticles. *J Pharm Investig.* 2017;47(4):287–96. [\[CrossRef\]](#)
23. Miatmoko A, Ayunin Q, Soeratri W. Ultradeformable vesicles: concepts and applications relating to the delivery of skin cosmetics. *Ther Deliv.* 2021;12:739–56. [\[CrossRef\]](#)
24. Rosita N, Meitasari VA, Rianti MC, Hariyadi DM. Enhancing skin penetration of epigallocatechin gallate by modifying partition coefficient using reverse micelle method. *Ther Deliv.* 2019;10(7):409–17. [\[CrossRef\]](#)
25. Benson HAE. Topical and Transdermal Drug Delivery. In: *Topical and Transdermal Drug Delivery.* A John Wiley & Sons, Inc., Publication; 2005. p. 5–37. [\[CrossRef\]](#)
26. Mohsenipour AA, Pal R. A Review of Polymer-Surfactant Interactions. In: *Handbook of Surface and Colloid Chemistry.* 4th ed. U.S.: CRC Press; 2016. p. 639. [\[CrossRef\]](#)
27. Owen SC, Doak AK, Wassam P, Shoichet MS, Shoichet BK. Colloidal aggregation affects the efficacy of anticancer drugs in cell culture. *ACS Chem Biol.* 2012;7(8):1429–35. [\[CrossRef\]](#)

28. Alvarez-Lorenzo C, Concheiro A. Effects of surfactants on gel behavior: design implications for drug delivery effects of surfactants on gel behavior design implications for drug delivery systems. *Am J Drug Deliv.* 2003;1(2):77–101. [[CrossRef](#)]
29. Sardar N, Kamil M, Kabir-Ud-Din. Interaction between non-ionic polymer hydroxypropyl methyl cellulose (HPMC) and cationic gemini/conventional surfactants. *Ind Eng Chem Res.* 2012;51(3):1227–35. [[CrossRef](#)]
30. Bao H, Li N, Gan LH, Zhang H. Interactions between ionic surfactants and polysaccharides in aqueous solutions. *Macromolecules.* 2008;41(23):9406–12. [[CrossRef](#)]
31. dos Santos MA, Grenha A. Polysaccharide nanoparticles for protein and peptide delivery: exploring less-known materials. 1st ed. Vol. 98, *Advances in protein chemistry and structural biology.* Elsevier Inc.; 2015. 223–261 p. [[CrossRef](#)]
32. Yoo J, Won Y. Phenomenology of the Initial Burst Release of Drugs from PLGA Microparticles. *ACS Biomater Sci Eng.* 2020;6:6053–62. [[CrossRef](#)]
33. Paul DR. Elaborations on the Higuchi model for drug delivery. *Int J Pharm.* 2011;418(1):13–7. [[CrossRef](#)]
34. Bruschi ML. 5 - Mathematical models of drug release. In: Bruschi MLBT-S to M the DR from PS, editor. *Strategies to Modify the Drug Release from Pharmaceutical Systems.* Woodhead Publishing; 2015. p. 63–86.
35. Yadav G, Bansal M, Thakur N, Khare P. Multilayer tablets and their drug release kinetic models for oral controlled drug delivery system. *Middle-East J Sci Res.* 2013;16(6):782–95.
36. Liechty WB, Kryscio DR, Slaughter B V, Peppas NA. Polymers for drug delivery systems. *Annu Rev Chem Biomol Eng.* 2010;1:149–73. [[CrossRef](#)]
37. Mazyed EA, Helal DA, Elkhoudary MM, Abd Elhameed AG, Yasser M. Formulation and optimization of nanospanlastics for improving the bioavailability of green tea epigallocatechin gallate. *Pharmaceuticals.* 2021 Jan 15;14(1):68. . [[CrossRef](#)]
38. Gomaa E, Abu Lila AS, Hasan AA, Ghazy F eldin S. Preparation and characterization of intravaginal vardenafil suppositories targeting a complementary treatment to boost in vitro fertilization process. *Eur J Pharm Sci.* 2018;111:113–20. [[CrossRef](#)]
39. Reddy RS, Kumar L, Pydi CR, Reddy MS, Verma R. Development of Fluconazole Suppositories for the Treatment of Candida Infection of Genitourinary Tract. *Indian J Pharm Educ Res.* 2018;52(4):S16–22.
40. Mohamed DFM, Mahmoud OAE, Mohamed FA. Preparation and Evaluation of Ketotifen Suppositories. *J Adv Biomed Pharm Sci.* 2020;3:10–22.
41. Hassan AS, Soliman GM, Ali MF, El-Mahdy MM, El-Gindy GEDA. Mucoadhesive tablets for the vaginal delivery of progesterone: in vitro evaluation and pharmacokinetics/pharmacodynamics in female rabbits. Vol. 44, *Drug Development and Industrial Pharmacy.* Taylor & Francis; 2018. 224–232 p. [[CrossRef](#)]



Understand societal impact

Analyze impact of research

Analyze and benchmark the impact of your institution's research for each of the UN SDGs

elsevier.com

OPEN

Journal of Research in Pharmacy

COUNTRY

Turkey



Universities and research institutions in Turkey

SUBJECT AREA AND CATEGORY

Medicine
 Pharmacology (medical)
 Pharmacology, Toxicology and Pharmaceutics
 Pharmacology, Toxicology and Pharmaceutics (miscellaneous)

PUBLISHER

Marmara University



Marmara University in Scimago Institutions Rankings

H-INDEX

16

PUBLICATION TYPE

Journals

ISSN

26306344

COVERAGE

2019-2021

INFORMATION

[Homepage](#)
[How to publish in this journal](#)
ikucukguzel@marmara.edu.tr

SCOPE

Journal of Research in Pharmacy is the official scientific journal of Marmara University Faculty of Pharmacy. The journal is the continuation of the former "Journal of Pharmacy of University of Marmara" which was published between 1985 and 1997. Since

CATCH EVERY DETAIL

300,000+ products & services to help you solve the toughest problems in life science

Disc


many different specialists. Journal of Research in Pharmacy publishes original research papers, review articles and scientific commentaries on all aspects of pharmaceutical sciences depending on their conceptual novelty and scientific quality. The journal welcomes articles in this multidisciplinary field, with a focus on topics relevant for drug action, drug discovery and development, conventional and emerging fields related to pharmaceutical sciences. Articles, which cannot be associated with pharmaceutical issues in any way, might be returned to authors without processing. Scientific commentaries and review articles are generally evaluated by invitation or assent of the Editors. Proceedings of scientific meetings may also be published as special issues or supplements to the Journal, upon decision by the editors.

Join the conversation about this journal

Quartiles

FIND SIMILAR JOURNALS ?






<p>1 Indian Journal of Pharmaceutical Sciences IND</p> <p>75% similarity</p>	<p>2 Turkish Journal of Pharmaceutical Sciences TUR</p> <p>73% similarity</p>	<p>3 Journal of Reports in Pharmaceutical Sciences IRN</p> <p>64% similarity</p>	<p>4 Acta Pharmaceu TUR</p> <p>63% similarity</p>
--	---	--	---



CATCH EVERY DETAIL

300,000+ products & services to help you solve the toughest problems in life science

Disc



Journal of Research in Pharmacy

Q3 Pharmacology, Toxicology and Pharmaceutics...
best quartile

SJR 2021
0.19

powered by scimagojr.com


← Show this widget in your own website

Just copy the code below and paste within your html code:


```
<a href="https://www.scimagojr.com" style="border: 1px solid #ccc; padding: 2px 5px; display: inline-block;">https://www.scimagojr.com
```

SCImago Graphica

Explore, visually communicate and make sense of data with our **new data visualization tool**.



Metrics based on Scopus® data as of April 2022



Source details

Journal of Research in Pharmacy

Formerly known as: Marmara Pharmaceutical Journal

Scopus coverage years: from 2019 to Present

Publisher: Marmara University

E-ISSN: 2630-6344

Subject area: Pharmacology, Toxicology and Pharmaceutics: General Pharmacology, Toxicology and Pharmaceutics

Medicine: Pharmacology (medical)

Source type: Journal

CiteScore 2021 **1.3**

SJR 2021 **0.189**

SNIP 2021 **0.372**

[View all documents >](#)

[Set document alert](#)

[Save to source list](#)


[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)

[Export content for category](#)

CiteScore rank 2021 In category: General Pharmacology, Toxicol...

☆	Rank	Source title	CiteScore 2021	Percentile
☆	#42	Journal of Research in Pharmacy	1.3	43rd percentile
	74			
☆	#1	Acta Pharmaceutica Sinica B	15.9	99th percentile
☆	#2	Drug Metabolism Reviews	8.8	97th percentile
☆	#3	Life Sciences	8.0	96th percentile
☆	#4	Expert Review of Clinical Pharmacology	6.8	95th percentile
☆	#5	Advanced Pharmaceutical Bulletin	6.1	93rd percentile
☆	#6	Journal of Zhejiang University: Science B	6.0	92nd percentile
☆	#7	Handbook of Experimental Pharmacology	5.9	91st percentile
☆	#8	ChemMedChem	5.6	89th percentile
☆	#9	Advances in Pharmacological and Pharmaceutical Sciences	5.5	88th percentile
☆	#10	Clinical and Translational Science	5.2	87th percentile
☆	#11	F1000Research	5.0	85th percentile

CiteScore trend



☆	Rank	Source title	CiteScore 2021	Percentile
☆	#12	Research in Pharmaceutical Sciences	4.7	84th percentile
☆	#13	Current Reviews in Clinical and Experimental Pharmacology	4.6	83rd percentile
☆	#14	Therapeutics and Clinical Risk Management	4.5	81st percentile
☆	#15	Pharmacogenetics and Genomics	4.5	80th percentile
☆	#16	Brazilian Journal of Medical and Biological Research	4.3	79th percentile
☆	#17	Drug Target Insights	4.3	77th percentile
☆	#18	Antibiotics	3.9	76th percentile
☆	#19	Revista Brasileira de Farmacognosia	3.9	75th percentile
☆	#20	Biomedical Reports	3.9	73rd percentile
☆	#21	Medicinal Chemistry Research	3.8	72nd percentile
☆	#22	Journal of Cardiovascular Development and Disease	3.6	70th percentile
☆	#23	Iranian Journal of Pharmaceutical Research	3.5	69th percentile
☆	#24	Bioanalysis	3.5	68th percentile
☆	#25	Pharmacology Research and Perspectives	3.3	66th percentile
☆	#26	ADMET and DMPK	3.2	65th percentile
☆	#27	American Journal of Pharmaceutical Education	2.9	64th percentile
☆	#28	Drug Metabolism and Personalized Therapy	2.9	62nd percentile
☆	#29	Mental Health Clinician	2.7	61st percentile
☆	#30	Brazilian Journal of Pharmaceutical Sciences	2.5	60th percentile
☆	#31	Farmacia	2.4	58th percentile
☆	#32	Journal of Applied Pharmaceutical Science	2.3	57th percentile
☆	#33	Currents in Pharmacy Teaching and Learning	2.3	56th percentile
☆	#34	Molecular and Cellular Toxicology	2.2	54th percentile
☆	#35	Journal of Anaesthesiology Clinical Pharmacology	2.0	53rd percentile
☆	#36	Rasayan Journal of Chemistry	2.0	52nd percentile
☆	#37	Current Bioactive Compounds	1.8	50th percentile

☆	Rank	Source title	CiteScore 2021	Percentile
☆	#38	ScienceRise: Pharmaceutical Science	1.7	49th percentile
☆	#39	Methods in Pharmacology and Toxicology	1.5	47th percentile
☆	#40	Pharmaceutical Sciences	1.5	46th percentile
☆	#41	Jundishapur Journal of Natural Pharmaceutical Products	1.3	45th percentile
☆	#42	Journal of Research in Pharmacy	1.3	43rd percentile
☆	#43	Journal of Pharmacy and Bioallied Sciences	1.1	42nd percentile
☆	#44	Handbook of Analytical Separations	1.1	41st percentile
☆	#45	Indian Journal of Pharmaceutical Education and Research	1.1	39th percentile
☆	#46	Current Drug Therapy	0.9	38th percentile
☆	#47	Journal of Reports in Pharmaceutical Sciences	0.9	37th percentile
☆	#48	Yaoxue Xuebao	0.9	35th percentile
☆	#49	Zhongguo Zhongyao Zazhi	0.9	34th percentile
☆	#50	Neuromethods	0.8	33rd percentile
☆	#51	Neuropsychopharmacologia Hungarica	0.7	31st percentile
☆	#52	Indian Journal of Chemistry - Section B Organic and Medicinal Chemistry	0.7	30th percentile
☆	#53	Nanomedicine Journal	0.6	29th percentile
☆	#54	Wikijournal of Medicine	0.6	27th percentile
☆	#55	Current Protocols	0.6	26th percentile
☆	#56	Progress in Drug Research	0.6	25th percentile
☆	#57	European Pharmaceutical Journal	0.6	23rd percentile
☆	#58	Iraqi Journal of Pharmaceutical Sciences	0.6	22nd percentile
☆	#59	Phytomedicine Plus	0.6	20th percentile
☆	#60	Pharmaceutical Sciences Asia	0.5	19th percentile
☆	#61	Defence Life Science Journal	0.5	18th percentile
☆	#62	Biomedical and Biopharmaceutical Research	0.3	16th percentile
☆	#63	African Journal Biomedical Research	0.3	15th percentile

☆	Rank	Source title	CiteScore 2021	Percentile
☆	#64	Azerbaijan Pharmaceutical and Pharmacotherapy Journal	0.3	14th percentile
☆	#65	Hacettepe University Journal of the Faculty of Pharmacy	0.3	12th percentile
☆	#66	Chinese Journal of New Drugs	0.2	11th percentile
☆	#67	Khyber Medical University Journal	0.1	10th percentile
☆	#68	Pharma Times	0.1	8th percentile
☆	#69	Acta Marisiensis - Seria Medica	0.1	7th percentile
☆	#70	Journal of Integrated Science and Technology	0.1	6th percentile
☆	#71	International Journal of Pharmaceutical Sciences and Nanotechnology	0.0	4th percentile
☆	#72	Health Biotechnology and Biopharma	0.0	2nd percentile
☆	#72	Revista Colombiana de Ciencias Quimico-Farmacéuticas(Colombia)	0.0	2nd percentile
☆	#72	World Economics	0.0	2nd percentile

About Scopus

[What is Scopus](#)

[Content coverage](#)

[Scopus blog](#)

[Scopus API](#)

[Privacy matters](#)

Language

[日本語版を表示する](#)

[查看简体中文版本](#)

[查看繁體中文版本](#)

[Просмотр версии на русском языке](#)

Customer Service

[Help](#)

[Tutorials](#)

[Contact us](#)

ELSEVIER

[Terms and conditions](#) ↗ [Privacy policy](#) ↗

Copyright © Elsevier B.V. ↗. All rights reserved. Scopus® is a registered trademark of Elsevier B.V.

We use cookies to help provide and enhance our service and tailor content. By continuing, you agree to the use of cookies ↗.

