

ISSN 1110 - 6131



**Egyptian Journal  
Of  
Aquatic Biology  
And  
Fisheries**

---

**Editor-in-Chief****Prof. Abdalla M. Ibrahim**

Zoology Dept., Faculty of Science, Ain Shams University, Cairo, Egypt.

Email Address: drabdallai@yahoo.com

---

**Co-Editor-in-Chief****Prof. Magdy T. Khalil**

Zoology Dept., Faculty of Science, Ain Shams University, Cairo, Egypt.

Email Address: mtkhalil52@hotmail.com

---

**Associate Editor****Prof. Waheed M. Emam**

Zoology Dept., Faculty of Science, Ain Shams University, Cairo, Egypt.

Email Address: eased2001@yahoo.com

---

**Prof. Abd El-Halim A. Saad**

Zoology Dept., Faculty of Science, Ain Shams University, Cairo, Egypt.

Email Address: halimsaad@ymail.com

---

**Prof. Nassrallah Abdel-Hameid,**

Zoology Dept., Faculty of Science, Banha University, Cairo, Egypt

Email Address: nassr65@gmail.com

---

**Prof. Sahar Fahmy Mehanna**

Professor of Fish Population Dynamics and stock assessment, National Institute of Oceanography and Fisheries, Suez. Egypt

Email Address: sahar\_mehanna@yahoo.com

---

**Prof. Alahmady Shafiq Alzahaby**

Department of zoology, Faculty of science, Zagazig University, Zagazig Egypt.

Email Address: alahmady\_alzahaby@yahoo.com

---

**Prof. Ahmad Mosaad Azab**

Zoology Dept., Faculty of Science Al-Azhar University, Cairo Egypt.

Email Address: amazab2000@yahoo.com

---

**Prof. Gamal Shebl Ramadan**

Prof. of Physiology, Zoology Dept., faculty of Science, Ain shams Univ. Egypt

Email Address: gamal\_ramadan@hotmail.com

---

**Advisory Board****Prof. Roger Flower**

University College London | UCL · Environmental Change Research Centre (ECRC), United Kingdom.

Email Address: rflower@geog.ucl.ac.uk

---

**Prof. Safwat H. Shakir**

Chemical Engineering Department, Roy G. Perry College of Engineering, Prairie View A&M University, Texas A&M University System, Prairie View, Texas, USA

Email Address: shshakir@pvamu.edu

---

**Prof. Richard Heckmann**

Zoology Department, Brigham Young University, USA.

Email Address: biosecretary@byu.edu

---



**Prof. Irvin W. Osborne-Lee**

Chemical Engineering Department, Roy G. Perry College of Engineering, Prairie View A&M University, Texas A&M University System, Prairie View, Texas, USA

Email Address: [oslee@pvamu.edu](mailto:oslee@pvamu.edu)

---

**Language Editor**



**Mrs. Mona S. Higazy**

English Department, Faculty of Alsun, Ain Shams University, Cairo, Egypt.

Email Address: [monahigazy0@gmail.com](mailto:monahigazy0@gmail.com)

---

Volume & Issue: Volume 26, Issue 5, September and October 2022 [XML](#)

- 
- 1 **Basic life-history parameters of the blue swimmer crab *Portunus pelagicus* (Linnaeus, 1758) along the Gulf of Suez, Red Sea, Egypt**  
Page 1-19  
Reem A. Mahmoud et al.  
[View Article](#) | [PDF 1.39 MB](#)

---

  - 2 **Perspectives of hotel and tourist village officials on the cultivation of wooden trees for shoreline stabilization on the northwestern coast, Egypt**  
Page 21-38  
Noura H.S. Hassan et al.  
[View Article](#) | [PDF 608.09 K](#)

---

  - 3 **Tilapia Lake Virus (TiLV): the aetiological agent of Tilapia Lake Virus Disease (TiLVD): A review article**  
Page 39-51  
Saleh Ahmed Rabeh  
[View Article](#) | [PDF 948.21 K](#)

---

  - 4 **Potential Application of Agriculture by-Products in Heavy Metals Bioremediation**  
Page 53-67  
Mohamed A. El-Sawy et al.  
[View Article](#) | [PDF 654.25 K](#)

---

  - 5 **Community structure and Spatio-Temporal Distribution of Chaetognatha near Corals Reefs on Hurghada coast, Red Sea, Egypt**  
Page 69-91  
Zeinab Abouelnaga et al.  
[View Article](#) | [PDF 1.42 MB](#)

---

  - 6 **Routes of Transmission of SARS CoV-2 through Aquatic Environment in Developing Countries**  
Page 93-116  
Waled M. El-Senousy et al.  
[View Article](#) | [PDF 721.05 K](#)

---

  - 7 **Risk assessment of heavy metals in mangrove trees (*Avicennia marina*) and associated seawater of Ras Mohammed Protectorate, Red Sea, Egypt**  
Page 117-135  
Donia H. Elnaggar et al.

[View Article](#) | [PDF 910.52 K](#)

---

8 **[Assessment of heavy metal concentration in water and the Nile tilapia of Lake Manzala, EL-Kapoty, Egypt](#)**

Page 137-147

Ahmed I. Abd El-kader et al.

[View Article](#) | [PDF 400.18 K](#)

---

9 **[Bacterial Load of Tilapia Fish Residing El-Ramla and Abu Simbel Khors, Lake Nasser, Egypt](#)**

Page 149-160

Sayed M. Ali

[View Article](#) | [PDF 615.74 K](#)

---

10 **[Captive breeding of the stinging catfish, \*Heteropneustes fossilis\* \(Bloch, 1794\) found in Brahmaputra River, Assam, India using inducing agent ovasis and its early embryogenesis](#)**

Page 161-173

Chandopal Saikia et al.

[View Article](#) | [PDF 808.64 K](#)

---

11 **[Post-mortem Biochemical, Microbiological and Sensory Quality Changes in the Grass Carp \(\*Ctenopharyngodon idella\*\) during Ice Storage](#)**

Page 175-187

Md. Golam Rasul et al.

[View Article](#) | [PDF 282.09 K](#)

---

12 **[Determination of sensory attributes, microbiological, and biochemical analysis, and pesticide contents of the dried shark \(\*Scoliodon sorrakowah\*\) in the Bengal Bay of Bangladesh](#)**

Page 189-206

Chakma Suprakash et al.

[View Article](#) | [PDF 584.35 K](#)

---

13 **[Antibacterial, antibiofilm, antioxidants and phytochemical profiling of \*Syzygium aromaticum\* extract](#)**

Page 207-218

Ahmed Mohamed Alghazzaly et al.

[View Article](#) | [PDF 605.95 K](#)

---

14 **[Age and growth of elephant-snout fish \*Mormyrus kannume\* \(Forsskål, 1775\) \(Family: Mormyridae\) from Lake Nasser, Egypt](#)**

Page 219-234

Mahmoud M. S. Farrag et al.

[View Article](#) | [PDF 962.56 K](#)

---

- 15 **Evaluation of dietary fish oil: soybean oil ratios on growth performance feed utilization, haematological indices and body fatty acid composition of Keeled mullet, *Liza carinata* fingerlings**

Page 235-256

Ahmed M. Aboseif et al.

[View Article](#) | [PDF 641.37 K](#)

---

- 16 **Effectiveness of dietary vitamin C on the performance of common carp (*Cyprinus carpio*).**

Page 257-287

Ahmed M. Aboseif et al.

[View Article](#) | [PDF 1.37 MB](#)

---

- 17 **Mosquitocidal activity of *Ophiocoma scolopendrina* extracts against *Culex pipiens* and their antimicrobial potential**

Page 289-305

Hussein A. El-Naggar et al.

[View Article](#) | [PDF 924.11 K](#)

---

- 18 **A study on shrimp fishing quantities offered in some markets of El Basrah, Iraq**

Page 307-317

Tariq H. Y. Al-Maliky

[View Article](#) | [PDF 543.26 K](#)

---

- 19 **Assessment of some Physico-chemical parameters of the fish farm water in Abengourou, Côte d'Ivoire**

Page 319-343

Ali Sanou et al.

[View Article](#) | [PDF 913.32 K](#)

---

- 20 **Bacteria forming histamine and shelf life of sardine (*Sardina Pilchardus*) at different temperatures and storage time with emphasis on histopathological changes in skeletal musculature**

Page 345-360

Wafaa A. Mohamed et al.

[View Article](#) | [PDF 829.52 K](#)

---

- 21 **Identification of productive miter squid fishing grounds using a generalized additive model in Indonesian Fisheries Management Area 711**

Page 361-384

Hadi Puspito et al.

[View Article](#) | [PDF 1.22 MB](#)

- 
- 22 **[A comparative evaluation of the effects of pollution on selected organs of \*Oreochromis niloticus\* \(Linnaeus, 1758\) from two subtropical lakes with different pollution levels in Zimbabwe: Histopathological comparison](#)**  
Page 385-404  
I. Nyagura, et al.  
[View Article](#) | [PDF 659.36 K](#)
- 
- 23 **[The Prospects and Problems of Rural Household Based Aquaculture: A Study of Selected Fish Farming Households in Kerala, India](#)**  
Page 405-427  
Mathews Plamoottil; Pradeep Kumar B.  
[View Article](#) | [PDF 552.24 K](#)
- 
- 24 **[Diet of The North African Green Frog \*Pelophylax saharicus\* in The Wetlands of Edough \(north- eastern of Algeria\)](#)**  
Page 429-443  
Fafani Driouche; Nadia Ziane  
[View Article](#) | [PDF 931.28 K](#)
- 
- 25 **[Phytochemical variation, antioxidant and antidiabetic capacity of extracts of common brown seaweeds from the Red Sea, Egypt](#)**  
Page 445-462  
Mona M. Ismail et al.  
[View Article](#) | [PDF 961.52 K](#)
- 

Records Per Page 



Volume & Issue: Volume 26, Issue 5, September and October 2022 [XML](#)

---

- 26 **Lignocellulosic substrate as a low-cost effective inducer for production of hydrolytic cellulases by marine halophilic *Aspergillus ochraceus***  
Page 463-481  
Taghreed S. Alnusaire; Aida M. Farag  
[View Article](#) | [PDF 725.01 K](#)
- 
- 27 **Microbial Remediation of some Heavy Metals in Wastewaters of Lake Manzala, Egypt**  
Page 483-493  
Ahmed, I. Abd El-kader et al.  
[View Article](#) | [PDF 661.57 K](#)
- 
- 28 **Study of the production performance by releasing carp fingerlings at selected floodplain of Kotalipara Upazilla, Gopalganj, Bangladesh**  
Page 495-512  
Md. A. Halim  
[View Article](#) | [PDF 639.54 K](#)
- 
- 29 **Ecological notes on the Red-eyed reef crab *Eriphia sebana* (Shaw & Nodder, 1803) from the intertidal zone of Kadmat island, Lakshadweep, India**  
Page 513-518  
N. M. Naseem; Nihal Davood  
[View Article](#) | [PDF 677.3 K](#)
- 
- 30 **Effect of commercial Nisin levels on hygienic quality aspects of the frozen cultured Nile tilapia (*Oreochromis niloticus*) fillets**  
Page 519-533  
M.K. Arafat, et al.  
[View Article](#) | [PDF 458.82 K](#)
- 
- 31 **Treatment of MAS caused by *Aeromonas sobria* in European seabass**  
Page 535-556  
Magdy Khalil Soliman et al.  
[View Article](#) | [PDF 1.53 MB](#)
- 
- 32 **Age, Growth and mortality of the rabbit fish *Siganus rivulatus* (Forsskål 1775) from the Egyptian Mediterranean coast off Alexandria, Egypt**  
Page 557-569  
Mohammed G. Desouky et al.

[View Article](#) | [PDF 833.16 K](#)

---

33 **[Some biological measurements on the European sea bass in Bardawil Lagoon, Egypt](#)**

Page 571-593

Heba E. Abd Elnabi et al.

[View Article](#) | [PDF 1.54 MB](#)

---

34 **[Liver toxicity and geospatial analysis as integrated indices of water quality deterioration in Idku Lake, Egypt](#)**

Page 595-612

Aml S. Said et al.

[View Article](#) | [PDF 2.61 MB](#)

---

35 **[A comparative study of fish production from Lake Manzalah before and during the cleansing and development operations based on surplus production modeling approach](#)**

Page 613-624

Sahar F. Mehanna et al.

[View Article](#) | [PDF 813.07 K](#)

---

36 **[GC-MS analysis of Anadara uropigmelana and screening of its in vitro antioxidant and antiproliferative effect against MCF-7 cells](#)**

Page 625-650

Aalaa M. El-Sadat et al.

[View Article](#) | [PDF 944.94 K](#)

---

37 **[Yersinia ruckeri infection and pathogenicity of the cultured Oreochromis niloticus](#)**

Page 651-666

Ashraf A. Abd El Tawab et al.

[View Article](#) | [PDF 799.98 K](#)

---

38 **[Toxicological effect of Boric Acid and Cadmium Chloride on the Nile Tilapia, Oreochromis niloticus](#)**

Page 667-680

Rana Abdel Aliem et al.

[View Article](#) | [PDF 1.08 MB](#)

---

39 **[Herbivorous Reef Fishes of South Sinai Marine Protected Areas \(Gulf of Aqaba, Egypt\): Influence of Fisheries Management and Protection Status](#)**

Page 681-705

Khaled M. El-Haddad et al.

[View Article](#) | [PDF 1.75 MB](#)

- 
- 40 **[Do marine synbiotics decrease estradiol impacts in early-weaned European seabass \(\*Dicentrarchus labrax\*\) larvae?](#)**  
Page 707-727  
Ahmed Md. Salem et al.  
[View Article](#) | [PDF 1.21 MB](#)
- 
- 41 **[Viruses as Indicators of Fecal Pollution in Aquatic Environment](#)**  
Page 729-764  
Mohammed K. Rashed et al.  
[View Article](#) | [PDF 692.4 K](#)
- 
- 42 **[Stock structure based on truss network analysis, length-weight relationship and condition factor of \*Decapterus macrostoma\* \(Bleeker, 1851\) from two fishing areas, Red Sea, Egypt](#)**  
Page 765-777  
Yassein A. Osman et al.  
[View Article](#) | [PDF 863.16 K](#)
- 
- 43 **[Monitoring land use/land cover spatiotemporal changes and its implications on the productivity of Idku Lake, Egypt](#)**  
Page 779-796  
Aml S. Said et al.  
[View Article](#) | [PDF 2.03 MB](#)
- 
- 44 **[The potential role of marine macroalgae \*Padina pavonica\* on growth performance, histological status, and resistance of the rabbitfish \*Siganus rivulatus\* to \*Pseudomonas anguilliseptica\* bacteria](#)**  
Page 797-819  
Mohamed N. Monier et al.  
[View Article](#) | [PDF 1.28 MB](#)
- 
- 45 **[Antifungal activity of volatiles emitted from living cultures of \*Chlorella vulgaris\*, \*Desertifilum tharense\*, and \*Navicula arenaria\*](#)**  
Page 821-840  
Asmaa Lotfi Alallaf et al.  
[View Article](#) | [PDF 1.04 MB](#)
- 
- 46 **[Ichthyofauna of River Chirchik at Tashkent region, Uzbekistan](#)**  
Page 841-850  
B.G. Kamilov et al.  
[View Article](#) | [PDF 522.3 K](#)
-

- 47 **Evaluation of metallic pollution of north eastern Algerian coasts using the demosponge *Chondrilla nucula* Schmidt, 1862**  
Page 851-878  
Islem Zerarguia et al.  
[View Article](#) | [PDF 703.17 K](#)
- 
- 48 **Efficacy of Essential Oils (Thyme and Laurel) on Maintaining the Quality Indices and Nutritional Value of the Frozen *Tilapia* (*Oreochromis niloticus*) Fillets**  
Page 879-895  
Shaban A. El-Sherif et al.  
[View Article](#) | [PDF 562.23 K](#)
- 
- 49 **Investigating technical and scale efficiencies of the rice-shrimp system in the Mekong Delta, Vietnam**  
Page 897-909  
Vo Thanh Toan et al.  
[View Article](#) | [PDF 599.55 K](#)
- 
- 50 **Biological aspects of the blue swimming crab, *Portunus segnis* (Brachyura: Decapoda: Portunidae), inhabiting the Suez Canal Lakes, Egypt**  
Page 911-937  
Arwa Abd-Allah et al.  
[View Article](#) | [PDF 1.04 MB](#)

Records Per Page

25 ▼



Volume & Issue: Volume 26, Issue 5, September and October 2022 [XML](#)

---

51 **[Systematic notes on some cyprinid fishes collected from Kasargod district of Kerala, India](#)**

Page 939-954

Mathews Plamoottil; Regy Johnson

[View Article](#) | [PDF 1.18 MB](#)

---

52 **[Phytoplankton functional groups structures and ecological quality of tropical lake \(Taabo, Côte d'Ivoire\)](#)**

Page 955-971

Maboundou Camara et al.

[View Article](#) | [PDF 849.96 K](#)

---

53 **[Single and Combined Effects of Bisphenol A Exposure and Schistosoma mansoni Infection: Biochemical and Histological Implications on Biomphalaria alexandrina Snails](#)**

Page 973-1001

Sherin K. Sheir et al.

[View Article](#) | [PDF 1.26 MB](#)

---

54 **[Component Analysis and Antimicrobial Activity of the Plocamium cartilagineum Extract](#)**

Page 1003-1017

Selma Mabrouki et al.

[View Article](#) | [PDF 517.25 K](#)

---

55 **[A surveillance study on motile Aeromonas Septicemia \(MAS\) affecting cultured Litopenaeus vannamei shrimp from Suez Governorate, Egypt](#)**

Page 1019-1032

Ismail A. Eissa et al.

[View Article](#) | [PDF 754.7 K](#)

---

56 **[Spatio-temporal variations in conservative and non-conservative properties of the surface seawater along the Red Sea coast, Egypt](#)**

Page 1033-1046

Ahmed Salah-Tantawy et al.

[View Article](#) | [PDF 894.93 K](#)

---

57 **[Water pollution evaluation in Aswan, Egypt, utilizing biochemical and molecular indicators in aquatic Chironomidae larvae in natural and laboratory settings](#)**

Page 1047-1067

Zeinab Mohamed et al.

[View Article](#) | [PDF 1.3 MB](#)

---

58 **Temperature variability and distribution of fish along the Suez Gulf, Red Sea as a reason of climate change**

Page 1069-1087

Ali Wheida et al.

[View Article](#) | [PDF 1.1 MB](#)

---

59 **The prevention impact of the green algal extract against genetic toxicity and antioxidant enzyme alteration in the Mozambique tilapia**

Page 1103-1118

Wagdy K. B. Khalil et al.

[View Article](#) | [PDF 775 K](#)

---

60 **Effects of Dietary Marine Bacillus subtilis HS1 Probiotic, Chitosan Prebiotic and Two Marine Synbiotics Mixtures on the Growth and Oxidative Stress of the European Seabass (Dicentrarchus labrax) Larvae**

Page 1119-1138

Ahmed Md. Salem; Hassan A. Ibrahim

[View Article](#) | [PDF 1.47 MB](#)

---

61 **Anthropic Impacts on Aquatic Mangrove Ecosystems Worldwide**

Page 1139-1163

Ana Grijalva-Endara et al.

[View Article](#) | [PDF 985.83 K](#)

---

62 **The Effect of Stocking Density on Growth Performance and Water Quality of the Silver rasbora (Rasbora argyrotænia) Fry Reared in Plastic-lined Pond**

Page 1165-1176

Muhammad Hanif Azhar et al.

[View Article](#) | [PDF 717.29 K](#)

---

63 **Weekly fluctuation in phytoplankton macromolecular composition in response to environmental changes, in Eastern Harbour, Alexandria (Egypt)**

Page 1089-1102

Mona M. Ismail et al.

[View Article](#) | [PDF 448.65 K](#)

---

64 **Growth of Kappaphcus alvarezii in vertical method of multi-trophic system based on feeding rate**

Page 1197-1210

et al. Heriansah

[View Article](#) | [PDF 970.38 K](#)

---

65 **Morphometrics Diversity and Phenotypic Relationship of the Red Snapper (*Lutjanus gibbus*) in Northern Papua Waters**

Page 1211-1227

Ridwan Sala et al.

[View Article](#) | [PDF 615.19 K](#)

---

66 **Winter Kills in Farmed European Seabass (*Dicentrarchus labrax*): Co- infected with *Shewanella putrefaciens* and *Aeromonas veronii***

Page 1177-1195

Mahmoud Abou-Okada et al.

[View Article](#) | [PDF 1015.93 K](#)

---

67 **Antibacterial effect of rhodanine derivatives against bacteria isolated from the River Nile, Egypt**

Page 1229-1239

Saleh Ahmed Rabeh

[View Article](#) | [PDF 511.8 K](#)

---

68 **Susceptibility of El-Alamein coast to contamination and future prediction of erosion using geostatistical analyses**

Page 1241-1271

Ahmed El-Zeiny et al.

[View Article](#) | [PDF 2.64 MB](#)

---

69 **Structure Community of Phytoplankton as a Bioindicator of Water Quality in Situ Rawa Dongkal, East Jakarta, Indonesia**

Page 1273-1288

Muhammad Ridwan et al.

[View Article](#) | [PDF 401.69 K](#)

---

70 **Water Quality Assessment for Irrigation Purposes in a Semi-Arid Region: Wadi Baghai (North East of Algeria)**

Page 1289-1302

Yamama Nedjar et al.

[View Article](#) | [PDF 583.52 K](#)

---

71 **Contribution to the study of the digenean trematode *Clinostomum complanatum* parasite of the Nile Tilapia (*Oreochromis niloticus*) in breeding basins in southern Mauritania**

Page 1303-1315

I. M'bareck et al.

[View Article](#) | [PDF 684.41 K](#)

---

72 **[Antimicrobial Activities of a Novel Marine Streptomyces sp. MMM2 Isolated from EI- Arish coast, Egypt](#)**

Page 1317-1339

Mona A. Malash et al.

[View Article](#) | [PDF 1.24 MB](#)

---

73 **[Heavy Metals Bioaccumulation in Liver and Muscle Tissues of the Common Sole Fish \(Solea solea\) Inhabiting Damietta Fishing Harbor, Egypt](#)**

Page 1359-1374

Ali A. Al-Halani et al.

[View Article](#) | [PDF 730.55 K](#)

---

74 **[Immune Response and Antischistosomal Effect of the Thymoquinone Bioactive Compound from Nigella Sativa Loaded with Chitosan Nanoparticles](#)**

Page 1375-1392

Hend M. El-Menyawy et al.

[View Article](#) | [PDF 606.38 K](#)

---

75 **[Morphological characteristics of Rhinogobius brunneus \(Perciformes: Gobiidae\) in some water basins of Uzbekistan](#)**

Page 1393-1401

Quvatov Asqar Qoraqulovich et al.

[View Article](#) | [PDF 625.17 K](#)

---

Records Per Page  ▼

⏪ ⏩ 1 2 3 ⏪ ⏩



## The Effect of Stocking Density on Growth Performance and Water Quality of the Silver rasbora (*Rasbora argyrotaenia*) Fry Reared in Plastic-lined Pond

Muhammad Hanif Azhar<sup>1,2\*</sup>, Wildan Kurnia Pratama<sup>1</sup>, Darmawan Setia Budi<sup>1</sup>,  
Akhmad Taufiq Mukti<sup>3</sup>, Muhammad Brawijoyo Santanumurti<sup>3</sup>, Devrim Memiş<sup>2</sup>

- <sup>1</sup>. Study Program of Aquaculture, Department of Health and Life Sciences, School of Health and Life Sciences (SIKIA), Universitas Airlangga, Jalan Wijaya Kusuma 113, 68423 Banyuwangi, Indonesia
- <sup>2</sup>. Department of Aquaculture, Faculty of Aquatic Sciences, Istanbul University, Istanbul 34134, Turkey
- <sup>3</sup>. Department of Aquaculture, Faculty of Fisheries and Marine, Universitas Airlangga, Surabaya 60115, Indonesia

\*Corresponding Author: hanifazhar@fpk.unair.ac.id

### ARTICLE INFO

#### Article History:

Received: Feb. 7, 2022

Accepted: Sept. 13, 2022

Online: Oct. 27, 2022

#### Keywords:

Silver rasbora,  
*Rasbora argyrotaenia*,  
growth,  
rearing,  
density

### ABSTRACT

Silver rasbora is one of the local fish species in Indonesia that gains a high public demand. To meet the increasing market demand, it is necessary to implement an optimal silver rasbora rearing system. The objective of the study was to determine the best stocking density for the growth and water quality of silver rasbora (*Rasbora argyrotaenia*) fry reared in plastic-lined ponds. The fries utilized had a mean initial body length of  $1.85 \pm 0.5$  cm and body weight of  $0.22 \pm 0.05$  g. The experiment lasted for 40 days and was carried out in a 500 L plastic-lined pond, with dimensions of 1 x 1 x 1 m and five treatments were evaluated (100, 150, 200, 250, 300 fish m<sup>-2</sup>). The feed used was commercial floating feed with 39% protein content, and feeding was conducted three times a day (08.00 AM, 12.00 PM, and 16.00 PM) on ad satiation basis. According to the results, the stocking density showed a significant effect on body weight, final biomass, and feed conversion ratio (FCR) in all treatments. T1 resulted in the highest body weight with a value of  $0.71 \pm 0.02$  g, and the lowest FCR with a value of  $0.90 \pm 0.01$ . While, T5 resulted in the highest final biomass with a value of  $102.81 \pm 5.62$  g. The pH concentration was the only parameter that showed a significantly different value, with the lowest value in T5 ( $7.02 \pm 0.0$ ). The best water quality parameter in T5 showed values of  $27.03 \pm 0.1^{\circ}\text{C}$  (temperature),  $7.02 \pm 0.03$  (pH),  $6.79 \pm 0.21$  mgL<sup>-1</sup>(DO),  $2.81 \pm 0.89$  mgL<sup>-1</sup> (nitrate),  $0.18 \pm 0.02$  mgL<sup>-1</sup> (nitrite), and  $0.05 \pm 0.01$  mgL<sup>-1</sup> (TAN). A stocking density of 300 fish showed suitable conditions for rearing silver rasbora fries in plastic-lined pond.

### INTRODUCTION

Silver rasbora (*Rasbora argyrotaenia*), a freshwater fish from the family Cyprinidae is commonly found in Southeast Asia, including Indonesia, and lives in flowing (rivers) to stagnant waters (rice fields and lakes) (Nelson, 2006; Said & Triyanto, 2013). Silver rasbora has a high economic value in Indonesia because it can be used as both a food consumption and an ornamental fish (Said *et al.*, 2011). Previous data states that in 2018,

rasbora production in Indonesia reached 9,523.96 tons at 60,000 IDR or 4.19 USD (**KKP, 2021; Rukayah & Lestari, 2021**). It is one of the popular freshwater fish for its delicious taste and high nutritional content (protein 47,54% and fat 2,9%) (**Said *et al.*, 2010; Sogandi *et al.*, 2019**), but unfortunately, rasboras cannot be optimally cultivated and still rely on wild harvests (**Budi *et al.*, 2020**). According to previous research, rasbora is an endangered species due to overfishing (**Muchlisin *et al.*, 2010**) and environmental pollution such as the use of pesticides, which affects the availability of rasbora in nature; therefore, aquaculture regulations should be carried out to prevent rasboras from extinction (**Prastika *et al.*, 2021**).

Since 2010, there has been an increase in efforts to cultivate rasbora, and several studies on its development have been conducted, including habitat characteristics, feeding habits and conservation systems (**Sulastri *et al.*, 2010; Sulistiyarto, 2013; Rosadi *et al.*, 2014**), reproductive aspects (**Said *et al.*, 2011**), mtDNA assay (**Kusuma *et al.*, 2017**), spawning induction using hormones (**Ningrum *et al.*, 2019**), sperm cryopreservation to support their breeding efforts (**Muchlisin *et al.*, 2020; Eriani *et al.*, 2022**); and also the use of different stocking densities in larval stages (**Budi *et al.*, 2020**). The majority of fish harvest has been currently directed to fulfill market demand. Hence, it is necessary to conduct appropriate fry-rearing techniques to fulfill the market demand. Information about the appropriate stocking density is very important to increase the productivity of silver rasbora fries and ensure a continuous supply.

Stocking density has a direct impact on biota growth, water quality, and production efficiency (**de Oliveira *et al.*, 2012; Tibile *et al.*, 2016**). According to previous studies, excessive stocking density reduces the oxygen content in waters causing the fish to become stressed, susceptible to disease, and witness a decrease in both growth performance, survival rate and feed efficiency (**Martos-Sitcha *et al.*, 2020; Muchlisin *et al.*, 2021**). Other studies stated that high stocking densities can cause high carbon dioxide levels due to fish respiration, thereby lowering pH and increasing acidity in the waters (**Hong *et al.*, 2019**). Furthermore, studies related to stocking density are important, because if the density is too high, it will cause a size disparity among fish, competition for food, and ammonia excretion in the waters, in addition to predation (**Ronald *et al.*, 2014; Tibile *et al.*, 2016**). However, if it is not enough, then the profit from aquaculture will not be optimal (**Szczepkowski *et al.*, 2011**). Therefore, this study aimed to obtain an appropriate method to produce quality fish and ensure a continuous supply of fries in silver rasbora hatchery activities using plastic-lined ponds by determining the effect of different stocking densities on growth parameters and water quality.

## MATERIALS AND METHODS

This study was conducted at the School of Health and Life Sciences (SIKIA) Banyuwangi Campus Laboratory of Airlangga University, Banyuwangi, Indonesia.

### **Fish stocking**

Silver rasbora fries were obtained from the Technical Services Unit of Freshwater Aquaculture Development in Umbulan (Pasuruan, East Java). Silver rasbora fries specimens were kept for 1 week in an adaptation pond before the experiment to minimize stress. During the adaptation period, the fries were fed the same type of commercial floating feed and at the same time (three times a day at 08.00 AM, 12.00 PM, and 16.00 PM) as will be carried out in the experimental process. However, before being placed in the experimental pond, the fries were selected and weighed individually, based on their length and size. The fries samples had a mean initial weight of  $0.22 \pm 0.05$  g and a mean length of  $1.85 \pm 0.05$  cm.

### **Experimental Design**

A completely randomized experimental design was used in this study. This study was conducted for 40 days, using 5 treatments of different stocking density levels (100, 150, 200, 250, and 300 fishes  $m^{-2}$ ) with 4 replicates. Silver rasbora fries were reared semi-outdoor using plastic-lined ponds measuring  $1 \times 1 \times 1$  m, with a water volume of 500 L and aerated with an air blower ( $140 \text{ L min}^{-1}$ ).

During the experiment, the feed was administered three times a day (at 08.00 AM, 12.00 PM, and 16.00 PM) using floating commercial feed with a protein content of 39%, lipid (5%), crude fiber (4%), moisture (10%), and ash (11%) (PF-500 Matahari Sakti Ltd Indonesia), with ad satiation basis. The feed was weighed before and after being given, and feeding was carried out gradually and stopped if the fish stops responding to the feed. The sampling of water quality and fish growth was conducted before morning feeding time and siphon process (carried out once every 10 days by changing the water to a maximum of 10% of the total volume of water in the fishpond).

### **Water Quality Measurement**

Temperature and pH were measured using a pH meter (Lutron YK2001) and DO (Dissolved oxygen) was measured using a DO meter (YSI Pro20). Nitrite and nitrate were measured using a test kit (Merck MColorTest 114658 and Hanna Instruments HI3874) and TAN (Total Ammonia Nitrogen) was analyzed using a spectrophotometer based on the APHA (1999) procedures. In addition, measurement of TAN, nitrite, and nitrate concentrations was carried out in the water quality analysis laboratory. The water samples were collected directly from four points in the water body using a 250 ml sample bottle and then sent to the laboratory for analysis. Meanwhile, the temperature and DO measurements were carried out in situ.

### **Observation Parameters**

Temperature, pH, and DO were measured daily (every morning) while TAN, Nitrite, and Nitrate were measured every 10 days. Water samples were taken in the morning before feeding time and the production parameters observed include Final Biomass (FB), Weight Gain (WG), Survival Rate (SR), Total amount of given feed (TF), and Feed Conversion Ratio (FCR). Furthermore, it can be formulated  $FCR = TF/WG$ ;  $WG = \text{Mean}$

final weight – mean initial weight; SR (%) = (Final number of fish/ Initial number of fish) x 100 (Sarker *et al.*, 2016). Growth sampling was also performed every 10 days in the morning before feeding and after measuring water quality by taking 10% of biomass (T1: 10; T2: 15; T3: 20; T4: 25; T5: 30 fish) from each treatment in the experimental pond and measuring the lengths and weights individually.

### Data analysis

The data obtained were processed using Microsoft Excel 2013 and then tested for normality and homogeneity using the Kolmogorov-Smirnov and Levene's test, followed by the analysis of the fish growth data using Analysis of Variance (ANOVA) at a 95% confidence interval ( $\alpha = 0.05$ ). Subsequently, if the results were statistically significant ( $P < 0.05$ ), the 95% Tukey Test was carried out. Water quality parameters were measured using repeated-measures ANOVA with two factors (water quality and time). Bonferroni test was used when differences between time and treatments were detected ( $P < 0.05$ ). Statistical analysis was also performed using IBM-SPSS version 18.

## RESULTS

### 1. Growth Performance Parameter

The use of different stocking densities had an effect on the growth parameters of silver rasbora seeds. The results were showed in Table 1.


**Table 1.** Growth performance of Silver rasbora fries for 40 days rearing period using the different stocking densitie

Variable	Treatment (Different Stocking Density)				
	T1 (100)	T2 (150)	T3 (200)	T4 (250)	T5 (300)
FB (g tank <sup>-1</sup> )	69.99±1.31 <sup>a</sup>	73.36±3.15 <sup>ab</sup>	78.91±1.73 <sup>b</sup>	90.58±3.68 <sup>c</sup>	102.81±5.62 <sup>d</sup>
TF (g tank <sup>-1</sup> )	45.43±0.01 <sup>a</sup>	47.75±0.02 <sup>a</sup>	65.56±0.01 <sup>b</sup>	79.15±0.01 <sup>c</sup>	97.83±0.02 <sup>d</sup>
FCR	0.90±0.01 <sup>a</sup>	0.93±0.03 <sup>a</sup>	1.20±0.01 <sup>b</sup>	1.20±0.14 <sup>b</sup>	1.30±0.01 <sup>b</sup>
WG (g)	50.3±1.87 <sup>a</sup>	51.6±2.37 <sup>ab</sup>	55.8±1.36 <sup>b</sup>	65.88±2.19 <sup>c</sup>	77.09±2.94 <sup>d</sup>
SR (%)	100±0.00	100±0.00	100±0.00	100±0.00	100±0.00


FB: Final Biomass; WG: Weight Gain; TF: Total amount of Feed given; FCR: Feed Conversion Ratio; SR: Survival Rate. Values are mean ± SD. Superscript letters denote significant differences ( $P < 0.05$ ) between treatments

The ANOVA test showed that stocking density had a significant effect ( $P < 0.05$ ) on production parameters (Final Biomass (FB), total amount of food given (TF), and Feed Conversion Ratio (FCR)) but had no significant effect ( $P > 0.05$ ) to Survival Rate (SR). The results of the study showed that the highest values of FB, TF, and FCR were found in T5 treatment using a stocking density of 300 fish m<sup>-2</sup> silver rasbora fries. Tukey test

showed that the FB and TF values in treatment T5 (300 fish m<sup>-2</sup>) were significantly higher than in other treatments. The FCR value in treatment T5 showed a significantly higher than in treatment T1 and T2 but not significant in treatment T3 and T4. The SR value showed no significant results among all the treatments (Table 1).



**Figure 1.** Body weight of Silver rasbora fries at 0, 10, 20, 30, and 40 days with different stocking density treatments. Superscript letters differ significantly ( $P < 0.05$ ) between all treatments



**Figure 2** Total length of Silver rasbora fries at 0, 10, 20, 30, and 40 days with different stocking density treatments. Total length of silver rasbora showed did not significantly ( $P > 0.05$ ) among all treatments

The results showed that the use of stocking density affected the weight of the fish (Figs. 1,2). Fish weight during the rearing period showed a significant value ( $P < 0.05$ ) between all treatments. However, the total length did not show a significant value

( $P > 0.05$ ) between all treatments. This indicates the higher the stocking density used, the lower the fish weight.

### Water Quality Parameters

The results of water quality analysis using a one-way repeated measure ANOVA showed in Table 2. The results were not significantly different ( $P > 0.05$ ) between the use of stocking density and water quality in the form of TAN, nitrite, nitrate, DO, and temperature on rearing media of silver rasbora. However, the use of stocking density gave a significant effect ( $P < 0.05$ ) on water quality in the form of pH value. The pH value between treatments showed a significant value ( $P < 0.05$ ) between T5 and T1, T2, and T4 treatments. However, it was not significant ( $P > 0.05$ ) from the T3 treatment. In general, water quality parameters during the rearing period of silver rasbora fry can be seen in Table 2.

**Table 2.** Water quality of Silver rasbora fry for 40 days rearing period using the different stocking densities

Variable	T1 (100)	T2 (150)	T3 (200)	T4 (250)	T5 (300)	P value <sup>†</sup>
TAN (mg L <sup>-1</sup> )	0.03±0.01	0.04±0.02	0.04±0.02	0.05±0.02	0.05±0.01	0.329
Nitrite (mg L <sup>-1</sup> )	0.09±0.01	0.18±0.03	0.28±0.01	0.18±0.08	0.18±0.02	0.317
Nitrate (mg L <sup>-1</sup> )	2.79±1.41	3.10±0.71	3.22±0.65	2.69±0.75	2.81±0.89	0.967
DO (mg L <sup>-1</sup> )	7.06±0.14	7.02±0.16	7.01±0.14	7.07±0.07	6.79±0.21	0.340
pH	7.19±0.07 <sup>bc</sup>	7.24±0.07 <sup>c</sup>	7.07±0.03 <sup>a</sup>	7.16±0.11 <sup>b</sup>	7.02±0.03 <sup>a</sup>	0.000
Temperature (°C)	27.5±0.4	27.4±0.4	27.5±0.1	27.6±0.1	27.3±0.1	0.114

TAN: Total ammonia nitrogen; DO: Dissolved oxygen. Values are mean ± SD. Superscript letters denote significant differences between treatments (Bonferroni test,  $P < 0.05$ ). <sup>†</sup>Results from one-way repeated measure ANOVA (Density x Times) and Bonferroni test.

### Inorganic Nitrogen

The results of the sampling every 10 days which were analyzed using a one-way repeated measure ANOVA showed that were not significantly different ( $P > 0.05$ ) between the use of stocking density and the concentration of inorganic nitrogen (TAN, nitrite, nitrate) on silver rasbora rearing media. The value of TAN concentration tends to increase at the beginning of the rearing period and tends to decrease at the end of the rearing period. While the value of Nitrite concentration tends to be high at the beginning of the maintenance period and tends to fluctuate until the end of the rearing period. Nitrate concentration values tend to increase until the end of the maintenance period in all treatments.

## DISCUSSION

The use of different stocking densities significantly affected the growth parameters of silver rasbora fry during the experimental period. The weight of silver rasbora fry

decreased with the increase in the stocking density of silver rasbora fry. The weight of the fish decreased with the increase in stocking density and this result is consistent with several studies using fish species from the family Cyprinidae (*Labeo bata*, *Puntius binotatus*, *Barbonymus gonionotus*) which showed a negative correlation between the stocking density and fish growth. In addition, the use of high stocking density could reduce the water quality, increase the competition for feed, and affect the aggressiveness of fish (**Rahman et al., 2015; Samad et al., 2016; Hayat et al., 2018**). However, several studies showed a positive correlation between stocking density and fish growth, such as the Sciaenidae family (*Argyrosomus japonicus* and *Argyrosomus regius*) (**Pirozzi et al., 2009; Millán-Cubillo et al., 2016**). This result is possible with animals who prefer to live in groups to avoid stress (**de las Heras et al., 2015**). Studies on stocking density must take into consideration a variety of factors, including species selection, social interactions, water quality, and environmental conditions (**Salari et al., 2012; Arifin et al., 2017**).

Furthermore, the stocking density also affected the FCR and the total amount of given feed as the higher the stocking density, the higher the FCR and TF value. However, the use of different stocking densities in the rearing of silver rasbora fry was still ideal because it showed an FCR value of less than 1.5 in all treatments. Feed conversion and efficiency are very important indicators of efficiency in the use of the feed provided because feed prices tend to rise (**Craig and Khun, 2017; Fry et al., 2018**). Moreover, the FCR value of less than 2 was still considered good in aquaculture industries (**Bag et al., 2016**). The use of optimal stocking densities in aquaculture is very necessary to reduce stress on fish and potential economic losses (**Arifin et al., 2017**).

The stocking density treatments did not affect the SR of silver rasbora fry until the end of the experimental period. The high SR in all treatments could be influenced by several factors, such as the species used and the influence of environmental factors (**Samad et al., 2016**). Stocking density is one of the most important factors in aquaculture activities because it directly affects SR, growth performance, water quality, and production efficiency in fish or shrimp farming (**Chakraborty and Mirza, 2007; Rahman and Verdegem, 2010; de Oliveira et al., 2012**). Several similar studies in the family Cyprinidae also showed results where stocking density did not affect SR (**Said and Triyanto, 2013; Rahman et al., 2015; Hayat et al., 2018**).

Silver rasbora is a fish of the genus *Rasbora* which is found on many islands in Indonesia (**Kusuma et al., 2017**). The result of water quality parameters in several *Rasbora* studies is shown in Table 3.

Based on observations made during the maintenance period, showed that the water quality parameters in the form of TAN, Nitrite, and Nitrate values during the maintenance period were higher than in the literature (Table 3). This could be caused by excess feed residue and the use of stocking density (**Hayat et al., 2018**). Concentrations of TAN, Nitrite, and Nitrate in the rearing medium fluctuated but increased until the end

of the rearing period. The increased concentration of TAN, Nitrite, and Nitrate in rearing ponds could be due to the higher organic matter content in the rearing media from uneaten feed and fish feces (Jiménez-Montealegre *et al.*, 2002; Torres-Beristain *et al.*, 2006; Bag *et al.*, 2016).

**Table 3.** Water quality parameters in the genus of *Rasbora*

Variable	During Experiment	Acceptable Range (Literature)	Notes
TAN (mg L <sup>-1</sup> )	0.029-0.061	0.010-0.013 <sup>c</sup>	Larva rearing
Nitrite (mg L <sup>-1</sup> )	0.101-0.310	0.002-0.012 <sup>a</sup>	Natural habitat
Nitrate (mg L <sup>-1</sup> )	1-3.8	0.398-3.272 <sup>a</sup>	Natural habitat
pH	6.5-8.0	6.7-8.8 <sup>ab</sup>	Natural habitat
DO (mg L <sup>-1</sup> )	6.2-7.2	6.2-7.5 <sup>ac</sup>	Natural habitat; larvae rearing
Temperature (°C)	27.1-27.7	24.5-32.0 <sup>d</sup>	Adaptation test

<sup>a</sup>Sentosa and Djumanto, (2010) ; <sup>b</sup>Sulastri *et al.*, (2010); <sup>c</sup> Budi *et al.*, (2020); <sup>d</sup> Said and Triyanto, (2013)

The siphoning process that was carried out every 10 days during the rearing period in all treatments did not affect the concentration of TAN, Nitrite, and Nitrate in the rearing pond. Furthermore, the fluctuating conditions of TAN, Nitrite, and Nitrate concentrations during the rearing period indicated the occurrence of nitrification and denitrification by natural bacteria found in rearing ponds (Durborow *et al.*, 1997; Turker *et al.*, 2003).

## CONCLUSION

Differences in stocking density significantly affected the growth performance of BW, FB, TF, FCR, WG, and water quality (pH value) of silver rasbora (*Rasbora argyrotænia*) fry in plastic-lined ponds. A stocking density of 100 fish 500 L<sup>-1</sup> resulted in optimal growth of the fry. However, based on all production parameters and overall water quality, the use of a stocking density of 300 fish 500 L<sup>-1</sup> still showed suitable conditions for cultivation activities with an FCR value of less than 1 and optimal water quality for rearing silver rasbora fries in plastic-lined ponds.

## AUTHORS' CONTRIBUTIONS

All authors contributed to the conception and design of the study. Wildan Kurnia Pratama and Darmawan Setia Budi prepared the materials, collected data, and analyzed the results. Muhammad Hanif Azhar wrote the first draft of the manuscript and Ahmad Taufiq Mukti, Devrim Memiş, and Muhammad Brawijoyo Santanumurti commented on previous versions of the manuscript.



## ACKNOWLEDGMENT

The authors are thankful that UPT PBAT Umbulan, Pasuruan and School of Health and Life Sciences (SIKIA) Airlangga University in Banyuwangi for providing us with research materials, areas and laboratories to conduct this research.

## CONFLICT OF INTEREST

The authors declare that there is no conflict of interest.

## REFERENCES

- Al Adawiyah, L.; Sulmartiwi, L.; Bodur T. and Budi D.S.** (2019). Induction of spermiation using Ovaprim™ with topical gill method in the silver rasbora (*Rasbora argyrotaenia*). *Theriogenology.*, 126: 172–176.
- APHA** (1999). Standard methods for the examination of the water and wastewater, 20th Edition, American Public Health Association, Washington, DC. 1268pp.
- Arifin, O.Z.; Subagja, J.; Prakoso, V.A. and Suhud E.H.** (2017). Effect of stocking density on growth performance of domesticated barb (*Barbonymus balleroides*). *Indonesia Aquac J.*, 12: 1-6.
- Bag, N.; Moulick, S. and Mal, B.C.** (2016). Effect of stocking density on water and soil quality, growth, production and profitability of farming Indian major carps. *Indian J Fish.*, 63: 39-49.
- Budi, D.S.; Rizkyan, I. and Prayogo, P.** (2020). The effect of stocking density on the survival and growth of silver rasbora (*Rasbora Argyrotaenia*) larvae. *Polish J Nat Sci.*, 35: 87–95.
- Chakraborty, B.K. and Mirza, M.J.A.** (2007). Effect of stocking density on survival and growth of endangered bata, *Labeo bata* (Hamilton-Buchanan) in nursery ponds. *Aquaculture.*, 265: 156–162.
- Craig, S. and Khun, D.D.** (2017). Understanding Fish Nutrition, Feeds, and Feeding. Virginia, Tech-Publication.
- de las Heras, V.; Martos-Sitcha, J.A.; Yúfera, M. and Mancera, J.M.** (2015). Influence of stocking density on growth, metabolism, and stress of thick-lipped grey mullet (*Chelon labrosus*) juveniles. *Aquaculture.*, 448:29–37.
- de Oliveira, E.G.; Pinheiro, A.B.; de Oliveira, V.Q.; da Silva Júnior, A.R.M.; de Moraes, M.G.; Rocha, I.R.C.B.; de sousa, R.R. and Costa, F.H.F.** (2012). Effects of stocking density on the performance of juvenile pirarucu (*Arapaima gigas*) in cages. *Aquaculture.*, 370-371:96-101.
- Durborow, R.M.; Crosby, D.M. and Brunson, M.W.** (1997). Nitrite in fish pond, Southern Regional Aquaculture Center, SRAC Publ, No. 463.
- Eriani, K.; Mustaqim, M.; Hasri, I.; Amalia, R.; Al Azhar.; Nur, F.M. and Muchlisin, Z.A.** (2022). Effect of fetal bovine serum on the sperm quality of depik (*Rasbora tawarensis*) after short-term cryopreservation. *Indian J. Anim. Res.*, 56(7): 843-847.
- Fry, J.P.; Mailloux, N.A.; Love, D.C.; Milli, M.C. and Cao, L.** (2018). Feed conversion efficiency in aquaculture: Do we measure it correctly? *Environ Res Lett.*, 13.2: 024017.

- Hayat, M.; Nugroho, R.A. and Aryani, R.** (2018). Influence of different stocking densities on the growth, feed efficiency, and survival of majalaya common carp (*Cyprinus carpio* linnaeus 1758). *F1000Research.*, 7: 1–10.
- Hong, J.; Chen, X.; Liu, S.; Fu, Z.; Han, M.; Wang, Y.; Gu, Z. and Ma, Z.** (2019). Impact of fish density on water quality and physiological response of golden pompano (*Trachinotus ovatus*) flingerlings during transportation. *Aquaculture.*, 507: 260-265.
- Jiménez-Montealegre, R.; Verdegem, M.C.J.; Van Dam, A. and Verreth, J.A.J.** (2002). Conceptualization and validation of a dynamic model for the simulation of nitrogen transformations and fluxes in fish ponds. *Ecol. Modell.*, 147: 123–152.
- Keshavanath, P.; Gangadhar, B.; Ramesha, J.; Priyadarshini, M.; Van Dam, A.A.; Verdegem, M.C.J. and Beveridge, M.C.M.** (2015). Impact of substrates and fish stocking densities on growth and production of the Indian major carp, *Labeo rohita* (HAM.). *J. Aqua. Trop.*, 30: 1-14.
- Kottelat, M.; Whitten, A.; Kartikasari, A.N. and Wirjoatmojo, S.** (1993). Freshwater fishes of Western Indonesia and Sulawesi: Ikan Air Tawar Indonesia Bagian Barat dan Sulawesi, Periplus, Hongkong. 428pp.
- KKP (Kementerian Kelautan dan Perikanan).** (2021). Produksi Perikanan. <https://statistik.kkp.go.id/home.php?m=total&i=2#panel-footer> (Accessed 12 December 2021).
- Kusuma, W.E.; Samuel, P.D.; Wiadnya, D.G.R.; Hariati, A.M. and Kumazawa, Y.** (2017). Complete mitogenome sequence of *Rasbora argyrotaenia* (Actinopterygii: Cyprinidae). *Mitochondrial DNA Part B. Resour.*, 2: 373–374.
- Lukman.** (2017). Perkembangan pemanfaatan dan penelitian ikan bada (*Rasbora argyrotaenia*) di danau maninjau. *Warta Iktiologi.*, 1: 24-27
- Martos-Sitcha, J.A.; Mancera, J.M.; Prunet, P. and Magnoni, L.J.** (2020). Welfare and stressors in fish: challenges facing aquaculture. *Frontiers in physiology.*, 11, 162.
- Millán-Cubillo, A.F.; Martos-Sitcha, J.A.; Ruiz-Jarabo, I.; Cárdenas, S. and Mancera, J.M.** (2016). Low stocking density negatively affects growth, metabolism, and stress pathways in juvenile specimens of meagre (*Argyrosomus regius*, Asso 1801). *Aquaculture.*, 451: 87–92.
- Muchlisin, Z.A.; Musman, M. and Siti Azizah, M.N.** (2010). Length- weight relationships and condition factors of two threatened fishes, *Rasbora tawarensis* and *Poropuntius tawarensis*, endemic to Lake Laut Tawar, Aceh Province, Indonesia. *J. Appl. Ichthyol.*, 26: 949-953.
- Muchlisin, Z.A.; Sarah, P.I.; Aldila, D.F.; Eriani, K.; Hasri, I.; Batubara, A.S.; Nur, F.M.; Mustaqim, M.; Mutmainnah, C.R.; Abinawanto, A. and Wilkes, M.** (2020). Effect of Dimethyl sulfoxide (DMSO) and egg yolk on sperm motility, fertility and hatching rates of depik *Rasbora tawarensis* (Pisces: Cyprinidae) eggs after short-term cryopreservation. *Aquac. Res.*, 51: 1700–1705.
- Muchlisin, Z.A.; Indrianti, Y.S.; Dewiyanti, I.; Lisra, F.; Yanti, R.Z.; Nur, F.M. and Batubara, A.S.** (2021). Effect of stocking density of the growth performance, survival rate and feed utilization of the eels *Anguilla bicolor* (pisces: anguillidae) Larvae. *IOP Conf. Ser. Earth Environ. Sci.*, 674: 012086.

- Nelson, J.S.** (2006). *Fishes of the World* 4<sup>th</sup> edition. John Willey & Sons. Inc, New Jersey. 622pp.
- Ningrum, D.R.K.; Budi, D.S. and Sulmartiwi, L.** (2019). Induksi pemijahan ikan wader pari (*Rasbora argyrotaenia*) menggunakan Ovaprim<sup>TM</sup> dengan dosis berbeda. *Depik.*, 8: 117–124.
- Pirozzi, I.; Booth, M.A. and Pankhurst, P.M.** (2009). The effect of stocking density and repeated handling on the growth of juvenile mulloway, *Argyrosomus japonicus* (Temminck & Schlegel 1843). *Aquac. Int.*, 17: 199–205.
- Prastika, A.B.; Sulmartiwi, L. and Lutfiyah, L.** (2021). Effect of sublethal dose organophosphate pesticides on embryogenesis and hatching rate of silver rasbora eggs (*Rasbora argyrotaenia*). *IOP Conf. Ser. Earth Envir. Sci.*, 679: 012065.
- Rahman, M.; Ferdous, Z.; Mondal, S. and Amin M** (2015.) Effects of stocking densities on growth and survival of Thai Sharpunti (*Barbonymus gonionotus*) in earthen ponds. *J. Bangladesh Agric. Univ.*, 9: 327–338.
- Rahman, M.M. and Verdegem, M.** (2010). Effects of intra- and interspecific competition on diet, growth and behaviour of *Labeo calbasu* (Hamilton) and *Cirrhinus cirrhosus* (Bloch). *Appl. Anim. Behav. Sci.*, 128: 103–108.
- Ronald, N.; Gladys, B. and Gasper, E.** (2014). The effects of stocking density on the growth and survival of Nile tilapia (*Oreochromis niloticus*) fry at Son Fish Farm, Uganda. *J. Aquac. Res. Development.*, 5: 1-7.
- Rosadi, E.; Yuli, H.E.; Setyohadi, D. and Bintoro, G.** (2014). Distribution, composition, and abiotic environment of silver rasbora (*Rasbora argyrotaenia* Blkr) Fish in Upstream Areas of Barito Watershed, South Kalimantan. *J. Environ Ecol.*, 5:117-131.
- Rukayah, S. and Lestari, W.** (2021) Upaya konservasi species asli melalui kajian reproduksi dan lingkungan ikan wader pari (*Rasbora argyrotaenia* Blker, 1854) di Waduk PB Soedirman Banjarnegara. *Prosiding SNPBS.*, 118-129
- Said, D.S.; Haryani, G.S.; Lukman.; Triyanto.; Mayasari, N.; Hamdani, A.; Sutrisno. and Sari, L.** (2010). Perkembangan ikan bada (*Rasbora Argyrotaenia*) Danau Maninjau-Sumatera Barat Pada Habitat Ex-Situ. *Prosiding Seminar Nasional Limnologi V.*
- Said, D.S.; Triyanto.; Lukman.; Sutrisno. and Hamdani, A.** (2011). Aspek biologi ikan bada (*Rasbora argyrotaenia*) di danau maninjau, sumatera barat. *Prosiding Forum Nasional Pemacuan Sumber Daya Ikan.*, III: 1-10
- Said D.S. and Triyanto.** (2013). Ketahanan hidup dan kemampuan tumbuh ikan bada (*Rasbora argyrotaenia*) pada suhu pemeliharaan bervariasi. *Limnotek.*, 20(2): 191-199.
- Said, D.S. and Hidayat.** (2015). 101 Ikan Hias Air Tawar, LIPI Press, Jakarta. 260pp.
- Salari, R.; Saad, C.; Kamarudin, M. and Zokaefar, H.** (2012). Effects of different stocking densities on tiger grouper juvenile (*Epinephelus fuscoguttatus*) growth and a comparative study of the flow-through and recirculating aquaculture systems. *African J. Agric. Res.*, 7: 3765–3771.
- Samad, M.A.; Khatun, A.; Reza, M.S.; Asrafuzzaman, Md. and Ferdaushy, M.H.** (2016). Effects of stocking density on growth, survival and production of mirror carp (*Cyprinus carpio* var. *specularis*) spawn in nursery pond. *Asian J. Med. Biol. Res.*, 2: 429–435.

- Sarker P.K.; Kapuscinski A.R.; Lanois A.J.; Livesey E.D.; Benhard, K.P. and Coley, M.L.** (2016) Towards sustainable aquafeeds: Complete substitution of fish oil with marine microalga *Schizochytrium* sp. improves growth and fatty acid deposition in juvenile Nile tilapia (*Oreochromis niloticus*). PLoS One 11: e0156684.
- Sentosa, A.A. and Djumanto.** (2010). Habitat Pemijahan Ikan Wader Pari (*Rasbora lateristriata*) di Sungai Ngrancah, Kabupaten Kulonprogo. Jurnal Iktiologi Indonesia., 10(1): 55-63.
- Sogandi.; Sanjaya, R.E.; Baity, N. and Syahmani.** (2019). Identification of nutritional content and profiles of amino acid from seluang fish (*Rasbora* sp.). Penelitian Gizi dan Makanan, 42: 73-80.
- Sulastri, Hartoto, D.I.; Yuniarti, I. and Nasution, S.H.** (2010). Karakteristik habitat, kebiasaan makan, dan sistem konservasi ikan bada *Rasbora Argyrotaenia* di Danau Maninjau. Pros Seminar Nasional Ikan., VI: 487-49
- Sulistiyarto, B.** (2013). Hubungan antara kelimpahan ikan saluang (*rasbora argyrotaenia* blkr) dengan populasi fitoplankton di dataran banjir Sungai Rungan Kalimantan Tengah. Jurnal Ilmu Hewani Tropika., 2: 27-30.
- Suryani, S.A.M.P; Wirawan, I.G.P.; Dwiyani, R. and Sritamin, M.** (2021). Genetic diversity and differentiation of cytochrome oxidase subunit I (COI) gene of *Rasbora lateristriata* bleeker in different habitat. IOP Conf. Ser. Mater. Sci. Eng. 1098: 052043.
- Szczepkowski, M.; Szczepkowska, B. and Piotrowska, I.** (2011). Impact of higher stocking density of juvenile Atlantic sturgeon, *Acipenser oxyrinchus Mitchell*, on fish growth, oxygen consumption, and ammonia excretion. Fish. Aquat. Life., 19(2): 59-67.
- Tibile, R.M.; Sawant, P.B.; Chada, N.K.; Lakra, W.S.; Prakash, C.; Swain S. and Bhagwati, K.** (2016). Effect of stocking density on growth, size variation, condition index and survival of discus, *Symphysodon aequifasciatus* Pellegrin, 1904. Turkish J Fish Aquat Sci., 16: 455–462.
- Torres-Beristain, B.; Verdegem, M.; Kerepeczki, E. and Verreth J.** (2006). Decomposition of high protein aquaculture feed under variable toxic conditions. Water Res., 40: 1341–1350.
- Turker, H.; Eversole, A.G. and Brune, D.E.** (2003). Comparative Nile tilapia and silver carp filtration rates of partitioned aquaculture system phytoplankton. Aquaculture., 220: 449–457.