

# CEK - HANIF - CEK

*by* Muhammad Hanif Azhar

---

**Submission date:** 21-Dec-2022 02:00PM (UTC+0800)

**Submission ID:** 1985374938

**File name:** 2.pdf (717.29K)

**Word count:** 5326

**Character count:** 27639

## The Effect of Stocking Density on Growth Performance and Water Quality of the Silver rasbora (*Rasbora argyrotaenia*) Fry Reared in Plastic-lined Pond

Muhammad Hanif Azhar<sup>1,2\*</sup>, Wildan Kurnia Pratama<sup>1</sup>, Darmawan Setia Budi<sup>1</sup>,  
Akhmad Taufiq Mukti<sup>3</sup>, Muhammad Brawijoyo Santanumurti<sup>3</sup>, Devrim Memiş<sup>2</sup>

- <sup>1</sup>. Study Program of Aquaculture, Department of Health and Life Sciences, School of Health and Life Sciences (SIKIA), Universitas Airlangga, Jalan Wijaya Kusuma 113, 68423 Banyuwangi, Indonesia
- <sup>2</sup>. Department of Aquaculture, Faculty of Aquatic Sciences, Istanbul University, Istanbul 34134, Turkey
- <sup>3</sup>. Department of Aquaculture, Faculty of Fisheries and Marine, Universitas Airlangga, Surabaya 60115, Indonesia

\*Corresponding Author: hanifazhar@fpk.unair.ac.id

### ARTICLE INFO

#### Article History:

Received: Feb. 7, 2022  
Accepted: Sept. 13, 2022  
Online: Oct. 27, 2022

#### Keywords:

Silver rasbora,  
*Rasbora argyrotaenia*,  
growth,  
rearing,  
density

### ABSTRACT

Silver rasbora is one of the local fish species in Indonesia that gains a high public demand. To meet the increasing market demand, it is necessary to implement an optimal silver rasbora rearing system. The objective of the study was to determine the best stocking density for the growth and water quality of silver rasbora (*Rasbora argyrotaenia*) fry reared in plastic-lined ponds. The fries utilized had a mean initial body length of  $1.85 \pm 0.5$  cm and body weight of  $0.22 \pm 0.05$  g. The experiment lasted for 40 days and was carried out in a 500 L plastic-lined pond, with dimensions of 1 x 1 x 1 m and five treatments were evaluated (100, 150, 200, 250, 300 fish  $m^{-2}$ ). The feed used was commercial floating feed with 39% protein content, and feeding was conducted three times a day (08.00 AM, 12.00 PM, and 16.00 PM) on ad satiation basis. According to the results, the stocking density showed a significant effect on body weight, final biomass, and feed conversion ratio (FCR) in all treatments. T1 resulted in the highest body weight with a value of  $0.71 \pm 0.02$  g, and the lowest FCR with a value of  $0.90 \pm 0.01$ . While, T5 resulted in the highest final biomass with a value of  $102.81 \pm 5.62$  g. The pH concentration was the only parameter that showed a significantly different value, with the lowest value in T5 ( $7.02 \pm 0.0$ ). The best water quality parameter in T5 showed values of  $27.03 \pm 0.1^{\circ}C$  (temperature),  $7.02 \pm 0.03$  (pH),  $6.79 \pm 0.21$   $mgL^{-1}$  (DO),  $2.81 \pm 0.89$   $mgL^{-1}$  (nitrate),  $0.18 \pm 0.02$   $mgL^{-1}$  (nitrite), and  $0.05 \pm 0.01$   $mgL^{-1}$  (TAN). A stocking density of 300 fish showed suitable conditions for rearing silver rasbora fries in plastic-lined pond.

### INTRODUCTION

Silver rasbora (*Rasbora argyrotaenia*), a freshwater fish from the family Cyprinidae is commonly found in Southeast Asia, including Indonesia, and lives in flowing (rivers) to stagnant waters (rice fields and lakes) (Nelson, 2006; Said & Triyanto, 2013). Silver rasbora has a high economic value in Indonesia because it can be used as both a food consumption and an ornamental fish (Said *et al.*, 2011). Previous data states that in 2018,

rasbora production in Indonesia reached 9,523.96 tons at 60,000 IDR or 4.19 USD (KKP, 2021; Rukayah & Lestari, 2021). It is one of the popular freshwater fish for its delicious taste and high nutritional content (protein 47,54% and fat 2,9%) (Said *et al.*, 2010; Sogandi *et al.*, 2019), but unfortunately, rasboras cannot be optimally cultivated and still rely on wild harvests (Budi *et al.*, 2020). According to previous research, rasbora is an endangered species due to overfishing (Muchlisin *et al.*, 2010) and environmental pollution such as the use of pesticides, which affects the availability of rasbora in nature; therefore, aquaculture regulations should be carried out to prevent rasboras from extinction (Prastika *et al.*, 2021).

Since 2010, there has been an increase in efforts to cultivate rasbora, and several studies on its development have been conducted, including habitat characteristics, feeding habits and conservation systems (Sulastri *et al.*, 2010; Sulistiyarto, 2013; Rosadi *et al.*, 2014), reproductive aspects (Said *et al.*, 2011), mtDNA assay (Kusuma *et al.*, 2017), spawning induction using hormones (Ningrum *et al.*, 2019), sperm cryopreservation to support their breeding efforts (Muchlisin *et al.*, 2020; Eriani *et al.*, 2022); and also the use of different stocking densities in larval stages (Budi *et al.*, 2020). The majority of fish harvest has been currently directed to fulfill market demand. Hence, it is necessary to conduct appropriate fry-rearing techniques to fulfill the market demand. Information about the appropriate stocking density is very important to increase the productivity of silver rasbora fries and ensure a continuous supply.

Stocking density has a direct impact on biota growth, water quality, and production efficiency (de Oliveira *et al.*, 2012; Tibile *et al.*, 2016). According to previous studies, excessive stocking density reduces the oxygen content in waters causing the fish to become stressed, susceptible to disease, and witness a decrease in both growth performance, survival rate and feed efficiency (Martos-Sittha *et al.*, 2020; Muchlisin *et al.*, 2021). Other studies stated that high stocking densities can cause high carbon dioxide levels due to fish respiration, thereby lowering pH and increasing acidity in the waters (Hong *et al.*, 2019). Furthermore, studies related to stocking density are important, because if the density is too high, it will cause a size disparity among fish, competition for food, and ammonia excretion in the waters, in addition to predation (Ronald *et al.*, 2014; Tibile *et al.*, 2016). However, if it is not enough, then the profit from aquaculture will not be optimal (Szczepkowski *et al.*, 2011). Therefore, this study aimed to obtain an appropriate method to produce quality fish and ensure a continuous supply of fries in silver rasbora hatchery activities using plastic-lined ponds by determining the effect of different stocking densities on growth parameters and water quality.

## MATERIALS AND METHODS

This study was conducted at the School of Health and Life Sciences (SIKIA) Banyuwangi Campus Laboratory of Airlangga University, Banyuwangi, Indonesia.

### Fish stocking

Silver rasbora fries were obtained from the Technical Services Unit of Freshwater Aquaculture Development in Umbulan (Pasuruan, East Java). Silver rasbora fries specimens were kept for 1 week in an adaptation pond before the experiment to minimize stress. During the adaptation period, the fries were fed the same type of commercial floating feed and at the same time (three times a day at 08.00 AM, 12.00 PM, and 16.00 PM) as will be carried out in the experimental process. However, before being placed in the experimental pond, the fries were selected and weighed individually, based on their length and size. The fries samples had a mean initial weight of  $0.22 \pm 0.05$  g and a mean length of  $1.85 \pm 0.05$  cm.

### Experimental Design

A completely randomized experimental design was used in this study. This study was conducted for 40 days, using 5 treatments of different stocking density levels (100, 150, 200, 250, and 300 fishes  $m^{-2}$ ) with 4 replicates. Silver rasbora fries were reared semi-outdoor using plastic-lined ponds measuring  $1 \times 1 \times 1$  m, with a water volume of 500 L and aerated with an air blower ( $140 \text{ L min}^{-1}$ ).

During the experiment, the feed was administered three times a day (at 08.00 AM, 12.00 PM, and 16.00 PM) using floating commercial feed with a protein content of 39%, lipid (5%), crude fiber (4%), moisture (10%), and ash (11%) (PF-500 Matahari Sakti Ltd Indonesia), with ad satiation basis. The feed was weighed before and after being given, and feeding was carried out gradually and stopped if the fish stops responding to the feed. The sampling of water quality and fish growth was conducted before morning feeding time and siphon process (carried out once every 10 days by changing the water to a maximum of 10% of the total volume of water in the fishpond).

### Water Quality Measurement

Temperature and pH were measured using a pH meter (Lutron YK2001) and DO (Dissolved oxygen) was measured using a DO meter (YSI Pro20). Nitrite and nitrate were measured using a test kit (Merck MColortest 114658 and Hanna Instruments HI3874) and TAN (Total Ammonia Nitrogen) was analyzed using a spectrophotometer based on the APHA (1999) procedures. In addition, measurement of TAN, nitrite, and nitrate concentrations was carried out in the water quality analysis laboratory. The water samples were collected directly from four points in the water body using a 250 ml sample bottle and then sent to the laboratory for analysis. Meanwhile, the temperature and DO measurements were carried out in situ.

### Observation Parameters

Temperature, pH, and DO were measured daily (every morning) while TAN, Nitrite, and Nitrate were measured every 10 days. Water samples were taken in the morning before feeding time and the production parameters observed include Final Biomass (FB), Weight Gain (WG), Survival Rate (SR), Total amount of given feed (TF), and Feed Conversion Ratio (FCR). Furthermore, it can be formulated  $FCR = TF/WG$ ;  $WG = \text{Mean}$

final weight – mean initial weight;  $SR (\%) = (\text{Final number of fish} / \text{Initial number of fish}) \times 100$  (Sarker *et al.*, 2016). Growth sampling was also performed every 10 days in the morning before feeding and after measuring water quality by taking 10% of biomass (T1: 10; T2: 15; T3: 20; T4: 25; T5: 30 fish) from each treatment in the experimental pond and measuring the lengths and weights individually.

### Data analysis

The data obtained were processed using Microsoft Excel 2013 and then tested for normality and homogeneity using the Kolmogorov-Smirnov and Levene's test, followed by the analysis of the fish growth data using Analysis of Variance (ANOVA) at a 95% confidence interval ( $\alpha = 0.05$ ). Subsequently, if the results were statistically significant ( $P < 0.05$ ), the 95% Tukey Test was carried out. Water quality parameters were measured using repeated-measures ANOVA with two factors (water quality and time). Bonferroni test was used when differences between time and treatments were detected ( $P < 0.05$ ). Statistical analysis was also performed using IBM-SPSS version 18.

## RESULTS

### 1. Growth Performance Parameter

The use of different stocking densities had an effect on the growth parameters of silver rasbora seeds. The results were showed in Table 1.


**Table 1.** Growth performance of Silver rasbora fries for 40 days rearing period using the different stocking densitie

| Variable                   | Treatment (Different Stocking Density) |                          |                         |                         |                          |
|----------------------------|--|--------------------------|-------------------------|-------------------------|--------------------------|
|                            | T1<br>(100)                            | T2<br>(150)              | T3<br>(200)             | T4<br>(250)             | T5<br>(300)              |
| FB (g tank <sup>-1</sup> ) | 69.99±1.31 <sup>a</sup>                | 73.36±3.15 <sup>ab</sup> | 78.91±1.73 <sup>b</sup> | 90.58±3.68 <sup>c</sup> | 102.81±5.62 <sup>d</sup> |
| TF (g tank <sup>-1</sup> ) | 45.43±0.01 <sup>a</sup>                | 47.75±0.02 <sup>a</sup>  | 65.56±0.01 <sup>b</sup> | 79.15±0.01 <sup>c</sup> | 97.83±0.02 <sup>d</sup>  |
| FCR                        | 0.90±0.01 <sup>a</sup>                 | 0.93±0.03 <sup>a</sup>   | 1.20±0.01 <sup>b</sup>  | 1.20±0.14 <sup>b</sup>  | 1.30±0.01 <sup>b</sup>   |
| WG (g)                     | 50.3±1.87 <sup>a</sup>                 | 51.6±2.37 <sup>ab</sup>  | 55.8±1.36 <sup>b</sup>  | 65.88±2.19 <sup>c</sup> | 77.09±2.94 <sup>d</sup>  |
| SR (%)                     | 100±0.00                               | 100±0.00                 | 100±0.00                | 100±0.00                | 100±0.00                 |


FB: Final Biomass; WG: Weight Gain; TF: Total amount of Feed given; FCR: Feed Conversion Ratio; SR: Survival Rate. Values are mean ± SD. Superscript letters denote significant differences ( $P < 0.05$ ) between treatments

The ANOVA test showed that stocking density had a significant effect ( $P < 0.05$ ) on production parameters (Final Biomass (FB), total amount of food given (TF), and Feed Conversion Ratio (FCR)) but had no significant effect ( $P > 0.05$ ) to Survival Rate (SR). The results of the study showed that the highest values of FB, TF, and FCR were found in T5 treatment using a stocking density of 300 fish m<sup>-2</sup> silver rasbora fries. Tukey test

showed that the FB and TF values in treatment T5 (300 fish m<sup>-2</sup>) were significantly higher than in other treatments. The FCR value in treatment T5 showed a significantly higher than in treatment T1 and T2 but not significant in treatment T3 and T4. The SR value showed no significant results among all the treatments (Table 1).



**Figure 1.** Body weight of Silver rasbora fries at 0, 10, 20, 30, and 40 days with different stocking density treatments. Superscript letters differ significantly ( $P < 0.05$ ) between all treatments



**Figure 2** Total length of Silver rasbora fries at 0, 10, 20, 30, and 40 days with different stocking density treatments. Total length of silver rasbora showed did not significantly ( $P > 0.05$ ) among all treatments

40

The results showed that the use of stocking density affected the weight of the fish (Figs. 1,2). Fish weight during the rearing period showed a significant value ( $P < 0.05$ ) between all treatments. However, the total length did not show a significant value

( $P > 0.05$ ) between all treatments. This indicates the higher the stocking density used, the lower the fish weight.

### Water Quality Parameters

The results of water quality analysis using a one-way repeated measure ANOVA showed in Table 2. The results were not significantly different ( $P > 0.05$ ) between the use of stocking density and water quality in the form of TAN, nitrite, nitrate, DO, and temperature on rearing media of silver rasbora. However, the use of stocking density gave a significant effect ( $P < 0.05$ ) on water quality in the form of pH value. The pH value between treatments showed a significant value ( $P < 0.05$ ) between T5 and T1, T2, and T4 treatments. However, it was not significant ( $P > 0.05$ ) from the T3 treatment. In general, water quality parameters during the rearing period of silver rasbora fry can be seen in Table 2.

**Table 2.** Water quality of Silver rasbora fry for 40 days rearing period using the different stocking densities

| Variable                      | T1<br>(100)             | T2<br>(150)            | T3<br>(200)            | T4<br>(250)            | T5<br>(300)            | P<br>value <sup>†</sup> |
|-------------------------------|-------------------------|------------------------|------------------------|------------------------|------------------------|-------------------------|
| TAN (mg L <sup>-1</sup> )     | 0.03±0.01               | 0.04±0.02              | 0.04±0.02              | 0.05±0.02              | 0.05±0.01              | 0.329                   |
| Nitrite (mg L <sup>-1</sup> ) | 0.09±0.01               | 0.18±0.03              | 0.28±0.01              | 0.18±0.08              | 0.18±0.02              | 0.317                   |
| Nitrate (mg L <sup>-1</sup> ) | 2.79±1.41               | 3.10±0.71              | 3.22±0.65              | 2.69±0.75              | 2.81±0.89              | 0.967                   |
| DO (mg L <sup>-1</sup> )      | 7.06±0.14               | 7.02±0.16              | 7.01±0.14              | 7.07±0.07              | 6.79±0.21              | 0.340                   |
| pH                            | 7.19±0.07 <sup>bc</sup> | 7.24±0.07 <sup>c</sup> | 7.07±0.03 <sup>a</sup> | 7.16±0.11 <sup>b</sup> | 7.02±0.03 <sup>a</sup> | 0.000                   |
| Temperature (°C)              | 27.5±0.4                | 27.4±0.4               | 27.5±0.1               | 27.6±0.1               | 27.3±0.1               | 0.114                   |

TAN: Total ammonia nitrogen; DO: Dissolved oxygen. Values are mean ± SD. Superscript letters denote significant differences between treatments (Bonferroni test,  $P < 0.05$ ). <sup>†</sup>Results from one-way repeated measure ANOVA (Density x Times) and Bonferroni test.

### Inorganic Nitrogen

The results of the sampling every 10 days which were analyzed using a one-way repeated measure ANOVA showed that were not significantly different ( $P > 0.05$ ) between the use of stocking density and the concentration of inorganic nitrogen (TAN, nitrite, nitrate) on silver rasbora rearing media. The value of TAN concentration tends to increase at the beginning of the rearing period and tends to decrease at the end of the rearing period. While the value of Nitrite concentration tends to be high at the beginning of the maintenance period and tends to fluctuate until the end of the rearing period. Nitrate concentration values tend to increase until the end of the maintenance period in all treatments.

## DISCUSSION

The use of different stocking densities significantly affected the growth parameters of silver rasbora fry during the experimental period. The weight of silver rasbora fry

decreased with the increase in the stocking density of silver rasbora fry. The weight of the fish decreased with the increase in stocking density and this result is consistent with several studies using fish species from the family Cyprinidae (*Labeo bata*, *Puntius binotatus*, *Barbonymus gonionotus*) which showed a negative correlation between the stocking density and fish growth. In addition, the use of high stocking density could reduce the water quality, increase the competition for feed, and affect the aggressiveness of fish (Rahman *et al.*, 2015; Samad *et al.*, 2016; Hayat *et al.*, 2018). However, several studies showed a positive correlation between stocking density and fish growth, such as the Sciaenidae family (*Argyrosomus japonicus* and *Argyrosomus regius*) (Pirozzi *et al.*, 2009; Millán-Cubillo *et al.*, 2016). This result is possible with animals who prefer to live in groups to avoid stress (de las Heras *et al.*, 2015). Studies on stocking density must take into consideration a variety of factors, including species selection, social interactions, water quality, and environmental conditions (Salari *et al.*, 2012; Arifin *et al.*, 2017).

Furthermore, the stocking density also affected the FCR and the total amount of given feed as the higher the stocking density, the higher the FCR and TF value. However, the use of different stocking densities in the rearing of silver rasbora fry was still ideal because it showed an FCR value of less than 1.5 in all treatments. Feed conversion and efficiency are very important indicators of efficiency in the use of the feed provided because feed prices tend to rise (Craig and Khun, 2017; Fry *et al.*, 2018). Moreover, the FCR value of less than 2 was still considered good in aquaculture industries (Bag *et al.*, 2016). The use of optimal stocking densities in aquaculture is very necessary to reduce stress on fish and potential economic losses (Arifin *et al.*, 2017).

The stocking density treatments did not affect the SR of silver rasbora fry until the end of the experimental period. The high SR in all treatments could be influenced by several factors, such as the species used and the influence of environmental factors (Samad *et al.*, 2016). Stocking density is one of the most important factors in aquaculture activities because it directly affects SR, growth performance, water quality, and production efficiency in fish or shrimp farming (Chakraborty and Mirza, 2007; Rahman and Verdegem, 2010; de Oliveira *et al.*, 2012). Several similar studies in the family Cyprinidae also showed results where stocking density did not affect SR (Said and Triyanto, 2013; Rahman *et al.*, 2015; Hayat *et al.*, 2018).

Silver rasbora is a fish of the genus *Rasbora* which is found on many islands in Indonesia (Kusuma *et al.*, 2017). The result of water quality parameters in several *Rasbora* studies is shown in Table 3.

Based on observations made during the maintenance period, showed that the water quality parameters in the form of TAN, Nitrite, and Nitrate values during the maintenance period were higher than in the literature (Table 3). This could be caused by excess feed residue and the use of stocking density (Hayat *et al.*, 2018). Concentrations of TAN, Nitrite, and Nitrate in the rearing medium fluctuated but increased until the end



of the rearing period. The increased concentration of TAN, Nitrite, and Nitrate in rearing ponds could be due to the higher organic matter content in the rearing media from uneaten feed and fish feces (Jiménez-Montealegre *et al.*, 2002; Torres-Beristain *et al.*, 2006; Bag *et al.*, 2016).

**Table 3.** Water quality parameters in the genus of Rasbora

| Variable                      | During Experiment | Acceptable Range (Literature) | Notes                           |
|-------------------------------|-------------------|-------------------------------|---------------------------------|
| TAN (mg L <sup>-1</sup> )     | 0.029-0.061       | 0.010-0.013 <sup>c</sup>      | Larva rearing                   |
| Nitrite (mg L <sup>-1</sup> ) | 0.101-0.310       | 0.002-0.012 <sup>a</sup>      | Natural habitat                 |
| Nitrate (mg L <sup>-1</sup> ) | 1-3.8             | 0.398-3.272 <sup>a</sup>      | Natural habitat                 |
| pH                            | 6.5-8.0           | 6.7-8.8 <sup>ab</sup>         | Natural habitat                 |
| DO (mg L <sup>-1</sup> )      | 6.2-7.2           | 6.2-7.5 <sup>ac</sup>         | Natural habitat; larvae rearing |
| Temperature (°C)              | 27.1-27.7         | 24.5-32.0 <sup>d</sup>        | Adaptation test                 |

<sup>a</sup>Sentosa and Djumanto, (2010) ; <sup>b</sup>Sulastri *et al.*, (2010); <sup>c</sup> Budi *et al.*, (2020); <sup>d</sup>Said and Triyanto, (2013)

The siphoning process that was carried out every 10 days during the rearing period in all treatments did not affect the concentration of TAN, Nitrite, and Nitrate in the rearing pond. Furthermore, the fluctuating conditions of TAN, Nitrite, and Nitrate concentrations during the rearing period indicated the occurrence of nitrification and denitrification by natural bacteria found in rearing ponds (Durborow *et al.*, 1997; Turker *et al.*, 2003).

## CONCLUSION

Differences in stocking density significantly affected the growth performance of BW, FB, TF, FCR, WG, and water quality (pH value) of silver rasbora (*Rasbora argyrotaenia*) fry in plastic-lined ponds. A stocking density of 100 fish 500 L<sup>-1</sup> resulted in optimal growth of the fry. However, based on all production parameters and overall water quality, the use of a stocking density of 300 fish 500 L<sup>-1</sup> still showed suitable conditions for cultivation activities with an FCR value of less than 1 and optimal water quality for rearing silver rasbora fries in plastic-lined ponds.

## AUTHORS' CONTRIBUTIONS

33

All authors contributed to the conception and design of the study. Wildan Kurnia Pratama and Darmawan Setia Budi prepared the materials, collected data, and analyzed the results. Muhammad Hanif Azhar wrote the first draft of the manuscript and Ahmad Taufiq Mukti, Devrim Memiş, and Muhammad Brawijoyo Santanumurti commented on previous versions of the manuscript.

### ACKNOWLEDGMENT

The authors are thankful that UPT PBAT Umbulan, Pasuruan and School of Health and Life Sciences (SIKIA) Airlangga University in Banyuwangi for providing us with research materials, areas and laboratories to conduct this research.

### CONFLICT OF INTEREST

The authors declare that there is no conflict of interest.

### REFERENCES

- Al Adawiyah, L.; Sulmartiwi, L.; Bodur T. and Budi D.S.** (2019). Induction of spermiation using Ovaprim™ with topical gill method in the silver rasbora (*Rasbora argyrotaenia*). *Theriogenology.*, 126: 172–176.
- APHA** (1999). Standard methods for the examination of the water and wastewater, 20th Edition, American Public Health Association, Washington, DC. 1268pp.
- Arifin, O.Z.; Subagja, J.; Prakoso, V.A. and Suhud E.H.** (2017). Effect of stocking density on growth performance of domesticated barb (*Barbonymus balleroides*). *Indonesia Aquac J.*, 12: 1-6.
- Bag, N.; Moulick, S. and Mal, B.C.** (2016). Effect of stocking density on water and soil quality, growth, production and profitability of farming Indian major carps. *Indian J Fish.*, 63: 39-49.
- Budi, D.S.; Rizkyan, I. and Prayogo, P.** (2020). The effect of stocking density on the survival and growth of silver rasbora (*Rasbora Argyrotaenia*) larvae. *Polish J Nat Sci.*, 35: 87–95.
- Chakraborty, B.K. and Mirza, M.J.A.** (2007). Effect of stocking density on survival and growth of endangered bata, *Labeo bata* (Hamilton-Buchanan) in nursery ponds. *Aquaculture.*, 265: 156–162.
- Craig, S. and Khun, D.D.** (2017). *Understanding Fish Nutrition, Feeds, and Feeding.* Virginia, Tech-Publication.
- de las Heras, V.; Martos-Sitcha, J.A.; Yúfera, M. and Mancera, J.M.** (2015). Influence of stocking density on growth, metabolism, and stress of thick-lipped grey mullet (*Chelon labrosus*) juveniles. *Aquaculture.*, 448:29–37.
- de Oliveira, E.G.; Pinheiro, A.B.; de Oliveira, V.Q.; da Silva Júnior, A.R.M.; de Moraes, M.G.; Rocha, I.R.C.B.; de Sousa, R.R. and Costa, F.H.F.** (2012). Effects of stocking density on the performance of juvenile pirarucu (*Arapaima gigas*) in cages. *Aquaculture.*, 370-371:96-101.
- Durborow, R.M.; Crosby, D.M. and Brunson, M.W.** (1997). Nitrite in fish pond, Southern Regional Aquaculture Center, SRAC Publ, No. 463.
- Eriani, K.; Mustaqim, M.; Hasri, I.; Amalia, R.; Al Azhar.; Nur, F.M. and Muchlisin, Z.A.** (2022). Effect of fetal bovine serum on the sperm quality of depik (*Rasbora tawarensis*) after short-term cryopreservation. *Indian J. Anim. Res.*, 56(7): 843-847.
- Fry, J.P.; Mailloux, N.A.; Love, D.C.; Milli, M.C. and Cao, L.** (2018). Feed conversion efficiency in aquaculture: Do we measure it correctly? *Environ Res Lett.*, 13.2: 024017.

- Hayat, M.; Nugroho, R.A. and Aryani, R.** (2018). Influence of different stocking densities on the growth, feed efficiency, and survival of majalaya common carp (*Cyprinus carpio* linnaeus 1758). *F1000Research.*, 7: 1–10.
- Hong, J.; Chen, X.; Liu, S.; Fu, Z.; Han, M.; Wang, Y.; Gu, Z. and Ma, Z.** (2019). Impact of fish density on water quality and physiological response of golden pompano (*Trachinotus ovatus*) flingerlings during transportation. *Aquaculture.*, 507: 260-265.
- Jiménez-Montealegre, R.; Verdegem, M.C.J.; Van Dam, A. and Verreth, J.A.J.** (2002). Conceptualization and validation of a dynamic model for the simulation of nitrogen transformations and fluxes in fish ponds. *Ecol. Modell.*, 147: 123–152.
- Keshavanath, P.; Gangadhar, B.; Ramesha, J.; Priyadarshini, M.; Van Dam, A.A.; Verdegem, M.C.J. and Beveridge, M.C.M.** (2015). Impact of substrates and fish stocking densities on growth and production of the Indian major carp, *Labeo rohita* (HAM.). *J. Aqua. Trop.*, 30: 1-14.
- Kottelat, M.; Whitten, A.; Kartikasari, A.N. and Wirjoatmojo, S.** (1993). Freshwater fishes of Western Indonesia and Sulawesi: Ikan Air Tawar Indonesia Bagian Barat dan Sulawesi, Periplus, Hongkong. 428pp.
- KKP (Kementerian Kelautan dan Perikanan).** (2021). Produksi Perikanan. <https://statistik.kkp.go.id/home.php?m=total&i=2#panel-footer> (Accessed 12 December 2021).
- Kusuma, W.E.; Samuel, P.D.; Wiadnya, D.G.R.; Hariati, A.M. and Kumazawa, Y.** (2017). Complete mitogenome sequence of *Rasbora argyrotaenia* (Actinopterygii: Cyprinidae). *Mitochondrial DNA Part B. Resour.*, 2: 373–374.
- Lukman.** (2017). Perkembangan pemanfaatan dan penelitian ikan bada (*Rasbora argyrotaenia*) di danau maninjau. *Warta Iktiologi.*, 1: 24-27
- Martos-Sittha, J.A.; Mancera, J.M.; Prunet, P. and Magnoni, L.J.** (2020). Welfare and stressors in fish: challenges facing aquaculture. *Frontiers in physiology.*, 11, 162.
- Millán-Cubillo, A.F.; Martos-Sittha, J.A.; Ruiz-Jarabo, I.; Cárdenas, S. and Mancera, J.M.** (2016). Low stocking density negatively affects growth, metabolism, and stress pathways in juvenile specimens of meagre (*Argyrosomus regius*, Asso 1801). *Aquaculture.*, 451: 87–92.
- Muchlisin, Z.A.; Musman, M. and Siti Azizah, M.N.** (2010). Length- weight relationships and condition factors of two threatened fishes, *Rasbora tawarensis* and *Poropuntius tawarensis*, endemic to Lake Laut Tawar, Aceh Province, Indonesia. *J. Appl. Ichthyol.*, 26: 949-953.
- Muchlisin, Z.A.; Sarah, P.I.; Aldila, D.F.; Eriani, K.; Hasri, I.; Batubara, A.S.; Nur, F.M.; Mustaqim, M.; Mutmainnah, C.R.; Abinawanto, A. and Wilkes, M.** (2020). Effect of Dimethyl sulfoxide (DMSO) and egg yolk on sperm motility, fertility and hatching rates of depik *Rasbora tawarensis* (Pisces: Cyprinidae) eggs after short-term cryopreservation. *Aquac. Res.*, 51: 1700–1705.
- Muchlisin, Z.A.; Indrianti, Y.S.; Dewiyanti, I.; Lisra, F.; Yanti, R.Z.; Nur, F.M. and Batubara, A.S.** (2021). Effect of stocking density of the growth performance, survival rate and feed utilization of the eels *Anguilla bicolor* (pisces: anguillidae) Larvae. *IOP Conf. Ser. Earth Environ. Sci.*, 674: 012086.

- Nelson, J.S.** (2006). *Fishes of the World* 4<sup>th</sup> edition. John Willey & Sons. Inc, New Jersey. 622pp.
- Ningrum, D.R.K.; Budi, D.S. and Sulmartiwi, L.** (2019). Induksi pemijahan ikan wader pari (*Rasbora argyrotaenia*) menggunakan Ovaprim<sup>TM</sup> dengan dosis berbeda. *Depik.*, 8: 117–124.
- Pirozzi, I.; Booth, M.A. and Pankhurst, P.M.** (2009). The effect of stocking density and repeated handling on the growth of juvenile mullet, *Argyrosomus japonicus* (Temminck & Schlegel 1843). *Aquac. Int.*, 17: 199–205.
- Prastika, A.B.; Sulmartiwi, L. and Lutfiyah, L.** (2021). Effect of sublethal dose organophosphate pesticides on embryogenesis and hatching rate of silver rasbora eggs (*Rasbora argyrotaenia*). *IOP Conf. Ser. Earth Envir. Sci.*, 679: 012065.
- Rahman, M.; Ferdous, Z.; Mondal, S. and Amin M** (2015.) Effects of stocking densities on growth and survival of Thai Sharpunti (*Barbonymus gonionotus*) in earthen ponds. *J. Bangladesh Agric. Univ.*, 9: 327–338.
- Rahman, M.M. and Verdegem, M.** (2010). Effects of intra- and interspecific competition on diet, growth and behaviour of *Labeo calbasu* (Hamilton) and *Cirrhinus cirrhosus* (Bloch). *Appl. Anim. Behav. Sci.*, 128: 103–108.
- Ronald, N.; Gladys, B. and Gasper, E.** (2014). The effects of stocking density on the growth and survival of Nile tilapia (*Oreochromis niloticus*) fry at Son Fish Farm, Uganda. *J. Aquac. Res. Development.*, 5: 1-7.
- Rosadi, E.; Yuli, H.E.; Setyohadi, D. and Bintoro, G.** (2014). Distribution, composition, and abiotic environment of silver rasbora (*Rasbora argyrotaenia* Blkr) Fish in Upstream Areas of Barito Watershed, South Kalimantan. *J. Environ Ecol.*, 5:117-131.
- Rukayah, S. and Lestari, W.** (2021) Upaya konservasi species asli melalui kajian reproduksi dan lingkungan ikan wader pari (*Rasbora argyrotaenia* Blker, 1854) di Waduk PB Soedirman Banjarnegara. *Prosiding SNPBS.*, 118-129
- Said, D.S.; Haryani, G.S.; Lukman.; Triyanto.; Mayasari, N.; Hamdani, A.; Sutrisno. and Sari, L.** (2010). Perkembangan ikan bada (*Rasbora Argyrotaenia*) Danau Maninjau-Sumatera Barat Pada Habitat Ex-Situ. *Prosiding Seminar Nasional Limnologi V.*
- Said, D.S.; Triyanto.; Lukman.; Sutrisno. and Hamdani, A.** (2011). Aspek biologi ikan bada (*Rasbora argyrotaenia*) di danau maninjau, sumatera barat. *Prosiding Forum Nasional Pemacuan Sumber Daya Ikan.*, III: 1-10
- Said D.S. and Triyanto.** (2013). Ketahanan hidup dan kemampuan tumbuh ikan bada (*Rasbora argyrotaenia*) pada suhu pemeliharaan bervariasi. *Limnotek.*, 20(2): 191-199.
- Said, D.S. and Hidayat.** (2015). 101 Ikan Hias Air Tawar, LIPI Press, Jakarta. 260pp.
- Salari, R.; Saad, C.; Kamarudin, M. and Zokaefar, H.** (2012). Effects of different stocking densities on tiger grouper juvenile (*Epinephelus fuscoguttatus*) growth and a comparative study of the flow-through and recirculating aquaculture systems. *African J. Agric. Res.*, 7: 3765–3771.
- Samad, M.A.; Khatun, A.; Reza, M.S.; Asrafuzzaman, Md. and Ferdaushy, M.H.** (2016). Effects of stocking density on growth, survival and production of mirror carp (*Cyprinus carpio* var. *specularis*) spawn in nursery pond. *Asian J. Med. Biol. Res.*, 2: 429–435.

- Sarker P.K.; Kapuscinski A.R.; Lanois A.J.; Livesey E.D.; Benhard, K.P. and Coley, M.L.** (2016) Towards sustainable aquafeeds: Complete substitution of fish oil with marine microalga *Schizochytrium* sp. improves growth and fatty acid deposition in juvenile Nile tilapia (*Oreochromis niloticus*). PLoS One 11: e0156684.
- Sentosa, A.A. and Djumanto.** (2010). Habitat Pemijahan Ikan Wader Pari (*Rasbora lateristriata*) di Sungai Ngrancah, Kabupaten Kulonprogo. Jurnal Iktiologi Indonesia., 10(1): 55-63.
- Sogandi.; Sanjaya, R.E.; Baity, N. and Syahmani.** (2019). Identification of nutritional content and profiles of amino acid from seluang fish (*Rasbora* sp.). Penelitian Gizi dan Makanan, 42: 73-80.
- Sulastri, Hartoto, D.I.; Yuniarti, I. and Nasution, S.H.** (2010). Karakteristik habitat, kebiasaan makan, dan sistem konservasi ikan bada *Rasbora Argyrotaenia* di Danau Maninjau. Pros Seminar Nasional Ikan., VI: 487-49
- Sulistiyarto, B.** (2013). Hubungan antara kelimpahan ikan saluang (*rasbora argyrotaenia* blkr) dengan populasi fitoplankton di dataran banjir Sungai Rungan Kalimantan Tengah. Jurnal Ilmu Hewani Tropika., 2: 27-30.
- Suryani, S.A.M.P.; Wirawan, I.G.P.; Dwiyani, R. and Sritamin, M.** (2021). Genetic diversity and differentiation of cytochrome oxidase subunit I (COI) gene of *Rasbora lateristriata* bleeker in different habitat. IOP Conf. Ser. Mater. Sci. Eng. 1098: 052043.
- Szczepkowski, M.; Szczepkowska, B. and Piotrowska, I.** (2011). Impact of higher stocking density of juvenile Atlantic sturgeon, *Acipenser oxyrinchus Mitchill*, on fish growth, oxygen consumption, and ammonia excretion. Fish. Aquat. Life., 19(2): 59-67.
- Tibile, R.M.; Sawant, P.B.; Chada, N.K.; Lakra, W.S.; Prakash, C.; Swain S. and Bhagwati, K.** (2016). Effect of stocking density on growth, size variation, condition index and survival of discus, *Symphysodon aequifasciatus* Pellegrin, 1904. Turkish J Fish Aquat Sci., 16: 455–462.
- Torres-Beristain, B.; Verdegem, M.; Kerepeczki, E. and Verreth J.** (2006). Decomposition of high protein aquaculture feed under variable toxic conditions. Water Res., 40: 1341–1350.
- Turker, H.; Eversole, A.G. and Brune, D.E.** (2003). Comparative Nile tilapia and silver carp filtration rates of partitioned aquaculture system phytoplankton. Aquaculture., 220: 449–457.

# CEK - HANIF - CEK

## ORIGINALITY REPORT

13%

SIMILARITY INDEX

8%

INTERNET SOURCES

8%

PUBLICATIONS

0%

STUDENT PAPERS

## PRIMARY SOURCES

|   |   |     |
|---|---|-----|
| 1 | <a href="https://creativecommons.org">creativecommons.org</a><br>Internet Source  | 1%  |
| 2 | Pei, Surui, Shuanglin Dong, Fang Wang, Qinfeng Gao, and Xiangli Tian. "Effects of stocking density and body physical contact on growth of sea cucumber, <i>Apostichopus japonicus</i> ", <i>Aquaculture Research</i> , 2012.<br>Publication | <1% |
| 3 | <a href="http://licenciamento.ibama.gov.br">licenciamento.ibama.gov.br</a><br>Internet Source   | <1% |
| 4 | <a href="http://microbiologyjournal.org">microbiologyjournal.org</a><br>Internet Source   | <1% |
| 5 | <a href="http://www.jeeng.net">www.jeeng.net</a><br>Internet Source   | <1% |
| 6 | Amany Mohamed Haroon. "Review on aquatic macrophytes in Lake Manzala, Egypt", <i>The Egyptian Journal of Aquatic Research</i> , 2022<br>Publication   | <1% |
| 7 | W A Nugrahani, R Kusdarwati, M F Ulkhaq. "Experimental infection of Streptococcus   | <1% |

agalactiae in silver rasbora (*Rasbora argyrotaenia*): Effect to hematological profile from infected fish", IOP Conference Series: Earth and Environmental Science, 2021

Publication

---

8

Nnenna Ugwu, Emma Love, Jo Murrell, Helen R Whay, Toby Knowles, Jo Hockenull.

"Development of an ethogram for hutch-housed dairy calves and determination of factors influencing their behaviour", Applied Animal Behaviour Science, 2021

Publication

---

<1 %

9

F P Putri, N N Dewi. " Growth monitoring of koi fish ( ) in natural hatchery techniques in Umbulan, Pasuruan, East Java ", IOP Conference Series: Earth and Environmental Science, 2019

Publication

---

<1 %

10

[journal.unhas.ac.id](http://journal.unhas.ac.id)

Internet Source

---

<1 %

11

"The Cladocera (Ctenopoda and Anomopoda) in rice fields during a crop cycle at Nakhon Si Thammarat province, southern Thailand", Crustaceana, 2010

Publication

---

<1 %

12

Munir, Mohammad Bodrul, Roshada Hashim, Yam Hok Chai, Terence L. Marsh, and Siti Azizah Mohd Nor. "Dietary prebiotics and

<1 %

probiotics influence growth performance, nutrient digestibility and the expression of immune regulatory genes in snakehead (*Channa striata*) fingerlings", *Aquaculture*, 2016.

Publication

13

[article.sciencepublishinggroup.com](http://article.sciencepublishinggroup.com)

Internet Source

<1 %

14

[globaljournals.org](http://globaljournals.org)

Internet Source

<1 %

15

[www.karger.com](http://www.karger.com)

Internet Source

<1 %

16

Ahmed Ismail. "Environmental Damage Caused by Wastewater Discharge into the Lake Manzala in Egypt", *American Journal of Bioscience and Bioengineering*, 2017

Publication

<1 %

17

Crexi, V.T.. "Production and refinement of oil from carp (*Cyprinus carpio*) viscera", *Food Chemistry*, 20100401

Publication

<1 %

18

V A Prakoso, K T Kim, J H Ryu, B H Min, Y J Chang. " Oxygen consumption of on several temperatures under brackish water conditions ", *IOP Conference Series: Earth and Environmental Science*, 2019

Publication

<1 %



|    |   |      |
|----|---|------|
| 19 | avmajournals.avma.org<br>Internet Source  | <1 % |
| 20 | ftp.academicjournals.org<br>Internet Source   | <1 % |
| 21 | mdpi.com<br>Internet Source   | <1 % |
| 22 | oaji.net<br>Internet Source   | <1 % |
| 23 | psasir.upm.edu.my<br>Internet Source  | <1 % |
| 24 | "Sustainable Aquaculture", Springer Science and Business Media LLC, 2018<br>Publication   | <1 % |
| 25 | Ester Santigosa, Denis Constant, Dauphinee Prudence, Thomas Wahli, Viviane Verlhac - Trichet. " A novel marine algal oil containing both and is an effective source of omega - 3 fatty acids for rainbow trout ( ) ", Journal of the World Aquaculture Society, 2020<br>Publication | <1 % |
| 26 | M. Jiang, H.H. Zhao, S.W. Zai, B. Shepherd, H. Wen, D.F. Deng. "A defatted microalgae meal (Haematococcus pluvialis) as a partial protein source to replace fishmeal for feeding juvenile yellow perch <i>Perca flavescens</i> ", Journal of Applied Phycology, 2018                | <1 % |

27 Pigment & Resin Technology, Volume 43, Issue 2 (2014-03-28) <1 %  
Publication

---

28 Shaharior Hossen, Soo Cheol Kim, Yusin Cho, Kang Hee Kho. "Vital Analysis of Cryopreserved Sperm of Marbled Flounder, Pseudopleuronectes yokohamae", Frontiers in Physiology, 2021 <1 %  
Publication

---

29 arhiva.nara.ac.rs <1 %  
Internet Source

---

30 media.neliti.com <1 %  
Internet Source

---

31 pakbs.org <1 %  
Internet Source

---

32 Irin Iriana Kusmini, Anang Hari Kristanto, Jojo Subagja, Vitas Atmadi Prakoso, Fera Permata Putri. "RESPONS DAN POLA PERTUMBUHAN BENIH ikan baung (Hemibagrus nemurus) DARI TIGA GENERASI DIPELIHARA PADA WADAH BUDIDAYA YANG BERBEDA", Jurnal Riset Akuakultur, 2018 <1 %  
Publication

---

33 Meidina Rizkita, Mia Rosmiati, Gede Suantika, Magdalena Lenny Sitomorang, Made Dendy Pratama, Syafira Rosefa. "Sustainability Status <1 %

# Analysis and Strategy Development for Common Carp (*Cyprinus carpio* L.) Hatchery Industry in Ciparay District, West Java, Indonesia", Research Square Platform LLC, 2022

Publication

34

[assets.researchsquare.com](https://assets.researchsquare.com)

Internet Source

<1 %

35

[d-nb.info](https://d-nb.info)

Internet Source

<1 %

36

[downloads.hindawi.com](https://downloads.hindawi.com)

Internet Source

<1 %

37

[edepot.wur.nl](https://edepot.wur.nl)

Internet Source

<1 %

38

[grouper.unisla.ac.id](https://grouper.unisla.ac.id)

Internet Source

<1 %

39

[habricentral.org](https://habricentral.org)

Internet Source

<1 %

40

[ispff.ir](https://ispff.ir)

Internet Source

<1 %

41

[moam.info](https://moam.info)

Internet Source

<1 %

42

[www.int-res.com](https://www.int-res.com)

Internet Source

<1 %

43

Paulo Adelino Medeiros, Edimar Lopes Costa, Elenice Martins Brasil, Eduardo Akifumi Ono, Elizabeth Gusmão Affonso. "Diets for grow-out of pirarucu in net cage: performance, physiological parameters, fillet composition and feeding cost", Boletim do Instituto de Pesca, 2019

Publication

<1 %

44

Helena Khatoon, Fatimah Yusoff, Sanjoy Banerjee, Mohamed Shariff, Japar Sidik Bujang. "Formation of periphyton biofilm and subsequent biofouling on different substrates in nutrient enriched brackishwater shrimp ponds", Aquaculture, 2007

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

# CEK - HANIF - CEK

GRADEMARK REPORT

FINAL GRADE

**/0**

GENERAL COMMENTS

**Instructor**

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10

PAGE 11

PAGE 12