



[Aquaculture, Aquarium, Conservation & Legislation](#)

You are here > [Home](#) · [AACL](#)

### AACL Bioflux

[Instructions to authors](#)

[Submission letter](#)

[Model of paper](#)

[Reviewer information pack](#)

[Editorial Board Expanded](#)

[Coverage / databases](#)

[Volume 13\(6\)/2020](#)

[Volume 13\(5\)/2020](#)

[Volume 13\(4\)/2020](#)

[Volume 13\(3\)/2020 \(June, 30\)](#)

[Volume 13\(2\)/2020 \(April, 30\)](#)

[Volume 13\(1\)/2020 \(February, 28\)](#)

[Volume 12\(6\)/2019 \(December, 30\)](#)

[Volume 12\(5\)/2019 \(October, 30\)](#)

[Volume 12\(4\)/2019 \(August, 30\)](#)

[Volume 12\(3\)/2019 \(June, 30\)](#)

[Volume 12\(2\)/2019 \(April, 30\)](#)

[Volume 12\(1\)/2019 \(February, 28\)](#)

[Volume 11\(6\)/2018 \(December, 30\)](#)

[Volume 11\(5\)/2018 \(October, 30\)](#)

[Volume 11\(4\)/2018 \(August, 30\)](#)

[Volume 11\(3\)/2018 \(June, 30\)](#)

[Volume 11\(2\)/2018 \(April, 30\)](#)

[Volume 11\(1\)/2018 \(February, 28\)](#)

[Volume 10\(6\)/2017 \(December, 30\)](#)

[Volume 10\(5\)/2017 \(October, 30\)](#)

[Volume 10\(4\)/2017 \(August, 30\)](#)

[Volume 10\(3\)/2017 \(June, 30\)](#)

# Aquaculture, Aquarium, Conservation & Legislation - International Journal of the Bioflux Society

ISSN 1844-9166 (online)

ISSN 1844-8143 (print)

Published by Bioflux - bimonthly -

in cooperation with The Natural Sciences Museum Complex (Constanta, Romania)

Peer-reviewed (each article was independently evaluated before publication by two specialists)

The journal includes original papers, short communications, and reviews on Aquaculture (Biology, Technology, Economics, Marketing), Fish Genetics and Improvement, Aquarium Sciences, Fisheries, Ichthyology, Aquatic Ecology, Conservation of Aquatic Resources and Legislation (in connection with aquatic issues) from wide world.

The manuscripts should be submitted to [zoobiomag2004@yahoo.com](mailto:zoobiomag2004@yahoo.com)

### Editor-in-Chief:

Petrescu-Mag I. Valentin: USAMV Cluj, Cluj-Napoca, University of Oradea (Romania)

Gavriloaie Ionel-Claudiu (reserve): SC Bioflux SRL, Cluj-Napoca (Romania).

### Editors:

Abdel-Rahim Mohamed M.: National Institute of Oceanography and Fisheries, Alexandria (Egypt)

Adascalitei Oana: Maritime University of Constanta, Constanta (Romania)

Amira Aicha Beya: Badji Mokhtar Annaba University, Annaba (Algeria)

Arockiaraj A. Jesu: SRM University, Chennai (India)


Appelbaum Samuel: Ben-Gurion University of the Negev (Israel)

Baharuddin Nursalwa: Universiti Malaysia Terengganu, Terengganu (Malaysia)

Boaru Anca: USAMV Cluj, Cluj-Napoca (Romania)


Volume 10(2)/2017 (April, 30)	Botha Miklos: Bioflux SRL, Cluj-Napoca (Romania)
Volume 10(1)/2017 (February, 28)	Breden Felix: Simon Fraser University (Canada)
Volume 9(6)/2016 (December, 30)	Burny Philippe: Universite de Liege, Gembloux (Belgium)
Volume 9(5)/2016 (October, 30)	Caipang Cristopher M.A.: Temasek Polytechnic (Singapore)
Volume 9(4)/2016 (August, 30)	Chapman Frank: University of Florida, Gainesville (USA)
Volume 9(3)/2016 (June, 30)	Coroian Cristian: USAMV Cluj, Cluj-Napoca (Romania)
Volume 9(2)/2016 (April, 30)	Creanga Steofil: USAMV Iasi, Iasi (Romania)
Volume 9(1)/2016 (February, 28)	
Volume 8(6)/2015 (December, 30)	Cristea Victor: Dunarea de Jos University of Galati, Galati (Romania)
Volume 8(5)/2015 (October, 30)	Das Simon Kumar: Universiti Kebangsaan Malaysia, Bangi, Selangor (Malaysia)
Volume 8(4)/2015 (August, 30)	Dimaggio Matthew A.: University of Florida (USA)
Volume 8(3)/2015 (June, 30)	Firica Cristian Manuel: Spiru Haret University Bucharest, Craiova (Romania)
Volume 8(2)/2015 (April, 30)	Georgescu Bogdan: USAMV Cluj, Cluj-Napoca (Romania)
Volume 8(1)/2015 (February, 28)	
Volume 7(6)/2014 (December, 30)	Karayucel Ismihan: University of Sinop, Sinop (Turkey)
Volume 7(5)/2014 (October, 30)	Khamesipour Faham: Shiraz University, Shiraz (Iran)
Volume 7(4)/2014 (August, 30)	Kosco Jan: Presov University, Presov (Slovakia)
Volume 7(3)/2014 (June, 30)	Kovacs Eniko: USAMV Cluj, Cluj-Napoca (Romania)
Volume 7(2)/2014 (April, 15)	Mehrad Bahar: Gorgan University of Agricultural Sciences and Nat. Res. (Iran)
Volume 7(1)/2014 (February, 15)	Miclaus Viorel: USAMV Cluj, Cluj-Napoca (Romania)
Volume 6(6)/2013 (November, 15)	Mihociu Tamara: R&D National Institute for Food Bioresources (Romania)
Volume 6(5)/2013 (September, 15)	Molnar Kalman: Hungarian Academy of Sciences, Budapest (Hungary)
Volume 6(4)/2013 (July, 25)	Muchlisin Zainal Abidin: Universiti Sains (Malaysia), Syiah Kuala University (Indonesia)
Volume 6(3)/2013 (May, 15)	Muntean George Catalin: USAMV Cluj, Cluj-Napoca (Romania)
Volume 6(2)/2013 (March, 15)	Nowak Michal: University of Agriculture in Krakow (Poland)
Volume 6(1)/2013 (January, 15)	
Volume 5(5)/2012 (December, 30)	Nyanti Lee: Universiti Malaysia Sarawak, Sarawak (Malaysia)
Volume 5(4)/2012 (September, 30)	Olivotto Ike: Universita Politecnica delle Marche, Ancona (Italy)
Volume 5(3)/2012 (July, 30)	Oroian Firuta Camelia: USAMV Cluj, Cluj-Napoca (Romania)
Volume 5(2)/2012 (June, 30)	Papuc Tudor: USAMV Cluj, Cluj-Napoca (Romania)
Volume 5(1)/2012 (March, 15)	Parvulescu Lucian: West University of Timisoara (Romania)
Volume 4(5)/2011 (December, 30)	Pasarin Benone: USAMV Iasi, Iasi (Romania)
Volume 4(4)/2011 (October, 30)	Pattikawa Jesaja Ajub: Pattimura University, Ambon (Indonesia)
Volume 4(3)/2011 (July, 30)	Petrescu Dacinia Crina: Babes-Bolyai University, Cluj-Napoca (Romania)

<b>Volume 4(2)/2011 (April, 30)</b>	Petrescu-Mag Ruxandra Malina: Babes-Bolyai University, Cluj-Napoca (Romania)
<b>Volume 4(1)/2011 (January, 30)</b>	Petrovici Milca: West University of Timisoara (Romania)
<b>Volume 3(5)/2010 (December, 5)</b>	Pratasik Silvester Benny: Sam Ratulangi University, Manado (Indonesia)
<b>Volume 3(4)/2010 (December, 1)</b>	Proorocu Marian: USAMV Cluj, Cluj-Napoca (Romania)
<b>Volume 3(3)/2010 (November, 15)</b>	Ray Sunuram: Khulna University (Bangladesh)
<b>Volume 3(2)/2010 (July, 30)</b>	Rhyne Andrew: Roger Williams University; New England Aquarium, Boston (USA)
<b>Volume 3(1)/2010 (February, 28)</b>	Ruchin Alexander B.: Joint Directorate of the Mordovia State Nature Reserve and National Park «Smolny», Saransk (Russia)
<b>Volume 2(4)/2009 (October, 30)</b>	Safirescu Calin: USAMV Cluj, Cluj-Napoca (Romania)
<b>Volume 2(3)/2009 (July, 30)</b>	Serrano Jr. Augusto E.: University of the Philippines Visayas (Philippines)
<b>Volume 2(2)/2009 (April, 30)</b>	Sima Nicusor Flaviu: USAMV Cluj, Cluj-Napoca (Romania)
<b>Volume 2(1)/2009 (January, 30)</b>	Tlusty Michael F.: New England Aquarium, Boston (USA)
<b>Volume 1(2)/2008 (December, 30)</b>	Vesa Stefan Cristian: Iuliu Hatieganu UMF, Cluj-Napoca (Romania)
<b>Volume 1(1)/2008 (September, 30)</b>	Vintila Iuliana: Dunarea de Jos University of Galati, Galati (Romania)
<b>Volume Pilot/2007 (December, 30) - available printed only</b>	Wariaghli Fatima: University Mohammed V in Rabat, Rabat (Morocco)
<b>Pontus Euxinus, Volume 1 (1980) - Parent Journal</b>	Yusli Wardiatno: Bogor Agricultural University, Bogor (Indonesia).



design: [www.simple-webdesign.com](http://www.simple-webdesign.com)

Home | Archive | Volume 1 (1) / 2008 | CEEX 140 | Volume 1 (2) / 2008 | Volume 2 (1) / 2009 | Volume 2 (2) / 2009 | Volume 2 (3) / 2009 | Volume 2 (4) / 2009 | Pilot | Volume 3 (1) / 2010 | Volume 3(2)/2010 (July, 30) | Volume 3(3)/2010 | Volume 3(4)/2010 | Volume 3(5)/2010 - ACVAPEDIA 2010 | Volume 4(1)/2011 | Volume 4(2)/2011 - ACVAPEDIA 2010 | Volume 4(3)/2011 | Volume 4(4)/2011 | Volume 4(5)/2011 | Volume 5(1)/2012 (March, 15) | Volume 5(2)/2012 | Volume 5(3)/2012 | Volume 5(4)/2012 | Volume 5(5)/2012 (December, 30) | Volume 6(1)/2013 - ACVAPEDIA 5th edn., Hungary, Szarvas (HAKI), 27-29th of November, 2012 | Volume 6(2)/2013 - ACVAPEDIA 5th edn., Hungary, Szarvas (HAKI), 27-29th of November, 2012 | Volume 6(3)/2013 | Volume 6(4)/2013 | Volume 6(5)/2013 | Volume 6(6)/2013 | Volume 7(1)/2014 | Volume 7(2)/2014 | Volume 7(3)/2014 | Volume 7(4)/2014 | Volume 7(5)/2014 | Volume 7(6)/2014 | Volume 8(1)/2015 | Volume 8(2)/2015 | Volume 8(3)/2015 | Volume 8(4)/2015 | Volume 8(5)/2015 | Volume 8(6)/2015 | Volume 9(1)/2016 | Volume 9(2)/2016 | Volume 9(3)/2016 | Volume 9(4)/2016 | Volume 9(5)/2016 | Volume 9(6)/2016 | Volume 10(1)/2017 | Volume 10(2)/2017 | Volume 10(3)/2017 | Volume 10(4)/2017 | Volume 10(5)/2017 | Volume 10(6)/2017 | Volume 11(1)/2018 | Volume 11(2)/2018 | Volume 11(3)/2018 | Volume 11(4)/2018 | Pontus Euxinus, Volume 1, 1980 | Volume 11(5)/2018 | Volume 11(6)/2018 | Volume 12(1)/2019 | Volume 12(2)/2019 | Volume 12(3)/2019 | Volume 12(4)/2019 | Volume 12(5)/2019 | Volume 12(6)/2019 | Volume 13(1)/2020 | Volume 13(2)/2020 | Volume 13(3)/2020 | Contact | Site Map





[Aquaculture, Aquarium, Conservation & Legislation](#)

You are here > [Home](#) · [Volume 13\(1\)/2020](#)

## AAACL Bioflux

<a href="#">Instructions to authors</a>
<a href="#">Submission letter</a>
<a href="#">Model of paper</a>
<a href="#">Reviewer information pack</a>
<a href="#">Editorial Board Expanded</a>
<a href="#">Coverage / databases</a>
<a href="#">Volume 13(6)/2020</a>
<a href="#">Volume 13(5)/2020</a>
<a href="#">Volume 13(4)/2020</a>
<a href="#">Volume 13(3)/2020 (June, 30)</a>
<a href="#">Volume 13(2)/2020 (April, 30)</a>
<a href="#">Volume 13(1)/2020 (February, 28)</a>
<a href="#">Volume 12(6)/2019 (December, 30)</a>
<a href="#">Volume 12(5)/2019 (October, 30)</a>
<a href="#">Volume 12(4)/2019 (August, 30)</a>
<a href="#">Volume 12(3)/2019 (June, 30)</a>
<a href="#">Volume 12(2)/2019 (April, 30)</a>
<a href="#">Volume 12(1)/2019 (February, 28)</a>
<a href="#">Volume 11(6)/2018 (December, 30)</a>
<a href="#">Volume 11(5)/2018 (October, 30)</a>
<a href="#">Volume 11(4)/2018 (August, 30)</a>
<a href="#">Volume 11(3)/2018 (June, 30)</a>
<a href="#">Volume 11(2)/2018 (April, 30)</a>
<a href="#">Volume 11(1)/2018 (February, 28)</a>
<a href="#">Volume 10(6)/2017 (December, 30)</a>
<a href="#">Volume 10(5)/2017 (October, 30)</a>
<a href="#">Volume 10(4)/2017 (August, 30)</a>
<a href="#">Volume 10(3)/2017 (June, 30)</a>

## Volume 13(1)/2020

**Nugraha E., Gunawan R., Danapraja S., Yusrizal, Kusdinar A., Waluyo A. S., Hutajulu J., Prayitno H., Halim S., Sutisna D. H., 2020 The sea surface temperature effect on the length and size of skipjack tuna (*Katsuwonus pelamis*) catches in the Banda Sea, Indonesia. AAACL Bioflux 13(1):1-18.**

**Asadi M. A., Pambudi G. S, 2020 Diversity and biomass of mangrove forest within Baluran National Park, Indonesia. AAACL Bioflux 13(1):19-27.**

**Sulistijowati R., Karim Z., Junianto, 2020 The inhibition of *Vibrio alginolyticus* by the flavonoid extract of *Sonneratia alba* fruit. AAACL Bioflux 13(1):28-35.**

**Trianto A., Ambariyanto, Sarjito, Pramesti R., Soenarjo N., Hartati R., Taufiq-Spj N., Endrawati H., Destio, Yusidarta I., 2020 Ecological study and preliminary culture of the sponge *Candidaspongia* a source of anticancer molecules. AAACL Bioflux 13(1):36-45.**

**Hartanto., Tjahjono A., 2020 Plankton indexes and heavy metal pollution in Kendal coastal waters, Indonesia. AAACL Bioflux 13(1):46-63.**

**Akbar H., Syari I. A., Suyatna I., Putriningtias A., Bahri S., Destilawaty, Putra S. A., 2020 Condition of coral reefs in East Belitung, Bangka Belitung Islands, Indonesia. AAACL Bioflux 13(1):64-70.**

**Herawati E. Y., Khasanah R. I., Ambarwati M., Sofarini D., 2020 The effect of hydro-oceanographic factors on the community structure of plankton in natural and artificial coral reefs in Paiton waters. AAACL Bioflux 13(1):71-85.**

**Ahmad A., Fahrudin A., Boer M., Kamal M. M., Wardiatno Y., 2020 Ecosystem approach reef fisheries management model in Ternate Island, North Maluku, Indonesia. AAACL Bioflux 13(1):86-99.**

**Adelina A., Feliatra F., Siregar Y. I., Suharman I., 2020 Utilization of feather meal fermented *Bacillus subtilis* to replace fish meal in the diet of silver pompano, *Trachinotus blochii* (Lacepede, 1801). AAACL Bioflux 13(1):100-108.**

**Koshinski R., 2020 Effect of *Taraxacum officinale* Weber ex Wiggers extract on growth performance, biochemical blood parameters and meat quality of rainbow trout (*Oncorhynchus mykiss* W.) cultivated in a recirculating system. AAACL Bioflux 13(1):109-117.**

**Radona D., Kusmini I. I., Prihadi T. H., Khasani I., Astuti D. N., 2020 Hormonal induction for maturation, ovulation and successful level of spawning on striped**

- Volume 10(2)/2017 (April, 30) **snakehead (*Channa striata*). AACL Bioflux 13(1):118-124.**
- Volume 10(1)/2017 (February, 28) **Kusmana C., Iswahyudi, Hidayat A., Noorchamat B. P., 2020 Sustainable status of mangrove forest ecosystem management in Langsa City, Aceh, Indonesia. AACL Bioflux 13(1):125-136.**
- Volume 9(6)/2016 (December, 30)
- Volume 9(5)/2016 (October, 30)
- Volume 9(4)/2016 (August, 30) **Serdiati N., Yonarta D., Pratama F. S., Faqih A. R., Valen F. S., Tamam M. B., Hamzah Y. I. G., Hasan V., 2020 *Andinoacara rivulatus* (Perciformes: Cichlidae), an introduced exotic fish in the upstream of Brantas River, Indonesia. AACL Bioflux 13(1):137-141.**
- Volume 9(3)/2016 (June, 30)
- Volume 9(2)/2016 (April, 30)
- Volume 9(1)/2016 (February, 28) **Silaban B., Wattimena M. L., Nanlohy E. E. E. M., Lewerissa S., Silaban R., 2020 Morphometric and proximate analysis of mole crabs (*Hippa* genus) in Maluku Province, Indonesia. AACL Bioflux 13(1):142-151.**
- Volume 8(6)/2015 (December, 30)
- Volume 8(5)/2015 (October, 30)
- Volume 8(4)/2015 (August, 30) **Nieves P. M., Mendoza Jr. A. B., Bradecina S. R. B., 2020 Occurrence and recurrence: the fish kill story in Lake Buhi, Philippines. AACL Bioflux 13(1):152-158.**
- Volume 8(3)/2015 (June, 30)
- Volume 8(2)/2015 (April, 30) **Kristiana V., Mukti A. T., Agustono, 2020 Increasing growth performance of Nile tilapia (*Oreochromis niloticus*) by supplementation of noni *Morinda citrifolia* fruit extract via diet. AACL Bioflux 13(1):159-166.**
- Volume 8(1)/2015 (February, 28)
- Volume 7(6)/2014 (December, 30)
- Volume 7(5)/2014 (October, 30) **Ndobe S., Rusaini, Masyahoro A., Serdiati N., Madinawati, Moore A. M., 2020 Reproductive and morphometric characteristics of climbing perch *Anabas testudineus* in Sigi, Central Sulawesi, Indonesia. AACL Bioflux 13(1):167-182.**
- Volume 7(4)/2014 (August, 30)
- Volume 7(3)/2014 (June, 30) **Sayed J. A., Iqbal H., Ayenuddin H., Al-Amin S., Samsad K., Amirun N., Kumar P. A., Shahanul I. M., 2020 A semi intensive approach on growth and profit margin of Indian major carps (*Catla catla*, *Labeo rohita* and *Cirrhinus cirrhosus*) with cost effective standard feed formulation. AACL Bioflux 13(1):183-193.**
- Volume 7(2)/2014 (April, 15)
- Volume 7(1)/2014 (February, 15)
- Volume 6(6)/2013 (November, 15)
- Volume 6(5)/2013 (September, 15) **Tjendanawangi A., Dahoklory N., 2020 The effect of macroalgal species and content of feed formulation on sea urchin *Tripneustes gratilla* gonad quality. AACL Bioflux 13(1):194-198.**
- Volume 6(4)/2013 (July, 25)
- Volume 6(3)/2013 (May, 15) **Rostika R., Azhima M. F., Ihsan Y. N., Andriani Y., Suryadi I. B. B., Dewanti L. P., 2020 The use of solid probiotics in feed to growth and survival rate of *mantap* common carp (*Cyprinus carpio*). AACL Bioflux 13(1):199-206.**
- Volume 6(2)/2013 (March, 15)
- Volume 6(1)/2013 (January, 15)
- Volume 5(5)/2012 (December, 30) **Endrizalova J., Didenko A., Pavlinsky S., Manko P., 2020 Diet and feeding niche of five invasive species in the Bodrog River watershed. AACL Bioflux 13(1):207-217.**
- Volume 5(4)/2012 (September, 30) **Doctolero J. S., Vera Cruz E. M., Velasco R. R., 2020 Morpho-behavioral and hematological responses of vaccinated proactive and reactive Nile tilapia (*Oreochromis niloticus*) subjected to handling stress. AACL Bioflux 13(1):218-229.**
- Volume 5(3)/2012 (July, 30)
- Volume 5(2)/2012 (June, 30)
- Volume 5(1)/2012 (March, 15)
- Volume 4(5)/2011 (December, 30) **Kurnialahi I. D., Husni A., Sinurat E., Isnansetyo A., 2020 Antioxidant activity of tropical seaweed *Sargassum muticum* fucoidan. AACL Bioflux 13(1):230-240.**
- Volume 4(4)/2011 (October, 30)
- Volume 4(3)/2011 (July, 30) **Suciyono, Anggara D. P., Azhar M. H., Ulkhaq M. F., Fasya A. H., 2020 Characteristics of *Acropora divaricata* and *Acropora nobilis* on different depths**




Volume 4(2)/2011 (April, 30)	<b>transplantation based on growth rate and zooxanthellae density. AACL Bioflux 13(1):241-249.</b>
Volume 4(1)/2011 (January, 30)	
Volume 3(5)/2010 (December, 5)	<b>Putri N. A., Sulmartiwi L., Pursetyo K. T., 2020 The histopathology of antique ark's mantle (<i>Anadara antiquata</i>) post-depuration with the shells' filtration. AACL Bioflux 13(1):250-260.</b>
Volume 3(4)/2010 (December, 1)	
Volume 3(3)/2010 (November, 15)	
Volume 3(2)/2010 (July, 30)	Pujiyati S., Hananya A., Retnoaji B., Lubis M. Z., 2020 Bioacoustic of <i>Anguilla bicolor</i> (Mc Clelland, 1844) feeding behaviour under controlled conditions. AACL Bioflux 13(1):261-267.
Volume 3(1)/2010 (February, 28)	
Volume 2(4)/2009 (October, 30)	Asriyana A., Halili H., Irawati N., 2020 Size structure and growth parameters of striped eel catfish ( <i>Plotosus lineatus</i> ) in Kolono Bay, Southeast Sulawesi, Indonesia. AACL Bioflux 13(1):268-279.
Volume 2(3)/2009 (July, 30)	
Volume 2(2)/2009 (April, 30)	...
Volume 2(1)/2009 (January, 30)	...
Volume 1(2)/2008 (December, 30)	...
Volume 1(1)/2008 (September, 30)	...
Volume Pilot/2007 (December, 30) - available printed only	
Pontus Euxinus, Volume 1 (1980) - Parent Journal	



design: [www.simple-webdesign.com](http://www.simple-webdesign.com)

Home | Archive | Volume 1 (1) / 2008 | CEEX 140 | Volume 1 (2) / 2008 | Volume 2 (1) / 2009 | Volume 2 (2) / 2009 | Volume 2 (3) / 2009 | Volume 2 (4) / 2009 | Pilot | Volume 3 (1) / 2010 | Volume 3(2)/2010 (July, 30) | Volume 3(3)/2010 | Volume 3(4)/2010 | Volume 3(5)/2010 - ACVAPEDIA 2010 | Volume 4(1)/2011 | Volume 4(2)/2011 - ACVAPEDIA 2010 | Volume 4(3)/2011 | Volume 4(4)/2011 | Volume 4(5)/2011 | Volume 5(1)/2012 (March, 15) | Volume 5(2)/2012 | Volume 5(3)/2012 | Volume 5(4)/2012 | Volume 5(5)/2012 (December, 30) | Volume 6(1)/2013 - ACVAPEDIA 5th edn., Hungary, Szarvas (HAKI), 27-29th of November, 2012 | Volume 6(2)/2013 - ACVAPEDIA 5th edn., Hungary, Szarvas (HAKI), 27-29th of November, 2012 | Volume 6(3)/2013 | Volume 6(4)/2013 | Volume 6(5)/2013 | Volume 6(6)/2013 | Volume 7(1)/2014 | Volume 7(2)/2014 | Volume 7(3)/2014 | Volume 7(4)/2014 | Volume 7(5)/2014 | Volume 7(6)/2014 | Volume 8(1)/2015 | Volume 8(2)/2015 | Volume 8(3)/2015 | Volume 8(4)/2015 | Volume 8(5)/2015 | Volume 8(6)/2015 | Volume 9(1)/2016 | Volume 9(2)/2016 | Volume 9(3)/2016 | Volume 9(4)/2016 | Volume 9(5)/2016 | Volume 9(6)/2016 | Volume 10(1)/2017 | Volume 10(2)/2017 | Volume 10(3)/2017 | Volume 10(4)/2017 | Volume 10(5)/2017 | Volume 10(6)/2017 | Volume 11(1)/2018 | Volume 11(2)/2018 | Volume 11(3)/2018 | Volume 11(4)/2018 | Pontus Euxinus, Volume 1, 1980 | Volume 11(5)/2018 | Volume 11(6)/2018 | Volume 12(1)/2019 | Volume 12(2)/2019 | Volume 12(3)/2019 | Volume 12(4)/2019 | Volume 12(5)/2019 | Volume 12(6)/2019 | Volume 13(1)/2020 | Volume 13(2)/2020 | Contact | Site Map



# Increasing growth performances of Nile tilapia (*Oreochromis niloticus*) by supplementation of noni *Morinda citrifolia* fruit extract via diet

<sup>1</sup>Vini Kristiana, <sup>2</sup>Akhmad T. Mukti, <sup>2</sup>Agustono

<sup>1</sup> Study Program of Aquaculture, Faculty of Fisheries and Marine, Universitas Airlangga, Surabaya 60115, Indonesia; <sup>2</sup> Department of Fish Health Management and Aquaculture, Faculty of Fisheries and Marine, Universitas Airlangga, Surabaya 60115, Indonesia. Corresponding author: A. T. Mukti, [akhmad-t-m@fpk.unair.ac.id](mailto:akhmad-t-m@fpk.unair.ac.id)

**Abstract.** High feed prices and its low nutritional quality are the inhibitions in the process of red tilapia fish cultivation. Therefore, the use of feed additives in feed becomes significant in order to optimize fish growth. Especially the ones made of natural ingredients, such as noni (*Morinda citrifolia*) fruit. The purpose of this study was to determine the effect of added noni fruit extract to the growth rate and protein retention of red tilapia fish. The method used was an experimental method with a completely randomized design, consisting of four replications and four treatments of noni fruit extract dosages given to red tilapia. The treatments used were the dosage amount of noni extraction given to red tilapia, namely P0: commercial feed as the control group without noni fruit extract + 2 g Progol (fish feed binder); P1: commercial feed 1 kg + noni fruit 100 mL kg<sup>-1</sup> feed + 2 g Progol; P2: commercial feed 1 kg + noni fruit 300 mL kg<sup>-1</sup> feed + 2 g Progol; P3: commercial feed 1 kg + noni fruit 500 mL kg<sup>-1</sup> feed + 2 g Progol. The parameters observed were specific growth rates, feed conversion ratio, survival rate, and protein retention of red tilapia. The result showed that the addition of noni fruit extract in commercial feed with different doses significantly affected the specific growth rate, daily length growth rate, feed conversion ratio, and protein retention of red tilapia. The survival rate from the preservation of red tilapia is not significantly different between treatments. The best result for the specific growth rate is P3 (500 mL kg<sup>-1</sup>) of 0.0150% day<sup>-1</sup>, daily length growth rate is P3 (500 mL kg<sup>-1</sup>) of 1.599% day<sup>-1</sup>, feed conversion ratio is P2 (300 mL kg<sup>-1</sup>) of 0.846%, survival rate of 100% in all treatments, and protein retention with the best results is P1 (10 mL kg<sup>-1</sup>) of 12.6246%. The addition of noni fruit extract in commercial feed can increase the specific growth rate and daily length, lowering the value of feed conversion ratio in red tilapia, and increasing the protein retention of red tilapia (*Oreochromis niloticus*).

**Key Words:** fish meal substitution, noni fruit (*Morinda citrifolia*) extract, red tilapia (*Oreochromis niloticus*).

**Introduction.** Red tilapia is one of the freshwater fish commodities that are currently in high demand for consumption and cultivation. The meat of red tilapia is delicious, solid, easy to serve, does not have many thorns, is a source of animal protein, and is relatively inexpensive (Siddiqui & Al-Harbi 1995; Zonneveld & Fadholi 1991). Red tilapia is popular within the community, making it an opportunity for fish farmers to increase the productivity of their ponds. Therefore, the availability of red tilapia for consumption needs to be increased. The demand for red tilapia continues to increase in the international market, mainly the American and European Union markets (Widanarni et al 2012). Red tilapia is one of the export fish which provided as a whole or in the form of a filet. The effort to increase the export of red tilapia is by increasing its production through cultivation.

The feed is one of the essential factors in the agriculture business. Feed is an energy source for fish to grow and develop (Younos et al 1990). The availability of adequate and balanced feed will affect the growth and survival of cultivated fish (Tiamiyu 2016). Feed must contain all necessary nutrients such as carbohydrates, fats, proteins, minerals, vitamins, and feed additives with sufficient and balanced amounts (Halver & Hardy 2003). High feed prices and low nutritional quality are inhibitions in the fish farming process. Feed as an energy source for growth is the most significant amount of

production costs component (Shepherd & Jackson 2013). One way to overcome this problem is to use feed additives. Feed additives in fish farming, especially red tilapia, are ingredients that are intentionally used to increase appetite, digestibility, endurance, reduce stress levels, and spur fish growth (Alemayehu et al 2018).

The provision of feed is considered necessary in fish farming, including the addition of feed additives to spur growth and production. Phytobiotics is one type of natural feed additive derived from plants (Vidanarachchi et al 2005). The purpose of adding natural feed additives to the feed is to spur growth, as well as improve the health and productivity of fish (Citarasu 2001; Citarasu et al 1998, 1999, 2002; Sivaram et al 2004). Noni fruit can be used as phytobiotic (feed additive) because it has antibacterial effects and bioactive substances such as alkaloids, flavonoids, and tannins which are very effective in suppressing the growth of pathogenic bacteria and improving intestinal morphological characteristics (Murdiati & Adiwinatai 2000). Bioactive substances in noni include xeronine and precursor of xeronine (proxeronine). Proxeronine will be converted to xeronine in the intestine by proxeronase enzymes and other substances. Xeronine increases the permeability of cell membranes, facilitates the transfer of peptides and increases nutrient absorption, as well as effectuates amino acids, vitamins, and minerals. Xeronine can activate several enzymes and regulate protein synthesis (Solomon 2001). Therefore, noni fruit extract can be used as an additional ingredient in feed to help the digestion process of fish, so that the consumed feed can be optimally utilized for growth and feed efficiency, as well as the protein content of fish. This study aims to determine the potential of given noni fruit (*Morinda citrifolia*) extract to the growth rate and protein retention of red tilapia (*Oreochromis niloticus*).

## Material and Method

**Preparation of red tilapia (*Oreochromis niloticus*) and rearing conditions.** This research was conducted from April to May 2017 at the Education Laboratory of the Airlangga University, Faculty of Fisheries and Marine, Surabaya, Indonesia. The proximate test of feed and protein retention test were conducted at the Laboratory of the Faculty of Veterinary Medicine, Universitas Airlangga, Surabaya, Indonesia. The instruments used in this study were an aquarium measuring 45 cm x 30 cm x 30 cm, a set of aerators, digital scales, pH pens, thermometers, rulers, plastic bags, plastic hoses with a diameter of 1 cm, dissolved oxygen (DO) meters, and ammonia kits. The fish used were 140 red tilapia fry in healthy condition, measuring 5-6 cm total length, weighing 2-5 g head<sup>-1</sup>. The red tilapia fry were obtained from the Gunung Sari fish market, Surabaya, East Java. Noni fruit extract was obtained from the result of squeezed noni fruit. The noni fruit used was ripe noni fruit, which is yellow and soft with a distinctive smell.

**Experimental design.** This research method was experimental. The primary test parameters in this study were protein retention, specific growth rate, feed conversion ratio, and survival rate of red tilapia. The variables observed were independent variables, dependent variables, and control variables. The independent variable is the dose of noni extract. The dependent variables are specific growth rate, feed conversion ratio, survival rate, and protein retention. Control variables are water quality (temperature, pH, DO, and ammonia). This study used a Completely Randomized Design (CRD) with four treatments and four replications, which treatments explained as follows:

P0: commercial feed as the control group without noni fruit extract + 2 g Progol (fish feed binder);

P1: commercial feed 1 kg + noni fruit 100 mL kg<sup>-1</sup> feed + 2 g Progol;

P2: commercial feed 1 kg + noni fruit 300 mL kg<sup>-1</sup> feed + 2 g Progol;

P3: commercial feed 1 kg + noni fruit 500 mL kg<sup>-1</sup> feed + 2 g Progol.

The feed given in this study was PF 1000 commercial feed from PT Matahari Sakti, with a protein content of 39-41%, 5% lipid, 6% crude fiber, 16% ash, and 10% moisture content.



**Preparation of the experimental diets.** First, ripe noni fruits were cleaned with water, then washed with warm water. The noni fruits were crushed and then squeezed by using strain cloth. The squeezed noni fruit extract was put in a bottle and stored at room temperature.

The commercial feed was mixed with noni fruit extract and 2 grams of Progol with a different amount in each treatment, as previously explained. The commercial feed and noni fruit extract were smoothly mixed. The mixed feed was dried in an open room and was aerated until dry. After that, the dried food was stored in a plastic jar. The mixed fish feed with different doses were then analyzed through the proximate test in the laboratory. The nutrient content of the feed used in this study can be seen in Table 1 as follows.

Table 1

Nutrient content of feed with various doses of noni fruit extract

Analysis	P0 (Control)	P1 (100 mL kg <sup>-1</sup> )	P2 (300 mL kg <sup>-1</sup> )	P3 (500 mL kg <sup>-1</sup> )
Dry ingredients (%)	90.1441	87.3484	86.7339	84.5036
Ash (%)	8.9591	8.7352	8.6875	8.5261
Crude protein (%)	29.1112	30.9015	30.2753	29.2539
Crude lipid (%)	8.3165	7.6176	6.9983	6.872
Crude fiber (%)	3.5026	2.488	2.6143	2.2593
Ca	2.5824	2.4109	2.4069	2.1481
Nitrogen-free extract (%)	40.2547	37.6061	38.1585	37.5923
Metabolism energy (Kcal kg <sup>-1</sup> )	3057.9622	2967.4828	2922.6623	2857.8501

**The analytical procedure of the diets.** The aquarium was then filled with 30 liters of water, aerated, and then left for 24 hours. The fish were selected based on a similar length and weight and its healthy condition. The fish were acclimatized for one week and were fasted for 24 hours before being treated. The fish were placed in fourteen aquariums according to the treatment given with a stocking density of 10 fish per aquarium. The fish were kept for thirty days and fed according to the treatment doses. Feeding was given three times a day: morning, afternoon, and evening, with a dose of 3% of body weight.

Below is the protein retention (PR) calculation based on a formula from Tung & Shiau (1991):

$$PR = \frac{\text{Ln final body weight} - \text{Ln initial body weight (g)}}{\text{Total protein intake (g)}} \times 100$$

The specific growth rate was calculated based on a formula from Effendie (1997):

$$SGR = \frac{\text{LnWt} - \text{LnWo}}{t} \times 100$$

where: SGR = specific growth rate (% day<sup>-1</sup>);

W0 = initial body weight (g);

Wt = final body weight (g);

T = treatment time (in days).

The daily length growth rate (LGR) was calculated based on a formula from Aggraeni & Abdulgani 2013):

$$LGR = \frac{\text{LnLt} - \text{LnLo}}{t} \times 100$$

where: LGR = daily length growth rate (% day<sup>-1</sup>);

Lt = average total length on day t (cm);

Lo = average total length on day 0 (cm);

t = total of observation time (in days).

The survival rate was calculated based on a formula from Wirabakti (2006):

$$SR = \frac{N_t}{N_o} \times 100$$

where: SR = survival rate (%);

N<sub>t</sub> = the number of survived fish at the end of the experiment (in fish);

N<sub>o</sub> = the number of survived fish at the start of the experiment (in fish).

The feed conversion ratio was calculated based on a formula from Council (1997):

$$FCR = \frac{F}{(W_t + D) - W_o}$$

where: FCR = feed conversion ratio;

W<sub>o</sub> = the biomass weight of fish at the start of the experiment (g);

W<sub>t</sub> = the biomass weight of fish at the end of the experiment (g);

D = total weight of dead fish (g);

F = feed intake (g).

**Statistical analysis.** Data analysis using Variant Analysis (ANOVA) with the study design was a Completely Randomized Design (CRD) to determine the differences in treatment. If there is a difference in treatment, Duncan's Multiple Range Test is applied.

**Results.** The calculation results of the average value of the specific growth rate, feed conversion ratio, and survival rate of red tilapia using statistical analysis after being treated for 30 days can be seen in Table 2.

Table 2

Average value of the specific growth rate, survival rate, feed conversion ratio, daily length growth, and protein retention of red tilapia

Treatment	Specific growth rate (% day <sup>-1</sup> )	Survival rate (%)	Feed conversion ratio (%)	Daily length growth rate (% day <sup>-1</sup> )	Protein retention (%)
P0	0.0105±0.0007 <sup>c</sup>	100	2.712±0.588 <sup>b</sup>	1.070±0.189 <sup>b</sup>	7.3345±0.3434 <sup>c</sup>
P1	0.0135±0.0006 <sup>b</sup>	100	1.157±0.226 <sup>a</sup>	1.145±0.050 <sup>b</sup>	12.6246±0.3074 <sup>b</sup>
P2	0.0147±0.0005 <sup>a</sup>	100	0.846±0.234 <sup>a</sup>	1.435±0.186 <sup>a</sup>	11.5692±0.4054 <sup>a</sup>
P3	0.0150±0.0009 <sup>a</sup>	100	0.918±0.436 <sup>a</sup>	1.599±1.312 <sup>a</sup>	11.4289±0.7356 <sup>a</sup>

P0 (0% noni fruit extract), P1 (100 mL kg<sup>-1</sup> noni fruit extract), P2 (300 mL kg<sup>-1</sup> commercial feed with noni fruit extract), P3 (500 mL kg<sup>-1</sup> commercial feed with noni fruit extract). Different superscripts show significant differences.

According to Table 2, the value of the specific growth rate (SGR) of red tilapia was 0.0105-0.0150% day<sup>-1</sup>. The results of the statistical analysis of added noni fruit extract on commercial feed to the SGR of red tilapia showed significantly different results (p < 0.05). Duncan's Multiple Range Test (DMTR) results showed that the highest growth rate in this study was P3 (0.0150% day<sup>-1</sup>) which was not significantly different from P2 (0.0147% day<sup>-1</sup>), while it was significantly different with P0 (0.0105% day<sup>-1</sup>) and P1 (0.0135% day<sup>-1</sup>). The lowest specific growth rate was P0 (0.0105% day<sup>-1</sup>) which was significantly different from P1 (0.0135% day<sup>-1</sup>), P2 (0.0147% day<sup>-1</sup>), and P3 (0.0150% day<sup>-1</sup>).

The feed conversion ratio (FCR) of red tilapia was 0.846-2.712%. The results of statistical analysis of the added noni fruit extract to commercial feed to FCR of red tilapia showed significantly different results (p < 0.05). Based on the results of DMTR, the highest FCR in this study was P0 (2.712%) which was significantly different from P1 (1.157%), P2 (0.846%), and P3 (0.918%). While the lowest feed conversion ratio was P2 (0.846%) and was not significantly different from P1 (1.157%), P3 (0.918%), and significantly different from P0 (2.712%). These results indicate that the addition of noni fruit extract to commercial feed can reduce feed conversion ratio of red tilapia.

The survival rate of red tilapia from all treatments was 100%. The average survival rate of red tilapia showed no significant difference ( $p > 0.05$ ). Based on the results of DMTR, the survival rates of P0, P1, P2, and P3 were not significantly different.

The value of the daily length growth rate of red tilapia for thirty days of preservation was 1.070-1.599% day<sup>-1</sup>. The results showed significant differences ( $p < 0.05$ ). The highest daily length growth value was at P3 (1.599% day<sup>-1</sup>) which was not significantly different from P2 (1.435% day<sup>-1</sup>) but was significantly different from P0 (1.070% day<sup>-1</sup>) and P1 (1.145% day<sup>-1</sup>). The lowest daily length growth rate was P0 (1.070% day<sup>-1</sup>) which was not significantly different from P1 (1.145% day<sup>-1</sup>) but significantly different from P2 (1.435% day<sup>-1</sup>) and P3 (1.599% day<sup>-1</sup>).

The results of the statistical analysis of the addition of noni fruit extract to commercial feed showed significantly different results ( $p < 0.05$ ) to the protein retention of red tilapia. The results of DMTR showed that the highest protein retention value was P1 (12.6246%) which was significantly different from P0 (7.3345%), P2 (11.5692%), and P3 (11.4289%). Protein retention value in P2 (11.5692%) was not significantly different from P3 (11.4289%) but was significantly different from P0 (7.3345%) and P1 (12.6246%). The lowest protein retention value was P0 (7.3345%), which was significantly different from P1 (12.6246%), P2 (11.5692%), and P3 (11.4289%).

Data on the range of water quality parameters for red tilapia preservation for thirty days can be seen in Table 3.

Table 3

Water quality range value of red tilapia in 30 days of preservation

<i>Parameter</i>	<i>Range</i>
Temperature (°C)	27.4-29.2
pH	7.2-7.9
DO (mg L <sup>-1</sup> )	3.3-5
Ammonia (mg L <sup>-1</sup> )	0-0.5

**Discussion.** The noni fruit extract mixed in the commercial feed had a significantly different effect on specific growth rate, daily length growth rate, feed conversion ratio, and protein retention of red tilapia for thirty days of preservation. These effects are due to the optimum absorption process or nutrient absorption of the feed in the digestive tract. Noni extract contains anthraquinone or acidic ingredients that can affect the pH of the digestive tract to be more acidic (Widianto et al 2015). The statement is right considering that in an acidic environment, protein-breaking enzymes (pepsin) can work optimally, so that protein intakes are absorbed more by the body which supports the optimal fish growth rate (Esminger et al 1990).

The extract of noni fruit contains proxeronine and xeronine. Proxeronine is converted to xeronine with the help of proxeronase enzymes. Xeronine is an active compound that has a role in facilitating metabolism inside the body of the animals and increasing nutrient absorption (Solomon 2001). Xeronine in noni fruit can function to expand the absorption in the intestine surface and facilitate the absorption process of food so the nutrients in the feed can be absorbed optimally (Wang et al 2008). The optimal absorption process will affect increased protein retention (Schlegel 1987; Frazier & Westhoff 1988; Cui et al 1992; Solomon 2001; Sjöfjan 2003; Sujono 2006; Widianto et al 2015). Increased protein retention in the treatment results was caused by adding noni fruit extract in commercial feed, which can suppress the growth of microbial pathogens and increase non-pathogenic microbes in the digestive tract, resulting in the ability to decompose feed ingredients, especially crude fiber. The use of noni fruit extract can inhibit the growth of pathogenic bacteria in the intestine so that the number of non-pathogenic bacteria in the intestine was increased (Solomon 2001). Noni extract can increase the absorption process in the digestive tract and help decompose feed ingredients, especially crude fiber (Widianto et al 2015). Non-pathogenic bacteria can produce enzymes that digest crude fiber, protein, and lipid (Sjöfjan 2003). *Bacillus* sp. bacteria in the digestive tract can break down proteins into amino acids (Sujono 2006).

These amino acids are used by bacteria to multiply themselves. Bacteria are a source of single-cell proteins so that the bacteria multiplication can increase protein feed and reduce crude fiber (Schlegel 1987).

Moreover, *Bacillus* sp. is a pectinolytic bacteria, which is an organism that can produce pectin or complex carbohydrates (Frazier & Westhoff 1988). The characteristics of *Bacillus* sp. bacteria are proteolytic, saccharolytic, and pectinolytic, which increases protein and carbohydrate feed. Protein retention indicates the amount of additional body protein from the consumed feed protein. The feed consumed was a source of protein used for preservation, metabolic activity, and growth (Cui et al 1992). The high protein retention in P1 treatment was caused by the increase in protein and carbohydrate content by the activity of non-pathogenic microbes that increase due to the addition of noni fruit extract in the feed. The only issue was the more noni fruit extract dosage in the feed cannot help at breaking down the crude fiber and result in higher crude fiber amount in the feed.

The feed conversion ratio is one of the standards to determine the quality of feed given to livestock in fulfilling the required nutrition. The higher the value of feed conversion, the worse the quality of the nutritional value of the feed. Noni fruit contains saponins which are bioactive in the growth of animals and digestive microbes. Giving saponins can increase the permeability of cell walls in the intestine, increase absorption of food substances hence result in higher conversion value (Önning et al 1996).

Survival rates of red tilapia were not significantly different between treatments. Thirty days preservation of red tilapia with the treatments given did not cause death in fish. This proves that adding noni fruit extract in commercial feed did not contain toxic ingredients that cause death. One study using noni fruit extract with a dose of 15,000 mg per kg body weight in experimental animals was not found any sign of toxic effects (Krishnakumar et al 2015). The study had performed surgery on experimental animals but still did not find any signs of toxicity in the animals' organs. Noni fruit is categorized as non-toxic substances (Wang et al 2008; Younos et al 1990).

Water quality in this study is an essential supporting parameter for the survival rate of red tilapia. Water as a living medium for fish that is preserved must meet the requirements. Water quality needs to be controlled in order to maintain its optimal conditions, thus creating an environment that is suitable for the habitat of red tilapia. The supporting parameters of this study are water quality parameters such as temperature, DO, pH, and ammonia. The temperature in this study averaged around 27.4-29.2°C. This temperature is good enough for the survival of red tilapia. The optimal water temperature for red tilapia is 25-30°C (Kungvakij & Pudadera 1984). Temperature is an environmental factor that influences the body metabolism speed. The optimum speed metabolism for fish will be achieved at optimal temperatures. The pH value has an essential influence on the survival of fish in the water. The pH in this study ranged from 7.2 to 7.9, which is quite adequate for the survival rate of red tilapia. The pH range required by fish is 6-9 (DeLong et al 2009). The average DO content in this study ranged from 3.3 to 5 mg L<sup>-1</sup>. Based on these results, the DO during the study indicated a suitable level for the survival of red tilapia (Yildiz et al 2017). Thus DO in the preservation of red tilapia met the requirements. Ammonia in the water is derived from fish feces and from the rest of the feed that is not consumed by fish. In general, the concentration of ammonia contained in the water must not exceed 1 mg L<sup>-1</sup> (BFAR-NFFTC 2000; Indian Council of Agricultural Research 2006). Ammonia during this study was 0-0.5 mg L<sup>-1</sup>, which is suitable for the preservation of red tilapia.

**Conclusions.** Based on this research, it can be concluded that the addition of noni fruit extract in commercial feed can increase specific growth rate, daily length growth rate, lower the feed conversion ratio, and increase protein retention of red tilapia (*Oreochromis niloticus*). Moreover, the addition of noni fruit extract in commercial feed did not show any toxic effect on the survival rate of red tilapia during the thirty days of preservation.

**Acknowledgements.** The authors would like to express their gratitude to all members of Study Program of Aquaculture, Faculty of Fisheries and Marine, Universitas Airlangga, Indonesia.

## References

- Aggraeni N. M., Abdulgani N., 2013 Pengaruh pemberian pakan alami dan pakan buatan terhadap pertumbuhan ikan betutu (*Oxyeleotris marmorata*) pada skala laboratorium. Jurnal Sains dan Seni Pomits 2(1):197-201. [in Indonesian]
- Alemayehu T. A., Geremew A., Getahun A., 2018 The role of functional feed additives in tilapia nutrition. Fisheries and Aquaculture Journal 9(2):249.
- BFAR-NFFTC, 2000 Basic biology of tilapia. NFFTC Aqua-Leaflet No. 2000-06 6, 2 pp..
- Citarasu T., Babu M. M., Punitha S. M. J., Venket Ramalingam K., Marian M. P., 2001 Control of pathogenic bacteria using herbal biomedicinal products in the larviculture system of *Penaeus monodon*. In: International conference on advanced technologies in fisheries and marine sciences. MS University, India, pp. 104.
- Citarasu T., Immanuel G., Marian M. P., 1998 Effects of feeding artemia enriched with stresstol and cod liver oil on growth and stress resistance in the Indian white shrimp *Penaeus indicus* postlarvae. Asian Fisheries Science 12:65-75.
- Citarasu T., Jayarani T., Babu M., Marian M., 1999 Use of herbal bio-medicinal products in aquaculture of shrimp. In: Aqua-Terr Annual Symposium. School of Biological Sciences, MK University, Madurai.
- Citarasu T., Babu M. M., Raja Jeya Sekar R., Peter Marian M., 2002 Developing artemia enriched herbal diet for producing quality larvae in *Penaeus monodon*, Fabricius. Asian Fisheries Science 15:21-32.
- Council N. R., 1977 Nutrient requirement of warmwater fishes. The National Academic Press, Washington DC, 87 pp.
- Cui Y., Liu X., Wang S., Chen S., 1992 Growth and energy budget in young grass carp, *Ctenopharyngodon idella* Val., fed plant and animal diets. Journal Fish Biology 41(2):231-238.
- DeLong D. P., Losordo T. M., Rakocy J. E., 2009 Tank culture of tilapia. SRAC Publication No. 282. North Carolina State University, Southern Regional Aquaculture Center.
- Effendie M. I., 1997. Biologi perikanan. Yayasan Pustaka Nusatama, Yogyakarta, 163 pp. [in Indonesian]
- Esminger M. E., Oldfield J. E., Heinemann W. W., 1990 Feed and nutrition. The Ensminger Publishing Company Clovis, California.
- Frazier W. C., Westhoff D. C., 1988 Food microbiology. 4th edition, McGraw-Hill, New York, 539 pp.
- Halver J. E., Hardy R. W., 2003 Fish nutrition. 3rd edition, Academic Press, San Diego, 500 pp.
- Indian Council of Agricultural Research, 2006 Handbook of fisheries and aquaculture. New Delhi, 755 pp.
- Krishnakumar N. M., Latha P. G., Suja S. R., Rajasekharan S., 2015 A review on the ethnomedicinal, therapeutic, and nutraceutical importance of 'noni' (*Morinda citrifolia* L.). International Journal of Medicinal Plants and Natural Products 1(3):1-14.
- Kungvakij P., Pudadera B., 1984 Physico-chemical factors for fish culture in pond. UNDP/FAO Network of Aquaculture Centers in Asia, Philippines, pp. 1-11.
- Murdiati T. B., Adiwinata G., Hildasari D., 2000 Penuluruhan senyawa aktif dari buah mengkudu (*Morinda citrifolia*) dengan aktivitas antelmintik terhadap *Haemonchus contortus*. Jurnal Ilmu Ternak dan Veteriner 5(4):255-259. [in Indonesian]
- Önning G., Wang Q., Weström B. R., Asp N. G., Karlsson B., 1996 Influence of oat saponins on intestinal permeability in vitro and in vivo in the rat. British Journal of Nutrition 76(1):141-151.
- Schlegel H., 1987 General microbiology. 6th edition, Cambridge University Press, Cambridge, 587 pp.
- Shepherd C. J., Jackson A. J., 2013 Global fishmeal and fish-oil supply: inputs, outputs and markets. Journal of Fish Biology 83(4):1046-1066.

- Siddiqui A. Q., Al-Harbi A. H., 1995 Evaluation of three species of tilapia, red tilapia and a hybrid tilapia as culture species in Saudi Arabia. *Aquaculture* 138(1-4):145-157.
- Sivaram V., Babu M. M., Immanuel G., Murugadass S., Citarasu T., Marian M. P., 2004 Growth and immune response of juvenile greasy groupers (*Epinephelus tauvina*) fed with herbal antibacterial active principle supplemented diets against *Vibrio harveyi* infections. *Aquaculture* 237(1-4):9-20.
- Sjofjan O., 2003 Kajian probiotik (*Aspergillus niger* dan *Bacillus* spp.) sebagai imbuhan ransum dan implikasi efeknya terhadap mikroflora usus serta penampilan produksi ayam petelur. PhD thesis, Universitas Padjajaran, Bandung, 100 pp. [in Indonesian]
- Solomon N., 2001 The Noni phenomenon. 21st edition, Direct Source Publishing, Utah, 296 pp.
- Sujono A. H., 2006 Pengaruh penggunaan tepung buah mengkudu (*Morinda citrifolia*) terhadap penambahan bobot badan dan tampilan pakan pada ayam pedaging. *Jurnal Protein* 13(1):10-16. [in Indonesian]
- Tiamiyu L. O., Okomoda V. T., Aende A., 2016 Growth performance of *Oreochromis niloticus* fingerlings fed *Moringa oleifera* leaf as replacement for soybean meal. *Journal of Aquaculture Engineering and Fisheries Research* 2(2):61-66.
- Tung P. H., Shiau S. Y., 1991 Effect of meal frequency on growth performance of hybrid tilapia *Oreochromis niloticus* × *O. aureus*, fed different carbohydrate diets. *Aquaculture* 92:343-350.
- Vidanarachchi J. K., Mikkelsen L. L., Sims I. M., Iji P., Choct M., 2005 Phytobiotics: alternatives to antibiotic growth promoters in monogastric animal feeds. *Recent Advances in Animal Nutrition* 15:131-144.
- Wang M. Y., Nowicki D., Anderson G., Jensen J., West B., 2008 Liver protective effects of *Morinda citrifolia* (Noni). *Plant Foods for Human Nutrition* 63(2):59-63.
- Widanarni, Ekasari J., Maryam S., 2012 Evaluation of biofloc technology application on water quality and production performance of red tilapia *Oreochromis* sp. cultured at different stocking densities. *HAYATI Journal of Bioscience* 19(2):73-80.
- Widianto B., Prayogi H. S., Nuryadi, 2015 Pengaruh penambahan tepung buah mengkudu *Morinda citrifolia* L. dalam pakan terhadap penampilan produk itik hibrida. *Jurnal Ilmu-Ilmu Peternakan* 25(2):28-35. [in Indonesian]
- Wirabakti M. C., 2006 Laju pertumbuhan ikan nila merah *Oreochromis niloticus* L. yang dipelihara pada perairan rawa dengan sistem keramba dan kolam. *Journal of Tropical Fisheries* 1:61-67. [in Indonesian]
- Yildiz H. Y., Robaina L., Pirhonen J., Mente E., Dominguez D., Parisi G., 2017 Fish welfare in aquaponic systems: its relation to water quality with an emphasis on feed and faeces: a review. *Water* 9(1):13.
- Younos C., Rolland A., Fleurentin J., Lanhers M. C., Misslin R., Mortier F., 1990 Analgesic and behavioural effects of *Morinda citrifolia*. *Planta Medica* 56(5):430-434.
- Zonneveld N., Fadholi R., 1991 Feed intake and growth of red tilapia at different stocking densities in ponds in Indonesia. *Aquaculture* 99(1-2):83-94.

Received: 26 August 2019. Accepted: 30 November 2019. Published online: 29 January 2020.

Authors:

Vini Kristiana, Study Program of Aquaculture, Faculty of Fisheries and Marine, Universitas Airlangga, Surabaya 60115, Indonesia, e-mail: vinikristiana1995@gmail.com

Akhmad Taufiq Mukti, Department of Fish Health Management and Aquaculture, Faculty of Fisheries and Marine, Universitas Airlangga, Surabaya 60115, Indonesia, e-mail: akhmad-t-m@fpk.unair.ac.id

Agustono, Department of Fish Health Management and Aquaculture, Faculty of Fisheries and Marine, Universitas Airlangga, Surabaya 60115, Indonesia, e-mail: agustono@fpk.unair.ac.id

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

How to cite this article:

Kristiana V., Mukti A. T., Agustono, 2020 Increasing growth performances of Nile tilapia (*Oreochromis niloticus*) by supplementation of noni *Morinda citrifolia* fruit extract via diet. *AAFL Bioflux* 13(1):159-166.