



ISSN: 0975-3575

PHARMACOGNOSY JOURNAL

An Official Journal of Phcog.net

Volume 14 | Number 02 | March - April 2022



Editorial Board (2020-21)

Editors & Editorial Board Members (2021)

Dr.Djemli Samir

Department of Biology , Applied Neuroendocrinology Laboratory
Badji Mokhtar Annaba University
Algeria

Dr. Raghava Naidu, Ph.D

Department of Human Oncology,
University of Wisconsin,
1111, Highland Ave, Madison,
Wisconsin 53705, USA

Dr.Karim Raafat

Associate Professor of Pharmacognosy and Phytochemistry,
Pharmaceutical Sciences Department,
Faculty of Pharmacy,
Beirut Arab University (BAU),
Beirut 115020, Lebanon

Ourlad Alzeus Tantengco, MD-PhD Molecular Medicine

College of Medicine, University of the Philippines Manila
Pedro Gil Street, Ermita, Manila, Philippines, 1000

Janib Achmad

Lecturer of Faculty of Fisheries and Marine Science,
University of Khairun Ternate
Kampus 2 JalanPertamina, KelurahanGambesi,
Ternate Selatan

Muammar Fawwaz, Ph.D

Department of Pharmaceutical Chemistry
Faculty of Pharmacy
Universitas Muslim Indonesia
Makassar 90231, South Sulawesi, Indonesia

Hany Ezzat Khalil

Associate Professor,
College of Clinical Pharmacy,
King Faisal University,
KSA

Emad Yousif

Department of Chemistry
College of Science
Al-Nahrain University
Baghdad,Iraq

Sughosh Upasani

R.C Patel Institute of pharmacy,
Shirpur,Dist-Dhule,Maharashtra,
India.

Gurusiddaiah suresh kumar

Scientist
Dept of biochemistry
CSIR-CFTRI
Mysore, Karnataka, INDIA

Arjun Patra

Assistant Professor
School of Pharmaceutical Sciences
Guru Ghasidas Central University
Koni, Bilaspur - 495 009
Chattisgarh, India

Francis O. Atanu, Ph.D

Department of Biochemistry
Faculty of Natural Sciences
Kogi State University
Anyigba, Nigeria.

Vijay Kumar Chattu

Faculty of Medical Sciences
University of the West Indies
St. Augustine, Trinidad & Tobago.

Dr.Kunle Okaiyeto, PhD

Applied and Environmental Microbiology Research Group (AEMREG)
Department of Biochemistry and Microbiology
University of Fort Hare

Alice campus
5700, Alice
South Africa.

Dr. Srisailam Keshetti, Ph.D

Principal, University College of Pharmaceutical Sciences, Satavahana University
Karimnagar 505001
Telangana
INDIA

Dr. Gayathri M Rao

Associate Professor
Department of Biochemistry
Kasturba Medical College, Mangaluru.

Shuge Tian

Experimental Teaching Demonstration Center of TCM in Xinjiang Medical University
Department of traditional medicine, TCM
Xinjiang Medical University
Xinjiang CHINA 830054

Dr. Ramachandra Setty Siddamsetty,

Professor, Govt College of Pharmacy,
Mission Road, Bengaluru, INDIA

Dr. (Mrs.) Sayyada Khatoon

HOD, Pharmacognosy Division
CSIR-National Botanical Research Institute,
Rana Pratap Marg, Post Box 436,
Lucknow-226001 (U.P.) India

Dr. A. Sajeli Begum

Department of Pharmacy
Birla Institute of Technology & Science
Hyderabad, India

Olga Silva

Department of Pharmacological Sciences,
Faculdade de Farmácia,
Universidade de Lisboa, Portugal

Xinwen Wang

Department of Clinical Pharmacy
University of Michigan
USA

Roman Lysiuk

Department of Pharmacognosy and Botany,
Danylo Halytsky Lviv National Medical University,
Pekarska, 69., Lviv 79010, Ukraine

Arif Nur Muhammad Ansori

Universitas Airlangga
Indonesia

Pharmacogn J. Vol 14, Issue 2, Mar-Apr, 2022

[View](#)[What links here](#)Submitted by [sys1](#) on Sat, 04/30/2022 - 10:59

The Activities of Pegagan Embun (*Hydrocotyle sibthorpioides* Lam.) on TNF- α , Macrophages and Leukocytes Male White Mice Exposed by H5N1 Virus Antigens

GC-MS Profiling, Antioxidants and Antimicrobial Activity of Prickly Pear (*Opuntia ficus-indica*) Pulp Extract

The Potential Effect of Nigericin from *Streptomyces hygroscopicus* subsp. *Hygroscopicus* Against the Syndemic of Malaria and COVID-19 through Molecular Docking Perspective

Anti-Ulcer Activity of Spray-dried Powders Prepared from Aerial Parts Extracts of *Ampelopsis cantoniensis*

Molecular Docking, Physicochemical and Drug-likeness Properties of Isolated Compounds from *Garcinia latissima* Miq. on Elastase Enzyme: In Silico Analysis

Antibacterial and Cytotoxic Activity of Selected Raw-Consumed Vegetables in West Java, Indonesia

Potential of Stem Cells in Overcoming Infertility Problems in Women

Screening of Secondary Metabolites and Antioxidant Activity of Wild Edible Termite Mushroom

Optimization of Ultrasound-Assisted Extraction of *Tinospora crispa* Stem, Phytochemical Screening, Total Phenolic Content and Anti Gout Potential Activity

Pasuchaca (*Geranium ruizii* Hieron.): A Medicinal Plant of the Geraniaceae Family with Hypoglycemic Effect on Alloxan-Induced Hyperglycemia in Mice

The Effect of Antioxidant activity, Total Phenols and Total Flavonoids on Arginase Inhibitory Activity on Plants of Genus *Sterculia*

Evaluation of Antibacterial and Antioxidant Activity of Endophytic Fungi Isolated from *CAPSICUM ANNUUM* L. and *ALLIUM CEPA* L.

Determination Antioxidant Activity of *Coffea Arabica*, *Coffea Canephora*, *Coffea Liberica* and Sunscreens Cream Formulation for Sun Protection Factor (SPF)

An In Silico Study of Examining Bioactive Compounds from *Azadirachta indica* Juss. (Neem) as Potential Death Receptor 5 Inductor in Hepatoma Cells

Preliminary Identification and Quantification of Four Secondary Metabolites, Total Tannin and Total Flavonoid Contents in Guava Fruit Ethanol Extract

Clinical Studies of Silymarin as a Protective Agent Against Liver Damage Caused by Anti-TB Drugs, Methotrexate, and in Cases of Chronic Hepatitis C and Diabetes Mellitus

Formulation and Characterization of Meniran (*Phyllanthus Niruri* Linn) Extract Nanoparticle on Antibacterial Activity Against *Salmonella Pullorum*

Cytotoxic Effects of Kirinyuh Herb (*Austroeupeatorium inulaefolium* (Kunth) R. d. King & H. Robinson) Extracts and Fractions on BSLT, MCF-7 Cells and T-47D Cells

The Potential of Moringa (*Moringa oleifera* Lamk) Seed Oil as Anti- Alopecia

Mechanism of the Bioactive *Sargassum cristaeifolium* in Inhibiting Inflammatory Mediators in a Nitroglycerin-Induced Migraine Model in Rats

Optimization of the Ultrasound Assisted Extraction of *Phaleria macrocarpa* (Scheff.) Boerl. Fruit Peel and its Antioxidant and Anti-Gout Potential

Potency of Antidiabetic Effects of the Combination of *Syzygium cumini* and *Andrographis paniculata* in Rats with High-Fat Diet and Streptozotocin-Induced Diabetes

Preliminary Data on the Antibacterial Activity of *Coffea Arabica*, *Coffea Canephora* and *Coffea Liberica*

Anxiolytic and Antidepressant-like Effects of *Foeniculum vulgare* Essential Oil

Optimization of Ultrasound-Assisted Extraction of *Andrographis paniculata* Nees Leaves, Phytochemical Screening, Total Phenolic Content and Anti-Gout Potential Activity

Coleus atropurpureus (L) Benth. Leaves as a New Promising Drug for Abscesses Caused by Methicillin-resistant *Staphylococcus aureus* and *Staphylococcus aureus*

Ten Commonly Available Medicinal Plants in Malaysia with Potential Sun Protection Factor and Antioxidant Properties – A Review

A Review of an Important Plants: *Annona squamosa* Leaf

Virgin coconut oil and tuberculosis: A mini-review

Inhibitory Capacity of Xanthine Oxidase in Antigout Therapy by Indonesian Medicinal Plants

Virgin coconut oil and tuberculosis: A mini-review

Fathul Djannah^{1,*}, Muhammad Nasrum Massi², Mochammad Hatta², Agussalim Bukhari³, Irda Handayani⁴, Muhammad Faruk⁵, Anny Setijo Rahaju^{6,7}

Fathul Djannah^{1,*}, Muhammad Nasrum Massi², Mochammad Hatta², Agussalim Bukhari³, Irda Handayani⁴, Muhammad Faruk⁵, Anny Setijo Rahaju^{6,7}

¹Department of Anatomical Pathology, Faculty of Medicine, Universitas Mataram, Mataram, West Nusa Tenggara INDONESIA.

²Department of Microbiology, Faculty of Medicine, Universitas Hasanuddin, Makassar South Sulawesi, INDONESIA.

³Department of Clinical Nutrition, Faculty of Medicine, Universitas Hasanuddin, Makassar South Sulawesi, INDONESIA.

⁴Department of Clinical Pathology, Faculty of Medicine, Universitas Hasanuddin, Makassar South Sulawesi, INDONESIA.

⁵Department of Surgeon, Faculty of Medicine, Universitas Hasanuddin, Makassar South Sulawesi, INDONESIA.

⁶Department of Anatomical Pathology, Faculty of Medicine, Universitas Airlangga, Surabaya, INDONESIA.

⁷Universitas Airlangga Academic Hospital, Dr. Soetomo General Academic Hospital, Surabaya, INDONESIA.

Correspondence

Fathul Djannah

Department of Anatomical Pathology, Faculty of Medicine, Universitas Mataram, Mataram, West Nusa Tenggara INDONESIA.

Email: fdjannah354@gmail.com

History

- Submission Date: 01-02-2022;
- Review completed: 11-02-2022;
- Accepted Date: 19-02-2022.

DOI : 10.5530/pj.2022.14.59

Article Available online

<http://www.phcogj.com/v14/i2>

Copyright

© 2022 Phcogj.Com. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International license.

ABSTRACT

Virgin coconut oil is widely promoted and used as healthy and beneficial oil. One of them is caused by antimicrobials. Caprylic, caproic acid, capric acid, lauric acid and tau glyceryl monolaurate are other VCO compositions. Furthermore, due to the non-heating manufacturing process, the content in VCO can reduce cholesterol levels of triglycerides, LDL, phospholipids, VLDL and increase HDL in blood serum. VCO consumption lowers the number of *Mycobacterium tuberculosis* colonies while increasing the conversion of BTA sputum. Until now, the prevalence of tuberculosis (TB) disease was extremely high. VCO can be used as a supplement to help TB patients recover faster.

Key words: Virgin coconut oil, *Mycobacterium tuberculosis*, Tuberculosis.

INTRODUCTION

Virgin coconut oil is widely promoted and used as healthy and beneficial oil. Some of the benefits of using coconut oil support overcoming various health problems.¹ Coconut oil is a vegetable oil staple for the food industry. Another term for coconut oil is tropical oil or lauric oil.² However, there are differences in the coconut oil used for cooking (coconut cooking oil) with VCO. The distinction is found in the extraction process. RBD or coconut oil is typically extracted using dried, ground, and chemically mixed parts of the coconut skin.

Meanwhile, VCO is produced from fresh, pure coconut milk, which undergoes a fermentation process by adding microbes such as lactobacillus fermentum and/or lactobacillus plantarum, separating coconut milk from oil (churning), and then adding an enzyme to separate oil and water. VCO is called virgin oil because no heating process is known as cold extraction.³ RBD or ordinary oil will produce a large amount of saturated fatty acids due to this process, whereas VCO produces no precipitate.⁴ In addition, without the heating process, the content in VCO can reduce cholesterol levels of triglycerides, LDL, phospholipids, VLDL and increase HDL in blood serum compared to coconut oil. Furthermore, VCO contains antioxidant enzymes and has antithrombotic effect.⁵ According to other research, to obtain the antimicrobial effect of coconut, triglycerides must first be broken down into monoglycerides also known as free fatty acids. Lipase-released free fatty acids, on the other hand, contain only weak antimicrobials.⁶

Some of the compositions of VCO have a positive impact on health. VCO can be antifungal, antibacterial, antioxidant, antiviral, prevent cardiovascular and liver disease, increase the immune system, anti-diabetic, and treat metabolic syndrome.⁷ Several scientific studies have shown that VCO can be used as an antibacterial agent to treat tuberculosis. Tuberculosis (TB) is a fatal airborne disease that frequently becomes

a significant problem in low and middle-income countries' health, economic, and social sectors.⁸ World case reports show around 10-20 million TB cases. 90% of cases are found in adults and 9% in people living with HIV. Ethiopians occupy the highest burden of TB disease globally, and Africa ranks third.⁹ The average TB incidence rate is 164 per 100,000 populations. 23% of them are latent TB.¹⁰

Tuberculosis is caused by a bacterium called *Mycobacterium tuberculosis*. Several species of these bacteria include MTBC (*mycobacterium tuberculosis* complex), which consists of *Mycobacterium tuberculosis* (Mtb), a common cause of TB in humans, and *M. africanum*, a type of TB, especially in the African region. Meanwhile, *M. caprae*, *M. bovis* and *M. pinnipedi* I are TB in mammals and *M. Microti* specific to mice.¹¹ The high number of sufferers and TB transmission rates and some literature show an increase in sputum conversion and nutritional status of TB patients after consuming VCO,¹² or other benefits. It is necessary to discuss the benefits of VCO for people with tuberculosis comprehensively.

Search strategies

The search for research literature was carried out through database journals such as Pubmed, ScienceDirect, Willey Online Library, and Google Scholar by using a combination of keywords MeSH (Medical Subject Headings) of "virgin", "coconut oil", "oil", tuberculosis, and infectious disease. Other additional references are also searched manually.

Virgin coconut oil (VCO)

Coconut oil is a necessary ingredient in food processing as well as the most important and primary dietary fat. In the food processing industry, the presence of coconut oil is vital. There are two types of coconut oil: RBD coconut oil or copra oil and VCO.¹³ RBD or copra is oil that is refined or purified using various chemical compounds (refined) and then purified (bleaching), and finally removing the unpleasant odour that appears (deodorised).¹⁴ Making RBD is to first dry the coconut either by using smoke, the sun, or both. Then after cleaning,

Cite this article: Djannah F, Massi MN, Hatta M, Bukhari A, Handayani I, Faruk M, et al. Virgin coconut oil and tuberculosis: A mini-review. Pharmacogn J. 2022;14(2): 464-469.

the grated coconut is pressed to get coconut oil. However, the three processes must still be carried out in the extraction process to get healthy coconut oil, such as refined, bleaching and deodorised.¹⁵

The other type is VCO. Unlike RBD, which is extracted using a dry process, VCO is extracted using a wet technique under carefully controlled temperature conditions, allowing better components of coconut oil to be preserved. Small components, such as vitamins A and E and polyphenols, can be preserved by avoiding UV radiation.

VCO composition

Currently, VCO has been widely used as a food supplement and moisturizer. Other VCO compositions are caprylic, caproic acid, capric acid and lauric acid.¹³ Based on other studies, it shows that the biological properties of VCO are much better than coconut oil. VCO contains chain fatty acids containing 45-50% lauric acid, a powerful antimicrobial and anti-inflammatory agent. With integrated or wet, the physicochemical structure showed the free fatty acid content (FFA) in the range of 0.13% ± 0.06%. The phenol content ranges from 16.02 ± 0.44 mg GAE in every 100 g VCO.¹⁶

VCO extraction

VCO is the result of coconut extraction, which has high economic value. In order to get the best VCO composition, there are several extraction techniques used, namely the fresh-wet technique (fermentation then centrifugation) and the fresh-dry technique (fresh-dry). These two techniques can maintain the antioxidant components in coconut oil, such as polyphenols, tocotrienols and tocopherols.¹⁷

VCO extracted by fermentation technique showed that the addition of yeast greatly affected the volume of VCO produced in water content, free fatty acid composition, and viscosity. The types of yeast used are *Saccharomyces cerevisiae*, baker's yeast, *Rhizopus* sp or tempeh yeast, and yeast tape.¹⁸ Meanwhile, other literature shows four ways to get good quality VCO: centrifugation, cooling, direct micro expelling-oven dried that does not use high temperatures, and direct micro expelling sun-dried.¹⁹ However, from other sources, it is described that the physicochemical properties of coconut oil's fatty acids with the best quality are obtained from processing methods by centrifugation or suspended solid-liquid separation.²⁰

Benefits of VCO

Fresh VCO contains a very beneficial component in lowering blood pressure, helping to lose excess weight, and is a biomarker in the inflammatory process. On the other hand, VCO that has been heated repeatedly, such as palm oil, can increase blood pressure. Therefore, VCO should be consumed fresh to obtain health benefits.¹⁷

In addition, other studies have shown that VCO treated with a new wet system has a good role as an endogenous antioxidant and paraoxonase one because it can prevent the oxidation of lipids and proteins.¹⁷ Another benefit is that VCO can be hepatoprotective with dry or fermentation techniques. The experimental results using paracetamol-induced liver damage showed that VCO significantly reduced liver damage or was hepatoprotective.²² Furthermore, consuming 30 mL of VCO daily can raise high-density lipoprotein cholesterol levels for healthy young adults. No problems or complications were discovered after eight weeks of daily VCO consumption.²³ Other research indicates that, while still in early stages, VCO's use effectively reduce the incidence of atopic dermatitis, and prevents dental caries and hair damage caused by UV exposure and treatment that results in protein loss.²⁴ VCO, a diet rich in unsaturated fatty acids, can reduce the incidence of cardiovascular disease and have local effects (CVD). However, some trials are still limited to animal trials, so further studies are needed in humans.²⁵ Others show that coconut oil contains a better concentration of palm

oil. Coconut oil does not affect body weight, body fat percentage, waist circumference, and plasma glucose. However, vegetable oil (coconut oil) cannot be said to reduce the risk of CHD.²⁶

The effect of VCO on TB patients


TB disease has been an epidemic for hundreds of years, with 5000 TB deaths per day. MTB, caused by *Mycobacterium tuberculosis*, is a parasitic disease in humans that can adapt and survive by transmitting it between humans. Therefore, a host maintaining systemic immunity is needed to control infection transmission.²⁷ Other literature states that TB is a disease that can induce the body's immunity, either as a protective immunity or by damaging tissues. Primary TB plays a role in mediating protective immunity to spread infection.

In contrast to primary TB, post-primary TB plays a role in forming tissue damage that results in the formation of cavities. Both components are indispensable for mycobacterium tuberculosis or MTB bacteria.²⁸ The process of MTB in multiplying is by making MTB-infected macrophages isolated into hypoxic tissues and then differentiation containing lipids containing triacylglycerol.²⁹ Other literature states that the MTB replication process is carried out by converting fatty acids into acyl-adenylate so that the inhibitor will be disrupted by the presence of various pathways.³⁰ The MTB structure plays a vital role in determining the survival of MTB bacteria, namely the plasma membrane, cell wall, mycomembranes and capsules.³¹

Based on the study results, giving VCO with different concentrations reduced the number of *Mycobacterium tuberculosis* colonies. When given VCO with different concentrations, 1:10 with a given concentration of 1:10,000 was able to reduce colonies from 46% to 100%. This is because different VCO preparations contain varying amounts of lauric acid.³² In addition, giving VCO combined with the administration of catfish albumin impacts the cure rate and shortens the length of treatment due to conversion of the sputum smear and interferon-gamma composition in the third month.³³ The positive impact of the use of VCO can be caused by the content of saturated fatty acid chains or glycerol monolaurate or also called monolaurin, which has chemical and biological effects or has strong antimicrobials in killing pulmonary tuberculosis bacteria by penetrating membranes and synthesizing, making efforts to inhibit the reproduction and replication of tuberculosis germs.³⁴

Other studies also show the results or benefits of VCO on TB. Pulmonary TB patients who were given MDT DOTS then combined with VCO tablets could accelerate the time needed to convert the sputum components of AFB. Accelerated conversion is caused by the content of long-chain saturated fat or also called glyceryl monolaurate, which functions as a potent bactericide for pulmonary TB, thereby inhibiting the replication and reproduction of mycobacterium TB.³⁵ In addition, from the results of the analysis of the VCO content, of the 84% composition of saturated fatty acids, 46.67% of them are lauric acid. Furthermore, when absorbed into the body, lauric acid will be converted or changed to glycerol monolaurate or monolaurin. In the study of detecting the resistance level to OAT or mycobacterial tuberculosis using the REMA (resazurin microtiter assay plate) technique, the results showed that VCO could be effective against mycobacterium tuberculosis. Monolaurin counteracts MTB at 313 mg/mL while rifampin at 0.0025 mg/mL.³⁶ While related to the dose of VCO, which can provide a good therapeutic effect, VCO should be given at a low dose first in order to improve hepatic structural, biochemical deviation or modify diet. The therapeutic effect of VCO will be more significant on LDH, SOD, MDH and total bilirubin.³⁷

The therapeutic effect of VCO begins with lipolysis in the mouth, continues to the stomach, and ends in the small intestine. Monolaurin and monocaprin are the two forms of VCO metabolites.³⁸ VCO fights

**Table 1: Summary of the study.**

Study	Outcome	Result
Dalmacion et al. ³²	The effect of VCO in reducing Mycobacterium tuberculosis	2 brands of VCO can reduce the number of mycobacterium tuberculosis colonies, 46-100% on Middle brook 7H10 media. There was a difference in the level of gamma interference in the intervention group given VCO and albumin and the control group at 3 months (mean SE 38.53-13.64). Meanwhile, in the 1st and 6th months, no difference was found. At 6 months there was no difference because the body no longer produced IFN-X cytokines after 6 months of treatment.
Montolalu et al. ³³	Effects of albumin and VCO supplementation with sputum and interferon conversion	The combination of VCO nutrition with albumin extracted from catfish showed significant results in increasing body immunity, accelerating the conversion of BTA VCO sputum with a P value <0.00 while the conversion of sputum with catfish extract showed a P value of 0.004. As for the combination of the two, VCO and albumin with P value <0.00.
Arifin et al. ³⁵	Accelerated conversion of BTA sputum	The results of clinical trials using the REMA (Resazurin microtiter assay) method, showed that VCO was active against MTB H 37 Rv at 625ug/ml and against clinical isolates from 78 to 625. While Monolaurin was active at 313 ug/mL against MTB H 37 Rv, and Rifampicin, at 0.0025 g/mL. These data indicate the high potential of VCO as an antimycobacterial tuberculosis.
Delia. ³⁶	Clinical trials of VCO as an antimycobacterial	The results of this study showed that the sputum smear conversion rate that received VCO and DOTS interventions had a mean = 4 weeks, 2 weeks faster than those who were only given DOTS with a mean = 6 weeks.
Kasman. ¹²	Increased conversion of BTA sputum and nutritional status of pulmonary TB patients with VCO	

microbial growth through three basic mechanisms: 1) disintegration of the lipid membrane, 2) inhibition of pathogenic organism maturation and 3) Preventing pathogens from entering the host cell membrane. Furthermore, the lipid breakdown process is the foundation for monolaurin activity.³⁹

From the table 1, we know that VCO can be an excellent supplement recommendation for TB sufferers. If described further, the mechanism or effect of VCO in overcoming TB is shown in Figure 1.

Based on the mechanism (Figure 1), it can be seen that the main antimicrobial content in VCO is glyceryl monolaurate/lauric acid/saturated fatty acids. Glyceryl monolaurate is a glycerol ester of lauric acid, often used in the food industry as a cosmetic, preservative, or emulsifier, and has been declared safe by the food and drug administration for topical use at a dose of 100 mg/ml.⁴⁰ However, glyceryl monolaurate is also known as a very potent antimicrobial against gram-positive bacteria. GML is also known as a topical microbicide used in a broad spectrum. Despite GML's superiority as an antimicrobial, other research indicates that its use is declining. This is because its water-soluble properties are still low, and its melting point is too high. However, the most recent nanoparticle formulation can effectively reduce the toxic effect against BTA germs.⁴¹

Apart from being an antimicrobial pathogen, Glyceryl monolaurate or monolaurin is an antiviral that can fight various viruses and improve the body's immune system. The mechanism of the role of glyceryl guacholate related to its function as an antiviral includes controlling the pro-inflammatory cytokine system, activating leukocytes, disintegrating membranes, and inhibiting the process of maturation and replication of the virus.⁴²

The antibacterial component of VCO that can increase sputum conversion is lauric acid. Lauric acid is a bioactive compound that can be an inhibitory factor for Mycobacterium tuberculosis. The lauric acid inhibitory concentration level was 22.2 -66.7 g/mL.⁴³ The way lauric acid works in inhibiting the performance of bacteria is by forming reactive oxygen species (ROS) and causing damage to the membrane, and inhibiting the formation of biofilms, thereby inhibiting pro-inflammatory cytokines.⁴⁴

As for getting the maximum antibacterial or antiviral effect from VCO, the thing that must be considered is the hydrolysis process of VCO. The higher the amount of NaOH used in the hydrolysis process, the antibacterial activity can increase. In addition, VCO, which is hydrolyzed with enzymes, is more effective than that which undergoes hydrolysis with sodium hydroxide [46].⁴⁵ Based on an antibacterial activity test of hydrolyzed VCO also known as VCOH, the hydraulic VCO process outperformed VCO without hydrolysis.⁴⁶ Furthermore, the length of fermentation and whether or not induction oil is used impact VCO quality.⁴⁷

A positive impact of giving VCO is an increase in sputum conversion BTA. Sputum conversion is the strongest predictor in determining the effectiveness of TB therapy. Mycobacteria characterize sputum conversion in the sputum culture of TB cases formed in the second month of the treatment process. The first month's conversion ranged from 60-80%, while in the second month, it reached 95%. Some of the supporting factors in increasing sputum conversion are the level of compliance in consuming drugs, smoking habits, nutritional status (BMI), and habits in consuming alcohol.⁴⁸

According to the previous reference, VCO, due to its glyceryl monolaurate content, can increase sputum conversion by inhibiting the replication and reproduction of mycobacterium bacteria. Another study found that glyceryl monolaurate inhibited the growth of Streptococcus aureus and Streptococcus epidermidis bacteria by inhibiting the production of lipase enzymes. Lipase deficiency can inhibit bacterial growth.⁴⁹ VCO

also has antibacterial properties and can be used as a skin protection material to protect against UV rays and radiation exposure. The test results of VCO creams show that they increase spreadability but have no effect on pH or adhesion.⁵⁰

Future direction

From the literature review described previously, we get an overview of the antibacterial benefits of VCO in increasing the conversion of BTA sputum. However, further scientific studies are needed to determine the best type of VCO, dosage and the effect of VCO consumption time.

CONCLUSION


TB is an infectious disease that is still becoming an epidemic with a high prevalence every year. Apart from medical drugs, nutrition or complementary supplements are needed to reduce the prevalence of events and accelerate and make the TB treatment process more effective. According to the literature, virgin coconut oil is one companion supplement that can increase the conversion of BTA sputum. Antimicrobials containing glyceryl monolaurate/saturated fatty acid/lauric acid are found in VCO. The use of VCO as an additional supplement for TB therapy is recommended.

REFERENCES

- Hartono, Suryawati B, Sari Y, Avicena A, Maryani, Sukmagautama C, et al. The effect of curcumin and virgin coconut oil towards cytokines levels in COVID-19 patients at Universitas Sebelas Maret Hospital, Surakarta, Indonesia. *Phcog J*. 2022;14(1).
- Eyres L, Eyres MF, Chisholm A, Brown RC. Coconut oil consumption and cardiovascular risk factors in humans. *Nutr Rev*. 2016;74(4):267-280.
- Tham YJ, Ng PE, Khoo KS, Cheng CK, Chew KW, Show PL. A comprehensive review on the techniques for coconut oil extraction and its application. *Bioprocess Biosyst Eng*. 2021;44(9):1807-1818.
- Suryani S. A Comparative Study of Virgin Coconut Oil, Coconut Oil and Palm Oil in Terms of Their Active Ingredients. *Processes*. 2020;8(4):402.
- Nevin KG, Rajamohan T. Influence of virgin coconut oil on blood coagulation factors, lipid levels and LDL oxidation in cholesterol fed Sprague–Dawley rats. *E-SPEN Eur E-J Clin Nutr Metab*. 2008;3(1):e1-e8.
- Sia CM, Yim HS, Lai CM. Commercial virgin coconut oil: assessment of antimicrobial potential. 2010;13.
- Deen A. Chemical composition and health benefits of coconut oil: an overview. *J Sci Food Agric*. 2021;101(6):2182-2193.
- Delogu G, Sali M, Fadda G. The Biology of *Mycobacterium Tuberculosis* Infection. *Mediterr J Hematol Infect Dis*. 2013;5(1):e2013070.
- FMOH. Guidelines for Clinical and Programatic Management TB, TB/HIV and Leprosy in Ethiopia, 6th edition. Ethiopia: Federal Ministry of Health Addis Ababa. 2016.
- World Health Organization and International Telecommunication Union, Be he@lthy, be mobile: a handbook on how to implement mTB-Tobacco. Geneva: World Health Organization. 2019.
- Brosch R. A new evolutionary scenario for the *Mycobacterium tuberculosis* complex. *Proc Natl Acad Sci*. 2002;99(6):3684-3689.
- Kasman. the influence of virgin coconut oil (VCO) intervention to increase conversion and nutrient status of patients with pulmonary TB in public lung health cancer of makassar city in 2013 / kasman. Universitas Hasanuddin: Fakultas Kesehatan Masyarakat. 2013.
- Holt B. Vegetable oil: properties, uses and benefits. New York Nova Science Publishers Inc. 2016.
- Adkins SW, Foale MA, Samosir YMS. Australian Centre for International Agricultural Research, and International Coconut Forum, Coconut revival: new possibilities for the 'tree of life': proceedings of the International Coconut Forum held in Cairns, Australia, 22-24 November 2005. Canberra. Australian Centre for International Agricultural Research. 2006.
- Fife B. Virgin coconut oil: nature's miracle medicine. Colorado Springs, CO: Piccadilly Books. 2006.
- Ahmad Z, Hasham R, Nor NFA, Sarmidi MR. Physico-Chemical and Antioxidant Analysis of Virgin Coconut Oil Using West African Tall Variety. 2015;13:11.
- Hamsi MA. Effect of consumption of fresh and heated virgin coconut oil on the blood pressure and inflammatory biomarkers: An experimental study in Sprague Dawley rats," *Alex J Med*. 2015;51(1):53-63.
- Agustine D, Gumilang MM, Komalasari N. The Effect of Yeast Starter Variations on the Quality of Virgin Coconut Oil (VCO) Using the Fermentation Method. 2021;6.
- Ghani NAA, Channip AA, Hwa CHP, Jafar F, Yasin HM, Usman A. Physicochemical properties, antioxidant capacities, and metal contents of virgin coconut oil produced by wet and dry processes. *Food Sci Nutr*. 2018;6(5):1298-1306.
- Ndife J, Obot D, Abasiokong K. Quality Evaluation of Coconut (*Cocos nucifera* L) Oils Produced by Different Extraction Methods. *Asian Food Sci J*. 2019;1-10.
- Arunima S, Rajamohan T. Effect of virgin coconut oil enriched diet on the antioxidant status and paraoxonase 1 activity in ameliorating the oxidative stress in rats – a comparative study. *Food Funct*. 2013;4(9):1402.
- Zakaria ZA. Hepatoprotective Activity of Dried- and Fermented-Processed Virgin Coconut Oil," *Evid. Based Complement. Alternat Med*. 2011;1-8.
- Chinwong S, Chinwong D, Mangklabruks A. Daily Consumption of Virgin Coconut Oil Increases High-Density Lipoprotein Cholesterol Levels in Healthy Volunteers: A Randomized Crossover Trial. *Evid Based Complement. Alternat Med*. 2017;1-8.
- Wallace TC. Health Effects of Coconut Oil-A Narrative Review of Current Evidence. *J Am Coll Nutr*. 2019;38(2):97-107.
- Ma ZF, Lee YY. Virgin Coconut Oil and its Cardiovascular Health Benefits. *Nat Prod Commun*. 2016;11(8):1934578X1601100.
- Nithya N, Seah JYH, Van Dam RM. The Effect of Coconut Oil Consumption on Cardiovascular Risk Factors: A Systematic Review and Meta-Analysis of Clinical Trials. *Circulations*. 2020;141(10).
- Lee RS. Population genomics of *Mycobacterium tuberculosis* in the Inuit. *Proc Natl Acad Sci*. 2015;112(44):13609-13614.
- Hunter RL. The Pathogenesis of Tuberculosis: The Early Infiltrate of Post-primary (Adult Pulmonary) Tuberculosis: A Distinct Disease Entity. *Front Immunol*. 2018;9.
- Daniel J, Maamar H, Deb C, Sirakova TD, Kolattukudy PE. *Mycobacterium tuberculosis* Uses Host Triacylglycerol to Accumulate Lipid Droplets and Acquires a Dormancy-Like Phenotype in Lipid-Loaded Macrophages. *PLoS Pathog*. 2011;7(6):e1002093.
- Arora P. Mechanistic and functional insights into fatty acid activation in *Mycobacterium tuberculosis*. *Nat Chem Biol*. 2009;5(3):166-173.
- Melly G, Purdy G. MmpL Proteins in Physiology and Pathogenesis of *M. tuberculosis*. *Microorganisms*. 2019;7(3):70.
- Dalmacion GV, Ortega AR, Pena IG, Ang CF. Preliminary study on the *in-vitro* Susceptibility of *Mycobacterium tuberculosis* Isolates to Virgin Coconut Oil. *Funct Foods Health Dis*. 2012;2(8):290.
- Montolalu FC, Hadju V, Lawrence GS, Wahyuni S, Maryunis M. The Effectiveness of Extract Snack Head Fish and Virgin Coconut Oil

- (VCO) on Sputum Conversion and Levels of Interferon- γ in Patients with Pulmonary TB Multi Drug Resistant (MDR) in Labuang Baji Hospital Makassar. *Int J Sci.* 2016;30(4):12.
34. Fife B, Fife B. The coconut oil miracle. 2021.
 35. Arifin M, Hadju V, Astuti N, Wahyuni S. Supplements Effect of Virgin Coconut Oil and Albumin Capsules (Catfish protein) on TB Patients Receiving Multi Drugs Therapy-DOTS Strategic in BBKPM Makassar, Indonesia. 2014;4(7):6.
 36. Delia GBD, Ontengco DC. Anti-Tuberculosis Activity of Virgin Coconut Oil Using the Resazurin Microtiter Assay. 2015;7.
 37. Adeyemi WJ. Investigation of the effects of dietary modification in experimental obesity: low dose of virgin coconut oil has a potent therapeutic value. *Biomed Pharmacother.* 2020;126:110110.
 38. Kp B, Av P. *Journal of Nutritional Biology.* 2021;14.
 39. Lafia, Nigeria, Abel AA. Antimicrobial Activity of Coconut Oil and its Derivative (Lauric Acid) on Some Selected Clinical Isolates. *Int J Med Sci Clin Invent.* 2017;4(8).
 40. Lin YC. Glycerol Monolaurate and Dodecylglycerol Effects on *Staphylococcus aureus* and Toxic Shock Syndrome Toxin-1 *in vitro* and *in vivo*. *PLoS ONE.* 2009;4(10):e7499.
 41. Lopes LQS, Santos GC, de Almeida R, Vaucher, Gende L, Raffin RP, *et al.* Evaluation of antimicrobial activity of glycerol monolaurate nanocapsules against American foulbrood disease agent and toxicity on bees. *Microb Pathog.* 2016;97:183-188.
 42. Subroto E, Indiaro R. Bioactive monolaurin as an antimicrobial and its potential to improve the immune system and against COVID-19: a review. *Food Res.* 2020;4(6):2355-2365.
 43. Muniyan R, Gurunathan J. Lauric acid and myristic acid from *Allium sativum* inhibit the growth of *Mycobacterium tuberculosis* H37Ra: *in silico* analysis reveals possible binding to protein kinase B. *Pharm Biol.* 2016;54(12):28142821.
 44. Yang HT. Lauric Acid Is an Inhibitor of *Clostridium difficile* Growth *in vitro* and Reduces Inflammation in a Mouse Infection Model. *Front Microbiol.* 2018;8:2635.
 45. Silalahi J, Permata YM, Putra EDL. Antibacterial Activity of Hydrolyzed Virgin Coconut Oil. 2014;7:5.
 46. Loung FS, Silalahi J, Suryanto D. Antibacterial activity of Enzymatic hydrolyzed of Virgin Coconut oil and Palm Kernel oil against *Staphylococcus aureus*, *Salmonella thypi* and *Escherichia coli*. *Intern J PharTech Research.* 2014;6(2):628-633.
 47. Anwar C, Irmayanti I, Ha U. Characteristics of Physical, Chemical, and Organoleptic Properties of Virgin Coconut Oil (VCO) By Studying the Ratio Between Coconut Cream with Inducement Oil and Length of Fermentation. *Acta Univ Agric Silv Mendel Brun.* 2020;68(3):473-482.
 48. Sari DK, Mega JY, Harahap J. Nutrition Status Related to Clinical Improvement in AFB-Positive Pulmonary Tuberculosis Patients in Primary Health Centres in Medan, Indonesia. *Open Access Maced J Med Sci.* 2019;7(10):1621-1627.
 49. Flanagan JL. Glycerol Monolaurate Inhibits Lipase Production by Clinical Ocular Isolates Without Affecting Bacterial Cell Viability. *Investig Ophthalmology Vis Sci.* 2016;57(2):544.
 50. Marpaung MP, Prasetyo D. Effect of virgin coconut oil (VCO) on the physical stability of sweet potato leaf extract (*Ipomoea batatas* (L.) Lam) creams and antibacterial activity test against *Staphylococcus aureus* ATCC 25923. *Pharmaciana.* 2021;11(1):109.

GRAPHICAL ABSTRACT



ABOUT AUTHORS



Fathul Djannah has completed a Bachelor of Medicine, from Faculty of Medicine Universitas Hang Tuah Surabaya and Anatomical Pathology Spesialistic degree, Faculty of Medicine Airlangga Indonesia. Currently, as a lecturer in Anatomical Pathology Department, Faculty of Medicine, Universitas Mataram Indonesia and student of doctoral Program from Faculty of Medicine, Universitas Hasanuddin Makassar Indonesia. Her research interests are related to tuberculosis, oncology, carcinoma mammae and molecular biology.



Prof. Mochammad Hatta, MD, PhD, Clinical Microbiologist (Cons) is professor of Molecular Biology and Immunology for Infectious Diseases in Medical Faculty, Hasanuddin University, Makassar, Indonesia since 1985. His MD degree from Medical Faculty, Hasanuddin University Makassar and completed PhD in Faculty of Medicine, Toyama Medical and Pharmaceutical Univ, Japan. He is holding a Chairman and Consultant as chairman of Post Graduate School, Medical Faculty, Hasanuddin University, Indonesia.



Irda Handayani has completed a MD, Clinical Pathology Spesialistic degree and Doctoral Program from Medical Faculty, Hasanuddin University Makassar Indonesia. Currently, She is a lecturer in Clinical Pathology Department, Medical Faculty, Hasanuddin University, Makassar, Indonesia. Her research interests are related to Molecular Biology and Immunology for Infectious Diseases.



Muh. Nasrum Massi has completed a MD, Clinical Microbiology Spesialistic degree from Medical Faculty, Hasanuddin University Makassar Indonesia. and Doctoral Program Spesialistic degree from Universitas Airlangga Indonesia. He completed PhD program in Kobe University Japan. Currently, He is a lecturer in Microbiology Department, Medical Faculty, Hasanuddin University, Makassar, Indonesia. His research interests are related to Molecular Biology and Immunology for Infectious Diseases.



Agussalim Bukhari has completed a Bachelor of Medicine from Medical Faculty Universitas Hasanuddin. His Master of Medicine degree from Monash University, Spesialistic degree from Medical Faculty, Universitas Indonesia and completed PhD in Faculty of Medicine, Toyama Medical and Pharmaceutical University, Japan.



Muhammad Faruk has completed a MD from Medical Faculty Universitas Wijaya Kusuma Indonesia and Surgeon Spesialistic degree from Medical Faculty, Hasanuddin University Makassar Indonesia. Currently, He is a lecturer in Medical Faculty, Universitas Al-Azhar Mataram Indonesia. His research interests are related to oncology, Vascular disease and molecular biology.




Anny Setijo Rahaju has completed a Bachelor of Medicine and Anatomical Pathology Specialist degree and Doctoral Program from Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia. Currently, She is a lecturer in Anatomical Pathology Department, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia. Her research interests are related to oncology, gastrointestinal, uropoetical pathology and molecular biology.

Cite this article: Djannah F, Massi MN, Hatta M, Bukhari A, Handayani I, Faruk M, et al. Virgin coconut oil and tuberculosis: A mini-review. *Pharmacogn J.* 2022;14(2): 464-469.

Pharmacognosy Journal

<p>COUNTRY</p> <p>India</p> <p>Universities and research institutions in India</p>	<p>SUBJECT AREA AND CATEGORY</p> <p>Pharmacology, Toxicology and Pharmaceutics</p> <ul style="list-style-type: none"> Drug Discovery Pharmacology 	<p>PUBLISHER</p> <p>EManuscript Technologies</p>	<p>H-INDEX</p> <h1>25</h1>
<p>PUBLICATION TYPE</p> <p>Journals</p>	<p>ISSN</p> <p>09753575</p>	<p>COVERAGE</p> <p>2009-2021</p>	<p>INFORMATION</p> <p>Homepage</p> <p>How to publish in this journal</p> <p>editor@phcogj.com</p>



SCImago Graphica

Explore, visually communicate and make sense of data with our **new data visualization tool**.

← Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scimagoj.com" data-bbox="412 888 494 904">
```

powered by scimagojr.com

Source details

Pharmacognosy Journal

Scopus coverage years: from 2009 to Present

Publisher: Pharmacognosy Network Worldwide

ISSN: 0975-3575

Subject area: Pharmacology, Toxicology and Pharmaceutics: Pharmacology

Pharmacology, Toxicology and Pharmaceutics: Drug Discovery

Source type: Journal

CiteScore 2021

1.9



SJR 2021

0.258




SNIP 2021

0.718



[View all documents >](#)

[Set document alert](#)

 [Save to source list](#)

[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)

i Improved CiteScore methodology

CiteScore 2021 counts the citations received in 2018-2021 to articles, reviews, conference papers, book chapters and data papers published in 2018-2021, and divides this by the number of publications published in 2018-2021. [Learn more >](#)

CiteScore 2021 

$$1.9 = \frac{1,760 \text{ Citations 2018 - 2021}}{946 \text{ Documents 2018 - 2021}}$$

Calculated on 05 May, 2022

CiteScoreTracker 2022 

$$1.6 = \frac{1,239 \text{ Citations to date}}{798 \text{ Documents to date}}$$

Last updated on 05 September, 2022 • Updated monthly

CiteScore rank 2021

Category	Rank	Percentile
Pharmacology, Toxicology and Pharmaceutics	#219/303	27th
Pharmacology		
Pharmacology, Toxicology and Pharmaceutics	#116/154	25th
Drug Discovery		

[View CiteScore methodology >](#) [CiteScore FAQ >](#) [Add CiteScore to your site !\[\]\(8ca028f9e2a49f208b9be4b48bf3bf2f_img.jpg\)](#)



UNIVERSITAS AIRLANGGA
FAKULTAS KEDOKTERAN

Kampus A Jl. Mayjen Prof. Dr. Moestopo 47 Surabaya 60131
Telp. (031) 5020251, 5030252-3 Fax. (031) 5022472
Laman : <http://www.fk.unair.ac.id> e-mail : dekan@fk.unair.ac.id

SURAT KETERANGAN
Nomor : 4499/UN3.1.1/TU/2023

Yang bertanda tangan di bawah ini :

Nama : Dr. Sulistiawati, dr., M.Kes
NIP : 196502281990032002
Pangkat/Golongan : Lektor Kepala/ IVA
Jabatan : Wakil Dekan III

Dengan ini menerangkan bahwa :

Nama : Dr. Anny Setijo Rahaju, dr., SpPA(K)
NIP : 197009202008122001
Pangkat/ Gol : Penata/ III-c
Jabata : Lektor


Telah melaksanakan publikasi sebagai berikut :

No	Judul Karya Ilmiah	Tahun Pelaksanaan Penelitian	Alasan
1	Tinjauan Literatur: Patogenesis Infeksi Human Papillomavirus Sebagai Dasar Dalam Pengembangan Pencegahan Dan Terapi Baru Pada Kanker Penis	2021	Penelitian merupakan tinjauan literatur
2	Virgin coconut oil and tuberculosis: A mini-review	2022	Penelitian merupakan mini review

Demikian surat keterangan ini kami buat untuk dapat dipergunakan sebagai persyaratan pengusulan kenaikan pangkat ke Lektor Kepala.

Surabaya, 17 April 2023

Wakil Dekan III,




Dr. Sulistiawati, dr., M.Kes.
NIP. 196502281990032002