

# Journal of Public Health in Africa

ISSN 1073-7364

Volume 52 Number 1

January 2013

Pages 1-100

Pages 101-200

Pages 201-300

Pages 301-400

Pages 401-500

Pages 501-600

Pages 601-700

Pages 701-800

Pages 801-900

Pages 901-1000



[HOME](#) / [ARCHIVES](#) /

Vol 14, No s1 (2023): 4th Joint Conference of UNAIR-USM, International Conference of Pharmacy and Health Sciences 2022 (ICPHS) | 11 October 2022

# Vol 14, No s1 (2023): 4th Joint Conference of UNAIR-USM, International Conference of Pharmacy and Health Sciences 2022 (ICPHS) | 11 October 2022

**Published:** 30-03-2023



## ORIGINAL ARTICLES



### *Eurycoma longifolia*: an overview on the pharmacological properties for the treatment of common cancer

Shankar Jothi, Thaigarajan Parumasivam, Noratiqah Mohtar

<https://doi.org/10.4081/jphia.2023.2495>

0 0 0 0

75 PDF: 81

PDF



### Challenges and strategies for collagen delivery for tissue regeneration

Lia Agustina, Andang Miatmoko, Dewi Melani Hariyadi

<https://doi.org/10.4081/jphia.2023.2505>

0 0 0 0

60 PDF: 78

PDF



### Allergic rhinitis behavioral changes after Indonesian house dust mites allergenic extract administration as immunotherapy

Yusuf Alif Pratama, Honey Dzikri Marhaeny, Lutfiatur Rohmah, Salsabilla Madudari Kasatu, Ahmad Dzulfikri

Nurhan, Mahardian Rahmadi, Junaidi Khotib

 <https://doi.org/10.4081/jphia.2023.2510> 0  0  0  0 71  PDF: 66 PDF

### ***In vitro* study of pinostrobin propionate and pinostrobin butyrate: cytotoxic activity against breast cancer cell T47D and its selectivity index**

Tri Widiandani, Tiffany Tandian, Bagus Dwi Zufar, Andi Suryadi, Bambang Tri Purwanto, Suko Hardjono, Siswandono Siswandono

 <https://doi.org/10.4081/jphia.2023.2516> 0  0  0  0 80  PDF: 80 PDF

### **Self-medication profiles in school-age adolescents in Surabaya city, Indonesia**

Mufarrihah Mufarrihah, Ana Yuda, Abhimata Paramanandana, Dini Retnowati, Devy Maulidya Cahyani, Retno Sari, Sugiyartono Sugiyartono, Tutiek Purwanti, Dewi Isadiartuti, Esti Hendradi, Andang Miatmoko

 <https://doi.org/10.4081/jphia.2023.2530> 0  0  0  0 75  PDF: 68 PDF

### **The effect of polysorbate 20 and polysorbate 80 on the solubility of quercetin**

Tristiana Erawati, Dewi Isadiartuti, Bintari Damartha Anggalih

 <https://doi.org/10.4081/jphia.2023.2503>

 0  0  0  0

 41  PDF: 74

 PDF



### ***In silico* screening of potential compounds from begonia genus as 3CL protease (3CL pro) SARS-CoV-2 inhibitors**

Saipul Maulana, Tutik Sri Wahyuni, Prihartini Widiyanti, Muhammad Sulaiman Zubair

 <https://doi.org/10.4081/jphia.2023.2508>

 0  0  0  0

 95  PDF: 54

 PDF



### **The activity of candlenut oil in the nanostructured lipid carrier system on hair growth in rats**

Tristiana Munandar Erawati, Noorma Rosita, Intan Rachmania

 <https://doi.org/10.4081/jphia.2023.2519>

 0  0  0  0

 94  PDF: 73

 PDF



### **Patterns of bronchodilator therapy in asthmatic outpatients**

Toetik Aryani, Riska Kholifatul Rahmawati, Ni Putu Cintyadewi, Arina Dery Puspitasari, Alfian Nur Rasyid, Samirah Samirah

 <https://doi.org/10.4081/jphia.2023.2533>

 0  0  0  0

 605  PDF: 80



## Chemoinformatics approach to design and develop vanillin analogs as COX-1 inhibitor

Norhayati Norhayati, Juni Ekowati, Nuzul Wahyuning  
Diyah, Bimo Ario Tejo, Samar Ahmed

<https://doi.org/10.4081/jphia.2023.2517>

0 0 0 0

600 PDF: 77

PDF



## The association between drug therapy problems and blood pressure control of patients with hypertension in public health center setting

I Nyoman Wijaya, Umi Athiyah, Fasich Fasich, Abdul  
Rahem, **Andi Hermansyah**

<https://doi.org/10.4081/jphia.2023.2531>

0 0 0 0

359 PDF: 78

PDF



## The sodium does not affect joint pain and functional activity of knee osteoarthritis patients

Anisyah Achmad, Suharjo Suharjo, Joewono  
Soeroso, Budi Suprapti, Siswandono Siswandono,  
Liza Pristianty, Mahardian Rahmadi, Jusak Nugraha,  
Cahyo Wibisono Nugroho, Yoki Surya, Satria Pandu  
Persada Isma, Erreza Rahadiansyah, Thomas Erwin  
C.J. Huwae, Bagus Putu Putra Suryana

<https://doi.org/10.4081/jphia.2023.2494>

0 0 0 0

156 PDF: 72

PDF



### The effect of isolated probiotics from Indonesian *Passiflora edulis sims.* on interferon gamma levels in peripheral blood mononuclear cell of adult tuberculosis patients *in vitro*

lif Hanifa Nurrosyidah, Ni Made Mertaniasih, Isnaeni Isnaeni

<https://doi.org/10.4081/jphia.2023.2504>

0 0 0 0

110 PDF: 60

PDF



### Moisturized and non-irritating hand gel based on sappan wood (*Caesalpinia sappan* L.) and limau citrus peel (*Citrus amblycarpa* (hassk.) ochse) extracts

Dina Yuspita Sari, Genta Windi Lestari, Habiba Fikri Farika Pulungan, Ira Remiyati, Ratna Widayarsi

<https://doi.org/10.4081/jphia.2023.2509>

0 0 0 0

101 PDF: 68

PDF



### The effect of peppermint oil addition on the physical stability, irritability, and penetration of nanostructured lipid carrier coenzyme Q10

Tristiana Erawati, Rizki Amalia Arifiani, Andang Miatmoko, Dewi Melani Hariyadi, Noorma Rosita, Tutiek Purwanti

<https://doi.org/10.4081/jphia.2023.2515>

0 0 0 0

76 PDF: 80

PDF



[HOME](#) / [Editorial Board](#)

## Editorial Board

### Editors-in-Chief

Prof. Nicaise Ndembi

Institute of Human Virology, *Nigeria*

Prof. Vittorio Colizzi

University of Rome Tor Vergata, *Italy*

Faculty of Science & Technology, Evangelic University of Cameroon, *Cameroon*

[vittorio.colizzi@publichealthinafrica.org](mailto:vittorio.colizzi@publichealthinafrica.org)

### Managing Editor

Mrs. Emanuela Fusinato

PAGEPress, *Italy*

### Board Members

Dr. John Nkengasong

Africa Centres for Disease Control and Prevention, *Cameroon*

Dr. Chikwe Ihekweazu

Nigeria Centres for Disease Control and Prevention, *Nigeria*

Dr. Alex Riolerus Ario

Uganda National Institute of Public Health, Ministry of Health, *Uganda*

Dr. Elvira Singh

National Cancer Registry, National Health Laboratory Service, *South Africa*

Prof. Abderrahmane Maaroufi

Institut Pasteur du Maroc [National Public Health Institute], *Morocco*

Dr. Ileshi Jani

National Institute of Health, *Mozambique*

Dr. Ebba Abate

Ethiopian Public Health Institute, Addis Ababa, *Ethiopia*

Dr. Mazyanga Lucy Mazaba Liwewe

Zambia National Public Health Institute, *Zambia*

Prof. Vincent Batwala

Mbarara University of Science & Technology, *Uganda*

Dr. Giacomo Paganotti

Botswana-University of Pennsylvania, *Botswana*

Dr. Georges Etoundi

Direction de la Lutte Contra la Maladie, les Epidemies et les Pandemies, *Cameroon*

Prof. Jacques Simpore

Joseph Ki-Zerbo University, *Burkina Faso*

Dr. Tolbert Nyenswah

Johns Hopkins University Bloomberg School of Public Health, *United States*

Dr. Souha Bougatef

National Observatory of New and Emerging Diseases, *Tunisia*

Dr. Raji Tajudeen

Africa Centres for Disease Control and Prevention, *Ethiopia*

Dr. Mohammed Abdulaziz

Africa Centres for Disease Control and Prevention, *Ethiopia*

Prof. Epée Emilienne

University of Yaoundé I, *Cameroon*

Dr. Nafisah Chotun

Africa Centres for Disease Control and Prevention, *Ethiopia*

The Journal of  
Public Health in  
Africa *is an Open  
Access, peer-  
reviewed journal  
published by Africa  
CDC, Addis Ababa,  
Ethiopia. All credits  
and honors to PKP  
for their OJS.*

About

About The Journal

Editorial Board

Info Submission

Submissions

Guidelines For  
Authors



pISSN: 2038-9922

eISSN: 2038-9930

FOR AUTHORS

SUBMIT YOUR PAPER

Guide for Authors

Benefits for Authors

How to write a scientific paper

How to write a Review article

Article Processing Charge

FOR REVIEWERS

Benefits for Reviewers

How to review

INDEXING

PubMed

PubMed Central

DOAJ

Scopus

ESCI

CABI Abstracts

MOST CITED ARTICLE

Global burden of diseases attributable to air pollution

” Cited: 43

📅 21 December 2018

© PAGEPress 2008-2023 • PAGEPress® is a registered trademark property of PAGEPress srl, Italy • VAT: IT02125780185 • Privacy

 Samuel Soledayo Babatola

---

Health care delivery in Malaysia:  
changes, challenges and champions

” Cited: 41

 5 September 2011

 Susan Thomas et al.

---

Investigation of medicinal plants  
traditionally used as dietary  
supplements: A review on Moringa  
oleifera

” Cited: 40

 21 December 2018

 Ivana Matic et al.

---

Last Updated: 3 April 2023



#### MOST READ LAST MONTH

Case based measles surveillance  
performance in 2010, littoral region of  
Cameroon

 229

---

Health economics in Africa from 1991  
to 2020: A systematic review

 183


---

Development of self-report  
assessment tool for anxiety among  
adolescents: Indonesian version of the  
Zung self-rating anxiety scale

 183

---

#### KEYWORDS



# The association between drug therapy problems and blood pressure control of patients with hypertension in public health center setting

I Nyoman Wijaya, Umi Athiyah, Fasich, Abdul Rahem, Andi Hermansyah

Department of Pharmacy Practice, Faculty of Pharmacy, Universitas Airlangga, Surabaya, Indonesia

## Abstract

**Background:** Patients with hypertension are at risk of experiencing Drug Therapy Problems (DTPs). However, few have studied the pattern of DTPs in Indonesian public health center (PHC) and how it affected the blood pressure control.

**Objective:** This study aims to identify DTPs and its associa-

Correspondence: I Nyoman Wijaya, Department of Pharmacy Practice, Faculty of Pharmacy, Universitas Airlangga, 60115, Surabaya, Indonesia.

Tel.: +62.315933150 - Fax: +62.315935249

E-mail: i-nyoman-w@ff.unair.ac.id

**Key words:** Drug therapy problem, Hypertension, Clinical outcome, Health care.

**Acknowledgments:** The authors highly appreciated the participation of respondents and pharmacists in PHC in Surabaya.

**Contributions:** INW, project leader, contributions to basic concept of the work, funding acquisition, manuscript writing and revision; UA, substantial contributions to the design of the experiments; F, project administration; AR, data acquisition and interpretation; AH, critically revising the manuscript for important intellectual content. All authors approved the final version to be published.

**Conflict of interest:** The authors declare no potential conflict of interest.

**Funding:** This research was funded by Hibah Mandat Grant Scheme of Universitas Airlangga.

**Ethical approval and consent to participate:** Ethical clearance was obtained from the Research Ethics Commission of the Faculty of Public Health, Universitas Airlangga (No. 200/EA/KEPK/2019).

**Availability of data and material:** Data and materials are available by the authors.

**Informed consent:** The manuscript does not contain any individual person's data in any form.

Received for publication: 28 December 2022.

Revision received: 5 February 2023.

Accepted for publication: 11 February 2023.

This work is licensed under a Creative Commons Attribution NonCommercial 4.0 License (CC BY-NC 4.0).

©Copyright: the Author(s), 2023

Journal of Public Health in Africa 2023; 14(s1):2531

doi:10.4081/jphia.2023.2531

tion to blood pressure control among patients with hypertension in Indonesian PHCs.

**Methods:** A descriptive, observational and cross-sectional study involving 150 selected persistent hypertensive patients was conducted from August to December 2019 in 63 PHCs in Surabaya. A questionnaire asking patients' experiences during treatment of hypertension was used. Chi-square test was used to analyze the association between DTPs and blood pressure control.

**Results:** The majority of respondents were aged 50-65 (54.7%), female (76%), only half has controlled blood pressure (52.7%) and most of them used three to four medicines at the same time (57.3%). A total 563 DTPs was identified with 15.6% was unnecessary drug use, 11.4% indicated need for additional drug therapy, one-fifth experienced dosage too low (21.5%) and non-adherence (19.2%) and 26% suffered from adverse drug reactions. There was a significant association between number of DTPs and blood pressure control ( $P < 0.05$ ).

**Conclusion:** Most patients experienced more than two DTPs and undertook more than three medicines at the same time. There is a significant association between the number of DTPs in hypertensive patient and the blood pressure control.

## Introduction

The shift of pharmacist orientation from product to services and patient's quality of life has been coined in the philosophy of pharmaceutical care.<sup>1</sup> Within this philosophy, pharmacists hold direct responsibility to assess, prevent and resolve drug therapy problems (DTPs).<sup>2</sup> DTPs have been defined as any undesirable events, actual or potential, that may prevent patients from the benefits of drug therapy.<sup>3</sup>

DTPs have been major concern in patients with chronic condition including hypertension. In fact, due to the possibility of polypharmacy and presence of multiple comorbidities, patients with hypertension are at high risk for DTPs. Costa *et al* (2019) highlighted that at least one DTP was identified from hypertensive patients undertaking polypharmacy.<sup>3</sup> This is not the only issue as battling against hypertension has been major homework globally.

World Health Organization mentioned that at least 1 of 3 people worldwide suffered from hypertension. This number might increase to 1.5 billion people by 2025 with fatality reaches 9.4 million people die from hypertension and its complications every year.<sup>4</sup> Indonesian population is also not immune to hypertension. There was a significant increase of hypertension prevalence among population aged 18 years from 25.8% (2013) to 34.1% (2018) with estimation of 63 million people living with hypertension.<sup>5</sup> Likewise, both the prevalence in East Java – the second largest province – and in Surabaya – the second largest city in Indonesia – have reached almost 20% of the population.<sup>6,7</sup>

Identifying DTPs in patients with hypertension is critical as it may lead to uncontrolled blood pressure particularly in resource-limited setting countries like Indonesia. For instance, a study in

Ethiopia determined the prevalence of DTPs ranged from 73-80% with poor selection of antihypertensive drugs, dosing issues and drug interactions were the major problems.<sup>9</sup> This number is not different to other countries including Brazil<sup>10</sup> and Nigeria.<sup>9</sup> Given that most of Indonesians have been commonly treated in public health center (PHC), it is therefore important to conduct this study in PHC setting.

## Materials and Methods

A descriptive, observational and cross-sectional study was conducted from August to December 2019 in 63 PHCs in Surabaya. Approximately two or three respondents were recruited using purposive sampling. Only patients who had been diagnosed with hypertension for more than six months, taking at least one antihypertensive drug, speak Indonesian, and providing consent for participation were recruited. A minimum sample of 150 respondents was required for this study. Patients completed interviewer-assisted questionnaire about the experience during therapy. Any presence of DTPs was recorded and subsequently analyzed for its association with blood pressure control using Chi-square test. Ethical clearance was obtained from the Research Ethics Commission of the Faculty of Public Health, Universitas Airlangga (No. 200/EA/KEPK/2019).

## Results and Discussion

### Characteristics of respondents

Table 1 shows that majority of respondents were aged 50-65 (54.7%) and female (76%). This result may slightly differ to some studies mentioning that females are less likely to have hypertension than male. However, after the age of 40 years, women are prone to hypertension than men due to changes in physiology and hormonal reducing the elasticity of blood vessels which may accelerate the increased of blood pressure.<sup>11</sup> This finding has strengthened other studies that women aged 40 years are susceptible to hypertension.<sup>12-15</sup>

### Respondents' experience with hypertension

Most respondents have suffered from hypertension for five years (26.7%). Half of the respondents had blood pressure of 120-139 mmHg and 80-89 mmHg (50.7%) indicating stage of pre-hypertension according to JNC VIII and they were having controlled blood pressure (52.7%) within the past 6 months. Two-third of the respondents took three to four medicines at the same time (Table 2).

The high portion of respondents with controlled blood pressure is in contrast to study in Lesotho with only 20% of hypertensive patients were in controlled state of blood pressure.<sup>14</sup> Understanding the state of blood pressure is important as hypertension requires long-term pharmacotherapy to prevent complications such as stroke and heart and kidney disorders.<sup>16</sup>

The fact that most respondents were taking polypharmacy is not surprising since anecdotal evidence illustrating that hypertensive patient visiting PHC in Surabaya often received more than one drug some of which were amlodipine, HCT, captopril, nifedipine, candesartan and lisinopril. In addition, other comorbidities and symptoms may present including pain, ulcers and cough, which makes patients receive more than one drug. This is in line to PHCs in Central Sulawesi stated that in addition to antihypertensive drugs, patients also used paracetamol (13%), mefenamic acid and

Ibuprofen (<2%).<sup>17</sup> The high number of medicines taken by patients may result non-compliance and possible drug interactions,<sup>2</sup> as indicated by a study in UK, showed that hypertensive patients who used three or more drug items were tended to incompliance to treatment.<sup>18</sup>

### Pattern of DTPs

A total 563 DTPs was recorded as highlighted in Table 3. Dosage too low (21.5%), adverse drug reaction (26.6%) and non-compliance (19.2%) are three DTPs recorded from one-fifth of respondents, respectively. This finding reflected that both actual and potential DTPs may exist in patients with hypertension. The most common adverse drug reaction as expressed by respondents was due to taking Captopril resulting dry cough. In addition, the use of HCT in patients may have contributed to hyperuricemia, hyperlipidemia, and hyperglycemia.

Another DTPs recorded was an unnecessary drug therapy that indicated that patient did not necessarily need the drug. As found

**Table 1. Characteristics of respondents.**

Characteristics (n=150)	Frequency	Percentage
Age (in years)		
<50	35	23.3
50-65	82	54.7
>65	33	22.0
Sex		
Male	36	24.0
Female	114	76.0
Education		
Illiterate	21	14.0
Elementary school	55	36.7
Junior high school	25	16.7
Senior high school	35	23.3
Diploma	3	2.0
Bachelor	10	6.7
Postgraduate	1	0.6
Occupation		
Unemployed	26	17.3
Government employee	4	2.7
Retired	9	6.0
Housewife	65	43.3
Private employee	46	30.7

**Table 2. Respondents' experience with hypertension.**

	Frequency	Percentage
Experience with hypertension (n=150)		
<1	31	20.6
1-5	40	26.7
5-10	39	26.0
10-20	37	24.7
>20	3	2.0
Outcome Therapy (Blood Pressure in mmHg)		
<120 and 80	3	2.0
120-139 and 80-89	76	50.7
140-159 and 90-99	30	20.0
160-180 and 100-110	39	26.0
>180 and >110	2	1.3
Blood pressure control		
Controlled blood pressure	79	52.7
Uncontrolled blood pressure	71	47.3
Number of Drug Items in Respondents		
<3 items	19	12.7
3-4 items	86	57.3
>4 items	45	30.0

in this study for example if patient received two painkillers at once such as paracetamol and mefenamic acid. Approximately 15% of respondents experienced unnecessary drug therapy. A few respondents experienced ineffective drug therapy (1.1%) since patients cannot achieve the desired therapeutic target after using the drug for a particular period. For instance, respondents received sympathomimetic drugs for reducing his or her common cold, which eventually increase blood pressure.<sup>19-24</sup>

Dosage too low occurred in more than 20% of respondents. For instance, a patient used 5mg amlodipine for several weeks, but his blood pressure was still above 160/100mmHg. This indicated that change to dose regimentation is necessary, yet nothing was done. Dosage too low also occurred in case patients using NSAIDs that constrict blood vessels.<sup>20,24</sup> In contrast, DTPs of dosage too high (4.6%) has been evident in patients taking two tablets of amlodipine 10 mg at once. This is against the maximum daily dose of amlodipine which is 10 mg. Doubling the dose may not necessarily need since combination of therapy of antihypertension, for instance with diuretics or other appropriate antihypertensive groups, can result better evidence.<sup>16,21</sup> In addition, a dosage too high can occur as a result of drug interactions. For example, concurrent use of simvastatin and amlodipine can cause muscle pain in patients.<sup>20</sup>

DTPs of non-compliance (19.2%) in this study illustrated by respondents did not refill drugs and forgot to use drugs as instructed. The overall DTPs in this study are similar to a study in Medan, North Sumatra mentioning unnecessary drug therapy (7.6%), needs additional drug therapy (47.0%), ineffective drug therapy (21.2%), adverse drug reaction (9,1%), dosage too low (1.5%), and non-compliance (13.6%) are the pattern of DTPs in PHC.<sup>22</sup>

Table 4 highlighted that most respondents had three to four

**Table 3. Pattern of DTPs in respondents.**

Number of types of DTPs (n=150)	Frequency	Percentage
Unnecessary drug therapy	88	15.6
Needs additional drug therapy	64	11.4
Ineffective drug therapy	6	1.1
Dosage too low	121	21.5
Adverse drug reaction	150	26.6
Dosage too high	26	4.6
Noncompliance	108	19.2
Total DTPs	563	100.0

**Table 4. Number of DTPs in respondents.**

Number of DTPs in respondents (n=150)	Frequency	Percentage
<3 DTPs	22	14.7
3-4 DTPs	86	57.3
>4 DTPs	42	28.0

**Table 5. Chi-square analysis.**

	Blood pressure control	Number of DTPs
Chi-Square	0.427 <sup>a</sup>	10,493 <sup>b</sup>
df	1	6
Asymp. Sig.	0.514	0.000

<sup>a</sup>0 cells (0,0%) have expected frequencies less than 5. The minimum expected cell frequency is 75,0. <sup>b</sup>0 cells (0,0%) have expected frequencies less than 5. The minimum expected cell frequency is 21,4. Chi-Square 103,493, Asymp. Sig. = 0.000 < 0.05, H0 is accepted (there is a relationship).

DTPs (57,3%) similar to a study by Redzuan *et al.* (2017) in Malaysia concluding on average 2.15±1.5 DTPs per patient with 88.8% of patients had at least one DTPs.<sup>23</sup>

### Association between DTPs and blood pressure control

The results of statistical analysis with chi-square test in Table 5 obtained *p-value* = 0.000 (<0.05) illustrating there is significant association between the number of DTPs and blood pressure control either controlled or uncontrolled.

### Conclusions

Most patients experienced more than two DTPs and undertook more than three medicines at the same time. There is a significant association between the number of DTPs in hypertensive patient and the blood pressure control.

### References

1. Ministry of Health of the Republic of Indonesia. Decree of the Minister of Health No. 1027/Menkes/SK/IX/2004 concerning Pharmaceutical Service Standards in Pharmacies. Jakarta: Government of the Republic of Indonesia 2004.
2. Cipolle JR, Strand ML, Morley CP. Pharmaceutical Care Practice: The Patient-centered Approach to Medication Management Services. 3rd Ed. New York: Mc. Graw Hills Inc 2012.
3. Costa FA, Mil JW, Risco AA. The Pharmacist Guide to Implementing Pharmaceutical Care. Switzerland: Springer International Publishing AG 2019.
4. World Health Organization. A Global Brief on Hypertension Silent Killer. Global Public Health Crisis 2015.
5. Ministry of Health of the Republic of Indonesia. Basic Health Research 2018. Jakarta: Ministry of Health of the Republic of Indonesia 2018.
6. East Java Provincial Health Office. Health Profile of East Java Province 2018, Surabaya 2019.
7. Surabaya City Health Office. Profile of the Surabaya City Health Office 2019. Surabaya 2019.
8. Dyker AG. Hypertension: Clinical Pharmacy and Therapeutics. Fifth edition. London: Churchill Livingstones 2012.
9. Weldegebreal AS, Tezeta F, Mehari AT, et al. Assessment of drug therapy problem and associated factors among adult hypertensive patients at Ayder comprehensive specialized hospital, Northern Ethiopia. *Afri Health Sci* 2019;19:2571-9.
10. Neves CM, Nascimento MMG, Silva DAM, Oliveira DR. Clinical Results of Comprehensive Medication Management Services in Primary Care in Belo Horizonte, Pharmacy. *Pharmacy* 2019;7:58.
11. Anggara FHD, Prayitno N. Factors related to blood pressure at Telaga Murni Health Center, West Cikarang in 2012, *Jurnal Ilmiah Kesehatan* 2013;5:20-5.
12. Desy A, Santi M. The Relationship between Characteristics and Central Obesity with the Incidence of Hypertension. *J Berkala Epidemiologi* 2018;6:43-50.
13. Mafaza RL, Wiratmadi B, Adriani M. Analysis of the relationship between abdominal circumference, fat intake, and the ratio of calcium-magnesium intake to hypertension. *Media Gizi Indonesia* 2016;11:127-34.
14. Thinyane KH, Mothebe T, Sooro M, et al. An observational study of hypertension treatment and patient outcomes in a

- primary care setting. *Pan Afr Med J* 2015; 20:424.
15. Tular GJ, Ratag BT, Kandou GD. The relationship between physical activity, family history and age with the incidence of hypertension in Tarabitan Village, Likupang Barat District, North Minahasa Regency. *Media Kesehatan* 2017; 9:1-6.
  16. Saseen JJ, Maclaughlin EJ. *Pharmacotherapy: A Pathophysiologic Approach*. 7th ed. New York: McGraw-Hill Companies, Inc 2012.
  17. Alaydrus S. Profile of Drug Use in Hypertensive Patients at Marawola Health Center January-March 2017. *J Mandala Pharmacon Indonesia* 2017;3:110-8.
  18. Hameed MA, Dasgupta I. Medication adherence and treatment resistant hypertension: a review. *Drugs in Context* 2019;8:212560.
  19. Katzung GB, Trevor JA. *Antihypertensive Agents: Basic and Clinical Pharmacology*. 13th ed. United State: McGraw-Hill Education 2014.
  20. Baxter K. *Stockley's Drug Interactions*. Eighth edition. London: Pharmaceutical Press; 2008.
  21. JNC-8 (The Eighth Joint National Committee). Evidence-Based Guideline for the Management of High Blood Pressure in Adults. *Guideline for Management of High Blood Pressure*. Ame Med Assoc 2014.
  22. Nasution A, Khairunnisa, Tanjung HR. Drug Therapy Problems In Management Of Hypertensive Outpatients Admitted To Four Indonesian Primary Health Centers. *Asian J Pharma Clin Res* 2016;9:87-90.
  23. Redzuan AM, Ramli AR, Pheng MTH. Drug-Related Problems in Hypertensive Patients with Multiple Comorbidities. *J Pharma Res* 2017;1:1-8.
  24. Ritter JM, Lewis LD, Mant TGK, Ferro AA *Textbook of Clinical Pharmacology and Therapeutics*. Fifth Edition. London: Hodder Arnold, an imprint of Hodden Education, part of Hachette Livre UK; 2008.

Non-commercial use only



## Journal of Public Health in Africa

### COUNTRY

Italy



Universities and research  
institutions in Italy



Media Ranking in Italy

### SUBJECT AREA AND CATEGORY

Medicine  
Public Health, Environmental and  
Occupational Health

### PUBLISHER

PagePress Publications

### H-INDEX

16

### PUBLICATION TYPE

Journals

### ISSN

20389922, 20389930

### COVERAGE

2010-2014, 2016-2021

### INFORMATION

[Homepage](#)

[How to publish in this journal](#)

[vittorio.colizzi@publichealthinfrica.org](mailto:vittorio.colizzi@publichealthinfrica.org)

### SCOPE

The Journal of Public Health in Africa (JPHiA) is a peer-reviewed, academic journal that focuses on health issues in the African continent. The journal editors seek high quality original articles on public health related issues, reviews, comments and more. The aim of the journal is to move public health discourse from the background to the forefront. The success of Africa's struggle against disease depends on public health approaches.



Join the conversation about this journal




Quartiles

FIND SIMILAR JOURNALS

options

1 <b>BMC Research Notes</b> GBR <b>54%</b> similarity	2 <b>Pan African Medical Journal</b> NGA <b>52%</b> similarity	3 <b>Open Public Health Journal</b> NLD <b>52%</b> similarity	4 <b>Malawi Medical Journal</b> MWI <b>49%</b> similarity	5 <b>International Health</b> GBR <b>49%</b> similarity
---	--	---	---	---



Journal of Public Health in Africa  
Public Health, Environmental and Occupational...  
Q3 best quartile  
SJR 2021 0.27  
powered by scimagojr.com

Show this widget in your own website  
Just copy the code below and paste within your html code:  
<a href="https://www.scimagojr.com/journal.php?q=21100781400&tip=si...">

SCImago Graphica  
Explore, visually communicate and make sense of data with our new data visualization tool.

Metrics based on Scopus® data as of April 2022

A **Anas** 2 years ago

Dear SJR team,  
I'm making this comment to let you know and the scientific community, that this journal is not professional. It has been now 13 months that we have submitted our paper, and the status remains the same. The journal does not respond to the authors' questions and reminders. They hold papers and do not answer no matter how many emails you send them. DON'T MAKE THE SAME MISTAKE AS WE DID.

reply

**Melanie Ortiz** 2 years ago

Dear Anas, thanks for your participation! Best Regards, SCImago Team

SCImago Team

P **Prof Oche Oche** 3 years ago

I sent a manuscript titled "Knowledge, Attitude and preventive Practices of Dengue fever among Health Care workers in a tertiary health institution in Northwest Nigeria" for publication over a year ago. since then I have sent several reminders with no response, I wonder why it is taking too long to respond and even review the paper as the authors are already thinking of withdrawing the paper. I hope you will let us know the current status of the paper.

prof. Oche .M.O  
corresponding author

reply



**Melanie Ortiz** 3 years ago

SCImago Team

Dear Oche,  
thank you for contacting us.  
We are sorry to tell you that SCImago Journal & Country Rank is not a journal. SJR is a portal with scientometric indicators of journals indexed in Elsevier/Scopus.  
Unfortunately, we cannot help you with your request, we suggest you contact the journal's editorial staff, so they could inform you more deeply.  
Best Regards, SCImago Team

N **Nduagubam** 3 years ago

What is the current publisher of JPHIA?

reply



**Melanie Ortiz** 3 years ago

SCImago Team

Dear Nduagubam,  
thank you for contacting us.  
It seems that the Publisher remains the same. However, we suggest you to contact the journal's editorial staff, so they could inform you more deeply.  
Best Regards, SCImago Team

D **Dr Sonopant Joshi** 5 years ago

Dear Sir / Madam

I have conducted study on " Model on Antenatal care on selected parameters related to Pregnancy outcome "

I would like to publis in your journal. What is processing fee to publish in INR

Thanks

reply



**Elena Corera** 5 years ago

SCImago Team

Dear Dr Sonopant Joshi,

thank you very much for your comment. Unfortunately, we cannot help you with your request, we suggest you contact journal's editorial staff so they could inform you more deeply.

Best Regards,  
SCImago Team

Leave a comment

Name

Email  
(will not be published)

Submit

The users of Scimago Journal & Country Rank have the possibility to dialogue through comments linked to a specific journal. The purpose is to have a forum in which general doubts about the processes of publication in the journal, experiences and other issues derived from the publication of papers are resolved. For topics on particular articles, maintain the dialogue through the usual channels with your editor.

Developed by:



Powered by:



Follow us on @ScimagoJR

Scimago Lab, Copyright 2007-2022. Data Source: Scopus®

EST MODUS IN REBUS  
Horatio (Satire 1.1.100)

[Cookie settings](#)

[Cookie policy](#)



# Source details

## Journal of Public Health in Africa

Open Access ⓘ

Scopus coverage years: from 2010 to 2014, from 2016 to Present

Publisher: PagePress

ISSN: 2038-9922 E-ISSN: 2038-9930

Subject area: Medicine: Public Health, Environmental and Occupational Health

Source type: Journal

CiteScore 2021

1.5 ⓘ

SJR 2021

0.268 ⓘ

SNIP 2021

0.434 ⓘ

[View all documents >](#)

[Set document alert](#)

[Save to source list](#) [Source Homepage](#)

[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)

### i Improved CiteScore methodology ⓘ

CiteScore 2021 counts the citations received in 2018-2021 to articles, reviews, conference papers, book chapters and data papers published in 2018-2021, and divides this by the number of publications published in 2018-2021. [Learn more >](#)

CiteScore 2021 ▾

$$1.5 = \frac{206 \text{ Citations } 2018 - 2021}{139 \text{ Documents } 2018 - 2021}$$

Calculated on 05 May, 2022

CiteScoreTracker 2022 ⓘ

$$0.8 = \frac{168 \text{ Citations to date}}{211 \text{ Documents to date}}$$

Last updated on 05 April, 2023 • Updated monthly

## CiteScore rank 2021 ⓘ

Category	Rank	Percentile
Medicine		
Public Health, Environmental and Occupational Health	#394/562	29th

[View CiteScore methodology >](#) [CiteScore FAQ >](#) [Add CiteScore to your site ↗](#)

---

## About Scopus

[What is Scopus](#)

[Content coverage](#)

[Scopus blog](#)

[Scopus API](#)

[Privacy matters](#)

## Language

[日本語版を表示する](#)

[查看简体中文版本](#)

[查看繁體中文版本](#)

[Просмотр версии на русском языке](#)

## Customer Service

[Help](#)

[Tutorials](#)

[Contact us](#)

---

## ELSEVIER

[Terms and conditions ↗](#) [Privacy policy ↗](#)

Copyright © Elsevier B.V. ↗. All rights reserved. Scopus® is a registered trademark of Elsevier B.V.

We use cookies to help provide and enhance our service and tailor content. By continuing, you agree to the use of cookies ↗.

