ISSN-0973-9122 (Print) . ISSN-0973-9130 (Electronic)

Volume 15 / Number 2 / April-June 2021

Indian Journal of Forensic Medicine & Toxicology

Website: www.ijfmt.com

Official Organ of Indian Association of Medico-Legal Experts (Regd.)

Indian Journal of Forensic Medicine & Toxicology

EDITOR in Chief

Prof. R K Sharma

Formerly at All India Institute of Medical Sciences, New Delhi, E-mail: editor.ijfmt@gmail.com

EDITOR

Prof. Dr. Adarsh Kumar

Forensic Medicine & Toxicology, AIIMS, New Delhi

INTERNATIONAL EDITORIAL ADVISORY BOARD

- 1. Prof Mete Gulmen Cukurova University, TURKEY
- Prof. Leandro Duarte De Carvalho, Minas Gerais, Belo Horizante, Brazil
- Prof. Donata Favretto (Full Professor) Forensic Toxicology at University of Padova, Italy
- Prof. Babak Mostafazadeh Department of Forensic Medicine & Toxicology, Shahid Beheshti University of Medical Sciences, Tehran-Iran
- Prof Halis Dokgoz, Mersin University, TURKEY
- 6. Prof Jozef Sidlo, Comenius University, Bratislava, SLOVAKIA
- Dr. Rahul Pathak (Lecturer) Forensic Science, Dept of Life Sciences Anglia Ruskin University, Cambridge, United Kingdom
- Dr. Hareesh (Professor & Head) Forensic Medicine, Ayder Referral Hospital, College of Health Sciences, Mekelle University, Mekelle Ethiopia East Africa
- Dr. Mokhtar Ahmed Alhrani (Specialist) Forensic Medicine & Clinical Toxicology, Director of Forensic Medicine Unit, Attorney General's Office, Sana'a, Yemen
- Dr. Sarathchandra Kodikara (Senior Lecturer) Forensic Medicine, Department of Forensic Medicine, Faculty of Medicine, University of Peradeniya, Sri Lanka
- Dr Noha A. Magdie El Rafie, Forensic Toxicology, Ain Shams University, Cairo, EGYPT

SCIENTIFIC COMMITTEE

- Prof Udai Pratap Singh, Department of Anthropology Lucknow University Lucknow
- Dr Anil Rahule (Associate Professor) Dept of Anatomy, Govt Medical College Nagpur
- Dr Shankar Bakkanwar (Associate Professor) Forensic Medicine, Kasturba Medical College, Manipal, Karnatakad
- 4. Dr K. Ravikumar Raksha Shakti University, Ahmedabad, Gujrat.
- Dr. Pragnesh Parmar (Associate Professor) Forensic Medicine, Valsad, Gujrat
- Dr Vandana Mudda (Awati) (Associate Prof) Dept of FMT, M.R.Medical College, Gulbarga, Karnataka,
- Dr. Asha Srivastava (Senior Scientific Officer) Forensic Psychology, Central Forensic Science Laboratory, CBI, Delhi
- Dr. Lav Kesharwani (Asst.Prof.) School of Forensic Science, Sam Higginbottom Institute of Agriculture Technology & Sciences, Allahabad U.P,
- Dr. Anu Sharma (Associate Prof) Dept of Anatomy, DMCH, Ludhiana (PB)
- Dr. Shalini Gupta (Prof) Oral Pathology and Microbiology, Dental Sciences King George Medical University, Lucknow, UP
- 11. Dr Rituja Sharma, Associate Prof, Law Banasthali Vidyapeeth Jaipur

"Indian Journal of Forensic Medicine & Toxicology" is peer reviewed quarterly journal. It deals with Forensic Medicine, Forensic Science, Toxicology, DNA fingerprinting, sexual medicine and environment medicine. It has been assigned International standard serial No. p-0973-9122 and e- 0973-9130. The Journal has been assigned RNI No. DELENG/2008/21789. The journal is indexed with Index Copernicus (Poland) and is covered by EMBASE (Excerpta Medica Database). The journal is also abstracted in Chemical Abstracts (CAS) database (USA. The journal is also covered by EBSCO (USA) database. The Journal is now part of UGC, DST and CSIR Consortia. It is now offical publication of Indian Association of Medico-Legal Experts (Regd.).

NATIONAL EDITORIAL ADVISORY BOARD

Prof Sudhir K Gupta - Head, Department of Forensic Medicine All India Institute of Medical Sciences, New Delhi

Members

- 1. Prof. SK Dhattarwal, Forensic Medicine, PGIMS, Rohtak, Haryana
- 2. Prof. N K Aggrawal Forensic Medicine, UCMS, Delhi
- Prof Ajay Ghangale Forensic Medicine Dr DY Patil Medical College, Pune, Maharashtra
- Dr. Amar Jyoti Patwory Professor, Forensic Medicine NEIGRIHMS, Shillong
- Dr S. Venkata Raghava Professor, Forensic Medicine, Banglore Medical College, Bengaluru
- Prof Praveen Arora, Professor Department of Forensic Medicine & Toxicology, SAIMS, Indore
- Dr. Pankaj Datta (Principal & Head) Department of Prosthodontics, Indraprastha Dental College & Hospital, Ghaziabad
- Dr. Mahindra Nagar (Head) Department of Anatomy, UCMS & GTB Hospital, Delhi
- Dr. Virender Kumar Chhoker Professor Forensic Medicine and Toxicology, Santosh Medical College, Ghaziabad, UP
- Dr. Dayanand G Gannur (Professor) Department of Forensic Medicine & Toxicology, Shri BM Patil Medical College, Hospital & Research centre, Bijapur, Karnataka
- Dr. Alok Kumar Professor Department of Forensic Medicine & Toxicology, UP Rural Institute of Medical Sciences and Research, Saifai, Etawah, U.P.

Print-ISSN:0973-9122 Electronic - ISSn: 0973-9130

Frequency: Quarterly, © All Rights reserved The views and opinions expressed are of the authors and not of the Indian Journal of Forensic Medicine & Toxicology, Indian Journal of Forensic Medicine & Toxicology does not guarantee directly or indirectly the quality or efficacy of any products or service featured in the advertisement in the journal, which are purely commercial.

Website: www.ijfmt.com

Editor

Dr. R.K. Sharma

Institute of Medico-legal Publications

Logix Office Tower, Unit No. 1704, Logix City Centre Mall, Sector- 32, Noida - 201 301 (Uttar Pradesh)

Printed, published and owned by

Dr. R.K. Sharma

Logix Office Tower, Unit No. 1704, Logix City Centre Mall, Sector- 32, Noida - 201 301 (Uttar Pradesh)

Published at

Institute of Medico-legal Publications

Logix Office Tower, Unit No. 1704, Logix City Centre Mall, Sector- 32, Noida - 201 301 (Uttar Pradesh)



Indian Journal of Forensic Medicine & Toxicology

Contents

Vo	lume 15, Number 2 April-June 2021
1.	Psychological Aspects in Terms of Physical, Mental and Sociological Factors During Worldwide Corona Crisis
2.	Anatomical Variation of Superficial Branch of Radial Nerve in Gujrat Region
3.	Combined Effect of Neural Tissue Mobilization and Deep Friction Massage in Piriformis Syndrome: A Research Protocol
4.	Police Brutality and Custodial Torture in Technological Era: Need for Anti-Torture Law in India - A Critical Analysis
5.	Study of Socio-Demographic Profile of Thoraco-Abdominal Injuries in Fatal Road Traffic Accidents in Mangalore City
6.	Study of Mortality Due to Thoraco-Abdominal Injuries in Road Traffic Accidents in Mangalore City 32 Abymon K K, Mohd Amjad Bhat, Azia Manzoor Bhat
7.	Pattern of Thoraco-Abdominal Injuries in Fatal Road Traffic Accidents in Mangalore City
8.	An Ayurvedic Approach for Treatment of Fissure in Ano: A Case Report of 32 Year Old Patient 43 Alok kumar Diwedi, Kiran Khandare, Rakesh Khatana, Harjodh Singh
9.	Ozone Therapy - A Boon to Dental Sciences
10.	Hyperammonaemia in Absence of Deranged Liver Enzymes with Valproate: A Case Report
11.	Assessment of Anxiety during COVID-19 Pandemic among General Population of India
12.	A Conceptual Study on Prevention & Management of Covid -19 Pandemic with Special Reference to Janapadodwamsa

VIII
V III

VIII		
95.	Surgical Management of Intestinal Obstruction	574
	Prateek Shaparia, Chiraag Hiran S, Vijoy Singh, Honeypalsinh H Maharaul, Mohammad Farooque Dudhwala	
96.	Nursing Management for Covid 19 Pandemics	
	Prerana Sakharwade, Khushbu Meshram, Sonali Kolhekar	
97.	A Protocol on Effect of Cryotherapy and Myofascial release Technique in Calf Muscle Spasticity	
	of Spastic Diplegic Cerebral Palsy Children	587
	Priya J. Tikhile, Ashish W. Bele	
98.	Successful Biological Treatment of Cutaneous wound Healing with Comparation of Aqueous	
	Humour Fluid and Fat Graft; Cattle as Model	592
	Qayes Taref Ali, Muslem Fahim Diwan, Amir I. Towfik	
<mark>99</mark> .	Sensitivity and Specificity of Several Malnutrition Screening Tools for Chronic Kidney Disease	
	Patient: A Narrative Review	598
	Qonita Rachmah, Dominikus Raditya Atmaka	
100.	Neonatal Transport - Review Article	603
	R. Abhilasha Singh Panwar, Sachin Damke	
<u>101.</u>	Atrial Myxomas: A Case Report on Rare Benign Cardiac Tumour and its Histopathological	
	Diagnosis	608
	R.K. Tandon, Charu Kiran Agrawal, Niti Adhwaryu	
102.	A Rare Case Report of Giant Cervical Angiomatous Polyp	613
	R.K.Tandon, Nidhi Patel, Shivani Vora	
103.	Toxicological Assessment of Ethanolic Leaves Extract of Kalanchoe pinnata in Rats	615
	V. Saravanan, S.S. Murugan, K.R. Navaneetha Krishnan, N. Mohana, K. Sakthivel, T. N. Sathya	
104.	Analysis of Risk Management Cycle on Capitation Mechanism at First Level Healthcare	
	Facilities in East Java	623
	Rachmad Suhanda, Usman Hadi, Widodo J. Pudjirahardjo, Djazuly Chalidyanto	
105.	Effectiveness of Cervical and Upper Shoulder Stabilization Exercises in Subjects With	
	Rheumatoid Arthritis Patients Associated with Neck Pain	629
	Raj Mhatre, Smita Patil	
106.	Changes in Intra Ocular Pressure after Cataract Surgery	638
	Rajni Sethia, Nirathya Anisha Pothuri, Aditya Desai, R.N. Kothari, Aarushi Shah, Jeet Patel	
107.	Trends of Fatal Poisoning in Jhalawar Region of Rajasthan: A 5 - Year Retrospective Study	647
	Ramakant Verma, Khushpal Singh Rathore, Bhavesh Bohra	
108.	A-b Ridge Count and Its Correlation with Epilepsy	653
	Rashmi C Goshi	

Analysis of Risk Management Cycle on Capitation Mechanism at First Level Healthcare Facilities in East Java

Rachmad Suhanda^{1,2}, Usman Hadi³, Widodo J. Pudjirahardjo⁴, Djazuly Chalidyanto⁴

¹ Doctoral Students, Faculty of Public Health, Universitas Airlangga, Indonesia, ²Lecturer at the Faculty of Medicine, Universitas Syiah Kuala, Indonesia, ³ Lecturer at Faculty of Medicine, Universitas Airlangga, Department of Internal Medicine, RSUD Dr. Soetomo, Indonesia, ⁴ Lecturer at Faculty of Public Health, Universitas Airlangga, Indonesia

Abstract

The implementation of risk management is an essential function in an organization to control risk. Based on the preliminary research that has been carried out, only 4.2% of FLHF in East Java implemented the complete risk management process, starting from establishing context to evaluating risks. Modifications of risk management cycle implementation are made by reviewing the opinion of Carroll (2001) and ISO 31000:2009. This research aims to analyze the influence between the stages of implementing modified risk management based at 95 FLHF in East Java. The results show that each stage of the capitation mechanism risk management has a strong significant influence on each other. The results provide recommendation that FLHF should improve the understanding and awareness of the risks and expected to be able to apply risk management cycle as a whole.

Keywords: risk management cycle, capitation mechanism, first level healthcare facilities

Introduction

The implementation of risk management is an essential function in an organization to control risk. This function will reduce the negative impact that may arise from various activities carried out by the organization. This function includes carrying out assessments and determining efficient ways to control risk.(1)(2)(3)(4) (5)(6) The implementation of risk management includes two major stages that are connected, namely the risk identification and analysis stage as well as the handling of such risks. Carrol (2001) describes the relationship between these two stages with the risk management process structure. The risk management process

Corresponding author: Prof. Dr. dr. Usman Hadi,

Faculty of Medicine, Department of Internal Medicine, RSUD Dr. Soetomo, Universitas Airlangga, Surabaya, East Java, Indonesia, email: usman.hadi2@fk.unair.ac.id structure explains that failure in risk management can result in decreases control over losses that may occur.(3)

Risk is a possibility of an unwanted loss with a certain severity. The existence of a severity level associated with the risk allows analysis and prevention of the risk. The element of uncertainty in risk requires effective control measures in order to reduce the negative impacts that may occur. Therefore, a management concept is formed to control risk, known as risk management.(2)(7)(8) Based on ISO 31000:2009, the risk management process has five main parts, namely: (a) communication and consultation, (b) establish context; (c) risk assessment (identification, analysis, and evaluation) (d) risk management and (e) monitoring as well as review. Those standards state that every part is a systematic application of management policies, procedures, and evaluation. (6)

First Level Health Facilities (FLHF) according to the Minister of Health Regulation (Permenkes) No. 71 of 2013 (9) consists of community health centre (Puskesmas), primary clinic (pratama clinic), class D hospital, and private dentist practice. FLHF is an institution that acts as the earliest health service provider (PPK). As an organization engaged in the health sector, FLHF certainly cannot avoid risks. Therefore, FLHF urgently needs the implementation of risk management.

Since 2014, Indonesia has implemented the National Health Insurance (JKN) policy. FLHF, in collaboration with Healthcare and Social Security Agency (BPJS Kesehatan), will act as gatekeepers in the program and could face risks that need to be managed. The capitation mechanism is a payment method used in JKN, which is implemented in Indonesia by adjusting a separate policy. Among the various risks faced, it is deemed necessary to analyze health services in the FLHF from the capitation mechanism approach as a payment method in FLHF.

The number of FLHFs working with BPJS Kesehatan throughout Indonesia until July 2020 is more than 27.000 FLHFs. In East Java, the number of FLHF who has worked with BPJS Kesehatan amounted to around 2.400 FLHFs.(10) Total FLHF cooperated with BPJS Kesehatan in East Java reached 11,4% out of the total FLHF in Indonesia. As many as 38.9% consisted of Puskesmas, and the rest consisted of individual practice and Pratama clinics. Based on the preliminary research that has been carried out, out of the total number of FLHFs in East Java, only 4.2% of FLHFs implemented the overall implementation of risk management, starting from establishing the context to evaluating risks.

Taking a statement from ISO 31000:2009(6) regarding the risk management process, this study then considers modifying the risk management process. Modifications are made by reviewing the opinion of Carroll (2001)(3) regarding the two stages in the process of implementing risk management and ISO 31000:2009 regarding the five main parts of risk management which are then adjusted to its application in FLHF. In conclusion, the risk management cycle in FLHF can be simplified into five components: (a) establish context; (b) risk identification; (c) risk analysis; (d) risk treatment, and (e) risk evaluation.

Establish context means determining the risk context of the capitation mechanism through socialization that affects the achievement of organizational goals. Risk identification is the study of potential sources of events and triggers that may cause harm to the organization due to the capitation mechanism. Risk analysis is the implementation of an investigation regarding possible causes, severity, urgency, and seriousness of losses related to the risks of the identified capitation mechanism. Risk treatment is the management of potential events that may result in organization losses due to the capitation mechanism. Risk evaluation can be interpreted as the implementation of activities to assess the management of events that may result in organization losses due to the capitation mechanism.

The modification results considered are to be more representative of the actual risk management cycle. Several essential points that make this modification different from existing concepts are found in the risk analysis and risk evaluation stages. In the risk analysis stage, there is a risk evaluation process in which risk is evaluated, so that the risk evaluation stage does not evaluate risk anymore. The risk evaluation stage is used to evaluate the overall risk management cycle. With the overall evaluation, it can be seen that the stages are still less than optimal in implementation. This research aims to analyze the influence between the stages of implementing modified risk management based on a combination of two statements.(3)(6) Modification of risk management implementation stages has been adjusted by its application in FLHF (Puskesmas and Pratama clinic) in collaboration with BPJS Kesehatan, especially in East Java Province.

Method

This research is an observational study with a cross-sectional approach using primary data from FLHF in East Java Province in collaboration with BPJS Kesehatan. The sampling method used was proportional stratified random sampling. The sample size after proportioning by type of FLHF has obtained as many as 54 community health centres and 28 primary clinics. Large sample proportionally distributed in districts or cities selected based on the number FLHF cooperation with BPJS Kesehatan. The results of the data collection obtained 95 FLHFs.

Research respondents were FLHF leaders and staff who were involved in the management of capitation funds and patient care for JKN participants. The minimum number of respondents for each FLHF was three people (according to FLHF conditions) consisting of 1 (one) leader and 2 (two) staff who are the key persons in the organization. The questionnaire used in data collection is the risk management implementation questionnaire. The assessment is carried out on a scale ranging from 1 (never implementing) to 5 (always implementing). The establish context variable was represented by 17 statements, risk identification variable 23 statements, risk analysis variable 22 statements, and risk treatment and risk evaluation variable 24 statements. In this statement, the

risk elements studied include financial, legal, social and performance aspects (performance, time, physical).

Results and Discussion

The implementation of capitation mechanism risk management in FLHF in this research consisted of five stages, namely establish context, risk identification, risk analysis, risk treatment, and risk evaluation. The results of research related to the implementation of capitation mechanism risk management are shown in Table 1.

No	The implementation of capitation mechanism risk management	n			Annotation			
			Mean	SD	Min	Max	р	
1	Establish Context	95	63,51	13,33	23	85	0,518	Normally distributed
2	Risk Identification	95	94,76	13,29	65	120	0,280	Normally distributed
3	Risk Analysis	95	87,64	14,12	57	115	0,797	Normally distributed
4	Risk Treatment	95	95,88	15,45	57	125	0,585	Normally distributed
5	Risk Evaluation	95	96,06	15,99	52	125	0,586	Normally distributed
	Total score	95	437,85	67,33	297	570	0,630	Normally distributed
	Factor analysis	95	0,001	1,00	-2,11	1,96	0,654	Normally distributed

 Table 1. Distribution of Mean, Standard of Deviation, and Normality Test Score of Capitation Mechanism

 Risk Management Implementation at FLHF in 4 Districts and Cities in East Java 2019

Based on Table 1, it is known that all measurement data are normally distributed. The result of the factor analysis from the total score of all sub-variables is the score that will be used for the frequency analysis of the risk management implementation intensity. The scores obtained are then grouped to make it easier to identify the intensity of risk management implementation in FLHF. Four groups of implementation of risk management intensity were obtained, which are very low, low, high, and very high. The results of the division of the capitation mechanism risk management implementation intensity by FLHF type can be seen in Table 2.

 Table 2. Frequency Distribution of Risk Management Implementation Intensity by Type of FLHF in 4

 Districts and Cities in East Java 2019

No	FLHF Туре	Very Low		Low		High		Very High		Total	
		n	%	n	%	n	%	n	%	n	%
1	Puskesmas	5	7,7%	24	36,9%	19	29,2%	17	26,2%	65	68,4%
2	Pratama clinics	7	23.3%	12	40,0%	6	20,0%	5	16,7%	30	31,6%
	Total	12	12,6%	36	37,9%	25	26,3%	22	23,2%	95	100,0%

The analysis result shows that in general, FLHF has not fully implemented the capitation mechanism risk management correctly. This result can be concluded from the number of FLHF that have implemented risk management with low and very low intensity, namely 48 FLHF (50,5%). This amount is slightly greater than FLHF that have implemented risk management with high and very high intensity (49,5%).

Meanwhile, based on the type of FLHF, risk management implementation with high and very high intensity is proportionally greater in Puskesmas (55,4%) rather than in Pratama clinics (36,7%). Organizational

leaders must be able to become "system thinkers" who demand in-depth analysis of security issues, replace punitive reactions to mistakes to be open and proactive in dealing with any risks. So it is hoped that there will be an opportunity to build a safer health organization.(11)

Furthermore, a continuation statistical analysis was carried out that observed the influence between stages of the capitation mechanism risk management implementation at FLHF in 4 districts and cities in East Java. The results of the statistical analysis can be seen in Table 3.

Tabel 3. Statitical Analysis of The Capitation Mechanism Risk Management Stages at FLHF in East Java2019

No	Independent variables	Dependent variables	В	b	р	R2	Annotation
1	Establish Context	Risk Identification	0,667	0,669	0,001	0,45	Significant
2	Risk Identification	Risk Analysis	0,986	0,928	0,001	0,86	Significant
3	Risk Analysis	Risk Treatment	1,062	0,971	0,001	0,94	Significant
4	Risk Treatment	Risk Evaluation	1,013	0,979	0,001	0,96	Significant
5	Risk Evaluation	Establish Context	0,572	0,687	0,001	0,47	Significant

After the statistical analysis has been done, it was found that all stages of risk management cycle influence each other. This result is consistent with research by Malilay, et al. (2014) on the role of applied epidemiology methods in disaster management cycle. In said research, it is stated that there are three stages of the disaster management cycle, namely predisaster, disaster and post-disaster. The three stages are interconnected in their implementation. If one stage is not appropriately implemented, it will have an impact on the other stages. (12)

Other research by Evans, Dalkir, dan Bidian (2014) also stated the same thing in a different field, namely knowledge management cycle (KMC). KMC consists of 7 phases, namely identify, store, share, use, learn,

improve, and create. The 7 phases are interrelated in their utilization and are grouped into the KMC model.(13) Research by Morita, Flynn, dan Ochiai (2011) on the strategic management cycle, also stated the same thing. The strategic management cycle consists of 4 stages, namely organizational visionary planning, formulating strategy, operations and practice, coordination and storing of practices. The success of the strategic management cycle is a function of how well each stage is carried out. An organization will survive and grow when management cycles are implemented effectively. (14)

Another research by Jorgensen (2007) regarding the sustainable environmental management system (EMS) stated a similar thing. Said research stated that to create an efficient and dynamic system, it takes synergies from several areas. In addition, implementation in each of the six stages of consisting of environmental policy, planning, implementation and operation, checking, management review, and continual improvement is also crucial. Each stage of an EMS is interconnected, so

compatibility between stages is essential.(15)

Based on the statistical analysis of the capitation mechanism risk management implementation cycle, the following results were obtained.



Figure 1. Statistical Analysis Results of Capitation Mechanism Risk Management Implementation Cycle

Figure 1 shows that each stage of the capitation mechanism risk management has a significant influence on each other. The strength of influence between stages of capitation mechanism risk management implementation is strong. This result is in line with research by Sendlhofer, et al. (2014)(16) related to the systematic implementation of clinical risk management in hospitals. The results indicate that clinical risk management plays a dominant role in enabling identification, analysis, and potential risks management. The application of clinical risk management plays can be challenging. Each stage in the clinical risk management application affects the quality of the risk management implementation cycle.

Based on the research, it can be concluded that various conditions will influence the risk management in the subsequent stages of implementation. This stage starts from establishing context to risk evaluating. Errors in the establishing context will affect every next step, up to the risk evaluation stage. The better the FLHF understanding and awareness of the risks faced, the better the implementation of identification, analysis, treatment and evaluation of these risks.

Conclusions and Suggestions

Conclusions

1. As many as 49.5 % of FLHF implemented risk

management with low and very low intensity, more than those with high and very high intensity. FLHF have not fully implemented the capitation mechanism risk management correctly.

2. There is a strong influence at each stage of risk management implementation (establish context, risk identification, risk analysis, risk treatment, and risk evaluation). It is important to note that understanding and awareness of the risks faced will determine how the FLHF manages each risk.

Suggestions

1. The implementation of risk management will improve if the understanding and awareness of the risks faced are well enough so that the enforcement of the risk context is known in the FLHF. It is expected to be able to apply risk management as a whole.

2. Organizational leaders need to monitor each stage of the risk management implementation in detail (starting from establish context, identification, analysis, treatment to evaluation) because they are closely related to one another.

Conflict of Interest : Nil

Ethical Clearance : This research has been proved by Health Research Ethics Committee, Faculty of Nursing Universitas Airlangga **Source of Funding** : Indonesia Endowment Fund for Education (LPDP)

References

- 1. Hubbard DW. The Failure of Risk Management: Why It's Broken and How to Fix It. New Jersey: John Wiley and Sons Inc.; 2009.
- Youngberg B. Principles of Risk Management and Patient Safety. Ontario: Jones & Bartlett Learning; 2011.
- Carroll R. Risk Management Handbook For Health Care Organizations: Third Edition. San Francisco: Jossey-Bass Inc.; 2001.
- 4. Rausand M. Risk Assessment: Theory, Methods, and Applications. New Jersey: John Wiley and Sons Inc.; 2011.
- 5. Culp CL. The Risk Management Process: Business Strategy and Tactics. New Jersey: John Wiley and Sons Inc.; 2001.
- International Standard. ISO 31000: Risk management — Principles and guidelines. First. Geneva: ISO copyright office; 2009.
- 7. Aven T. Quantitative Risk Assessment: The Scientific Platform. Cambridge: Cambridge University Press; 2011.
- 8. Webb A. The Project Manager's Guide to Handling Risk. England: Gower Publisher Ltd; 2013.
- 9. Kemenkes RI. PMK No. 71 Tahun 2013 Tentang Pelayanan Kesehatan pada JKN. 71.

- BPJS Kesehatan. BPJS Kesehatan [Internet]. Badan Penyelenggara Jaminan Sosial Kesehatan. 2019 [cited 2020 Sept 25]. Available from: https:// faskes.bpjs-kesehatan.go.id/aplicares/#/ap
- 11. Alam AY. Steps in the Process of Risk Management in Healthcare. Journal of Epidemiology and Preventive Medicine. 2016;02(02):118–23.
- Malilay J, Heumann M, Perrotta D, Wolkin AF, Schnall AH, Podgornik MN, et al. The Role of Applied Epidemiology Methods in The Disaster Management Cycle. American Journal of Public Health. 2014;104(11):2092–102.
- Evans M, Dalkir K, Bidian C. A Holistic View of The Knowledge Life Cycle: The Knowledge Management Cycle (KMC) Model. Electrical Journal of Knowledge Management. 2014;12(2):85–97.
- Morita M, James Flynn E, Ochiai S. Strategic Management Cycle: The Underlying Process Building Aligned Linkage among Operations Practices. International Journal of Production Economics. 2011;133(2):530–40.
- Jørgensen TH. Towards More Sustainable Management Systems: Through Life Cycle Management and Integration. Journal of Cleaner Production. 2008;16(10):1071–80.
- 16. Sendlhofer G, Brunner G, Tax C, Falzberger G, Smolle J, Leitgeb K, et al. Systematic Implementation of Clinical Risk management in A Large University Hospital: The Impact of Risk Managers. The Central European Journal of Medicine. 2014;127(1–2):1–11.