



## Editorial Team



**Viskasari P. Kalanjati**

**Editor-in-Chief**

Department of Anatomy, Histology, and Pharmacology, Faculty of  
Medicine, Universitas Airlangga, Indonesia

International Federation of Associations of Anatomists (IFAA)

 0000-0002-7005-0025

Google Scholar

54388384000

 5975075



**Kuntaman Kuntaman**

**Editorial Board**

Faculty of Medicine, Universitas Wijaya Kusuma  
Surabaya, Indonesia

0000-0003-4897-8879

Google Scholar

8700386400

 6086035



**Muhammad Miftahussurur**

**Editorial Board**

Division of Gastroentero-Hepatology, Department of Internal  
Medicine, Faculty of Medicine, Universitas Airlangga, Indonesia;

Division of Gastroenterology and Hepatology, Baylor College  
Medicine, US

 0000-0003-1415-6033

Google Scholar

56323903000

 6031037



**Aryati Aryati**

**Editorial Board**

Department of Clinical Pathology, Faculty of Medicine, Universitas  
Airlangga; Indonesian Society of Clinical Pathology and Laboratory  
Medicine, Indonesia

 0000-0001-9055-0177

Google Scholar

56054058600

 6043865



**Irwanto Irwanto**

**Editorial Board**

Department of Child Health, Faculty of Medicine, Universitas Airlangga; Dr. Soetomo General Academic Hospital, Indonesia

 0000-0002-7573-8793

Google Scholar

-

 255759



**Bambang Purwanto**

**Editorial Board**

Department of Physiology and Medical Biochemistry, Faculty of Medicine, Universitas Airlangga, Indonesia

 0000-0002-3786-7068

Google Scholar

57152859900

 5987193



**Andrew Paul Smith**

**Editorial Board**

Centre for Occupational and Health Psychology, School of Psychology, Cardiff University, United Kingdom

 0000-0001-8805-8028

-

35429650000

 -



**Yoshio Yamaoka**

**Editorial Board**

Department of Environmental and Preventive Medicine, Faculty of Medicine, Oita University, Japan

 0000-0002-1222-5819

":

Google Scholar

55183784100

 -



**Franco Servadei**

**Editorial Board**

Department of Neurosurgery, Humanitas University, Italy

 0000-0002-3595-3464



Google Scholar  
7006077203



**Marten J Postma**  
**Editorial Board**

Faculty of Science, Swammerdam Institute for Life Sciences,  
University of Amsterdam, Netherlands

0000-0002-3167-3633  
Google Scholar



**Christianto Lumenta**  
**Editorial Board**

Bogenhausen Academic Teaching Hospital, Technical University,  
Germany

0000-0002-4030-0708



55397690000



**Dirk Jan Marie de Ridder**  
**Editorial Board**

Department of Development and Regeneration, Katholieke  
Universiteit Leuven, Belgium

-

Google Scholar



**Horie Shigeo**  
**Editorial Board**

Department of Urology, Faculty of Medicine, Juntendo University,  
Japan

0000-0002-8612-8368



57222920684



**Yusuke Suzuki**  
**Editorial Board**

Department of Nephrology, Faculty of Medicine, Juntendo University,  
Japan

-



-  
8388474600  
 -

**Arend Frederik Bos**  
**Editorial Board**

Division of Neonatology, Faculty of Medical Sciences, University of Groningen, Netherlands

ID -  
-



36839156800  
 -

**Hiroaki Kimura**  
**Editorial Board**

Department of Physical Medicine and Rehabilitation, Hiroshima University Hospital, Japan

ID 0000-0003-0005-5805  
-



49763362000  
 -

**Jitti Hanprasertpong**  
**Editorial Board**

Division of Gynecologic Oncology, Department of Obstetrics and Gynecology, Faculty of Medicine, Prince of Songkla University, Thailand

ID 0000-0002-0640-6824  
Google Scholar  
21740891300



-

**Anucha Thatrimontrichai**  
**Editorial Board**

Department of Pediatrics, Faculty of Medicine, Prince of Songkla University, Thailand

ID 0000-0003-2363-6362  
Google Scholar  
35088990700



-

**Surasak Sangkhathat**  
**Editorial Board**

Division of Surgery, Faculty of Medicine, Prince of Songkla University, Thailand



 0000-0003-3622-3233

-

6602156542



### **Delvac Oceandy** **Editorial Board**

Department of Medicine and Health, Faculty of Biology, University of Manchester, United Kingdom

 0000-0002-6242-6491

-

6506557120



### **Brahmaputra Marjadi** **Editorial Board**

School of Medicine, Western Sydney University, Australia

 0000-0003-0898-3737

Google Scholar

28267975300



### **Purwo Sri Rejeki** **Editorial Board**

Department of Physiology and Medical Biochemistry, Faculty of Medicine, Universitas Airlangga, Indonesia

 0000-0003-2363-6362

Google Scholar

35088990700



6067303

### **Siti Khaerunnisa** **Editorial Board**

Department of Physiology and Medical Biochemistry, Faculty of Medicine, Universitas Airlangga, Indonesia; American Association for Biochemistry and Molecular Biology, US

 0000-0002-6358-8265

Google Scholar

57205438945




6072589

### **Wihasto Suryaningtyas** **Editorial Board**

Indonesian Neurosurgical Society; Dr. Soetomo General Academic Hospital, Indonesia





 0000-0002-1187-3777

Google Scholar

57216140907

 5987008

**Asra Al Fauzi**

**Editorial Board**

Indonesian Neurosurgical Society; Surabaya Neuroscience Institute (SNeI), Indonesia

 0000-0002-5155-2476

Google Scholar

57215857858

 5984620

**Azimatul Karimah**

**Editorial Board**

Department of Psychiatry, Faculty of Medicine, Universitas Airlangga; Dr. Soetomo General Academic Hospital, Indonesia

 0000-0002-0261-7878

Google Scholar

55640202700

 6058132

**Lucky Prasetiowati**

**Editorial Board**

Department of Anatomy, Histology, and Pharmacology, Faculty of Medicine, Universitas Airlangga; Indonesian Association of Anatomist, Indonesia

 0000-0002-8929-9816

Google Scholar

57192906307

 5995382

**Reny I'tishom**

**Editorial Board**

Department of Biomedical Sciences, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia

 0000-0002-9971-7786

Google Scholar

57209243610

 5984521






**Anggie Kusumawardhani**

**Assistant Editor**

Unit Konsorsium Jurnal dan Folia Medica Indonesiana, Indonesia

 **0000-0002-7998-8589**

Google Scholar



-



**Atika Wardah**

**Assistant Editor**

Unit Konsorsium Jurnal dan Folia Medica Indonesiana, Indonesia

 **0000-0002-8065-4922**

Google Scholar



-

## Login

Username \*

Password \*

[Forgot your password?](#)

Keep me logged in

**Login**

[Register](#)

Focus and Scope

Section Policies

Publication Frequency

Publication Ethics

Article Processing Charge

Peer Reviewers

Peer Review Process

Editorial Process Flowchart

Open Access Statement	Archiving
Reference Manager	Repository Policy
External Link	Plagiarism Check
Copyright Notice	ORCID ID Policy
License Term	Contact
Old Website	

## Meet Our Editorial Team



**Viskasari P. Kalanjati**  
 Editor-in-Chief  
 Universitas Airlangga, Indonesia  
 54388384000



**Muhammad Miftahussurur**  
 Editorial Board  
 Universitas Airlangga, Indonesia  
 56323903000

[Read More](#)

## Instruction For Author

How to Submit	Guidelines for Authors
Online Submission	Document Template
Patient Consent for Publication	Authors' Declaration & CTA

**ISSN**





Printed ISSN

Electronic ISSN

## Journal Standing



## Reference Manager



## Plagiarism Checker



## External Links



## In Collaboration with



## Accreditation Certificate

# SERTIFIKAT

Direktorat Jenderal Pendidikan Tinggi, Riset dan Teknologi  
Kementerian Pendidikan, Kebudayaan, Riset dan Teknologi Republik Indonesia



Kutipan dari Keputusan Direktorat Jenderal Pendidikan Tinggi, Riset dan Teknologi  
Kementerian Pendidikan, Kebudayaan, Riset, dan Teknologi Republik Indonesia

Nomor IS8/E/KPT/2021  
Peringkat Akreditasi Jurnal Ilmiah Periode 1 Tahun 2021

Nama Jurnal Ilmiah  
**Folia Medica Indonesiana**  
E-ISSN: 2599056X

Penerbit: Fakultas Kedokteran Universitas Airlangga

Ditetapkan Sebagai Jurnal Ilmiah

### TERAKREDITASI PERINGKAT 2

Akreditasi Berlaku selama 5 (lima) Tahun, yaitu  
Volume 56 Nomor 4 Tahun 2020 Sampai Volume 61 Nomor 3 Tahun 2025

Jakarta, 09 December 2021  
Plt. Direktur Jenderal Pendidikan Tinggi,  
Riset, dan Teknologi



Prof. Ir. Nizam, M.Sc., DIC, Ph.D., IPU, ASEAN Eng  
NIP. 19610706198710101



## Visitors



[View My Stats](#)

## Indexed By





# publons

---

## Information

---

[For Readers](#)

[For Authors](#)

[For Librarians](#)

---

## Keywords



Diabetes  
diabetes mellitus  
BMI  
motility  
disease  
LDL  
health risk  
obesity  
pediatric  
MDA  
NT-proBNP  
Escherichia coli  
virus  
CRP  
human & health  
age  
ESBL  
children  
mice  
quality of life

---

## Current Issue

RTOM 1.0

RSS 2.0

RSS 1.0

---

## Address

Folia Medica Indonesiana  
Unit of Journal Consortium and Folia Medica Indonesiana,  
Faculty of Medicine, Universitas Airlangga  
Jl. Prof. Dr. Moestopo 47, Surabaya, 60131, Indonesia

## Contact Info:

Email: [fmi@journal.unair.ac.id](mailto:fmi@journal.unair.ac.id)



Lembaga Inovasi, Pengembangan Jurnal,  
Penerbitan dan Hak Kekayaan Intelektual

**LIPJPHKI**

Gedung AUP, Kampus C, Universitas Airlangga, Kota Surabaya, Jawa Timur, 60115



This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike  
4.0 International License.

## PROCALCITONIN AS BIOMARKER OF ANTIBIOTIC EFFECTIVITY IN COMMUNITY-ACQUIRED PNEUMONIA

Ana Khusnul Faizah<sup>1</sup>, Winariani<sup>2</sup>, Yulistiani<sup>3</sup>, Samirah<sup>3</sup>

<sup>1</sup>Post graduate Student of Master of Clinical Pharmacy Program, Faculty of Pharmacy, Airlangga University

<sup>2</sup>Pulmonology Department Dr. Soetomo Teaching Hospital Surabaya

<sup>3</sup>Clinical Pharmacy Department, Faculty of Pharmacy, Airlangga University

### ABSTRAK

*Procalcitonin merupakan marker spesifik infeksi akut yang ditandai peningkatan kadar procalcitonin sehingga procalcitonin digunakan sebagai marker efektifitas terapi antibiotika. Penelitian ini bertujuan mengkaji procalcitonin sebagai marker efektifitas terapi antibiotika pada pasien pneumonia komunitas. Penelitian ini merupakan penelitian kohort prospektif pada pasien pneumonia komunitas di Instalasi Rawat Inap SMF Pulmonologi dan Ilmu Kedokteran Respirasi RSUD Dr. Soetomo Surabaya. Procalcitonin dan parameter SIRS sebelum, hari ke-3 dan hari ke-5 terapi antibiotika dianalisis secara deskriptif. Didapatkan 19 pasien pneumonia komunitas dengan penyakit penyerta TB paru, sepsis, pneumothorax dan efusi pleura. Procalcitonin pada pasien pneumonia komunitas sebelum terapi antibiotika adalah  $0,51 \pm 0,19$  (0,28-0,89) ng/ml, sesudah terapi antibiotika hari ke-3  $0,53 \pm 0,24$  (0,08-0,93) ng/ml dan pada hari ke-5 mencapai kadar  $0,43 \pm 0,24$  (0,00-0,83) ng/ml. Procalcitonin pada pasien pneumonia komunitas dengan sepsis adalah  $0,48 \pm 0,24$  (0,29-0,89) ng/ml;  $0,40 \pm 0,26$  (0,08-0,74) dan  $0,22 \pm 0,13$  (0,00-0,32) ng/ml. Procalcitonin pada pasien pneumonia komunitas dengan TB paru adalah  $0,52 \pm 0,18$  (0,38-0,83) ng/ml;  $0,55 \pm 0,25$  (0,23-0,89) ng/ml dan  $0,57 \pm 0,16$  (0,41-0,82) ng/ml. Procalcitonin pasien pneumonia komunitas dengan TB paru dan sepsis sebelum terapi antibiotika, hari ke-3 dan hari ke-5 terapi antibiotika adalah  $0,55 \pm 0,12$  (0,38-0,70) ng/ml,  $0,56 \pm 0,15$  (0,42-0,73) ng/ml dan  $0,47 \pm 0,15$  (0,33-0,64) ng/ml. Terdapat 60% pasien pneumonia komunitas dengan sepsis, 20% pasien pneumonia komunitas dengan efusi pleura dan tidak ada pasien pneumonia komunitas dengan TB Paru yang mencapai kadar procalcitonin  $< 0,26$  ng/ml pada hari ke-5 sesudah terapi antibiotika. Procalcitonin tidak dapat digunakan sebagai marker efektifitas terapi antibiotika pada pasien pneumonia komunitas. (FMI 2015;51:96-100)*

**Kata kunci:** procalcitonin, pneumonia komunitas, antibiotika, biomarker

### ABSTRACT

*Major therapy for community-acquired pneumonia (CAP) is antibiotics. Procalcitonin used as the measure of antibiotics effectiveness. Procalcitonin is specific biomarker that elevates in acute bacterial infection. This study aimed to analyze procalcitonin as effective antibiotics biomarker in CAP patients hospitalized at Pulmonology Department Dr. Soetomo Teaching Hospital. This study was performed by cohort prospective data and analyzed by descriptively. Data were assessed at before antibiotics, day 3 and day 5 after antibiotics therapy. Nineteen community-acquired pneumonia were enrolled. Serum procalcitonin before antibiotic therapy, day 3 and day 5 antibiotics therapy in CAP are  $0,51 \pm 0,19$  (0,28-0,89) ng/ml,  $0,53 \pm 0,24$  (0,08-0,93) ng/ml and  $0,43 \pm 0,24$  (0,00-0,83) ng/ml. Serum procalcitonin in CAP with sepsis patients are  $0,51 \pm 0,19$  (0,28-0,89) ng/ml (before antibiotic therapy),  $0,40 \pm 0,26$  (0,08-0,74) ng/ml (day 3) and  $0,22 \pm 0,13$  (0,00-0,32) ng/ml (day 5). Serum Procalcitonin in CAP with pulmonary TB are  $0,52 \pm 0,18$  (0,38-0,83) ng/ml,  $0,55 \pm 0,25$  (0,23-0,89) ng/ml dan  $0,57 \pm 0,16$  (0,41-0,82) ng/ml. Serum procalcitonin in CAP with sepsis and pulmonary TB are  $0,55 \pm 0,12$  (0,38-0,70) ng/ml,  $0,56 \pm 0,15$  (0,42-0,73) ng/ml dan  $0,47 \pm 0,15$  (0,33-0,64) ng/ml. Procalcitonin of 60% CAP with sepsis patients, 20% CAP with pleural effusion patients and no patients of CAP with pulmonary TB reach  $< 0,26$  ng/ml. In conclusion, procalcitonin is not useful biomarker in CAP. (FMI 2015;51:96-100)*

**Keywords:** procalcitonin, community-acquired pneumonia, antibiotic therapy, biomarker

**Correspondence:** Ana Khusnul Faizah, Royal Ketintang Regency E-02 Ketintang Madya Surabaya, +6285655458944, goodmorningana@yahoo.com

### INTRODUCTION

Community-acquired pneumonia is an infection of the pulmonary parenchyma that develops outside the hospital (Marrie 2008). The clinical manifestations are frequently febrile with a tachycardic response and may have chills or sweats and cough that are either

nonproductive or productive of mucoid, purulent, or blood-tinged sputum (Wunderink 2010). Many physiological changes occur in the host in order to restore the impaired homeostasis during bacterial and viral. The release of inflammatory mediators, such as interleukin (IL) 1 and tumor necrosis factor (TNF), results in fever. Chemokines, such as IL-8 and

granulocyte colony-stimulating factor, stimulate the release of neutrophils and their attraction to the lung, producing both peripheral leukocytosis and increased purulent secretions (Wunderink 2010).

White blood cell (leukocyte) count and other parameter SIRS are the most frequently used markers of acute phase responses in clinical practice. Classical diagnostic instruments including leukocyte count, chest x ray and sputum culture do not have sufficient specificity in differentiating between bacterial infections, non-infectious systemic inflammations or viral infections. Therefore, more specific and reliable markers that might be helpful in deciding the treatment are needed in these community-acquired pneumonia patients.

Procalcitonin, a protein of 116 amino-acids with molecular weight of 13 kDa, is prohormone of calcitonin produced by C-cells of thyroid gland and intracellularly cleaved by proteolytic enzymes into the active hormone. The production of procalcitonin during inflammation is linked with a bacterial endotoxin and with inflammatory cytokines (TNF, IL-6) (Meissner 2000). The aim of this study was to analyze procalcitonin as effective biomarker of antibiotics community-acquired pneumonia patients who hospitalized at the Pulmonology Department Dr. Soetomo Teaching Hospital.

**MATERIALS AND METHODS**

A cohort prospective study was conducted at Pulmonology Department Dr. Soetomo Teaching Hospital during September to December 2014. Patient selection based on inclusion and exclusion criteria. Inclusion criteria (1) patient diagnose with community-acquired pneumonia aged 18-75 years old; (2) patient or their family agreed to follow the study. Patient hospital-acquired pneumonia, thyroid carcinoma, pulmonary carcinoma, received corticosteroids or antibiotic therapies before come to Emergency Department were excluded.

Blood samples were collected from all patients before antibiotic therapy in order to perform complete blood count and routine biochemistry, as well as to analyze serum procalcitonin. Procalcitonin levels were reassessed after 3 and 5 d. culturing of the sputum specimens that were obtained from patients with sputum production were performed. The blood collected for serum PCT measurement was centrifuged and kept at -20°C until the time of the measurement. PCT measurement was performed by the Bio-Rad iMark device. In this study, we used Biotech Human PCT ELISA Kit made in the China. Descriptive analyses were performed

to determine the demographic patients, profile of procalcitonin and parameter SIRS at before and after antibiotic therapy. Correlation between procalcitonin and parameter SIRS (pulse, respiratory rate, body temperature and leukocyte) was measured by the Spearman’s correlation test and Pearson’s correlation test. A probability value of 0.05 was regarded as significant.

**RESULTS**

During the study period, 24 consecutive patients with CAP were screened for eligibility. Of these, 19 were eligible. Baseline characteristic on admission are shown in Table 1. The most age of the patient was 19-59 years and included 9 females and 10 males. The classic triad of dyspnea, cough, and sputum, as reported by the patients, was present in 90%, 90% and 58% of cases. The most coexisting illnesses in patients are pulmonary TB (63%) and sepsis (57%).

Eleven patients had purulent sputum. In 8 of those with purulent sputum, a causative microorganism was identified in 6 patients. Bacterial growth was shown in sputum culture. The results were as follows: Diplococcus in 4 patient, *Acinobacter baumannii* in 1 patients and *Aeromonas hydrophila* in 1 patient. On the day of admission, serum procalcitonin levels were 0.26 to 0.5 ng/mL in 12 patients and >0.5 ng/mL in 7 patients. The results of PCT before and after antibiotic therapy are shown in Figure 1. Serum PCT levels in all patients were >0.25 ng/ml. Results of PCT after 3 days and 5 days of antibiotic therapy shows that the number of patients seemed to have decreased from >0.25 at diagnosis to <0.1 ng/mL (2 patients).

There was no statistically significant correlation between procalcitonin and parameter SIRS (leukocyte, pulse, respiratory rate and the body temperature) in all community-acquired pneumonia patients coexisting illnesses group.

Table 1. Characteristics of CAP patients

	Characteristic (n=19)	n (%)
Sex	Female	9 (47)
	Male	10 (53)
Age	13 – 18 years	1 (5)
	19 – 59 years	12 (63)
	>59 years	6 (32)
	<0.1	0 (0)
Procalcitonin (ng/ml)	0.1 – 0.25	0 (0)
	0.26 – 0.5	12 (63)
	>0.5	7 (37)
	4 – 10	3 (16)
Leukocyte counts	>10	16 (84)
	Pulmonary TB	12 (63)
Coexisting illnesses*	Sepsis	11 (57)

	Hypertension	2 (11)
	Diabetes mellitus	2 (11)
	Pleural effusion	2 (11)
	Fluidopneumothorax	2 (11)
	Pyopneumothorax	1 (5)
	Heart failure	1 (5)
Clinical manifestations	Dyspnea	17 (90)
	Cough	17 (90)
	Sputum	11 (58)
Antibiotics	Ceftazidime	11 (58)
	Levofloxacin	11 (58)
	Anti TB	10 (53)
	Ceftriaxone	4 (21)

\* One patient may have one or more coexisting illnesses  
As expected, high levels of WBC, pulse, respiratory rate and temperature were decrease after 3 and 5 days antibiotic therapy (Figure 2 and 3).




Figure 1. Serum procalcitonin before and after antibiotic therapy

Table 2. Correlation in CAP with PTB

Parameter	Procalcitonin			
	Day 3		Day 5	
	p	r	p	r
Pulse	1.000	0.000	0.946	0.042
RR	0.004	0.979	0.511	-0.395
Temp	0.586	-0.331	0.627	0.297
Leukocyte	0.121	-0.779	0.588	0.330




Figure 2. Pulse and RR before and after antibiotic therapy




Figure 3. Temperature and WBC before and after antibiotic therapy

Table 3. Correlation in CAP with sepsis

Parameter	Procalcitonin			
	Day 3		Day 5	
	p	r	p	r
Pulse	0.926	0.058	0.355	0.533
RR	0.638	-0.289	0.105	0.799
Temp	0.880	-0.94	0.810	-0.150
Leukocyte	0.320	-0.565	0.405	-0.487

Table 4. Correlation in CAP with PTB and sepsis

Parameter	Procalcitonin			
	Day 3		Day 5	
	p	R	p	r
Pulse	0.199	0.689	0.741	0.205
RR	0.149	0.744	0.347	-0.541
Temp	0.870	0.102	0.391	-0.500
Leukocyte	0.332	-0.555	0.180	0.709

## DISCUSSION

PCT is a protein having a molecular weight of 13 kD and it consists of 116 amino acid residues (Meissner 2002). In healthy humans, its normal serum level is 0.1 ng/ml (Carrol et al 2002). In a previous study, administration of bacterial endotoxin to healthy individuals resulted in an increase in PCT levels starting two hours after administration, with a peak value

Table 5. Procalcitonin before and after antibiotics therapy

PCT (ng/ml)	Criteria*	Patients		
		Before Antibiotics Therapy	After Antibiotics Therapy	
		Day 0	Day 3	Day 5
		n (%)	n (%)	n (%)
< 0.1	Bacterial infection very unlikely	0 (0)	0 (0)	2 (10)
0.1 – 0.25	Bacterial infection unlikely	0 (0)	1 (6)	0 (0)
0.26 – 0.5	Bacterial infection likely	12 (63)	9 (47)	10 (59)
> 0.5	Bacterial infection very likely	7 (37)	9 (47)	7 (41)
	Total	19 (100)	19 (100)	19 (100)

\*Based on algorithm for procalcitonin-guided antibiotic therapy (Crain et al 2010, Albrich et al 2012)

reached in 12 hours (Linscheid et al 2004). Consequently, the serum level remains constant for another 12 hours and decreases back to normal level in 20-24 hours. PCT gives rapid response to bacterial infections

In this study, we have demonstrated that serum PCT levels increase in day 3 antibiotic therapies and decrease in day 5 antibiotic therapy. Serum procalcitonin levels remain high after 5 days antibiotic therapy because of pulmonary TB. Studies have shown that after 2 months of intensive phase of treatment, the number of pulmonary TB patients with an increase in PCT was more than those with decrease or no change in the procalcitonin levels. Although a clearly conclusive statement on the prognostic utility of PCT cannot be made at the end of 2 months of intensive treatment, the procalcitonin seems to be a good prognostic marker for the efficacy of the 6 months treatment, because at the end of 6 months, PCT in all patients had decreased (Rohini et al 2013). A similar result was found in another study that PCT level reduced significantly ( $P < 0.001$ ) after 6 months treatment anti-TB (Ghobadi et al 2014). PCT levels increase with the increasing severity of the inflammatory response to infection and may help in assessing the severity of infection, the prognosis of disease, and the response to therapeutic measures. In this study, procalcitonin is not useful biomarker biomarker of antibiotic effectivity in community-acquired pneumonia with pulmonary TB.

Serum procalcitonin levels decrease until  $< 0.26$  ng/mL in community-acquired pneumonia with sepsis. Procalcitonin decrease as parameter SIRS to response antibiotic therapy. Serum PCT levels are elevated in patients with sepsis. A decrease in PCT concentrations was associated with a favorable outcome in patients with sepsis. Based on this data, procalcitonin is useful biomarker of antibiotic effectivity in community-acquired pneumonia with sepsis.

Studies have shown that the diagnostic efficacy of leukocyte count is not high; it has either moderate or low efficacy. In this study, we also were unable to find

(Müller et al 2007, Kibe et al 2011). Following the administration of bacterial endotoxin, PCT levels increase faster and return to normal range more rapidly compared to the levels of CRP (Linscheid et al 2004). any significant relationship between PCT and leukocyte count. This suggests that WBC count is not a significant factor in identifying a bacterial infection. In another study, it was shown that PCT has no significant correlation with leukocyte (TA ÇI 2008, Ugajin et al 2011).

Our study has several limitations. First, since outpatients were excluded, the enrolled patients might have had relatively severe conditions. As a result, serum PCT was elevated compared with previous studies. Secondly, community-acquired pneumonia patients could be associated with bacterial coinfection (pulmonary TB and sepsis). Third limitations of this study are the low sample size (because of lesser prevalence of community-acquired pneumonia that meets inclusion). A further study with a larger sample size and no coexisting illness is suggested.

## CONCLUSION

The results of procalcitonin in 19 patients of community-acquired pneumonia with coexisting illness suggest that procalcitonin is not useful biomarker in community-acquired pneumonia.

## REFERENCES

Albrich WC, Dusemund F, Bucher B, Meyer S, Thomann R, Kühn F, Bassetti S, Sprenger M, Bachli E, Sigrist T, Schwietert M, Amin D, Hausfater P, Carre E, Gaillat J, Schuetz P, Regez K, Bossart R, Schild U, Mueller B; ProREAL Study Team (2012). Effectiveness and safety of procalcitonin-guided antibiotic therapy in lower respiratory tract infections in "real life": an international, multicenter poststudy survey (ProREAL). Arch Intern Med 172, 715-722



- Carrol ED, Thomson APJ, Hart CA (2002). Procalcitonin as a marker of sepsis. *International Journal of Antimicrobial Agents* 20, 1-9
- Christ-Crain M, Stolz D, Bingisser R, Müller C, Miedinger D, Huber PR, Zimmerli W, Harbarth S, Tamm M, Müller B (2006). Procalcitonin guidance of antibiotic therapy in community-acquired pneumonia. *Am J Respir Crit Care Med* 174, 84-93
- Ghobadi H, Lari SM, Amani F, Habibzadeh S, Karimi A, Pourfarzi F (2014). The impact of treatment on serum level of procalcitonin in patients with active pulmonary tuberculosis. *Journal of Cardio-Thoracic Medicine* 2, 238-242
- Kibe S, Adams K, Barlow G (2011). Diagnostic and prognostic biomarkers of sepsis in critical care. *J Antimicrob Chemother* 66, ii33-ii40
- Linscheid P, Seboek D, Schaer DJ, Zulewski H, Keller U, Müller B (2004). Expression and secretion of procalcitonin and calcitonin gene-related peptide by adherent monocytes and by macrophage-activated adipocytes. *Crit Care Med* 32, 1715-1721
- Marrie T (2008). Acute bronchitis and community-acquired pneumonia. In: Fishman AP, Elias JA, Fishman JA, Grippi MA, Senior RM, Pack AI (eds). *Fishman's Pulmonary Diseases and Disorders*, 4th ed, New York, McGraw-Hill Education/Medical, p 2097
- Meisner M (2002). Pathobiochemistry and clinical use of procalcitonin. *Clin Chim Acta* 323, 17-29
- Meisner M (2000). Procalcitonin (PCT): A New, Innovative Infection Parameter Biochemical and Vlinical Aspects, 3 revised and expanded edition, Stuttgart, Georg Thieme Verlag, p 529-539
- Müller B, Harbarth S, Stolz D, Bingisser R, Mueller C, Leuppi J, Nusbaumer C, Tamm M, Christ-Crain M (2007). Diagnostic and prognostic accuracy of clinical and laboratory parameters in community-acquired pneumonia. *BMC Infect Dis* 2, 7-10
- Rohini K, Bhat S, Srikumar PS, Kumar AM (2013). Diagnostic and prognostic value of procalcitonin in tuberculosis patients. *British Journal of Medicine & Medical Research* 3, 2189-2196
- TA ÇI C (2008). The importance of serum procalcitonin levels in patients with chronic obstructive pulmonary disease exacerbations. *Turkish Journal of Medical Sciences* 38, 139-144
- Ugajin M, Miwa S, Shirai M, Ohba H, Eifuku T, Nakamura H, Suda T, Hayakawa H, Chida K (2011). Usefulness of serum procalcitonin levels in pulmonary tuberculosis. *Eur Respir J* 37, 371-375
- Wunderink R (2010). Pneumonia. In: Loscalzo J (ed). *Harrison's Pulmonary and Critical Care Medicine*, 1st ed, New York, McGraw-Hill Education, p 99



FOLIA MEDICA INDONESIA

UNIVERSITAS AIRLANGGA

P-ISSN : 23558393 <> E-ISSN : 2599056X Subject Area : Education

**0.6**  
Impact Factor

**1121**  
Google Citations

**Sinta 2**  
Current Accreditation

[Google Scholar](#) [Garuda](#) [Website](#) [Editor URL](#)

History Accreditation

2018 2020 2021 2022 2023 2024 2025

Garuda Google Scholar

Longer Lag Time in Early-Stage Retinoblastoma

Faculty of Medicine, Universitas Airlangga [Folia Medica Indonesiana Vol. 58 No. 2 \(2022\): June 103-107](#)

2022 DOI: 10.20473/fmi.v58i2.24975 Accred : Sinta 2

The AKT Pathway and Satellite Cell Activation in Skeletal Muscle Mass Regulation

Faculty of Medicine, Universitas Airlangga [Folia Medica Indonesiana Vol. 58 No. 1 \(2022\): March 68-73](#)

2022 DOI: 10.20473/fmi.v58i1.13354 Accred : Sinta 2

Viral Load And Cd4+ among Hiv/Aids Patients Receiving Antiretroviral Therapy In Jayawijaya District, Papua Province, Indonesia

Faculty of Medicine, Universitas Airlangga [Folia Medica Indonesiana Vol. 58 No. 1 \(2022\): March 10-14](#)

2022 DOI: 10.20473/fmi.v58i1.18247 Accred : Sinta 2

Back Matter Vol.58 No.1 March 2022


Faculty of Medicine, Universitas Airlangga [Folia Medica Indonesiana Vol. 58 No. 1 \(2022\): March](#)

2022 DOI: 10.20473/fmi.v58i1.34145 Accred : Sinta 2

Acute Diarrhea Patients among Children Under Five Hospitalized in a Tertiary Hospital in East Java, Indonesia

Faculty of Medicine, Universitas Airlangga [Folia Medica Indonesiana Vol. 58 No. 1 \(2022\): March 33-38](#)

2022 DOI: 10.20473/fmi.v58i1.31423 Accred : Sinta 2



Journal By Google Scholar

	All	Since 2018
Citation	1121	820
h-index	14	12
i10-index	31	19

### [Radiological Aspects of HR-CT Scan on Temporal Bone](#)

Faculty of Medicine, Universitas Airlangga [Folia Medica Indonesiana Vol. 58 No. 1 \(2022\): March 88-102](#)

2022 [DOI: 10.20473/fmi.v58i1.23836](#) [Accred : Sinta 2](#)

### [Profile of Skin Biopsy Patients in Dr. Soetomo General Academic Hospital, Surabaya, Indonesia](#)

Faculty of Medicine, Universitas Airlangga [Folia Medica Indonesiana Vol. 58 No. 1 \(2022\): March 1-9](#)

2022 [DOI: 10.20473/fmi.v58i1.16811](#) [Accred : Sinta 2](#)

### [Front Matter Vol.58 No.1 March 2022](#)

Faculty of Medicine, Universitas Airlangga [Folia Medica Indonesiana Vol. 58 No. 1 \(2022\): March](#)

2022 [DOI: 10.20473/fmi.v58i1.34148](#) [Accred : Sinta 2](#)

### [Prostate-Specific Antigen and Time to PSA Nadir as Prognostic Significance in Castration-Resistant Prostate Cancer](#)

Faculty of Medicine, Universitas Airlangga [Folia Medica Indonesiana Vol. 58 No. 1 \(2022\): March 50-55](#)

2022 [DOI: 10.20473/fmi.v58i1.32649](#) [Accred : Sinta 2](#)

### [Pre-Operative Hormonal Administration in Hypospadias Patients Undergoing Urethroplasty](#)

Faculty of Medicine, Universitas Airlangga [Folia Medica Indonesiana Vol. 58 No. 1 \(2022\): March 80-87](#)

2022 [DOI: 10.20473/fmi.v58i1.32182](#) [Accred : Sinta 2](#)

[View more ...](#)

Get More with  
SINTA Insight

[Go to Insight](#)

#### Citation Per Year By Google Scholar

#### Journal By Google Scholar

	All	Since 2018
Citation	1121	820
h-index	14	12
i10-index	31	19