

Ghrelin and Serotonin as Indicators of Obesity due to The Influence of Circadia on Wistar Rats

by Anis Irmawati

Submission date: 02-Apr-2023 11:46AM (UTC+0800)

Submission ID: 2053194793

File name: Jurnal_1.pdf (203.48K)

Word count: 3630

Character count: 18253

ORIGINAL ARTICLE

Ghrelin and Serotonin as Indicators of Obesity due to The Influence of Circadia on Wistar Rats

*Retno Indrawati¹, Anis Irmawati¹, Indeswati Diyatri¹, Ninuk Hariyani², Mukhammad Fauzi³

1

¹ Department of Oral Biology, Faculty of Dental Medicine, Universitas Airlangga, Surabaya 60132, Indonesia

² Department of Public Health, Faculty of Dental Medicine, Universitas Airlangga, Surabaya 60132, Indonesia

³ Undergraduate Student, Faculty of Dental Medicine, Universitas Airlangga, Surabaya 60132, Indonesia

ABSTRACT

Introduction: One of the main factors supporting obesity is the disruption in the performance of the circadian rhythm which results in a decrease in the quality and quantity of sleep, which triggers stress which is regulated by the hormone serotonin in the body. This increase in serotonin is related to factors that trigger the risk of obesity, besides that melatonin and leptin in the body can decrease, thereby increasing ghrelin which stimulates appetite. The aim of this study was to prove that ghrelin and serotonin can be used as indicators of obesity in Wistar rats as experimental animals. **Materials and Methods:** This study involved 3 groups with each group consisting of 6 samples. Group 1 was the normal group (12 hours light, 12 hours dark), group 2 was the dark group (24 hours dark) and group 3 was the light group (24 hours light). Each group was treated with circadian and modified feed until obesity was found and then the blood measured using the Rat ELISA Kit. **Results:** The results of the correlation test of body weight of Wistar rats with ghrelin and serotonin show that there is a strong relationship between body weight and ghrelin with $p\text{-value} = 0.006$ ($p < 0.05$), $r = 0.609$. The correlation between body weight and serotonin was moderate with $p\text{-value} = 0.023$ ($p < 0.05$), $r = 0.517$. **Conclusion:** Ghrelin and Serotonin can be used as indicators of obesity in Wistar rats.

Keywords: Obesity, Circadian rhythm, Ghrelin, Serotonin

3

Corresponding Author:

Retno Indrawati, DDS, MDS

Email: retnoindrawati@fkg.unair.ac.id

Tel: +62315030255

INTRODUCTION

Obesity is a pathological condition in which excess fat is deposited in subcutaneous tissue, as well as a low-level chronic inflammatory condition in white adipose tissue. Obesity is classified as a chronic disease due to excessive food consumption, according to WHO (1). Obesity cases in the world increase threefold between 1975 and 2016. In 2016, more than 650 million people were obese (1). The percentage of obesity is about 13% of the world's adult population, which is 11% in men and 15% in women. Indonesians who are obese are not less than 12% and are expected to continue to increase every year (2). In addition, people who are obese have a 7 to 11 times higher risk of diabetes. In addition, if the person has diabetes, the risk of heart disease increases up to 8-fold. Obesity is a very serious health problem, not only for individuals but has become a global health problem.

The disruption of circadian rhythms, which act as

regulators of biological functions and behaviors, is one of the risk factors that contribute to obesity (3). This messed-up circadian rhythm causes a drop in sleep quality and quantity, triggering the onset of stress, which is controlled by the hormone serotonin in the body (4). The increase in serotonin is linked to obesity risk factors, on the other hand, melatonin and leptin levels in the body can drop, causing ghrelin levels to rise, stimulating appetite (5–7).

Current technological developments have caused 62% of the earth's population to be exposed to artificial light at night (8). Increased exposure to light at night is known to lead to increased metabolic disorders, impaired reproductive function and neuroendocrine (8). Changes in the duration of light or treatment of certain photoperiods are also a triggering factor for stress response (9).

10

This study was conducted to determine the effect of ghrelin and serotonin levels on obesity in experimental animals treated with photoperiod, so as to determine the relationship between obesity caused by changes in circadian rhythm, which is known through the influence of ghrelin and serotonin levels.

MATERIALS AND METHODS

Samples

This experimental laboratory research adopted true experimental design with (post test control group design). The study was carried out at The Institute Tropical Disease (ITD) Universitas Airlangga and has received ethical approval from the Dental Committee of Airlangga University, Indonesia (No. 320/HRECC.FODM/VII/2020). Federer's formula was used to calculate the sample size. About 18 Wistar rats were randomly sampled and divided into three groups: a normal control group that received 12 hours of light, a dark treatment group that received 0 hours of light, and a light treatment group that received 24 hours of light. The inclusion criteria of the samples were 8-week-old male wistar rat, rat weighing 140–200 grams, with no anatomical defects, and active movement.

Preparation of the light exposure to animal model

All sample groups, namely the control group and the treatment group, were subjected to circadian treatment and dietary treatment. The circadian treatment was carried out with each group using a bright period placed in a lighting room that had a fluorescent lamp along with a timer switch for setting light duration in each group, namely the normal control group with 12 hours of light (12T) and the light treatment group with 24 hours of light (24T). The distance between the floor of the lighting room and the light source is ± 75 cm, so that the light received by the test animals ranges from 100-110 lux. Circadian treatment in the dark treatment group was a group with 0 hours of light or no light at all in the room. Each group will be given treatment for 8 weeks.

Diet treatment was carried out in the control group and treatment group by way of rats induced into obesity by increasing fat intake by 40-60% of total calories, feed, and drink given to rats ad libitum. Modified foods containing 65-70 percent carbohydrates, 20-25 percent protein, and 5-12 percent fat were given to rats. For 8 weeks, Wistar rats are given a drink containing 10-15% fructose solution in drinking water, which causes obesity (Wulansari, 2018). During the study, an analytical balance sheet was used to measure animal weight every 7 days with a precision of 0.1 grams.

Measurement of ghrelin and serotonin levels

After the circadian treatment and diet treatment, the samples' ghrelin and serotonin levels were measured by taking the rats serum, done by taking blood through the heart (intracardiac) as much as at least 3 mL, then accommodated into the vacutainer. The blood is then centrifuged at a speed of 3000 rpm for 10 minutes. Serum results from blood collected, then tested using Rat ELISA kit (Elab Science, USA) in accordance with factory rules.

Statistical analysis

Data analysis was tested by data normality test using Saphiro Wilk test, correlation test using Spearman Rho Correlation and different test using Repeated Measured Anova and Mann Whitney U test.

RESULTS

This study used experimental methods of in vivo laboratory with true experimental design with posttest control design to observe differences in initial data and results of control group experiments and treatment groups. This study was conducted with the aim of knowing that ghrelin and serotonin can be used as indicators of obesity in Wistar rat using experiments that have been divided into 3 circadian treatment groups, namely normal, dark and light groups with each group consisting of 6 sample numbers.

Before the test was conducted between groups, each group conducted a different test each group using statistical test Repeated Measured Anova. The results of the Normal Anova Repeated Measured test group showed no meaningful difference in Wistar rat weight from week 1 to week 9, due to overall $p > 0.05$ values.

Based on Figure, the results of the Repeated Measured Anova dark group test showed in week 2, there was weight loss but not significant ($p > 0.05$). Significant weight loss ($p < 0.05$) occurred in the 3rd week of 4 grams. This meant that during the 3 weeks the rat's weight was reduced by 4 grams. Week 4 of the rat's weight still experienced a significant decrease ($p < 0.05$) of 4.33 grams. This meant that during the 4 weeks there was a significant weight loss of 4.33 grams. Week 5 and the 6th weight of the rat began to increase although insignificant ($p > 0.05$). The 7th week later, there was a significant increase in weight ($p < 0.05$) of 33.67 grams of initial weight. In the 9th week the body weight of rat still experienced a significant decrease ($p < 0.05$) of 4.00 grams. This means that there is a significant weight loss of 4.00 grams judging by the weight of the 7th week against the 9th week.




Fig. 1: The average of body weight over 8 weeks.

Repeated Measured Anova test results of the bright group showed from week 1 to week 8 there was a decrease in weight but not significant ($p > 0.05$), whereas if observations were observed from week 8 experienced significant weight loss on the 9th day ($p < 0.05$), amounting to 2,857 grams.

After conducting different tests each group then conducted tests between groups using statistics test Mann Whitney U test. The result was that there was a meaningful difference in Wistar rat weight between normal and dark group ($p < 0.05$). In the normal and light group ($p > 0.05$) it was found that there was no meaningful difference in Wistar rat weight. In the dark and light group ($p < 0.05$) there was a meaningful difference in the weight of Wistar rat (Table I).

Table I: The significant levels of body weight on circadian rhythm

Type of Data	Normal-Dark- (p-value)	Nomal-Bright (p-value)	Dark-Bright (p-value)
Weight	0,004	0,181	0,003

The average levels of ghrelin in Wistar rats showed that the dark group had higher levels than the light and normal groups. The average ghrelin level in the dark group rats is 923.673 ng/L, the light group rats are 837.014 ng/L and 768.390 ng/L in the normal group (Table II). Based on Table II, the average serotonin levels in Wistar rats showed that the dark group had higher levels than the light group and normal group. The average ghrelin level in rats with dark treatment is 30,412 ng/ml, in light group rats it is 28.291 ng/ml and in normal group rats is 27.299 ng/ml.

Table II: The average of ghrelin and serotonin levels

Sample groups	N	Mean ± SD	
		Ghrelin	Serotonin
Normal	6	768.390 ± 108.89	27.299 ± 0.797
Dark	6	923.673 ± 55.05	30.412 ± 1.220
Bright	6	837.011 ± 63.06	28.291 ± 1.634

Noted: SD: Standard Deviation; N: replication

After a different test was conducted Wistar mouse weight correlation test with ghrelin and serotonin using statistic test Spearman Rho Correlation. The result of Wistar rat weight correlation test with ghrelin obtained was p-value = 0.006 ($p < 0.05$), meaning that there was a significant relationship between weight and Ghrelin, with a value of $r = 0.609$ indicating that the relationship between the two was strong. The direction of the relationship is straight / positive, meaning the higher the weight, the higher ghrelin, and vice versa. The results of the Wistar mouse weight correlation test with serotonin obtained were p-value = 0.023 ($p < 0.05$), meaning that there was a significant relationship between weight and serotonin, with a value of $r = 0.517$ indicating that the relationship

between the two was moderate. The direction of the relationship is straight/positive, meaning the higher the BB, the higher the serotonin, and vice versa (Table III).

Table III: The correlation between body weight to ghrelin and serotonin levels

Type of Test	Weight -Ghrelin		Weight -Serotonin	
	p	r	p	r
Correlation Test Spearman's Rho	0.006	0.609	0.023	0.517

DISCUSSION

This study was conducted on Wistar rats, where the animal is a group of nocturnal animals that are active in the dark (10). Rats are nocturnal rodents that make the time of gene expression in contrast to 180o with diurnal animals (11, 12). Humans belong to the diurnal group who are accustomed to doing activities in the morning and rest at night which causes differences in the circadian rhythm of the mouse body with humans. Exposure to light at night received by humans causes circadian rhythm disturbances, as well as in rat the disorder can occur when obtained in groups that do not get any light at all or dark groups. Literature study conducted Tähkämä (13) states that many studies were found on the impact of light on human health, especially with regard to circadian rhythms.

5-HTP is a precursor compound in the production of the neurotransmitter serotonin. Serotonin is a hormone that is responsible for carrying messages between cells in the brain. Serotonin in the brain regulates anxiety, happiness, and mood (14). Elevated serotonin levels in the dark group occur due to circadian rhythm disturbances that cause serotonin that should be converted into melatonin to return to serotonin so that melatonin continues to decline. Melatonin decreases causing the body to be awake and experience a decrease in blood pressure, so it may experience sleep deprivation (10). Rat in the dark for a period of several weeks will show a change in behavior due to the absence of light to the center of the brain that controls mood disorders so that they may experience stress. Long-term lack of light can also exacerbate stress (15). Stress can trigger an increase in the hormone cortisol that begins with the release of stress hormones that secrete corticotrophin releasing factor (CRF) released from the hypothalamus so that it reaches the pituitary gland that is just below the hypothalamus which then stimulates the release of adenocorticotrophin hormone (ACTH) and when cortisol levels in the blood fall, the hypothalamus releases corticotropin-releasing hormone (CRH), which directs the pituitary gland to produce adrenocorticotrophic hormone (ACTH). ACTH then stimulates the adrenal glands to produce and release cortisol (16).

Rat in the dark tend to do activities continuously, so an increase in cortisol can lead to an increase in

the hormone ghrelin that motivates the body to eat. Research conducted by (17) said that ghrelin levels in serum are higher in stressful conditions compared to normal conditions (18), this is evidenced by measuring the weight of feed given to rat every day, that in the dark group has a more intake of 100 grams / day with the normal weight of feed given is 50 grams / day in each cage. This habit can be referred to as Dinner Syndrome (SMM) (19), according to the theory that one of the risk factors that support the occurrence of obesity is the disruption of the performance of circadian rhythms that act as regulators of the body's biologis function as well as behavior (20). Tryptophan amino acids that are abundant in the body due to the presence of food intake can also be the cause of increased serotonin levels. Serotonin is a chemical in the brain that can affect mood. Eating foods that contain the essential amino acid known as tryptophan can help the body to produce more serotonin (21).

Serotonin is a precursor to melatonin, where serotonin increases it is also in line with the increase in melatonin. In bright conditions serotonin levels were also found to be increased but not higher compared to the dark group because the rat in the light state also experienced circadian rhythm disturbances. Exposure to received light can lead to decreased stimulation of norepinephrine in the pineal gland so that serotonin activity decreases (22). Over-available serotonin can lead to serotonin syndrome that can lead to excessive anxiety disorders that lead to stress to depression (23). Rat in a bright state will tend to do less activity so that the increase in ghrelin is not higher compared to the dark group, this is evidenced by the intake that is not always consumed in some cages in the light group.

CONCLUSION

Increased levels of ghrelin and serotonin due to circadian effects can be used as an indicator of obesity in Wistar rats.

ACKNOWLEDGEMENTS

We would like thank you for the financial support from the Government of Indonesia (DIKTI) which has been received for this research.

REFERENCES

- Susantiningsih T. Obesitas dan stres oksidatif. *JuKe Unila*. 2015;5:89–93.
- Su Y, Liu Y, Xu W, Wang J. The situation analysis of the network public opinion based on the variable-weighted Choquet fuzzy integral. *Proc - 2011 Int Conf Inf Technol Comput Eng Manag Sci ICM 2011*. 2011;1:219–25. doi:10.1109/ICM.2011.379.
- Petersen PE. Challenges to improvement of oral health in the 21st century—the approach of the WHO Global Oral Health Programme. *Int Dent J*. Elsevier; 2004;54:329–43. doi:10.1111/j.1875-595x.2004.tb00009.x.
- Weljie AM, Meerlo P, Goel N, Sengupta A, Kayser MS, Abel T, et al. Oxalic acid and diacylglycerol 36:3 are cross-species markers of sleep debt. *Proc Natl Acad Sci. National Acad Sciences*; 2015;112:2569–74. doi:10.1073/pnas.1417432112.
- Lane JM, Chang A-M, Bjornnes AC, Aeschbach D, Anderson C, Cade BE, et al. Impact of common diabetes risk variant in MTNR1B on sleep, circadian, and melatonin physiology. *Diabetes. Am Diabetes Assoc*; 2016;65:1741–51. doi:10.2337/db15-0999.
- Pejovic S, Vgontzas AN, Basta M, Tsaoussoglou M, Zoumakis E, Vgontzas A, et al. Leptin and hunger levels in young healthy adults after one night of sleep loss. *J Sleep Res*. 2010;19:552–8. doi:10.1111/j.1365-2869.2010.00844.x.
- Simpson NS, Banks S, Dinges DF. *Sleep restriction is associated with increased morning plasma leptin concentrations, especially in women*. Biol Res Nurs. Sage Publications Sage CA: Los Angeles, CA; 2010. doi:10.1177/1099800410366301.
- Navara KJ, Nelson RJ. The dark side of light at night: Physiological, epidemiological, and ecological consequences. *J Pineal Res*. 2007;43:215–24. doi:10.1111/j.1600-079X.2007.00473.x.
- Ridwan A, Zakaria Z, Barlian A. Pengaruh fotoperiode terhadap respon stres dan parameter reproduksi pada mencit jantan (*Mus musculus* L.) galur swiss webster. *J Mat Sains*. 2012;17:39–45.
- Haris RA, Tendean L, Turalaki G, Smf B, Fakultas B, Universitas K, et al. Pengaruh pemberian kopi terhadap kualitas spermatozoa tikus wistar (*Rattus norvegicus*) yang terpapar stres. *Kandidat Skripsi Fakultas Kedokteran Universitas Sam Ratulangi Manado motilitas spermatozoa yang tidak motil dengan cara menghambat siklus nukleot*. 2016;4:0–5.
- Schweinfurth MK. The social life of Norway rats (*Rattus norvegicus*). *Elife*. 2020;9:e54020. doi:10.7554/eLife.54020.
- Núñez AA, Yan L, Smale L. The cost of activity during the rest phase: Animal models and theoretical perspectives. *Front Endocrinol (Lausanne)*. Frontiers; 2018;9:72. doi:10.3389/fendo.2018.00072.
- Tähkämu L, Partonen T, Pesonen AK. Systematic review of light exposure impact on human circadian rhythm. *Chronobiol Int*. 2019;36:151–70. doi:10.1080/07420528.2018.1527773.
- Yabut JM, Crane JD, Green AE, Keating DJ, Khan WI, Steinberg GR. Emerging roles for serotonin in regulating metabolism: New implications for an ancient molecule. *Endocr Rev*. 2019;40:1092–107. doi:10.1210/er.2018-00283.
- Gonzalez M. Dim light at night and constant darkness: Two frequently used lighting conditions

- that jeopardize the health and well-being of laboratory rodents. *Front Neurol*. *Frontiers*; 2018;9:609. doi: 10.3389/fneur.2018.00609.
16. Debi Z. Pengaruh Stress Terhadap Kadar Kortisol Dan Perkembangan Kelenjar Mammae Tikus (*Rattus Novergicus*) Hamil. Universitas Andalas; 2017.
 17. Ozsoy S, Besirli A, Abdulrezzak U, Basturk M. Serum ghrelin and leptin levels in patients with depression and the effects of treatment. *Psychiatry Investig*. 2014;11:167. doi:10.4306/pi.2014.11.2.167.
 18. Zarouna S, Wozniak G, Papachristou AI. Mood disorders: A potential link between ghrelin and leptin on human body? *World J Exp Med*. 2015;5:103. doi:10.5493/wjem.v5.i2.103.
 19. Afriani AE, Margawati A, Dieny FF. Tingkat stres, durasi dan kualitas tidur, serta sindrom makan malam pada mahasiswi obesitas dan non obesitas fakultas kedokteran. *Sport Nutr J*. 2019;1:63–73. doi:10.15294/spnj.v1i2.35014.
 20. Potter GDM, Skene DJ, Arendt J, Cade JE, Grant PJ, Hardie LJ. Circadian rhythm and sleep disruption: Causes, metabolic consequences, and countermeasures. *Endocr Rev*. 2016;37:584–608. doi:10.1210/er.2016-1083.
 21. Moriska M, Lubis G, Herman RB, Moriska M, Lubis G, Herman RB. Hubungan antara kadar total triptofan plasma dan indeks massa tubuh dengan gejala depresi dan skor. 2016;17:373–8. doi:14238/sp17.5.2016.373-378.
 22. Abdel-Rahman M, Rezk MM, Ahmed-Farid OA, Essam S, Moneim AEA. *Saussurea lappa* root extract ameliorates the hazards effect of thorium induced oxidative stress and neuroendocrine alterations in adult male rats. *Environ Sci Pollut Res*. 2020;27:13237–46. doi:10.1007/s11356-020-07917-y.
 23. Scotton WJ, Hill LJ, Williams AC, Barnes NM. Serotonin syndrome: Pathophysiology, clinical features, management, and potential future directions. *Int J Tryptophan Res*. 2019;12:1–14. doi:10.1177/1178646919873925.

Ghrelin and Serotonin as Indicators of Obesity due to The Influence of Circadia on Wistar Rats

ORIGINALITY REPORT

18%

SIMILARITY INDEX

14%

INTERNET SOURCES

11%

PUBLICATIONS

2%

STUDENT PAPERS

PRIMARY SOURCES

1	www.dovepress.com Internet Source	1%
2	epubs.surrey.ac.uk Internet Source	1%
3	medic.upm.edu.my Internet Source	1%
4	my.clevelandclinic.org Internet Source	1%
5	www.medicalnewstoday.com Internet Source	1%
6	Evi Hanizar, Yung-Sen Huang, Tri Agus Siswoyo, Mohamad Syaifudin Aswan. "Light Environment Effect in the Sperm and Ribonucleic Acid Quality and Body Weight of Male <i>Mus musculus</i> ", Open Access Macedonian Journal of Medical Sciences, 2021 Publication	1%
7	psasir.upm.edu.my Internet Source	1%

8	Impact of Sleep and Sleep Disturbances on Obesity and Cancer, 2014. Publication	1 %
9	repository-tnmgrmu.ac.in Internet Source	1 %
10	repository.unair.ac.id Internet Source	1 %
11	garuda.kemdikbud.go.id Internet Source	1 %
12	www.researchgate.net Internet Source	1 %
13	www.wjgnet.com Internet Source	1 %
14	etheses.whiterose.ac.uk Internet Source	1 %
15	nmbu.brage.unit.no Internet Source	<1 %
16	hdl.handle.net Internet Source	<1 %
17	academic.oup.com Internet Source	<1 %
18	Å–nkol, F. Lemis, Rengin Zembat, and GÅ¼lden UyanÅ±k Balat. "Computer use attitudes, knowledge and skills, habits and	<1 %

methods of preschool teachers", Procedia
Computer Science, 2011.

Publication

19

Nike Hendrijantini, Poedjo Hartono,
Muhammad Dimas Aditya Ari, Fedik Abdul
Rantan. "Human Umbilical Cord Mesenchymal
Stem-Cell Therapy to Increase the Density of
Osteoporotic Mandibular Bone", European
Journal of Dentistry, 2019

Publication

20

Rachael Horne, Jane A. Foster. "Metabolic and
Microbiota Measures as Peripheral
Biomarkers in Major Depressive Disorder",
Frontiers in Psychiatry, 2018

Publication

21

Submitted to Universitas Airlangga

Student Paper

22

www.researchsquare.com

Internet Source

23

journal.frontiersin.org

Internet Source

24

link.springer.com

Internet Source

25

rjpbcs.com

Internet Source

26

www.cureus.com

Internet Source

<1 %

<1 %

<1 %

<1 %

<1 %

<1 %

<1 %

<1 %

27

www.degruyter.com

Internet Source

<1 %

28

www.falw.vu.nl

Internet Source

<1 %

29

Fuller, Grace, Christopher W. Kuhar, Patricia M. Dennis, and Kristen E. Lukas. "A Survey of Husbandry Practices for Lorisid Primates in North American Zoos and Related Facilities : Lorisid Husbandry Survey", Zoo Biology, 2013.

Publication

<1 %

30

Tuti Kusumaningsih, Anis Irmawati, Diah Savitri Ernawati, Chiquita Prahasanti, Mohammed Aljunaid, Sarah Amelia. "The differences in the number of fibroblasts and blood vessels after the topical and systemic administration of Lactobacillus casei Shirota probiotics for the treatment of traumatic ulcers in Wistar rats (Rattus norvegicus)", Veterinary World, 2021

Publication

<1 %

31

bmcpregnancychildbirth.biomedcentral.com

Internet Source

<1 %

32

pjmhsonline.com

Internet Source

<1 %

33

www.mdpi.com

Internet Source

<1 %

34

Abraham Haim, Sinam Boynao, Abed Elsalam Zubidat. "Chapter 4 Consequences of Artificial Light at Night: The Linkage between Chasing Darkness Away and Epigenetic Modifications", IntechOpen, 2019

Publication

<1 %

35

Andrew W. McHill, Joseph T. Hull, Ciaran J. McMullan, Elizabeth B. Klerman. "Chronic Insufficient Sleep Has a Limited Impact on Circadian Rhythmicity of Subjective Hunger and Awakening Fasted Metabolic Hormones", Frontiers in Endocrinology, 2018

Publication

<1 %

36

Aubé, Martin, Johanne Roby, and Miroslav Kocifaj. "Evaluating Potential Spectral Impacts of Various Artificial Lights on Melatonin Suppression, Photosynthesis, and Star Visibility", PLoS ONE, 2013.

Publication

<1 %

37

Fatma Yasmin Mahdani, Ulyasari Rachmaningayu, Elva Anjani, Asri Dewanti Adhi Nugrahaeni et al. "Topical Application of *Citrus limon* Peel Essential Oil Increased Pro-Inflammatory Cytokines Expression on Type IV Hypersensitivity Reaction", International Journal of Medical and Dental Sciences, 2023

<1 %

38

Gregory D. M. Potter, Debra J. Skene, Josephine Arendt, Janet E. Cade, Peter J. Grant, Laura J. Hardie. "Circadian Rhythm and Sleep Disruption: Causes, Metabolic Consequences, and Countermeasures", *Endocrine Reviews*, 2016

<1 %

Publication

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

Ghrelin and Serotonin as Indicators of Obesity due to The Influence of Circadia on Wistar Rats

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5
