



International Journal of Psychosocial Rehabilitation

ISSN:1475-7192

[Login/Register \(/register-login\)](#)

Journal Information

This peer reviewed Journal was created in 1996 by practitioners, mental health program managers and mental health consumers to provide international practitioners, scholars and consumers with a forum to *publish and discuss their work in program development, evaluation research, policy innovations, and therapeutic practices that have been successful in their particular region and cultures.* IJPR is not associated with any university or governmental institution, nor is it part of any 'old boy' or other professional network. It was created to provide information to an international readership about issues related to psychosocial rehabilitation and associated topics.

Articles on psychosocial interventions, psychopharmacotherapy, mental health primary care, institutional and community care innovations, decentralization, policy changes, community & regionally based systems, and program evaluation are given particular attention. However, all articles that relate to psychosocial rehabilitation will be considered.

We invite comment from all readers on any and all subjects published in this journal, including the journal format itself. Feel free to comment on the Bulletin Board as well.

Current Editorial Board

Dr. Nira Hariyatie Hartani – Malaysia

Asst. Prof. Dr. Mohd Armi Abu Samah – Malaysia

Rathi Aarti, BPTTh, MPTh, – Cardiorespiratory – India

editor@psychosocial.com

Associate Editors

Ass. Prof. Dr. I Wayan Suryasa, Indonesia

Publishing Information

The International Journal of Psychosocial Rehabilitation is published by ADG, S..A .(ADG) ADG is a private non-profit, limited corporation in Honduras and operating in a limited capacity in the United

Kingdom. All information in IJPR is provided for personal use. Reproduction for publication or other use is strictly prohibited. Written permission must be obtained ADG, S.A., to reprint any information contained within the Journal, either in part or its entirety .

submission@psychosocial.com

For authors

Scope (<https://www.psychosocial.com/scope/>)

Track Your Paper (<https://www.psychosocial.com/track-your-paper/>)

Submit Online (<https://www.psychosocial.com/submit-online/>)

Editorial Overview (<https://www.psychosocial.com/editorial-overview/>)

Instructions for Authors (<https://www.psychosocial.com/instructions-for-authors/>)

Privacy & Cookie Policy (<https://www.psychosocial.com/privacy-cookie-policy/>)

Terms & Conditions (<https://www.psychosocial.com/terms-conditions-of-use/>)



Quick Links

About Publisher (<https://www.psychosocial.com/about-publisher/>)

Copy Rights (<https://www.psychosocial.com/text-book/>)

Contact Us

editor@psychosocial.com (<mailto:editor@psychosocial.com>)

Hampstead Psychological Associates,
Suite B19, 110 Gloucester Road,
London, NW1 8JA.
United Kingdom

Copyrights © 2022 SDA, LTD. All Rights Reserved.



International Journal of Psychosocial Rehabilitation

ISSN:1475-7192

[Login/Register \(/register-login\)](#)

Issue 2

A Framework on Health Smart Home Using IoT and Machine Learning for Disabled People (<https://www.psychosocial.com/article/PR200304/9743/>)**Authors:** S.L. Rakshanasri, J. Naren, Dr.G. Vithya, S. Akhil, K. Dinesh Kumar and S. Sai Krishna Mohan Gupta**DOI:** 10.37200/IJPR/V24I2/PR200304 (<https://doi.org/10.37200/IJPR/V24I2/PR200304>)**Pages:** 1-9**Keywords:** Smart Environment, IoT, Machine Learning, Fuzzy Logic[Abstract \(https://www.psychosocial.com/article/PR200304/9743/\)](https://www.psychosocial.com/article/PR200304/9743/)[Download \(/register-login\)](#)**Ontological Framework for Analyzing Student's Emotional behavior Performance Enhancement Using Fuzzy Logic (<https://www.psychosocial.com/article/PR200305/9745/>)****Authors:** Kirthika Ashokkumar, J. Naren, S.L. Rakshanasri and Dr.G. Vithya**DOI:** 10.37200/IJPR/V24I2/PR200305 (<https://doi.org/10.37200/IJPR/V24I2/PR200305>)**Pages:** 10-18**Keywords:** Personality Traits, Semantic Feedback, Ontology, Fuzzy Logic[Abstract \(https://www.psychosocial.com/article/PR200305/9745/\)](https://www.psychosocial.com/article/PR200305/9745/)[Download \(/register-login\)](#)**Health Smart Home with IoT – A State of Art Survey (<https://www.psychosocial.com/article/PR200306/9747/>)****Authors:** S.L. Rakshanasri, J. Naren, Dr.G. Vithya, S. Akhil, K. Dinesh Kumar and S. Sai Krishna Mohan Gupta**DOI:** 10.37200/IJPR/V24I2/PR200306 (<https://doi.org/10.37200/IJPR/V24I2/PR200306>)**Pages:** 19-28**Keywords:** Smart Health System, IOT, Art Survey[Abstract \(https://www.psychosocial.com/article/PR200306/9747/\)](https://www.psychosocial.com/article/PR200306/9747/)[Download \(/register-login\)](#)

Economic Dimensions of Blockchain Technology: In the Context of Extention of Cryptocurrencies (<https://www.psychosocial.com/article/PR200307/9749/>)**Authors:** Valliappan Raju and Siew Poh Phung**DOI:** 10.37200/IJPR/V24I2/PR200307 (<https://doi.org/10.37200/IJPR/V24I2/PR200307>)**Pages:** 29-39**Keywords:** Blockchain Technology, FinTech, Distributed Ledger Technology, Smart Contracts, Blockchain RegulationAbstract (<https://www.psychosocial.com/article/PR200307/9749/>)

Download (/register-login)

Observation of Bitcoin Trade for the Purpose of Hedge Trade in Global Stock Market (<https://www.psychosocial.com/article/PR200308/9751/>)**Authors:** Valliappan Raju and Siew Poh Phung**DOI:** 10.37200/IJPR/V24I2/PR200308 (<https://doi.org/10.37200/IJPR/V24I2/PR200308>)**Pages:** 40-52**Keywords:** Bitcoin Trade, Global Stock ExchangeAbstract (<https://www.psychosocial.com/article/PR200308/9751/>)

Download (/register-login)

Development of Auditing Compliance by Implementation of Blockchain Technology: An Explanatory Study based on Observations (<https://www.psychosocial.com/article/PR200309/9753/>)**Authors:** Valliappan Raju and Siew Poh Phung**DOI:** 10.37200/IJPR/V24I2/PR200309 (<https://doi.org/10.37200/IJPR/V24I2/PR200309>)**Pages:** 53-60**Keywords:** Blockchain Auditing, Blockchain TechnologyAbstract (<https://www.psychosocial.com/article/PR200309/9753/>)

Download (/register-login)

Utility of Blockchains in Publishing Sector: Focus on Academic Publishing (<https://www.psychosocial.com/article/PR200310/9755/>)**Authors:** Siew Poh Phung and Valliappan Raju**DOI:** 10.37200/IJPR/V24I2/PR200310 (<https://doi.org/10.37200/IJPR/V24I2/PR200310>)**Pages:** 61-66**Keywords:** Blockchain, Cryptocurrency, Bitcoin, Scholarly Communication, PublishingAbstract (<https://www.psychosocial.com/article/PR200310/9755/>)

Download (/register-login)

[Download \(/register-login\)](#)

Association between Arterial Stiffness and Left Ventricular Diastolic Dysfunction in Stable Coronary Heart Disease (<https://www.psychosocial.com/article/PR200719/10761/>)

Authors: Diah Masita Cahyani, Achmad Lefi, Budi Utomo, Agus Soebagjo

DOI: 10.37200/IJPR/V24I2/PR200719 (<https://doi.org/10.37200/IJPR/V24I2/PR200719>)

Pages: 4006-4013

Keywords: Arterial stiffness, Cardio-ankle vascular index, Diastolic, Left ventricular

[Abstract \(https://www.psychosocial.com/article/PR200719/10761/\)](https://www.psychosocial.com/article/PR200719/10761/)

[Download \(/register-login\)](#)

CORRELATION BETWEEN SOCIAL ENGAGEMENT AND COGNITIVE FUNCTION IN ELDERLY (<https://www.psychosocial.com/article/PR200720/10763/>)

Authors: Margareth Haryanto, Eko Wahono, Riani Wisnujono, Mudjjiani Basuki

DOI: 10.37200/IJPR/V24I2/PR200720 (<https://doi.org/10.37200/IJPR/V24I2/PR200720>)

Pages: 4014-4019

Keywords: social engagement, cognitive function, elderly, index social engagement

[Abstract \(https://www.psychosocial.com/article/PR200720/10763/\)](https://www.psychosocial.com/article/PR200720/10763/)

[Download \(/register-login\)](#)

DIAGNOSTIC TEST USING NEUROLOGICAL DEPRESSION DISORDERS INVENTORY FOR EPILEPSY COMPARED TO HAMILTON DEPRESSION RATING SCALE -17 AS A GOLD STANDARD (<https://www.psychosocial.com/article/PR200721/10765/>)

Authors: Andi Prasetiawan, Kurnia Kusumastuti

DOI: 10.37200/IJPR/V24I2/PR200721 (<https://doi.org/10.37200/IJPR/V24I2/PR200721>)

Pages: 4020-4029

Keywords: Diagnostic Test, Neurological Disorders Depression Inventory for Epilepsy, Hamilton Depression Rating Scale-17

[Abstract \(https://www.psychosocial.com/article/PR200721/10765/\)](https://www.psychosocial.com/article/PR200721/10765/)

[Download \(/register-login\)](#)

PAIN REDUCTION AFTER WET AND DRY CUPPING THERAPIES: ROLES OF $\alpha 2\beta 1$ INTEGRIN AND μ -OPIOID RECEPTOR IN ANIMAL MODELS (<https://www.psychosocial.com/article/PR200722/10767/>)

Authors: Imam Subadi, Boya Nugraha, Abdulhadi Suwandi, Nur Sulastri, Imam Susilo

DOI: 10.37200/IJPR/V24I2/PR200722 (<https://doi.org/10.37200/IJPR/V24I2/PR200722>)

Abstract (<https://www.psychosocial.com/article//39220/>)

Download (/register-login)

Innovative personality traits and their relationship to the skills of standing on the hands and card will on the floor of the floor movements in your women's artistic gym. (<https://www.psychosocial.com/article//39222/>)

Authors: Sanarya Jabbar Mahmood

DOI: 10.37200/IJPR/V24I2/PR2024464 (<https://www.psychosocial.com/wp-content/uploads/2022/04/PR2024465.pdf>)

Pages: 9891-9896

Keywords: Innovative personality traits, hands-on skill, kart will skill in women's art

Abstract (<https://www.psychosocial.com/article//39222/>)

Download (/register-login)

Quantify the contribution of personality traits according to the risky cognitive method versus caution in performing the skills of rolling up and landing a human wheel on the balance model device in your artistic gym for female students. (<https://www.psychosocial.com/article//39224/>)

Authors: Sanarya Jabbar Mahmood. Basma Naeem Mohsen. Hanan Adnan Abaaab

DOI: 10.37200/IJPR/V24I2/PR2024465 (<https://www.psychosocial.com/wp-content/uploads/2022/04/PR2024465.pdf>)

Pages: 9897-9908

Keywords: .

Abstract (<https://www.psychosocial.com/article//39224/>)

Download (/register-login)

Physical abilities and their relationship to some physiological variables of the 100-meter freestyle swimmers for youth (<https://www.psychosocial.com/article//39507/>)

Authors: Ahmed Mohammed Farhan

Pages: 9908-9914

Keywords: physical abilities, functional variables, free-swimming

Abstract (<https://www.psychosocial.com/article//39507/>)

Download (/register-login)

For authors

Scope (<https://www.psychosocial.com/scope/>)

Track Your Paper (<https://www.psychosocial.com/track-your-paper/>)

Submit Online (<https://www.psychosocial.com/submit-online/>)

Editorial Overview (<https://www.psychosocial.com/editorial-overview/>)

[Instructions for Authors \(https://www.psychosocial.com/instructions-for-authors/\)](https://www.psychosocial.com/instructions-for-authors/)

[Privacy & Cookie Policy \(https://www.psychosocial.com/privacy-cookie-policy/\)](https://www.psychosocial.com/privacy-cookie-policy/)

[Terms & Conditions \(https://www.psychosocial.com/terms-conditions-of-use/\)](https://www.psychosocial.com/terms-conditions-of-use/)



Quick Links

[About Publisher \(https://www.psychosocial.com/about-publisher/\)](https://www.psychosocial.com/about-publisher/)

[Copy Rights \(https://www.psychosocial.com/text-book/\)](https://www.psychosocial.com/text-book/)

Contact Us

editor@psychosocial.com (mailto:editor@psychosocial.com)

Hampstead Psychological Associates,
Suite B19, 110 Gloucester Road,
London, NW1 8JA.
United Kingdom

Copyrights © 2022 SDA, LTD. All Rights Reserved.

PAIN REDUCTION AFTER WET AND DRY CUPPING THERAPIES: ROLES OF $\alpha 2\beta 1$ INTEGRIN AND μ -OPIOID RECEPTOR IN ANIMAL MODELS

¹Imam Subadi, ²BoyaNugraha, ³AbdulhadiSuwandi, ^{*1}Nur Sulastri^{1*}, ⁴Imam Susilo

ABSTRACT---Background: Wet cupping therapy (WCT) and dry cupping therapy (DCT) are complementary therapies in pain management. Stretching keratinocyte by cupping seems to stimulate the expression of $\alpha 2\beta 1$ -integrin. Its benefit in pain reduction could be mediated by the expression of μ -opioid receptor (MOR).

Objective: This study aims to determine pain thresholds, $\alpha 2\beta 1$ -integrin and MOR after WCT and DCT.

Method: Thirty-two adult male Wistar rats were divided into negative control, positive control, DCT, and WCT groups. In DCT group and WCT group, cupping was conducted at left and right paralumbar region 48h after CFA injection. Hot plate test was conducted 24h after therapy to assess pain threshold and immunohistochemistry were conducted from the skin for $\alpha 2\beta 1$ -integrin and the spinal cord for MOR.

Result: WCG showed the highest expression of $\alpha 2\beta 1$ -integrin in the keratinocytes and MOR in the lamina II spinal cord than the other groups. On immunohistochemical examination, the expression of $\alpha 2\beta 1$ integrin showed positive cells (keratinocytes) with brown stain on dry cupping and wet cupping. Significant differences were shown between negative control group showed significant differences with positive control group, dry cupping therapy group, and wet cupping therapy group.

Conclusion: WCT is more effective than DCT in reducing pain. The benefit of both therapies in reducing pain in rats could be mediated by the expression of $\alpha 2\beta 1$ -integrin and MOR.

Keywords---Dry cupping, pain reduction, wet cupping,

I. INTRODUCTION

Cupping therapy is one of the complementary and alternative therapies that is widely used to reduce pain (Kim *et al.*, 2011). Previous study showed that cupping therapy can be used for acute and chronic pain (Cao *et al.*, 2004). In Indonesia there are two methods of cupping therapy, the dry cupping therapy and wet cupping therapy. Dry cupping therapy cups the skin without bleeding. In wet cupping therapy, the skin is injured and cupped, resulting in blood loss (Arslan, Gökgöz and Dane, 2016). The difference in the effectiveness of each cupping therapy method is not

¹Department of Physical Medicine and Rehabilitation, Faculty of Medicine/Universitas Airlangga Hospital, Universitas Airlangga, Surabaya 60131, Indonesia

²Department of Rehabilitation Medicine, Hannover Medical School, Hannover 30625, Germany

³Institute of Medical Microbiology and Hospital Epidemiology, Hannover 30625 Medical School, Hannover, Germany

⁴Department of Pathology Anatomy, Faculty of Medicine, Universitas Airlangga, Surabaya 60131, Indonesia

*Corresponding Author: Nur Sulastri, MD

Department of Physical Medicine and Rehabilitation, Hospital of Airlangga University, Campus C Mulyorejo, Surabaya 60115, Indonesia

Phone: +62315916290

E-mail: nuri.rehab@gmail.com

widely known. In both cupping methods negative pressure is applied which causes the skin to stretch. Stretched cells cause stimulation of mechanoreceptors of degenerin/epithelial sodium channel (DEG/ENaC), acid-sensing ion channels ASICs, transient receptor potential vanilloid 4 TRPV4 and integrins (Delmas, Hao and Rodat-Despoix, 2011).

Integrins are transmembrane receptors that connect the extracellular matrix with the cytoskeleton (Hynes, 2004). Integrins function as mechanoreceptors which transmit forces between the extracellular matrix and the cytoskeleton to maintain tissue structure integrity (Baker and Zaman, 2010). Cell strain causes activation of signal pathways which include mitogen-activated protein (MAP) kinase, Rho guanosine triphosphate (GTP)ase, increased cytoplasmic calcium, and reactive oxygen. These events cause changes in gene expression (Schwartz, 2010). Puncture on the skin on wet cupping is a stressor that can stimulate the expression of endorphin (Subadi *et al.*, 2017). The hypothesis of this study was that wet cupping is more effective than dry cupping because there is an addition of puncture which causes endorphin expression. This study aims to determine pain thresholds, $\alpha 2\beta 1$ -integrin and MOR after WCT and DCT.

II. METHODS

This study used post-test control group design. Thirty-two Wistar, 3 months, 250 ± 50 grams, male rats were randomized into 4 groups: negative control (n=8); positive control (n=8); dry cupping (n=8); wet cupping (n=8). In the negative control group, the rats had no exposure at all. In the positive control group, the rats were given complete Freund's adjuvant (CFA) (Sigma-Aldrich, Saint Louis, Missouri, USA). In the dry cupping group, the rats received CFA and dry cupping. In the wet cupping therapy group, the rats received CFA and wet cupping therapy.

Adult Wistar rats were put into plastic cage (4-5 rats per cage). They were acclimatized for 7 days with 12:12 h (dark/light; the light was turned off every 19:30) with temperature maintained at 24°C . CFA (100 μL) was injected in the center of the right plantar footpad in the three groups of the rats (Fehrenbacher, Vasko and Duarate, 2012). In the groups of dry cupping and wet cupping, 48 hours after the injection, the rats were treated with dry cupping or wet cupping therapy (Subadi *et al.*, 2017). In the dry cupping group, the skin of the right and left paralumbal areas was cupped (2 cm in diameter) with a negative pressure of 200 mmHg for 5 minutes (Subadi *et al.*, 2017). In the wet cupping group, the rats were pricked with lancet 28G before being cupped. Twenty-four hours after wet and dry cupping, all rats were examined for pain threshold using a hot plate (Hot/Cold Plate Cat 35100, Ugo Basile, Varese, Italy). The temperature of the hot plate was 51°C [15]. After examination of the pain threshold, the rats were terminated by cervical dislocation and the paralumbal skin was examined with immunohistochemistry.

Pain was examined using hot plate in bright conditions. The pain threshold was calculated from the moment the rats were placed on the hot plate surface until the nociceptive response occurred. Response showing discomfort (Nocifensive) behaviors include forepaw withdrawal or licking, hind paw withdrawal or licking, stamping, leaning posture and jumping (Deuis, Dvorakova and Vetter, 2017), with a maximum time limit of 30 seconds. One measurement used a stopwatch for each rat to obtain the average value. All the nocifensive behavior responses were conducted by a trained and experienced person in order to have homogenous results.

Expression of $\alpha 2\beta 1$ integrin in keratinocytes and μ -opioid receptors in nerve cells in the spinal cord lamina II was immunohistochemically examined using anti-integrin monoclonal antibody $\alpha 2\beta 1$ (Cat. No MAB2141Z, Chemicon, Merck Millipore co., Darmstadt, Germany) and μ -opioid anti-receptor monoclonal antibody (ab64746, Abcam,

Cambridge, USA). Cells that were positive for $\alpha 2\beta 1$ integrins and μ -opioid receptors were calculated using a microchemical light (Olympus CX21, New York, USA).

III. RESULT

Pain threshold was determined 24 hours after dry cupping therapy and wet cupping therapy, while in positive control group it was performed 72 hours after CFA injection. Figure 1 demonstrates the results of pain threshold experiment. Significant differences were observed between negative control and dry cupping therapy; and negative control and wet cupping therapy. Similar significant differences were also observed between positive control and dry cupping therapy; and positive control and wet cupping therapy. Interestingly, wet cupping therapy and dry cupping therapy showed significant difference. In this study there were two samples in wet cupping group which had to be stopped because it exceeded 30 seconds.

On immunohistochemical examination, the expression of $\alpha 2\beta 1$ integrin showed positive cells (keratinocytes) with brown stain on dry cupping and wet cupping (Figure 2). Negative reactions of $\alpha 2\beta 1$ integrin expression did not show brown stain. Number of keratinocytes expressing $\alpha 2\beta 1$ integrin is demonstrated on Figure 3. Significant differences were shown between negative control group and dry cupping therapy group, negative control group and wet cupping therapy group, and positive control group and wet cupping therapy.

On immunohistochemical examination, the expression of μ -opioid receptors showed positive cells (neurons) with brown stain on dry cupping and wet cupping (Figure 4). Negative reactions of μ -opioid receptors did not show brown stain. Number of nerve cells in the spinal cord lamina II expressing μ -opioid receptors is demonstrated on Figure 4. Significant differences were shown between negative control group showed significant differences with positive control group, dry cupping therapy group, and wet cupping therapy group. Similar trend was also observed in positive control group. Interestingly, dry cupping therapy group showed significant difference with wet cupping therapy group.

IV. DISCUSSION

In this study, dry cupping therapy and wet cupping therapy increased the expression of $\alpha 2\beta 1$ integrins in keratinocytes and μ -opioid receptors in spinal cord nerve cells. It seems that the negative pressure applied in dry cupping therapy and wet cupping therapy triggers the expression of $\alpha 2\beta 1$ integrins. In this study, $\alpha 2\beta 1$ integrin expression in wet cupping therapy was higher than that in dry cupping therapy. The difference between wet and dry cupping therapy methods is the puncture. The puncture causes damage to the keratinocyte membrane. Cell damage causes inflammatory reactions and secretion of inflammatory mediators such as bradykinin, prostaglandin, leukotriene, serotonin, histamine, and substance P (Amaya *et al.*, 2013). The integrins which contained subunit $\beta 1$ play a role in nociceptor sensitization by inflammatory mediators (Dina *et al.*, 2004, 2005). Additionally, they also modulate neuronal excitability and injury after mechanical stimulation to the surrounding extra cellular matrix (Wang *et al.*, 2006; Hemphill *et al.*, 2011). Mechanical stimulation is resulted by both dry and wet cupping therapy.

This study shows that dry cupping therapy and wet cupping therapy can reduce pain. These findings were in accordance with those of other researchers that dry cupping therapy and wet cupping therapy reduced pain (Dalton and BJ, 2017). Pain threshold test in this study was limited to 30 seconds. This study was conducted to prevent tissue

damage to the rat's paws. There were 25% (2 out of 8) samples in wet cupping therapy group exceeded 30 seconds. It could be caused by effectiveness of wet cupping therapy in increasing pain thresholds.

The expression of μ -opioid receptors in wet cupping therapy was higher than that in dry cupping therapy. μ -opioid receptor activation inhibits pain transmission in the spinal cord (Al-Hasani and Bruchas, 2011). In a previous study, β -endorphin expression was found after wet cupping therapy (Subadi *et al.*, 2017). β -endorphin causes analgesia through bonding with μ -opioid receptors. In central nervous system, β -endorphin binds with μ -opioid receptor, causing a cascade of interactions that inhibits the expression of substance P and GABA that contribute to pain transmission (Sprouse-Blum *et al.*, 2010).

In this study, the expression of opioid receptor in wet cupping therapy was higher than that in dry cupping therapy. It was allegedly because puncture was applied in wet cupping therapy, which caused cell injury. Cells affected by lesion will express inflammatory cytokines that can trigger endorphins through the POMC in the hypothalamus (Slominski *et al.*, 2000).

V. CONCLUSION

This study provided evidence-based data that wet cupping therapy is more effective in reducing pain as compared to dry cupping therapy. Pain reduction mechanism in both therapies was mediated by the expression of $\alpha 2\beta 1$ integrin and μ -opioid receptors which are higher in wet cupping.

REFERENCE

- [1] Al-Hasani, R. and Bruchas, M. (2011) 'Molecular mechanisms of opioid receptor-dependent signaling and behavior', *Anesthesiology*, 115(6), pp. 1363–81.
- [2] Amaya, F. *et al.* (2013) 'Tissue Injury and Related Mediators of Pain Exacerbation', *Curr Neuropharmacol*, 11(6), pp. 592–597.
- [3] Arslan, M., Gökğöz, N. and Dane, S. (2016) 'The effect of traditional wet cupping on shoulder pain and neck pain: A pilot study', *Complement Ther Clin Pract*, 23, pp. 30–3.
- [4] Baker, E. and Zaman, M. (2010) 'The biomechanical integrin', *J Biomech*, 43(1), pp. 38–44.
- [5] Cao, H. *et al.* (2004) 'Cupping therapy for acute and chronic pain management: a systematic review of randomized clinical trials', *J Tradit Chinese Med Sci*, 1(1), pp. 49–61.
- [6] Dalton, E. and BJ, V. (2017) 'Cupping Therapy: An Alternative Method of Treating Pain', *Public Health - Open Journal*, 2(2), pp. 59–63.
- [7] Delmas, P., Hao, J. and Rodat-Despoix, L. (2011) 'Molecular mechanisms of mechanotransduction in mammalian sensory neurons', *Nat Rev Neurosci*, 12(3), pp. 139–53.
- [8] Deuis, J., Dvorakova, L. and Vetter, I. (2017) 'Methods Used to Evaluate Pain Behaviors in Rodents', *Front Mol Neurosci*, 10, p. 284.
- [9] Dina, O. *et al.* (2004) 'Integrin signaling in inflammatory and neuropathic pain in the rat', *Eur J Neurosci*, 19(3), pp. 634–42.
- [10] Dina, O. *et al.* (2005) 'Primary afferent second messenger cascades interact with specific integrin subunits in producing inflammatory hyperalgesia', *Pain*, 115(1–2), pp. 191–203.
- [11] Fehrenbacher, J., Vasko, M. and Duarate, D. (2012) 'Models of inflammation: Carrageenan- or complete Freund's Adjuvant (CFA)-induced edema and hypersensitivity in the rat', *Curr Protoc Pharmacol*.
- [12] Hemphill, M. *et al.* (2011) 'A Possible Role for Integrin Signaling in Diffuse Axonal Injury', *PLOS ONE*, 6(7), p. e22899.
- [13] Hynes, R. (2004) 'The emergence of integrins: a personal and historical perspective', *Matrix Biol*, 23(6), pp. 333–40.
- [14] Kim, J. *et al.* (2011) 'Cupping for treating pain: a systematic review', *Evid Based Complement Alternat Med*, 2011, p. 467014.
- [15] Schwartz, M. (2010) 'Integrins and extracellular matrix in mechanotransduction', *Cold Spring Harb Perspect Biol*, 2(12), p. a005066.

- [16] Slominski, A. *et al.* (2000) 'Corticotropin releasing hormone and proopiomelanocortin involvement in the cutaneous response to stress', *Physiol Rev*, 80(3), pp. 979–1020.
- [17] Sprouse-Blum, A. *et al.* (2010) 'Understanding Endorphins and Their Importance in Pain Management', *Hawaii Med J*, 69(3), pp. 70–71.
- [18] Subadi, I. *et al.* (2017) 'Pain Relief with Wet Cupping Therapy in Rats is Mediated by Heat Shock Protein 70 and β -Endorphin', *Iran J Med Sci*, 42(4), pp. 384–391.
- [19] Wang, X. *et al.* (2006) 'Role of TGF β -mediated inflammation in cutaneous wound healing', *J Investig Dermatol Symp Proc*, 11(1), pp. 112–7.




Figure 1: Results of Pain Threshold Experiment




Figure 2: Expression of $\alpha 2\beta 1$ Integrin with Keratinocytes




Figure 3: Number of Keratinocytes Expressing $\alpha 2\beta 1$ Integrin




Figure 4: Result of Immunohistochemical Examination The Expression of μ -opioid Receptors with Neurons