

Clinical and ultrasonography evaluation of thyroid tumor screening in symptomatic patient of Bajulmati primary care center, Banyuwangi, East Java, Indonesia

by Rosy Setiawati

Submission date: 02-Apr-2023 04:59AM (UTC+0800)

Submission ID: 2053013121

File name: ulmati_primary_care_center,_Banyuwangi,_East_Java,_Indonesia.pdf (500.58K)

Word count: 4410

Character count: 23935

Clinical and ultrasonography evaluation of thyroid tumor screening in symptomatic patient of Bajulmati primary care center, Banyuwangi, East Java, Indonesia

Rosy Setiawati, MD, PhD^{a,*}, Tri Wulanhandarini, MD^a, Fierly Hayati, MD^a, Dyah Erawati, MD^a, Merlin Guntur Jaya, MD^a, Andi Ahmad Thoriq, MD^a, Triana Mediyawati Wijaya, MD^a, Galih Nur Ismiyati, MD^a, Dyan Wahyu Kusumaningrum, MD^a, Belinda Koesmarsono, MD^a, Agnes Triana Basja, MD^a, M. Ikhsan Nugroho, MD^a, Silvi Yuliana, MD^a, Syadza Zahrah Shedyta, MD^a, Hendra Boy Situmorang, MD^a

Abstract

This study aims to assess the prevalence, clinical, and ultrasonography (US) in thyroid screening in healthy subjects with general symptoms of thyroid abnormality in low iodine intake in Bajulmati primary care center, East Java Indonesia. We retrospectively reviewed US thyroid examination of 74 subjects with symptoms of mass in the neck, shaky, sleep difficulties, over sweating, and chronic fatigue on September 15th, 2021. Following the WHO guidelines, subjects also underwent physical examination in which the result were classified into 3 categories, that is, no palpable nor visible goiter, palpable but no visible goiter, as well as palpable and visible goiter. We evaluate US thyroid characteristics following Korean Society of Thyroid Radiology guidelines. Image analysis was reviewed by 4 general radiologists with 2 to 13 years' experience. Categorical variables were compared using chi-squared or Fisher exact tests. Correlation between variables was measured with gamma statistics. Statistical analyses were conducted using IBM SPSS Statistics 23.0. A *P*-value < .05 was considered to indicate statistical significance. Of the 74 subjects, 32 (43.2%) show abnormalities. Statistical analysis showed no significant differences in the result of thyroid US in subjects with complaint fatigue (*P* = .464), insomnia (*P* = .777), over sweating (*P* = .158), and tremor (*P* = .778), but there were significant differences with the complaint of mass in the neck (*P* = .008). Furthermore, there was also a strong correlation between goiter palpation and US thyroid result (*R* = 0.773, *P* = .00). We conclude there were significant differences in US result of patients with and without complaint of mass in the neck. We also found a strong correlation between goiter palpation and US examination. Clinical findings, laboratory examination, cytology and molecular markers, patients' age, nodules size, and ultrasound features should be considered for the treatment planning.

Abbreviations: FNA = fine needle aspiration, TSH = thyroid stimulating hormone, US = ultrasonography.

Keywords: primary health care, screening, thyroid cancer, ultrasonography

1. Introduction

Thyroid cancer in 2020 had an incidence rates of 10.1 per 100,000 women and 3.1 per 100,000 men with age-standardized mortality rates 0.5 per 100,000 women and 0.3 per 100,000 men globally.^[1] There is increase of incidence reported with findings: nearly all registered increase is limited to papillary histotype which is the least aggressive variant of thyroid cancer, and it is more marked in population groups with higher socioeconomic standard and better healthcare access.^[2] The true thyroid cancer increase might be due to environmental factors, female genetic and hormonal background of people who are more prone to thyroid diseases.^[3]

The submission is appropriate for publication in PPH because we raised the topic of screening of thyroid in the primary health care for the importance of early detection of thyroid abnormalities. This manuscript is original and has not been submitted elsewhere in part or in whole. The authors have no funding and conflicts of interest to disclose.

All data generated or analyzed during this study are included in this published article [and its supplementary information files].

^a Radiology Department, Universitas Airlangga, Faculty of Medicine, Dr. Soetomo Academic General Hospital, Surabaya, Indonesia.

* Correspondence: Rosy Setiawati, Radiology Consultant, Universitas Airlangga, Faculty of Medicine, Dr. Soetomo Academic General Hospital, Moestopo No. 47, Surabaya, Indonesia (e-mail: rosy-s@fk.unair.ac.id).

The early finding of the malignant thyroid cancer can help decreasing mortality, reduce morbidity and to avoid unnecessary tests and surgery in benign nodule, hence the nodule detection is crucial.^[4] Routine examination of thyroid include physical examination inspection and palpation of the neck. A more detailed, noninvasive, inexpensive could be achieved with ultrasonography.^[5] Ultrasonography (US) becomes a choice for the screening tools in thyroid abnormality for its non-invasiveness and its availability.^[6] The advances of the US had resulted in the identification of the thyroid nodules whether in the asymptomatic or symptomatic patient.^[7] Thyroid nodules detected by the US were 65%, while in the palpations were 4%.^[8]

Copyright © 2022 the Author(s). Published by Wolters Kluwer Health, Inc. This is an open access article distributed under the Creative Commons Attribution License 4.0 (CCBY), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Setiawati R, Wulanhandarini T, Hayati F, Erawati D, Jaya MG, Thoriq AA, Wijaya TM, Ismiyati GN, Kusumaningrum DW, Koesmarsono B, Basja AT, Nugroho MI, Yuliana S, Shedyta SZ, Situmorang HB. Clinical and ultrasonography evaluation of thyroid tumor screening in symptomatic patient of Bajulmati primary care center, Banyuwangi, East Java, Indonesia. *Medicine* 2022;101:52(e32546).

Received: 14 August 2022 / Received in final form: 11 December 2022 / Accepted: 12 December 2022

<http://dx.doi.org/10.1097/MD.00000000000032546>

Thyroid nodules may depend on several factors, including low iodine intake in the area with iodine insufficiency.^[9] Iodine intake is a crucial determinant of the prevalence of thyroid abnormality in a population where thyroidal hyperplasia, hypertrophy, and nodules formation may happen within months of the iodine deficiency from the free radical oxygen mutagenic effects.^[10] Low iodine intake may increase the thyroid-stimulating hormone (TSH) stimulation and causing increased thyroid cell responsiveness to TSH, thus resulting in thyroid cell epidermal growth factor beta 1 production and increased angiogenesis related to promotion of tumor growth.^[11] East Java is one of Indonesia's regions with low median urinary iodine concentration ($<150 \mu\text{f/L}$), showing deficiency of iodine in the East Javanese population.^[12]

Currently, no studies had evaluated the use of ultrasonography screening to provide clear guidance and to improve clinicians' judgment, especially in Indonesia with high rates of low iodine intake.^[13] Therefore, early detection of thyroid abnormality could benefit the treatment in the future. This study aims to assess the prevalence, clinical, and US in thyroid screening in healthy subjects with general symptoms of thyroid abnormality in Bajulmati primary care center, East Java Indonesia.

12

2. Materials and Methods

2.1. Patients

This retrospective study was approved by the ethical board of our institution, and the necessity for informed consent for research was waived. A total of 76 consecutive symptomatic subjects underwent screening thyroid US at the primary care center Bajulmati with community services on September 15th, 2021. We include subjects with symptoms of mass in the neck, shaky, sleep difficulties, over sweating, and chronic fatigue. There were 2 subjects with a history of thyroid surgery excluded from this study. The subjects were asked about their symptoms then underwent neck palpation examination. Following the WHO guidelines, subjects were then separated into 3 classifications: subjects with no palpable nor visible goiter, palpable but no visible goiter, as well as palpable and visible goiter.^[14]

2.2. US imaging

US thyroid examination was performed at the primary care center in Bajulmati Banyuwangi Indonesia using portable US equipment Samsung HM70A and Midray D10 with linear transducer by 4 board-certified radiologists. Based on the recommendation by the Korean society of thyroid radiology, thyroid nodules were described by the features of the nodule. The features mentioned in our study include composition, echogenicity, margins, axis, calcification, size. The composition includes cystic ($>90\%$ of the cystic portion) or predominantly cystic ($>50\%$ of the cystic portion and $<90\%$ of the cystic portion) nodules with reverberating artifacts and spongiform nodules. The echogenicity includes hypoechoic, isoechoic, and hyperechoic. The size of the nodule can be $>1 \text{ cm}$, $>2 \text{ cm}$. The margin mentioned could be smooth, spiculated, and ill-defined. While shapes mentioned were ovoid to round, taller than wider, and irregular.

2.3. Imaging analysis, reference standards, and statistical analysis

Image analysis was reviewed by 4 general radiologists with 2 to 13 years' experience. In multiple nodules, the most suspicious nodules were the 1 to be described. Discrepancies in US results will be resolved by the opinion of the most experienced radiologist. Categorical variables were compared

using chi-squared or Fisher exact tests, while the correlation between variables was measured with gamma statistics. Statistical analyses were conducted using IBM SPSS Statistics 23.0. A P -value <0.05 was considered to indicate statistical significance.

3. Results

A total of 74 subjects included in this study were 68 women and 6 men with a mean age of 42.71 years (range from 1 year to 87 years). Among them, 15 subjects (20.3%) with complaint of tremor, 15 subjects (20.3%) with a complaint of over sweating, 16 subjects (21.6%) with a complaint of insomnia, 26 subjects (35.1%) with a complaint of chronic fatigue, and 20 subjects (17%) with a complaint of a neck mass. (Table 1)

Among the included subjects, 32 subjects (43.2%) have nodule abnormality. The nodule involvement were 17 subjects (23%) unilateral and 15 patients (20.3%) bilateral. Subjects with single nodule were 17 subjects (23%) while with multiple nodules were 15 subjects (20.3%). The shape of the nodules for 30 subjects (40.5%) was ovoid to round or wider than taller while 2 subjects (2.7%) were taller than wider. The contents of the right thyroid nodules were solid in 6 subjects (8.1%), predominant solid in 6 subjects (8.1%), predominant cystic in 7 subjects (9.5%), cystic in 5 subjects (6.8%), and spongiform in 3 subjects (4.1%). While the contents of the left thyroid nodules were solid in 1 subject (1.4%), predominant solid in 8 subjects (10.8%), predominant cystic in 7 subjects (9.5%), cystic in 3 subjects (4.1%), and spongiform in 2 subjects (2.7%). The echogenicity of the nodules in the right thyroid was hypoechoic in 13 subjects (17.6%), isoechoic in 2 subjects (2.7%), hyperechoic in 5 subjects (6.8%), and mixed echo in 2 subjects (2.7%). While in the left side, the nodules were hypoechoic in 17 subjects (23%), isoechoic in 3 subjects (4.1%), hyperechoic in 5 subjects (6.8%), and mixed echo in 1 subject (1.4%). The margin of the nodules in the right side of the thyroid were smooth in 24 subjects (32.4%), ill-defined in 3 patients (4.1%) while in the left side of the thyroid 19 subjects (25.7%) were smooth and 2 subjects (2.7%) were ill-defined. Subjects with nodules calcification shaped stippled, fine coarse and nonlinear were 2 subjects (2.7%), and 2 other subjects (2.7%) were shaped curvilinear, smooth margin (Table 2).

Statistical analysis showed there were significant differences in US result of patients with and without complaint of mass in the neck ($P = .008$). (Table 3) There was also a significant strong correlation between goiter palpation and US thyroid result ($R = 0.773$, $P = .00$). (Table 4)

Table 1

Patients' demography.

Patient demography	
Age (range) (yr)	42.71 (0.88–87)
Gender	
Female	68 (91.9%)
Male	6 (8.1%)
Patients' complaint	
Palpitation	22 (29.7%)
Shaky	15 (20.3%)
Over sweating	15 (20.3%)
Insomnia	16 (21.6%)
Chronic fatigue	26 (35.1%)
Neck mass	20 (17.0%)
Neck physical examination	
Not palpable nor seen	52 (70.3%)
Palpable but not seen	5 (6.8%)
Seen enlarged	17 (23%)

Table 2
US Features of screening-detected thyroid cancer.

US result	
Normal	42 (56.8%)
With nodul	32 (43.2%)
Involvement	
Unilateral	17 (23%)
Bilateral	15 (20.3%)
Amount of nodul	
Single	17 (23%)
Multiple	15 (20.3%)
Shape of the nodul	
Ovoid to round	30 (40.5%)
Taller than wider	2 (2.7%)
Content of the right thyroid nodul	
Normal	47 (63.5%)
Solid	6 (8.1%)
Predominant solid	6 (8.1%)
Predominant cystic	7 (9.5%)
Cystic	5 (6.8%)
Spongioform	3 (4.1%)
Content of the left thyroid nodul	
Normal	53 (71.6%)
Solid	1 (1.4%)
Predominant solid	8 (10.8%)
Predominant cystic	7 (9.5%)
Cystic	3 (4.1%)
Spongioform	2 (2.7%)
Echogenicity of right thyroid nodul	
Normal	53 (71.6%)
Hypoechoic	13 (17.6%)
Isoechoic	2 (2.7%)
Hyperechoic	5 (6.8%)
Mixed	2 (2.7%)
Echogenicity of left thyroid nodul	
Normal	47 (63.5%)
Hypoechoic	17 (23.0%)
Isoechoic	3 (4.1%)
Hyperechoic	5 (6.8%)
Mixed	1 (1.4%)
Margin of the right mass	
Normal	47 (63.5%)
Smooth	24 (32.4%)
Ill defined	3 (4.1%)
Margin of the left mass	
Normal	53 (71.6%)
Smooth	19 (25.7%)
Ill defined	2 (2.7%)
Mass Calcification	
No calcification	70 (94.6)
Stipplen, fine coarse, non linear	2 (2.7%)
Curvilinear, smooth margin	2 (2.7%)

US = ultrasonography.

4. Discussion

Thyroid incidentalomas can be found in the imaging study for non-thyroid neck disease, on the US in the screening procedure, or in the histopathologic examination in the surgical specimens for non-nodular disease. The prevalence data from ultrasonographic studies were revealed ranging from 19% to 46% in the general population.^[15] The result was similar to this study of 43.2% (32 subjects) nodule detection among subjects with symptoms. Similar results of incidentalomas findings showed in the Choi study that was 37%.^[16] That being said, study of 2079 patients in Korea showed high sensitivity and specificity 96.2% and 51.7%, respectively.^[17] In the 253 randomly selected adults, 5.1% had abnormality of the neck with sensitivity and specificity to detect thyroid nodules were 11.6% and 97.3%, respectively.^[18] Of 1845 subjects, one of the study show cancer rate detection of 1.6% with sensitivity and specificity 100% and

Table 3
Comparison of patient complaint in normal patient and in patient with nodul.

	Normal (n = 42)	Nodul (n = 32)	p-value
Patient complaints			
Mass in the neck	6	14	< .01
Fatigue	13	13	> .05
Insomnia	10	6	> .05
Over sweating	6	9	> .05
Shaky	8	7	> .05

Table 4
Correlation of neck physical examination to US result.

Goitre palpation	Normal (n = 42)	Nodule (n = 32)	Coef. Correlation	P value
No palpable/ visible mass	37	15	0.773	< .01
Palpable but not visible mass	2	3		< .01
Palpable and visible mass	3	14		< .01

US = ultrasonography.

98.7%, respectively, showing good diagnostic performance of thyroid screening US.^[19]

The value of thyroid US study is still a debate due to several reasons including overdiagnosis which potentially cause overtreatment.^[20] This may lead to unnecessary procedures and expose the patient to the treatment side effect.^[21] The increased incidence and low mortality interpreted as a result of overdiagnosis.^[22] Moreover, study from Jun et al shows no difference in the mortality between the patient who went through screening and those who don't.^[23] The risk of overdiagnosis could be diminished by taking account into other factor such as size of thyroid nodules, presence of lymphadenopathy, cytology and molecular markers results, TSH measurement, US characteristics, patients' age, family history of thyroid cancer, and history of head and neck irradiation.^[6]

Routine screening population of low risk for thyroid cancer individual were unlikely to be cost effective and more useful in higher-risk individuals such as patient with radiation exposure, positive family history, lesser than 14 years or greater than 70 years of age with palpable thyroid nodule, and with suspicious clinical signs.^[24] With the advancement of the artificial intelligence, some models can be used to minimize false positives although with the cost of increasing false negative.^[25] Long-term controlled trials are a necessity to be performed to study the efficacy of US thyroid screening.

Chen et al stated the proper use of ultrasonography is when there are clinically supported features such as examination finding of a palpable thyroid nodule, large thyroid or a goiter, thyroid nodule findings from another imaging test, and new-onset hoarseness or compressive symptoms.^[26] Even so, the definition of inappropriate use of thyroid ultrasound still varies. It could be due to deemed unnecessary by reviewing endocrinologists, functional disease or nonspecific symptoms without specifying the presence of a palpable mass/ nodule, hypothyroidism, no history of thyroid nodule or neck irradiation.^[27] In this study, there was a significant difference in the symptoms of mass in the neck of the subjects ($P = .008$). Moreover, there was a strong correlation between goiter palpation dan US thyroid result ($R = 0.773$, $P = .00$). This result is in line with a study by Germano et al in which the incidentalomas were found almost as frequently as the palpable nodules demonstrating the important role of imaging.^[28]

Arpana et al showed USG study results of texture, size, margin, echogenicity, and vascularity are important factors for predicting thyroid malignancy but are not to be used independently as a screening tool.^[29] The combination of suspicious findings can help differentiate nodules that may require fine needle aspiration (FNA).^[30,31]

Iodine deficiency was one of the risk factors that must be taken into account in associating with increased rates of thyroid diseases.^[6] Indonesia was included in the category of adequate iodine nutrition (UIC 100–299 $\mu\text{g/L}$) based on national iodine status in 2014 through median urinary iodine concentration.^[32] The distribution of iodine status in Indonesian province, however, was not equally adequate. The risk of nodular goiter is only increased at the low intake due to the promotion of cell growth and DNA mutagenesis causing clusters of autonomous thyrocytes.^[10,32,33] However, iodine enrichment has been associated with increased thyroid cancer and shift from the follicular to papillary histotype.^[34]

There are several limitations in our study. Our study includes a small number of samples due to the short duration of the community services and the retrospective nature. Secondly, we did not perform further examination such as FNA or treatment of the subjects. There was no blinding in the subjects, and the distribution of the patients is mostly female due to the timing of the examination. The study was also being conducted in the rural area at a primary care facility, thus there was no laboratory facility to support this study. One of the goals of this study was to help screening of thyroid abnormality, especially at rural area without laboratory facility to predict thyroid function without the need of thyroid tests such as biochemical marker. This is also showed the importance of this study, since the result showed that there were significant differences in US result of patients with and without complaint of mass in the neck. We also found a strong correlation between goiter palpation and US examination. Therefore, it is important to do US examination for patient with complaint of mass in the neck and with goiter. Ultrasound can also be an alternative to laboratory limitation in the rural area because the modality of ultrasound is mobile, cheap, and noninvasive. Our suggestion for future study is there should be a screening thyroid study on the nation scale with a larger sample and correlating with the FNA or biopsy results and biochemical marker of thyroid function. The mortality rate of the thyroid cancer should be take into consideration to avoid overdiagnosis.

We conclude there were significant differences in US result of patients with and without complaint of mass in the neck. We also found a strong correlation between goiter palpation and US examination. Although US examination proves to be beneficial in nodule thyroid detection, other factors namely clinical, laboratory examination, cytology and molecular markers, patients' age, nodules size, and ultrasound features should be held into account before taking the further step of treatment.

13

Acknowledgments

The authors would like to thank all who have contributed to the process and completion of this report, including the teaching staffs and fellow residents of Department of Radiology of Faculty of Medicine, Universitas Airlangga, Dr Soetomo Academic General Hospital, Surabaya, Indonesia. We also want to thank Bajulmati primary care center, Banyuwangi, East Java, Indonesia.

Author contributions

Conceptualization: Rosy Setiawati, Tri Wulanhandarini, Fierly Hayati, Dyah Erawati, Merlin Guntur Jaya, Agnes Triana Basja.
Data curation: Rosy Setiawati, Fierly Hayati, Dyah Erawati, Merlin Guntur Jaya, Andi Ahmad Thoriq, Triana Mediyawati Wijaya, Galih Nur Ismiyati, Dyan Wahyu Kusumaningrum,

Belinda Koesmarsono, M. Ikhsan Nugroho, Silvi Yuliana, Syadza Zahrah Shedyta, Hendra Boy Situmorang.

Investigation: Rosy Setiawati, Fierly Hayati, Ahmad Thoriq, Triana Mediyawati Wijaya, Galih Nur Ismiyati, Dyan Wahyu Kusumaningrum, Belinda Koesmarsono, Agnes Triana Basja, M. Ikhsan Nugroho, Silvi Yuliana, Syadza Zahrah Shedyta, Hendra Boy Situmorang.

Methodology: Rosy Setiawati, Merlin Guntur Jaya.

Supervision: Dyah Erawati.

Validation: Rosy Setiawati.

Writing – original draft: Rosy Setiawati, Merlin Guntur Jaya.

Writing – review & editing: Rosy Setiawati, Tri Wulanhandarini, Merlin Guntur Jaya.

References

- [1] Pizzato M, Li M, Vignat J, et al. The epidemiological landscape of thyroid cancer worldwide: GLOBOCAN estimates for incidence and mortality rates in 2020. *Lancet Diabetes Endocrinol.* 2022;10:264–72.
- [2] Vigneri R, Malandrino P, Vigneri P. The changing epidemiology of thyroid cancer: why is incidence increasing?. *Curr Opin Oncol.* 2015;27:1–7.
- [3] Marcello MA, Malandrino P, Almeida JFM, et al. The influence of the environment on the development of thyroid tumors: a new appraisal. *Endocr Relat Cancer.* 2014;21:T235–54.
- [4] Frates MC, Benson CB, Charboneau JW, et al. Management of thyroid nodules detected at US: society of radiologists in ultrasound consensus conference statement. *Ultrasound Q.* 2006;22:231–8.
- [5] Wiest PW, Hartsborne MF, Inskip PD, et al. Thyroid palpation versus high-resolution thyroid ultrasonography in the detection of nodules. *J Ultrasound Med.* 1998;487:96.
- [6] Singh Ospina N, Iñiguez-Ariza NM, Castro MR. Thyroid nodules: diagnostic evaluation based on thyroid cancer risk assessment. *BMJ.* 2020;368:1–20.
- [7] Kang HW, No JH, Chung JH, et al. Prevalence, clinical and ultrasonographic characteristics of thyroid incidentalomas. *Thyroid.* 2004;14:29–33.
- [8] Ezzat S, Sarti DA, Cain DR, et al. Thyroid incidentalomas prevalence by palpation and ultrasonography background: thyroid nodules are commonly identified on autopsy examination. There are relatively few descriptions, however, of the frequency with which thyroid nodules are encountered. *2015;154:23–5.* Available at: <http://archinte.jamanetwork.com/>.
- [9] Carlé A, Krejbjerg A, Laurberg P. Epidemiology of nodular goitre. Influence of iodine intake. *Best Pract Res Clin Endocrinol Metab [Internet].* 2014;28:465–79.
- [10] Krohn K, Führer D, Bayer Y, et al. Molecular pathogenesis of euthyroid and toxic multinodular goiter. *Endocr Rev.* 2005;26:504–24.
- [11] Knobel M, Medeiros-Neto G. Relevance of iodine intake as a reputed predisposing factor for thyroid cancer. *Arq Bras Endocrinol Metabol.* 2007;51:701–12.
- [12] Kartono D, Atmarita A, Jahari AB, et al. The situation of urinary iodine concentration (Uic) among school age children, women at reproductive age and pregnant women in Indonesia: the analysis of Riskesdas 2013. *Gizi Indones.* 2017;39:49.
- [13] Bibbins-Domingo K, Grossman DC, Curry SJ, et al. Screening for thyroid cancer: US preventive services task force recommendation statement. *JAMA.* 2017;317:1882–7.
- [14] World Health Organization. Goitre as a determinant of the prevalence and severity of iodine deficiency disorders in populations. Geneva, Switzerland: World Health Organization. 2014.
- [15] Delivannis DA, Regina Castro M. Thyroid incidentalomas. *Contemp Endocrinol.* 2018;29:153–67.
- [16] Choi YJ, Park YL, Koh JH. Prevalence of thyroid cancer at a medical screening center: pathological features of screen-detected thyroid carcinomas. *Yonsei Med J.* 2008;49:748–56.
- [17] Kim SJ, Moon WK, Cho N. Sonographic criteria for fine-needle aspiration cytology in a Korean female population undergoing thyroid ultrasound screening. *Acta radiol.* 2010;51:475–81.
- [18] Brander A, Vilkinoski P, Nickels J. Thyroid gland: US screening in random adult population. *Radiology.* 1991;181:683–7.
- [19] Yoo J, Ahn HS, Kim SJ, et al. Evaluation of diagnostic performance of screening thyroid ultrasonography and imaging findings of screening-detected thyroid cancer. *Cancer Res Treat.* 2018;50:11–8.

- [20] Ali A, Alawad MA. Thyroid Cancer Screening: Is It Beneficial?. *Am J Otolaryngol Head Neck Surg*. 2018;1:1021. Available at: https://www.researchgate.net/publication/327944142_Thyroid_Cancer_Screening_Is_It_Beneficial.
- [21] Nys P, Cordray JP, Sarafian V, et al. Screening for thyroid cancer according to French recommendations with thyroid ultrasound in newly diagnosed Graves' disease without palpable nodule is not useful. *Ann Endocrinol (Paris)* [Internet]. 2015;76:13–8.
- [22] Grani G. Inappropriate use of thyroid ultrasound is common in clinical practice. *Clin Thyroidol*. 2022;34:23–5.
- [23] Jun JK, Hwang SY, Hong S, et al. Association of screening by thyroid ultrasonography with mortality in thyroid cancer: a case-control study using data from two national surveys. *Thyroid*. 2020;30:396–400.
- [24] Li Q, Lin X, Shao Y, et al. Imaging and screening of thyroid cancer. *Radiol Clin North Am* [Internet]. 2017;55:1261–71.
- [25] Idarraga AJ, Luong G, Hsiao V, et al. False negative rates in benign thyroid nodule diagnosis: machine learning for detecting malignancy. *J Surg Res* [Internet]. 2021;268:562–9.
- [26] Chen DW, Reyes-Gastelum D, Radhakrishnan A, et al. Physician-reported misuse of thyroid ultrasonography. *J Am Med Assoc*. 2020;155:984–6.
- [27] Edwards MK, Iñiguez-Ariza NM, Singh Ospina N, et al. Inappropriate use of thyroid ultrasound: a systematic review and meta-analysis. *Endocrine* [Internet]. 2021;74:263–9.
- [28] Germano A, Schmitt W, Almeida P, et al. Ultrasound requested by general practitioners or for symptoms unrelated to the thyroid gland may explain higher prevalence of thyroid nodules in females. *Clin Imaging* [Internet]. 2018;50:289–93.
- [29] Arpana PO, Gurung G, Pradhan S. Ultrasound findings in thyroid nodules: a radio-cytopathologic correlation. *J Med Ultrasound*. 2018;26:90–3.
- [30] Kim JY, Lee CH, Kim SY, et al. Radiologic and pathologic findings of nonpalpable thyroid carcinomas detected by ultrasonography in a medical screening center. *J ultrasound Med* [Internet]. 2008;27:215–23.
- [31] Hoang JK, Wai KL, Lee M, et al. US features of thyroid malignancy: pearls and pitfalls. *Radiographics*. 2007;27:847–60.
- [32] Zimmermann MB, Boelaert K. Iodine deficiency and thyroid disorders. *Lancet Diabetes Endocrinol*. 2015;3:286–95.
- [33] Laurberg P, Cerqueira C, Ovesen L, et al. Iodine intake as a determinant of thyroid disorders in populations. *Best Pract Res Clin Endocrinol Metab*. 2010;24:13–27.
- [34] Dong W, Zhang H, Zhang P, et al. The changing incidence of thyroid carcinoma in Shenyang, China before and after universal salt iodization. *Med Sci Monit*. 2013;19:49–53.

Clinical and ultrasonography evaluation of thyroid tumor screening in symptomatic patient of Bajulmati primary care center, Banyuwangi, East Java, Indonesia

ORIGINALITY REPORT

18%

SIMILARITY INDEX

12%

INTERNET SOURCES

17%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1	mafiadoc.com Internet Source	1%
2	www.researchsquare.com Internet Source	1%
3	Meyer Knobel, Geraldo Medeiros-Neto. "Relevance of iodine intake as a reputed predisposing factor for thyroid cancer", Arquivos Brasileiros de Endocrinologia & Metabologia, 2007 Publication	1%
4	Wei Yang, Ghaneh Fananapazir, Jennifer LaRoy, Machel Wilson, Michael J. Campbell. "Can the American Thyroid Association, K-Tirads, and Acr-Tirads Ultrasound Classification Systems Be Used to Predict Malignancy in Bethesda Category IV Nodules?", Endocrine Practice, 2020 Publication	1%

5	Won-Jin Moon, Jung Hwan Baek, So Lyung Jung, Dong Wook Kim et al. "Ultrasonography and the Ultrasound-Based Management of Thyroid Nodules: Consensus Statement and Recommendations", Korean Journal of Radiology, 2011 Publication	1 %
6	cardiothoracicsurgery.biomedcentral.com Internet Source	1 %
7	Zimmermann, Michael B.. "Thyroid gland: Iodine deficiency and thyroid nodules", Nature Reviews Endocrinology, 2014. Publication	1 %
8	www.ncbi.nlm.nih.gov Internet Source	1 %
9	Matthew K. Edwards, Nicole M. Iñiguez-Ariza, Naykky Singh Ospina, Eddy Lincango-Naranjo, Spyridoula Maraka, Juan P. Brito. "Inappropriate use of thyroid ultrasound: a systematic review and meta-analysis", Endocrine, 2021 Publication	1 %
10	file.scirp.org Internet Source	1 %
11	zagan.unizar.es Internet Source	1 %

12 Liguang Zhou, Bo Chen, Miaoqing Zhao, Huawei Zhang, Bo Liang. "Sonographic Features of Medullary Thyroid Carcinomas According to Tumor Size", Journal of Ultrasound in Medicine, 2015
Publication 1 %

13 f1000research.com
Internet Source 1 %

14 Hahn Wook Kang, Jung Heyun No, Jae Hoon Chung, Yong-Ki Min, Myung-Shik Lee, Moon-Kyu Lee, Jung Hyun Yang, Kwang-Won Kim. "Prevalence, Clinical and Ultrasonographic Characteristics of Thyroid Incidentalomas", Thyroid, 2004
Publication <1 %

15 Schwarz, Nicolas T., and Karl-Heinz Reutter. "General Treatment Approach", General and Visceral Surgery Review, 2012.
Publication <1 %

16 worldwidescience.org
Internet Source <1 %

17 doaj.org
Internet Source <1 %

18 www.hindawi.com
Internet Source <1 %

19

Ana Germano, Willian Schmitt, Pedro Almeida, Rui Mateus-Marques, Valeriano Leite.

"Ultrasound requested by general practitioners or for symptoms unrelated to the thyroid gland may explain higher prevalence of thyroid nodules in females", *Clinical Imaging*, 2018

Publication

<1 %

20

surgicalneurologyint.com

Internet Source

<1 %

21

www.scopus.com

Internet Source

<1 %

22

Fangqiong Luo, Lei Huang, Xiuqun Gong, Zhiyu Han, Fangyi Liu, Zhigang Cheng, Jianping Dou, Xiaoling Yu, Ping Liang, Jie Yu.

"Microwave ablation of benign thyroid nodules: 3 - year follow - up outcomes", *Head & Neck*, 2021

Publication

<1 %

23

Hyobin Seo, Kwang Nam Jin, Ji Sang Park, Koung Mi Kang, Eun Kyung Lee, Ji Ye Lee, Roh-Eul Yoo, Young Joo Park, Ji-hoon Kim. "Risk of thyroid cancer in a lung cancer screening population of the National Lung Screening Trial according to the presence of incidental thyroid nodules detected on low-dose chest CT", *Ultrasonography*, 2023

Publication

<1 %

24

Respati S. Dradjat, Panji Sananta, Rizqi Daniar Rosandi, Lasa Dhakka Siahaan. "Osteocalcin biomarker level evaluation on fracture healing with bone defect after stromal vascular fraction application in murine model", *Annals of Medicine and Surgery*, 2021

Publication

<1 %

25

www.unboundmedicine.com

Internet Source

<1 %

26

Maria Mathew D'Souza, Raman Kumar Marwaha, Rajnish Sharma, Abhinav Jaimini et al. "Prospective evaluation of solitary thyroid nodule on 18F-FDG PET/CT and high-resolution ultrasonography", *Annals of Nuclear Medicine*, 2010

Publication

<1 %

27

jamanetwork.com

Internet Source

<1 %

28

Luu-Ngoc Do, Hyo-Jae Lee, Chaeyeong Im, Jae Hyeok Park, Hyo Soon Lim, Ilwoo Park. "Predicting Underestimation of Invasive Cancer in Patients with Core-Needle-Biopsy-Diagnosed Ductal Carcinoma In Situ Using Deep Learning Algorithms", *Tomography*, 2022

Publication

<1 %

29

revistasylibrosmedicos.com

Internet Source

<1 %

30

www.bmj.com

Internet Source

<1 %

31

www.imrpress.com

Internet Source

<1 %

32

www.tandfonline.com

Internet Source

<1 %

33

www1.racgp.org.au

Internet Source

<1 %

34

zums.ac.ir

Internet Source

<1 %

35

Giorgio Grani. "Inappropriate Use of Thyroid Ultrasound Is Common in Clinical Practice", *Clinical Thyroidology*, 2022

Publication

<1 %

36

Hayashida, Naomi, Misa Imaizumi, Hiroki Shimura, Noriyuki Okubo, Yasushi Asari, Takeshi Nigawara, Sanae Midorikawa, Kazuhiko Kotani, Shigeyuki Nakaji, Akira Otsuru, Takashi Akamizu, Masafumi Kitaoka, Shinichi Suzuki, Nobuyuki Taniguchi, Shunichi Yamashita, and Noboru Takamura. "Thyroid Ultrasound Findings in Children from Three Japanese Prefectures: Aomori, Yamanashi and Nagasaki", *PLoS ONE*, 2013.

<1 %

37 S. Gaengler, X.D. Andrianou, A. Piciu, P. Charisiadis, C. Zira, K. Aristidou, D. Piciu, K.C. Makris. "Iodine status and thyroid nodules in females: a comparison of Cyprus and Romania", *Public Health*, 2017

Publication

38 Yukitoshi Shimamura, Masahiro Kanayama, Fumihiro Oha, Takeru Tsujimoto et al. "Pre-existing adjacent level foraminal stenosis does not affect the outcome of a single level lumbar interbody fusion", *Journal of Orthopaedic Science*, 2022

Publication

39 repository.unair.ac.id

Internet Source

40 www.scielo.br

Internet Source

41 "Management of Thyroid Nodules and Differentiated Thyroid Cancer", Springer Science and Business Media LLC, 2017

Publication

42 "Thyroid Cancer", Springer Nature, 2016

Publication

43 Congqing Lu, Yan Wang, Ming Yu. "Is ultrasonographic evaluation sensitive enough

to detect multicentric papillary thyroid carcinoma?", Gland Surgery, 2020

Publication

44

Endocrine Updates, 2012.

Publication

<1 %

45

Tingting Li, Zirui Jiang, Man Lu, Shibin Zou, Minggang Wu, Ting Wei, Lu Wang, Juan Li, Ziyue Hu, Xueqing Cheng, Jifen Liao.

"Computer-aided diagnosis system of thyroid nodules ultrasonography", Medicine, 2020

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Clinical and ultrasonography evaluation of thyroid tumor screening in symptomatic patient of Bajulmati primary care center, Banyuwangi, East Java, Indonesia

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5
