



Source details

Indian Journal of Forensic Medicine and Toxicology

Scopus coverage years: from 2008 to 2021

(coverage discontinued in Scopus)

Publisher: Medico Legal Society

ISSN: 0973-9122 E-ISSN: 0973-9130

Subject area: Social Sciences: Law Medicine: Pathology and Forensic Medicine

Pharmacology, Toxicology and Pharmaceutics: Toxicology Environmental Science: Health, Toxicology and Mutagenesis

Source type: Journal

CiteScore 2020

0.1



SJR 2020

0.115



SNIP 2021

0.078



[View all documents >](#)

[Set document alert](#)

[Save to source list](#) [Source Homepage](#)

[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)

i Improved CiteScore methodology



CiteScore 2020 counts the citations received in 2017-2020 to articles, reviews, conference papers, book chapters and data papers published in 2017-2020, and divides this by the number of publications published in 2017-2020. [Learn more >](#)

CiteScore 2020

$$0.1 = \frac{387 \text{ Citations } 2017 - 2020}{3,509 \text{ Documents } 2017 - 2020}$$

Calculated on 05 May, 2021

CiteScore rank 2020

Category	Rank	Percentile
Social Sciences		
└ Law	#639/722	11th
Medicine		
└ Pathology and Forensic Medicine	#183/191	4th
Pharmacology,		

[View CiteScore methodology >](#) [CiteScore FAQ >](#) [Add CiteScore to your site](#)

About Scopus

[What is Scopus](#)

[Content coverage](#)

[Scopus blog](#)

[Scopus API](#)

[Privacy matters](#)

Language

[日本語版を表示する](#)

[查看简体中文版本](#)

[查看繁體中文版本](#)

[Просмотр версии на русском языке](#)

Customer Service

[Help](#)

[Tutorials](#)

[Contact us](#)

ELSEVIER

[Terms and conditions](#) ↗ [Privacy policy](#) ↗

Copyright © Elsevier B.V. ↗. All rights reserved. Scopus® is a registered trademark of Elsevier B.V.

We use cookies to help provide and enhance our service and tailor content. By continuing, you agree to the use of cookies ↗.





ESCI & PubMed Indexed

Publish in Orthopedic Research and Reviews & get 25% off with code DIGIT

Dove Medical Press

[Learn More](#)

Indian Journal of Forensic Medicine and Toxicology

Discontinued in Scopus as of 2020

COUNTRY

India



Universities and research institutions in India

SUBJECT AREA AND CATEGORY

Environmental Science
Health, Toxicology and Mutagenesis

Medicine
Pathology and Forensic Medicine

Pharmacology, Toxicology and Pharmaceutics
Toxicology

Social Sciences
Law

PUBLISHER

Institute of Medico-legal Publications

H-INDEX

12

PUBLICATION TYPE

Journals

ISSN

09739122, 09739130

COVERAGE

2008-2021

INFORMATION

[Homepage](#)

[How to publish in this journal](#)

editor.ijfmt@gmail.com

ESCI & PubMed Indexed

Publish in Orthopedic Research and Reviews & get 25% off with code DIGIT

Quick Turnaroun Times

International Journal of General Medicine. Indexed on PubMed and has an Impact Factor



Dove

SCOPE

“Indian Journal of Forensic Medicine & Toxicology ” is a double-blind peer reviewed international journal. The frequency is quarterly. It deals with Forensic Medicine, Forensic Science, Toxicology, DNA fingerprinting, sexual medicine, environmental medicine, Forensic Pathology, legal medicine and public health laws.



Join the conversation about this journal

IF & PubMed Indexed

International Journal of General Medicine. Indexed on PubMed and has an Impact Factor

Dove Medical Press

[Learn More](#)



Quartiles




IF & PubMed Indexed

International Journal of General
Medicine. Indexed on PubMed and
has an Impact Factor


Dove Medical Press

[Learn More](#)


FIND SIMILAR JOURNALS ?




SJR



Total Documents



Total Cites Self-Cites




Citations per document



External Cites per Doc Cites per Doc




% International Collaboration




Citable documents

Non-citable documents



Cited documents

Uncited documents



Indian Journal of Forensic Medicine and Toxicology

Not yet assigned quartile

SJR 2021

0

powered by scimagojr.com


← Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scimagojr.com" data-bbox="612 318 757 332">
```

SCImago Graphica

Explore, visually communicate and make sense of data with our **new data visualization tool**.



IF & PubMed Indexed

International Journal of General Medicine. Indexed on PubMed and Impact Factor

Dove Medical Press

[Learn More](#)

Metrics based on Scopus® data as of April 2022



Editorial Team

Editor in Chief

Prof S K Dhattarwal

Forensic Medicine, PGIMS, Rohtak, Haryana

E-mail: editor.ijfmt@gmail.com

EDITORIAL ADVISORY BOARD

1. Prof Sudhir K Gupta, Head, FMT. AIIMS, New Delhi , India
2. Prof Mete Gulmen ,Cukurova University, TURKEY
3. Prof. Leandro Duarte De Carvalho , Minas Gerais, Belo Horizonte, BRAZIL
4. Dr. Valery Gunas, National Pirogov Memorial Medical University,Vinnytsya, UKRAINE
5. Dr. Rahul Pathak Forensic Science, Dept of Life Sciences ,Anglia Ruskin University, Cambridge, UNITED KINGDOM
6. Prof Emilo Nuzzalese, University of Turin , Italy
7. Dr Noha A. Magdy Elrafie, Forensic Toxicology, Ain Shams University, Cairo, EGYPT
8. Dr Rituja Sharma, Associate Prof, Law, Banasthali Vidyapeeth Jaipur
9. Dr Shankar Bakkanwar (*Associate Professor*) Forensic Medicine, Kasturba Medical College, Manipal, Karnataka
10. Dr K. Ravikumar , Raksha Shakti University, Ahmedabad, Gujrat.
11. Dr C. Behera (*Addl. Prof*) Dept of FMT, AIIMS, New Delhi
12. Dr. Kanak Lata Verma, Deputy Director, Toxicology ,RFSL, Chanakyapuri New Delhi
13. Dr. Asha Srivastava (*Senior Scientific Officer*) Forensic Psychology,Central Forensic Science Laboratory, CBI, Delhi
14. Dr. Raghvendra Kumar Vidua, (Associate Prof), FMT, AIIMS Bhopal
15. Dr. Vaibhav Saran (*Asst.Prof.*) School of Forensic Science, Sam Higginbottom Institute of Agriculture Technology & Sciences,Allahabad
16. Ms Aparna R. Asst. Prof. Forensic Serology & Biology, Jain University, Bengaluru
17. Dr. Deepali Jain, Asst Prof, Forensic Science ,BB Ambedkar University, Lucknow
18. Prof. NK Aggrawal Forensic Medicine, UCMS, Delhi
19. Prof. Manoj Kumar Mohanty, Forensic Medicine, AIIMS, Bhuvneshwar, Odisha
20. Prof. Amar Jyoti Patowary, Forensic Medicine, NEIGRIHMS, Shillong, Meghalaya
21. Prof S. Venkata Raghava , Forensic Medicine, Banglore Medical College, Bengaluru

22. Prof. Shalini Gupta Oral Pathology and Microbiology, Faculty of Dental Sciences, King George Medical University, Lucknow
23. Prof. Virender Kumar Chhoker Forensic Medicine, Santosh Medical College, Ghaziabad, UP
24. Prof. Dayanand G Gannur , Forensic Medicine , Shri BM Patil Medical College, Hospital & Research centre, Bijapur, Karnataka
25. Prof Praveen Arora, Forensic Medicine, SAIMS, Indore, MP
26. Prof Barkha Gupta , Saraswathi Institute of Medical Sciences Hapur, Uttar Pradesh India
27. Prof M Prashant Apollo Medical College Hyderabad
28. Prof Dimple Patel , Anatomay, AMC MET Medical College , Ahmedabad , Gujarat
29. Dr Mohammed Nader Shalaby, Associate Professor of Biological Sciences and Sports Health Department, Faculty of Physical Education, Suez Canal University, Egypt

30 Dr. Avinash Harishchandra Waghmode

Professor and Head, Dept of Forensic Medicine and Toxicology, BKL Walawalkar Rural
Medical College Chiplun Ratnagiri

31 Dr. Anita Yadav Assistant Professor Forensic Science, SBAS Galgotias University, Greater
Noida, UP

32 Dr. Risha Jasmine Nathan *Lecturer in Forensic Chemistry Anglia Ruskin University*

East Road, Cambridge, CB1 1PT England, United Kingdom

Current Issue

ATOM 1.0

RSS 2.0

RSS 1.0

[Make a Submission](#)

Browse

[Open Journal Systems](#)

ISSN-0973-9122 (Print) • ISSN-0973-9130 (Electronic)

Volume 16

Number 1

January-March 2022



Indian Journal of Forensic Medicine & Toxicology

Website: www.ijfmt.com

Official Organ of Indian Association of Medico-Legal Experts (Regd.)

Indian Journal of Forensic Medicine & Toxicology

Editor in Chief

Prof. SK Dhattarwal

Forensic Medicine, PGIMS, Rohtak, Haryana

INTERNATIONAL EDITORIAL ADVISORY BOARD

1. **Prof Mete Gulmen** Cukurova University, TURKEY
2. **Prof. Leandro Duarte De Carvalho**, Minas Gerais, Belo Horizonte, Brazil
3. **Prof. Donata Favretto** (Full Professor) Forensic Toxicology at University of Padova, Italy
4. **Prof. Babak Mostafazadeh** Department of Forensic Medicine & Toxicology, Shahid Beheshti University of Medical Sciences, Tehran-Iran
5. **Prof Halis Dokgoz**, Mersin University, TURKEY
6. **Prof Jozef Sidlo**, Comenius University, Bratislava, SLOVAKIA
7. **Dr. Rahul Pathak** (Lecturer) Forensic Science, Dept of Life Sciences Anglia Ruskin University, Cambridge, United Kingdom
8. **Dr. Hareesh** (Professor & Head) Forensic Medicine, Ayder Referral Hospital, College of Health Sciences, Mekelle University, Mekelle Ethiopia East Africa
9. **Dr. Mokhtar Ahmed Alhrani** (Specialist) Forensic Medicine & Clinical Toxicology, Director of Forensic Medicine Unit, Attorney General's Office, Sana'a, Yemen
10. **Dr. Sarathchandra** Kodikara (Senior Lecturer) Forensic Medicine, Department of Forensic Medicine, Faculty of Medicine, University of Peradeniya, Sri Lanka
11. **Dr Noha A. Magdie El Rafie**, Forensic Toxicology, Ain Shams University, Cairo, EGYPT

SCIENTIFIC COMMITTEE

1. **Prof Udai Pratap Singh**, Department of Anthropology Lucknow University Lucknow
2. **Dr Anil Rahule** (Associate Professor) Dept of Anatomy, Govt Medical College Nagpur
3. **Dr Shankar Bakkanwar** (Associate Professor) Forensic Medicine, Kasturba Medical College, Manipal, Karnataka
4. **Dr K. Ravikumar** Raksha Shakti University, Ahmedabad, Gujrat.
5. **Dr. Pragnesh Parmar** (Associate Professor) Forensic Medicine, Valsad, Gujrat
6. **Dr Vandana Mudda** (Awati) (Associate Prof) Dept of FMT, M.R.Medical College, Gulbarga, Karnataka.
7. **Dr. Asha Srivastava** (Senior Scientific Officer) Forensic Psychology, Central Forensic Science Laboratory, CBI, Delhi
8. **Dr. Lav Kesharwani** (Asst.Prof.) School of Forensic Science, Sam Higginbottom Institute of Agriculture Technology & Sciences, Allahabad U.P.

"Indian Journal of Forensic Medicine & Toxicology" is peer reviewed quarterly journal. It deals with Forensic Medicine, Forensic Science, Toxicology, DNA fingerprinting, sexual medicine and environment medicine. It has been assigned International standard serial No. p-0973-9122 and e- 0973-9130. The Journal has been assigned RNI No. DELENG/2008/21789. The journal is covered by EMBASE (Excerpta Medica Database). The journal is also abstracted in Chemical Abstracts (CAS) database (USA). The journal is also covered by EBSCO (USA) database. It is now official publication of Indian Association of Medico-Legal Experts (Regd.).

9. **Dr. Anu Sharma** (Associate Prof) Dept of Anatomy, DMCH, Ludhiana (PB)
10. **Dr. Shalini Gupta** (Prof) Oral Pathology and Microbiology, Dental Sciences King George Medical University, Lucknow, UP
11. **Dr Rituja Sharma**, Associate Prof, Law Banasthali Vidyapeeth Jaipur

NATIONAL EDITORIAL ADVISORY BOARD

Chairman

Prof Sudhir K Gupta - Head, Department of Forensic Medicine
All India Institute of Medical Sciences, New Delhi

Members

1. **Prof. N K Aggrawal** Forensic Medicine, UCMS, Delhi
2. **Prof Ajay Ghangale** Forensic Medicine Dr DY Patil Medical College, Pune, Maharashtra
3. **Dr. Amar Jyoti Patwory** Professor, Forensic Medicine NEIGRIHMS, Shillong
4. **Dr S. Venkata Raghava** Professor, Forensic Medicine, Bangalore Medical College, Bengaluru
5. **Prof Praveen Arora**, Professor Department of Forensic Medicine & Toxicology, SAIMS, Indore
6. **Dr. Pankaj Datta** (Principal & Head) Department of Prosthodontics, Indraprastha Dental College & Hospital, Ghaziabad
7. **Dr. Mahindra Nagar** (Head) Department of Anatomy, UCMS & GTB Hospital, Delhi
8. **Dr. Virender Kumar Chhoker** Professor Forensic Medicine and Toxicology, Santosh Medical College, Ghaziabad, UP
9. **Dr. Dayanand G Gannur** (Professor) Department of Forensic Medicine & Toxicology, Shri BM Patil Medical College, Hospital & Research centre, Bijapur, Karnataka
10. **Dr. Alok Kumar** Professor Department of Forensic Medicine & Toxicology, UP Rural Institute of Medical Sciences and Research, Saifai, Etawah, U.P.

Print-ISSN:0973-9122 Electronic - ISSN: 0973-9130

Frequency: Quarterly, © All Rights reserved The views and opinions expressed are of the authors and not of the Indian Journal of Forensic Medicine & Toxicology. Indian Journal of Forensic Medicine & Toxicology does not guarantee directly or indirectly the quality or efficacy of any products or service featured in the advertisement in the journal, which are purely commercial.

Website: www.ijfmt.com

Published at

Institute of Medico-legal Publications

Logix Office Tower, Unit No. 1704, Logix City Centre Mall,
Sector- 32, Noida - 201 301 (Uttar Pradesh)



1.	Deleted.....	1
2.	Antibacterial Activity of (+) Usnic Acid against Multi Drug Resistant <i>Acinetobacter baumannii</i> from Clinical Isolates <i>B. Nagaraju, K. Fathimunnisa, R. Vijayaraghavan, B. Sreekanth</i>	11
3.	A Comparative study to Evaluate the Efficacy of Supervised Exercise Program and Cyriax Physiotherapy on Pain and Function in Lateral Epicondylitis..... <i>Gouri Kalaskar, Pratik Phansopkar</i>	18
4.	Deleted	25
5.	A Study to Assess Respiratory Health Status among Bricks Workers <i>Jyot B. Darji, Siddaram Sarate</i>	29
6.	Study of Ossification Centres Around the Elbow and Wrist of Adolescent Age Group 15-18 Years..... <i>Kishore Kumar.G, R.Jenifer Jacquiline</i>	33
7.	Cheiloscopy - A Tool For Identification in Twins <i>M.Sugatha, R.Jenifer Jacquiline, M.Taqiuddin Khan</i>	41
8.	Synthesis, Characterization, Molecular Docking of Sulphamethoxazole Schiff Base Metal Complexes and Its Antibacterial, Anti-Inflammatory and Anti Depressant Activity <i>R.Gandhimathi, Magdivari Sangeetha</i>	47
9.	Aortic Atherosclerosis and Risk Factors: A Clinico-Demographic Autopsy-Based Study..... <i>Mainak Tarafder, Suparna Datta, Ranjan Biswas, Prabir Chakraborty, Saptarshi Chatterjee</i>	58
10.	Estimation of Stature & Gender from Thumb Indices in Indian Population <i>Mohite Hemlata S, Kakade Satish V, Mohite Sandeep S, Roy Priya P, Doshi Megha A</i>	66
11.	Sudden Death with Dual Organ Pathologies–An Autopsy -based Case Report <i>Mukul Sharma, Kimi Soumya Padhi, Govinda Balmuchu, Prachi Nemani, ManasRanjan Sahu</i>	76
12.	Demographic Study of Unnatural Deaths in Paediatric Age Group in General Hospital, Khammam <i>N. Devchand, A. Krishnanjaneyulu, Raja Sekhar Uppu, Uday Pal Singh</i>	83

II

13. Emergence of Synthetic Cannabinoids as Drugs of Abuse..... 89
Nupur Joshi, Mahipal Singh Sankhla
14. A Study of Metacognition in Adults with Type II Diabetes Mellitus – A Cross Sectional Study..... 98
S.Dhiviyashri, A.Sangeetha, Kumaresan M, A.V.Anuja, Karthika M
15. Anti-Smoking Campaign – An Innovative Initiative against Smoking by Tertiary Care Teaching Institute of North India 104
Shweta Talati, Pankaj Arora I, Anil Kumar Gupta, Prem Chand Sharma
16. The Economic Impact of Government Sponsored Health Insurance Schemes on a Tertiary Care Multispeciality Hospital in South India 110
Sri Theja Karra, Rajesh Kamath, Charan Tej Koyi, Renu Mirchandani, Kamath Madhusudana, Biju Soman
17. Hate Speech Against Disabled Persons: A Forensic Analysis of Legal Framework in India and Beyond 120
Suvidutt M.S., Aditya Tomer
18. Assessment and Evaluation of Implantology Courses for Post Graduate Students 126
Swati M Devani, Arvind I Moldi, Shrikar Desai, Amol Sangewar, Gaurav Khemaria, Vijay Namdevrao Yannawar, Shubhangi Lokhande
19. Effectiveness of Learning Software for Teaching Curricula and their Influence on Developing Thinking Skills and Achievement among Senior Students 133
Israa Yaseen Abdul-Kareem, Ban Abdul-Rahman Ibraheem, Iman Hamad Shihab
20. Health Satisfaction in Health Care between Patients of 3rd Tertiary Allergy Center in Kirkuk with Salam Primary Health Care Center Patients in Kirkuk..... 140
Abdulameer Anwar Samad, Fakher Abobaker Ahmed Gli, Chinar Baqir Kanbar, Diyar Mohammed Majeed
21. Strengthening Human Resources as Lecturers Through the E-Learning Learning Method..... 149
Agus Supinganto, Suharmanto, Irwan Budiana, Kusniyati Utami
22. Study of Lipid Profile, Lipid Peroxidation and DNA Damage in Patients with Coronary Artery Disease 155
Ahmed Jabbar AL Jabbiry, Kadhum J. Gattia, Hazim I. Abd Albari
23. The Effect of Aquatic Exercises on Special Endurance (Strength and Speed) and Achievement Run (800m) for Youth..... 162
Ahmed Elewy Arhem, MazinEnhier Lami
24. Association of Adipoq +45 T>G Gene Polymorphism with Insulin Resistance and Icam-1 Value in Obese Adolescents 168
Alexander Leonard Caesar Josediputra, Nur Aisiyah Widjaja, Roedi Irawan
25. Hemorrhagic Varicella in a 41-Year-Old Woman with Evans Syndrome : Case Report..... 176
Alfina Rahma, Eka Devinta Novi Diana, Frieda, Wibisono Nugraha, Muhammad Eko Irawanto, Moerbono Mochtar

26.	Estimation of Interleukin-10 and Interleukin-22 Levels in the Advances of Breast Cancer	184
	<i>Ali Jabbar Abd Zaid Al-Kilabi, Ghada Basil Ali Alomashi</i>	
27.	Occupational Health Hazards among Workers at Government Garment Factory in-Al Najaf City.....	191
	<i>Ali Sadoon Almusawy, Murtadha K. Adea Aljebory</i>	
28.	Relationship between Parenting Stress and Risk of Attention Deficit Hyperactivity Disorder (ADHD) in Elementary School Children.....	196
	<i>Alma Rossabela Setyanisa, Yunias Setiawati, Irwanto, Izzatul Fithriyah, Satria Arief Prabowo</i>	
29.	Quantitative Risk Assesment of Benzene Exposure in Printing Industry X Surabaya City	204
	<i>Amanda Fithri Habibati, Abdul Rohim Tualeka, Juliana Jalaludin, Syamsiar Russeng, Ahsan, Puji Rahmawati</i>	
30.	Age Estimation with Cemental Incremental Lines in Normal And Periodontally Diseased Teeth Using Phase Contrast Microscope	213
	<i>Ameena Sultana, Heena Zainab, Pramod Jahagirdar, Deepa Hugar, Shaimaa</i>	
31.	Vascular Endothelial Growth Factor (VEGF) Expression on Placenta Accreta Spectrum (PAS) FIGO Grading	224
	<i>Ana Puji Rahayu, Agus Sulistyono, Ernawati, Grace Ariani, Rozi Aditya Aryananda</i>	
32.	Sexual Dimorphism and Clinical Relevance of Thickness and Angulation of Lateral Lamella of Cribriform Plate-A Multidetector Computed Tomographic Study.....	230
	<i>Anandagouda V Naikanur, Balappa M Bannur, Manjula Patil, Sanjeev I Kolagi</i>	
33.	Effectiveness of Pisang Raja Peel Extract (Musa Paradisiaca L) on Bacterial Growth of Porphyromonas Gingivalis as the Cause of Periodontitis.....	237
	<i>Andy Fairuz Zuraida Eva, Masriadi, Nur Rahmah Hasanuddin, Andi Tenri Biba Mallombasang, Muhammad Ajis</i>	
34.	Measurement of Maximal Oxygen Uptake (VO ₂ Max) as a Cardiorespiratory Physiological Fitness Parameter Using Sensor Technology-Based Device Development.....	244
	<i>Anton Komaini, Wiyonna Gusvominesia, Syahrial Bakhtiar, Yendrizar, Novadri Ayubi</i>	
35.	Alpha-Mangostin and Gamma-Mangostin Isolated from Mangosteen (Garcinia mangostana L.) as Promising Candidates against SARS-CoV-2: A Bioinformatics Approach	251
	<i>Arif Nur Muhammad Ansori</i>	
36.	Prevalence of Musculoskeletal Disorder in Wrist and Fingers among Amateur Piano Players in Vellore.....	259
	<i>Ashish Mathew A, S.F.Mariyam Farzana</i>	
37.	Estimate of Biological Activity of Parsley extract on the Isolated Pathogenic Bacteria in Baghdad City.....	267
	<i>Ashwak Jasim Kzar Shammari</i>	
38.	Psychological Autopsy: A Lead to the Truth Untold	274
	<i>Mohd Asrarul Haque</i>	

IV

39. Diplopia as the Initial Manifestation of Cerebral Vasculitis in a Patient with Systemic Lupus Erythematosus: Diagnostic Approach and Challenges 278
Audy Ariana, Awalia
40. The Levels of Salivary IgA and Lactoferrin and Some Salivary Parameters in Waterpipe Smokers and Cigarette Smokers 283
Ausama Ahmed Fathallah, Maha Adel Mahmood
41. Incidence of Abruption and Placenta Previa in Pregnant with Previous Caesarian Section..... 291
Baidaa Abdulkareem Alwan, Areej Sabah Abdulridha
42. Levels of Suicidal Ideations and Intents among the Inmates in the Correctional Institutions in Baghdad City - Iraq..... 298
Maan Hameed Ibrahim Al-Ameri
43. Relationship between Vitamin D AND IL6 in Convalescent Healthcare Workers with Covid-19 in Baquba Hospitals in Diyala Province 306
Maysam Abbas, Luma T. Ahmed, Mohammed A. Alkarkhi
44. A Clinical Study of Parasitical Leishmaniasis in Al-Kut Province..... 313
Bushra Qasim Dhumad Al Edhari
45. Determinant Factors affecting Quality of Life of Children with HIV/AIDS..... 318
Candra Maulana, Irwanto, Dominicus Husada
46. Study of Psychosocial Risks in the Professional Environment of Health Care Workers in Morocco... 327
Chakhtoura Khalid, Chaib Yassine, Elanssari Anas, Aboussaleh Youssef, Ahmed O.T.Ahami
47. The Epistemological Aspects of Investigative Hypotheses 337
David Petretei
48. CT Scan Finding Characteristics of Confirmed Covid-19 Patients Based on Clinical Symptom Onset Patterns 343
Dewi Roziqo, Anita Widyoningroem, Prijambodo, Rosy Setiawati
49. Isolation and Diagnosis of the Bacteria Causing Corneal Ulceration Associated With Ocular Myiasis Infection 352
Dhuha Mahdi Jabir, Esraa Fadhel Wathah
50. Study the Effect of Many Medications in Ophthalmomyiasis Treatment and the Possibility of Using these Medications Instead of Mechanical Removal 358
Dhuha Mahdi Jabir, Ghada Basil Ali Alomashi
51. Physical and Psychological Stressor Exposure during Pregnancy Impacts the Expression of Synapsin and Neuronal Cells Number of MUS Musculus Offspring..... 366
Diana Estu Rumahastuti, Hermanto Tri Joewono, Widjiati
52. Distribution of Virulence genes in Streptococcus pneumoniae Isolated from Different Baghdad Hospitals..... 373
Dina Muthana Faghan, Basima Qasim Hasan Al-Saadi

CT Scan Finding Characteristics of Confirmed Covid-19 Patients Based on Clinical Symptom Onset Patterns

Dewi Roziqo¹, Anita Widyoningroem², Prijambodo², Rosy Setiawati²

¹Resident, ²Radiologist, Department of Radiology, Faculty of Medicine, Airlangga University, Surabaya, Indonesia

Abstract

Chest computed tomography (CT) scan is one of the main modalities to detect COVID-19 infection. Several CT scan findings are the hallmark to rapidly detect suspected COVID-19 infection, therefore timely treatment could be administered. This study aims to describe chest CT scan findings of confirmed COVID-19 patients based on clinical symptom onset patterns.

This study is a descriptive study with a retrospective design in confirmed COVID-19 patients. Data regarding characteristics of chest CT scan findings and the patient's history including clinical symptoms onset were collected.

There were 123 subjects in this study. The study data were categorized into modified clinical symptom onset patterns which were early, progressive, peak, absorption, and advanced phases. Most of male (57%) and female (43%) patients that were admitted to the hospital were in absorption phase of the disease with fever, cough, and dyspnea as the most prevalent symptoms. Chest CT scan findings of absorption phase included GGO (ground glass opacity) in 28 patients (76%), crazy paving pattern in 18 patients (49%), consolidation in 22 patients (59%), and fibrosis in 31 patients (84%). Chest CT scan findings of confirmed COVID-19 patients vary depending on the phase of the disease. Risk factors and secondary bacterial infection may contribute to long-term persistence of symptoms.

Keywords: COVID-19, chest CT scan, clinical symptom onset patterns

Introduction

The COVID-19 disease has symptoms similar to the common cold infection with an incubation period of about 2-14 days, therefore most cases go undetected and causes fast-growing transmission. Prompt diagnosis is essential to help prevent this

disease transmission.⁽¹⁾ WHO and Center for Disease Control (CDC) guidelines stated that chest computed tomography (CT) scan is one of the main modalities in detecting COVID-19 infection. The typical chest CT scan findings of a patient infected with COVID-19 causing pneumonia are multiple areas of bilateral lobular and subsegmental consolidation and ground-glass opacity (GGO).⁽²⁾ These findings are the key to rapidly detect a suspected COVID-19 infection, therefore timely treatment could be administered and could prevent further transmission.

Studies regarding correlation between symptom onset in COVID-19 patients and CT scan findings has been conducted by several researchers. A study by Pan

Corresponding author:

Anita Widyoningroem

Department of Radiology, Medical Faculty, Airlangga University Jl. Mayjen Prof. Dr. Moestopo No.47, Surabaya, Indonesia

Email: anita.widyo75@gmail.com

et al.,2020, examined the time course of pulmonary changes on chest CT scans during recovery from COVID-19 pneumonia and divided them into four stages of chest CT scan findings based on the quartiles of the time course from day 0 to day 26. In stage 1 (early stage: 0-4 days) GGO was found; in stage 2 (progressive stage: 5-8 days) GGO was scattered with enhanced crazy-paving pattern findings; in stage 3 (peak stage: 9-13 days) consolidated consolidation, scattered GGO, crazy-paving pattern, and residual parenchymal bands were found; and in stage 4 (absorption stage: >14 days) the consolidation was gradually absorbed and no longer crazy-paving pattern was found.⁽³⁾ This study aims to describe chest CT scan findings of confirmed COVID-19 patients based on clinical symptom onset patterns.

Methods and Materials

Study Design

This is a descriptive study with a retrospective design that used medical records and the examination results of non-contrast chest CT scan from patients with confirmed COVID-19 who were treated at Dr. Soetomo Surabaya General Hospital, Surabaya, Indonesia from June to October 2020.

Data Collection

Researchers identified medical record and history data of confirmed COVID-19 patients. The collected data were clinical symptoms, comorbid, and symptom onset patterns, which were calculated from the start of the patient's symptom until a non-contrast chest CT scan was performed. Evaluation of CT scan finding was conducted by 2 experimented radiologists senior (mean 10 years of experience) and one resident (mean 3 years of experience) using standardized CT reports. For cases with controversial results, two radiologists with more than 10 years experience jointly reviewed and evaluated the result.

Imaging Protocols

All patients were adults and non-pregnant who

underwent non-contrast chest CT scan to detect COVID-19 pneumonia in supine position during full inspiration. A total of 69 patients were subjected to CT scans in the Emergency Room using multi-detector 128 slice CT scan (Toshiba Aquilion Vision) and 54 patients had CT scans at the Radiodiagnostic Installation using multi-detector 128 slice CT scan (Philips) and 16 slice CT scan (Hitachi). The protocol used was 120 kV voltage tube, automatic tube current (120–380) mA, 5 mm thickness, 0.5 mm slice interval, 0-5 second rotation speed, and 1-0.828 helical distance.

Clinical and Imaging Interpretation

Identification of clinical symptom data categorized by clinical syndrome according to WHO, which includes: mild disease/non-specific symptoms, mild pneumonia, severe pneumonia, ARDS, sepsis and septic shock.⁽⁴⁾ The analysis results were reported using The Covid-19 Reporting and Data System (CO-RADS) category.⁽⁵⁾ Severity was assessed using a visual scoring system of each lobe involved. The scores for all lung lobes are summed to show lung involvement severity.⁽⁶⁾ This study used a modified assessment of per-lobe severity. Score 0: if there is no lesion or 0%, score 1: when the estimated number of lesions visually is 1% to 25%, score 2: when the estimated number of lesions visually is 26% to 50%, score 3: when the estimated number of lesions is visual 51% to 75%, score 4: when the estimated number of lesions visually is more than 75%. The final results were categorized into mild (total score 7 or less), moderate (total score between 8 and 17), and severe (total score 18 or more).

Data Analysis

The collected data were categorical, which were processed using SPSS 21 software and then presented in tabular form with percentages (%).

Results

The study was conducted on 123 study subjects of COVID-19 pneumonia patients. The study data were

categorized into modified clinical symptom onset patterns. The categories included initial (0-4 days), progressive (5-8 days), peak (9-13 days), absorption (14-30 days), and advanced phase (>30 days).

Subject Characteristics Based on Clinical Symptom Onset Patterns

The age range of study subjects was 17-91 years, with a mean age of 52 years. Based on the characteristics of clinical symptom onset patterns, most patients were included in absorption phase with an age range of 32-70 years (mean= 55±9 years) with 37 patients. The number of subjects who were male was 70 people (57%), which was more than female subjects with 53 people (43%). In initial, peak, absorption, and advanced phases most patients were male with 18 (56%), 9 (64%), 21 (57%), and 10 (67%) patients, respectively. While in progressive phase most patients were female with 13 (52%) patients.

The 3 most common clinical symptoms were fever (66%), cough (70%), and dyspnea (58%). Fever was found mostly in initial and progressive phase, with 24 (75%) and 21 (84%) patients, respectively. Cough was

mostly found in peak phase with 11 (79%) patients, while in absorption and advanced phase the number of patients who experienced cough and dyspnea were similar, with 28 (76%) and 10 (67%) patients in each phase, respectively. The study subjects in this study had variable comorbid, although some patients had no concomitant comorbid. In early, absorption, and peak phase, the most frequent comorbid was Diabetes Mellitus (DM) with 10 (31%), 11 (30%), and 3 (20%) patients, respectively. In progressive and peak phase, the most frequent comorbid was hypertension (HT), with 10 (40%) and 6 (43%) patients, respectively.

The study subjects were categorized into clinical syndromes based on WHO criteria. The results showed that most patients in early and progressive phase had mild pneumonia with 13 (41%) and 11 (44%) patients, respectively. In peak phase, there were the same number of patients for mild and severe pneumonia, which were 4 (29%) patients. Most patients in absorption and advanced phase had severe pneumonia, with 10 (27%) and 5 (33%) patients, respectively.

The subject characteristics are outlined in Table 1.

Table 1. Subject Characteristics

Characteristics	Early phase N (%)	Progressive phase N (%)	Peak phase N (%)	Absorption phase N (%)	Advanced phase N (%)	Total N (%)
Gender						
Male	18 (56)	12 (48)	9 (64)	21 (57)	10 (67)	70 (57)
Female	14 (44)	13 (52)	5 (36)	16 (43)	5 (33)	53 (43)
Age (Years)						
Mean \pm Standard Deviation	43 \pm 15	49 \pm 16	59 \pm 13	55 \pm 9	59 \pm 8	52 \pm 14
Age Range	17 - 74	32 - 91	35 - 85	32 - 70	45 - 70	17 - 91
Symptoms						
Fever	24 (75)	21 (84)	8 (57)	19 (51)	9 (60)	81 (66)
Cough	17 (53)	20 (80)	11 (79)	28 (76)	10 (67)	86 (70)
Dyspnea	13 (41)	14 (56)	6 (43)	28 (76)	10 (67)	71 (58)
No Symptom	1 (3)	0 (0)	0 (0)	0 (0)	0 (0)	1 (0.8)
Comorbid						
Chronic Kidney Disease (CKD)	2 (6)	3 (12)	0 (0)	2 (5)	2 (13)	9 (7)
DM	10 (31)	7 (28)	5 (36)	11 (30)	3 (20)	36 (29)
HT	7 (22)	10 (40)	6 (43)	7 (19)	2 (13)	32 (26)
Obesity	1 (3)	5 (20)	0 (0)	1 (3)	1 (7)	8 (6)
Malignancy	3 (9)	2 (8)	0 (0)	0 (0)	1 (7)	6 (5)
Others	13 (41)	4 (16)	2 (14)	8 (22)	5 (33)	32 (26)
No Comorbid	9 (28)	9 (36)	7 (50)	15 (40)	6 (40)	46 (37)
Clinical Syndrome (WHO)						
None	1 (3)	0 (0)	0 (0)	0 (0)	0 (0)	1 (0.8)
Mild Disease / Non-Specific Symptoms	10 (31)	2 (8)	3 (21)	4 (11)	1 (7)	20 (16)
Mild Pneumonia	13 (41)	11 (44)	4 (29)	6 (16)	3 (20)	37 (30)
Severe Pneumonia	0 (0)	8 (32)	4 (29)	10 (27)	5 (33)	27 (22)
ARDS	3 (9)	0 (0)	2 (14)	7 (19)	4 (27)	16 (13)
Sepsis	3 (9)	1 (4)	0 (0)	6 (16)	0 (0)	10 (8)
Septic Shock	2 (6)	3 (12)	1 (7)	3 (8)	1 (7)	10 (8)

Characteristics of Chest CT Scan Findings by Clinical Symptom Onset Pattern

The results of this study showed that the most common characteristics in early phase were GGO in

26 (81%) patients, crazy paving pattern on 8 (25%) patients, consolidation on 14 (44%) patients, and fibrosis on 23 (72%) patients. In progressive phase there were GGO in 17 (68%) patients, crazy paving pattern in 11 (44%) patients, consolidation in 15 (60%)

patients, and fibrosis in 22 (88%) patients. In peak phase, there were GGO in 11 (79%) patients, crazy paving pattern in 4 (29%) patients, consolidation in 7 (50%) patients, and fibrosis in 10 (71%) patients. In absorption phase there were GGO in 28 (76%) patients, crazy paving pattern in 18 (49%) patients, consolidation in 22 (59%) patients, and fibrosis in 31 (84%) patients. In advanced phase there were

GGO in 8 (53%) patients, crazy paving patterns and consolidation in 1 (7%) patient each, and fibrosis in 12 (80%) patients. Bronchiectasis and bronchiolectasis were commonly found in absorption and advanced phases. The most common distribution for chest CT scan findings were bilateral predominantly peripheral in all clinical symptom onset patterns. Table 2 outlines these chest CT scan findings.

Table 2. Characteristics of Chest CT Scan Findings by Clinical Symptom Onset Pattern

Chest CT Scan Findings	Early phase N (%)	Progressive phase N (%)	Peak phase N (%)	Absorption phase N (%)	Advanced phase N (%)	Total N (%)
Characteristics						
GGO	26 (81)	17 (68)	11 (79)	28 (76)	8 (53)	90 (73)
Crazy Paving Pattern	8 (25)	11 (44)	4 (29)	18 (49)	1 (7)	42 (34)
Consolidation	14 (44)	15 (60)	7 (50)	22 (59)	1 (7)	59 (48)
Halo sign	4 (12)	2 (8)	2 (14)	1 (3)	1 (7)	10 (8)
Reverse halo sign	2 (6)	1 (4)	1 (7)	2 (5)	1 (7)	7 (6)
Air bubble sign	0 (0)	0 (0)	1 (7)	3 (8)	0 (0)	4 (3)
Reticular pattern	2 (6)	3 (12)	0 (0)	1 (3)	2 (13)	8 (6)
Nodule	1 (3)	1 (4)	2 (14)	3 (8)	3 (20)	10 (8)
Fibrosis	23 (72)	22 (88)	10 (71)	31 (84)	12 (80)	98 (80)
Bronchiectasis	2 (6)	4 (16)	2 (14)	20 (54)	8 (53)	36 (29)
Bronchiolectasis	2 (6)	3 (12)	3 (21)	22 (59)	7 (47)	37 (30)
Pleural effusion	3 (9)	2 (8)	1 (7)	4 (11)	2 (13)	12 (10)
Pleural thickening	2 (6)	5 (20)	2 (14)	7 (19)	2 (13)	18 (15)
Distribution						
Bilateral predominantly peripheral	20 (62)	20 (80)	13 (93)	35 (95)	12 (80)	100 (81)
Unilateral predominantly peripheral	8 (25)	3 (12)	0 (0)	0 (0)	0 (0)	11 (9)
Bilateral predominantly central	1 (3)	1 (4)	0 (0)	0 (0)	0 (0)	2 (2)
No predominance	1 (3)	1 (4)	0 (0)	0 (0)	1 (7)	3 (2)
No lesion	2 (6)	0 (0)	1 (7)	2 (5)	2 (13)	7 (6)

Characteristics of Chest CT Scan Findings by CO-RADS Categories

In CO-RADS 1 group, most patients showed normal lung features and only 1 (14%) had fibrosis. In CO-RADS 2 category group, most patients had fibrosis and pleural thickening, with 7 (78%) and 4 (44%) patients, respectively. In CO-RADS 3 category group, most patients had consolidation and GGO with 4 (80%) and 3 (60%) patients, respectively.

In CO-RADS 4 group, most patients had GGO and consolidation with 11 (85%) and 8 (61%) patients, respectively. In CO-RADS 5 group, most patients had GGO and fibrosis, with 35 (85%) and 33 (80%) patients, respectively. Finally, in CO-RADS 6 group, most patients had fibrosis and GGO, with 45 (94%) and 40 (83%) patients, respectively. All CO-RADS categories mostly had bilateral predominantly peripheral in their finding’s distribution. Table 3 outlines these findings.

Table 3. Characteristics of Chest CT Scan Findings by CO-RADS Categories

Chest CT Scan Findings	CO-RADS						Total N (%)
	C1 N (%)	C2 N (%)	C3 N (%)	C4 N (%)	C5 N (%)	C6 N (%)	
Mean severity score ± standard deviation	0 ± 0	2 ± 1	5 ± 3	5 ± 4	10 ± 5	10 ± 5	8 ± 5
Characteristic							
GGO	0 (0)	1 (11)	3 (60)	11(85)	35 (85)	40 (83)	90 (73)
Crazy Paving Pattern	0 (0)	1 (11)	0 (0)	3 (23)	21 (51)	17 (35)	42 (34)
Consolidation	0 (0)	1 (11)	3 (60)	8 (61)	23 (56)	24 (50)	59 (48)
Halo sign	0 (0)	0 (0)	0 (0)	2 (15)	6 (15)	2 (4)	10 (8)
Reverse halo sign	0 (0)	1 (11)	0 (0)	1 (8)	4 (10)	1 (2)	7 (6)
Air bubble sign	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	4 (8)	4 (3)
Reticular pattern	0 (0)	1 (11)	0 (0)	2 (15)	3 (7)	2 (4)	8 (6)
Nodule	0 (0)	1 (11)	0 (0)	1 (8)	2 (5)	6 (12)	10 (8)
Fibrosis	1 (14)	7 (78)	4 (80)	8 (61)	33 (80)	45 (94)	98 (80)
Bronchiectasis	0 (0)	2 (22)	0 (0)	1 (8)	6 (15)	27 (56)	36 (29)
Bronchiolectasis	0 (0)	0 (0)	1 (20)	0 (0)	3 (7)	33 (69)	37 (30)
Pleural efusion	0 (0)	3 (33)	1 (20)	4 (31)	1 (2)	3 (6)	12 (10)
Pleural thickening	1 (14)	4 (44)	1 (20)	1 (8)	2 (5)	9 (19)	18 (15)
Distribution							
Bilateral predominantly peripheral	2 (25)	1 (12)	1 (20)	9 (69)	40 (98)	47 (98)	100 (81)
Unilateral predominantly peripheral	0 (0)	5 (62)	3 (60)	3 (23)	0 (0)	0 (0)	11 (9)
Bilateral predominantly central	0 (0)	0 (0)	1 (20)	0 (0)	1 (2)	0 (0)	2 (2)
No predominance	1 (12)	0 (0)	0 (0)	1 (8)	0 (0)	1 (2)	3 (2)
No lesion	7 (100)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	7 (6)

Discussion

The typical findings of a patient infected with COVID-19 that causes pneumonia on a chest CT scan are multiple areas of bilateral lobular and subsegmental consolidation and ground-glass opacity (GGO).⁽²⁾

These findings are the key to rapidly detect a suspected COVID-19 infection, therefore a timely treatment could be provided and prevent further infection transmission. CT scan findings in this study included GGO, crazy paving pattern, consolidation, halo sign,

inverted halo sign, bubble air sign, reticular pattern, nodule, fibrosis, bronchiectasis, bronchiolectasis, pleural effusion and pleural thickening. These findings were in accordance to several previous studies.^(3,7,8) Based on clinical symptom onset patterns, the findings in progressive, peak, and absorption phases in this study were mostly GGO, crazy paving pattern, consolidation, and fibrosis, with mostly bilateral predominantly peripheral distribution. These were similar to several previous studies which reported the presence of GGO with and without consolidation in subpleural visceral with peripheral ($p < 0.001$) or bilateral multifocal distribution.^(3,8,9) The virus ability to reach terminal bronchi and alveoli might be the main cause of this peripheral distribution.⁽¹⁰⁾ Fibrosis was found in the early phase of this study. However, fibrosis due to COVID-19 usually occurs in chronic or healing phase.⁽⁸⁾ It could be assumed that this finding is not a COVID-19 infection process and existed before COVID-19 infection. Statistically, Indonesia is a tuberculosis endemic area with a prevalence of bacteriological confirmed tuberculosis in 2013-2014 of 759 per 100,000 population aged 15 years and over.⁽¹¹⁾

In this study, the lung severity was in line with the CO-RADS category. Lung severity scoring progresses with clinical onset. Lung severity scoring in initial phase was mild, then increased in progressive, peak, and up to absorption phase, and then decreased in advanced phase. Study by Guan et al., 2019 compared chest CT scan findings between progressive and recovery phase. From this study, several alterations were found in progressive phase: rounded image of GGO developed into uneven GGO, increased consolidation, and sometimes pleural effusions. In recovery phase, the crazy paving pattern and air bronchogram findings decreased significantly.⁽¹²⁾ In this study, there were prolonged clinical symptoms and CT scan findings that still showed GGO, consolidation and crazy-paving patterns in absorption (14-30 days) and advanced (> 30 days). We assumed that there are several risk factors for these events. In this study, it

was found that the average age of study subject was 52 years ($SD \pm 14$), with mostly male patients in all phases of clinical symptom onset pattern. We also noted the presence of comorbid factors such as DM, HT, CKD, obesity and other comorbidities. In absorption and advanced phases, it was found that most of patients still had cough and dyspnea. This is in accordance to a study by Miyazato et al., 2020 who reported prolonged or delayed symptoms of coronavirus disease. The findings of this study were dyspnea, fatigue, cough, dysosmia, and dysgeusia, which lasted more than 120 days after symptom onset.⁽¹³⁾

A meta-analysis study by Biswas et al., 2021 estimated risk factors such as sex, age, or comorbidity with mortality in COVID-19 patients, and proved that male patients aged ≥ 50 years or that had comorbidities was significantly associated with an increased mortality risk. Age ≥ 50 years and male sex have higher expression of angiotensin-converting enzyme 2 (ACE2) regulated by male sex hormones, making men more prone of SARS-CoV-2 infection and had worse symptoms. Another hypothesis stated that women could fight the development of SARS-CoV-2 infection due to X-linked heterozygous alleles, which is called sex dimorphisms.⁽¹⁴⁾ Comorbidity is associated with decreased immune function, as in DM patients. Natural immune function is substantially reduced, which could restrict the body from producing antibodies against any infection. ACE-2 is highly expressed in lungs, intestines, kidneys, and blood vessels epithelial cells and is predominantly upregulated in DM or HT patients treated with ACE inhibitors (ACEIs) and angiotensin II type-I receptor blockers (ARBs).⁽¹⁵⁾

Secondary bacterial infection might also be the cause of prolonged symptoms. A study by Zhang et al., 2020 reported that COVID-19 patients who received invasive mechanical ventilation or were in critical condition had a higher likelihood of developing secondary infection ($p < 0.0001$). The most common infections are respiratory, haematogenous and urinary tract infections. Secondary infection will

lead to a higher mortality rate.⁽¹⁵⁾

This study has some limitations. Medical intervention as well as comorbid histories of confirmed patients with concomitant infection might interfere the CT scan findings. A history of comorbid such as malignancy, CKD, and others could also raise a bias with overlapped CT findings. We have a lack of standardized timeline of evaluation in overall patients, since the CT scan has not been as the routine protocol in patient's evaluation. CT scanning is performed mainly to follow up the disease's progression. Some cases of this study obtained were in the state of prolonged disease onset, approximately from 30-120 days.

Conclusion

CT scans are able to reveal distinct confirmed COVID-19 findings that are variable according to clinical symptom onset. The typical finding in early phase is GGO with a bilateral predominantly peripheral distribution. The next phases have variable findings, with GGO, crazy paving pattern, and consolidation as the most common findings with bilateral predominantly peripheral distribution. This study also showed that most patients were in absorption phase with dominant findings of fibrosis. However, GGO, consolidation, and crazy paving patterns were still found. Cough and dyspnea were still found in this phase. These results show long-term COVID-19 findings from both the symptoms and CT scan findings.

Acknowledgement: We would like to thank Department of Radiology of Dr. Soetomo Hospital Surabaya, Indonesia in providing the data for analysing.

Ethical Clearance: This study was approved by the ethical committee of Dr. Soetomo General Hospital, Surabaya, Indonesia (Registration Number: 0025 / KEPK / VII / 2020).

Source of Funding: Self

Conflict of Interest: Nil

References

1. Mahdi Asghari Ozma, et all. Clinical Manifestation, Diagnosis, Prevention and Control of SARS-CoV-2 (COVID-19) During The Outbreak Period. *Le Infezioni in Medicine*, n. 2, 153-165, April 2020, <https://www.researchgate.net/publication/340417798>
2. Qionjie HU, et all. Early CT Features and Temporal Lung Changes in COVID-19 Pneumonia in Wuhan, China. *European Journal of Radiology*, 128 (2020)109017, April 2020, doi: <https://doi.org/10.1016/j.ejrad.2020.109017>
3. Feng Pan, MD, et all. Time Course of Lung Changes On Chest CT During Recovery From 2019 Novel Coronavirus (COVID-19) Pneumonia. *RSNA*, Vol. 295, No. 3, June 2020, <https://doi.org/10.1148/radiol.2020200370>
4. Stephen A. Lauer, MS, PHD et all. The Incubation Period of Coronavirus Disease 2019 (COVID-19) From Publicly Reported Confirmed Cases: Estimation and Application. *Ann Intern Med*, 10 March 2020, doi: <https://doi.org/10.7326/M20-0504>
5. Mathias Prokop, MD et all. CO-RADS – A Categorical CT Assessment Scheme for Patients With Suspected COVID-19: Definition and Evaluation. *RSNA*, org, 2020.
6. Saeed, G. A. *et al.* (2020) 'Correlation between Chest CT severity scores and the clinical parameters of adult patients with COVID-19 pneumonia', *medRxiv*, 2021. doi: [10.1101/2020.10.15.20213058](https://doi.org/10.1101/2020.10.15.20213058).
7. Haixia Chen, et all. Clinical and Imaging Features of COVID-19. *Radiology of Infectious Disease*, April 2020, <https://doi.org/10.1016/j.jrid.2020.04.003>
8. Zheng Ye, et all. Chest CT Manifestations of New Coronavirus Disease 2019 (COVID-19): A Pictorial Review. *European Society of Radiology*,

- 11 March 2020, <https://doi.org/10.1007/s00330-020-06801-0>
9. Prashant Nagspal, et al. Imaging of COVID-19 Pneumonia: Pattern, Pathogenesis, and Advances. *BJR*, July 2020, <https://doi.org/10.1259/bjr.20200538>
 10. Xiang, C. *et al.* (2020) 'CT Findings in a Novel Coronavirus Disease (COVID-19) Pneumonia at Initial Presentation', *BioMed Research International*, 2020. doi: 10.1155/2020/5436025.
 11. Kementerian Kesehatan RI (2018). Pusat Data dan Informasi Kesehatan RI: Tuberculosis. ISSN 2442-7659. Available from: <https://pusdatin.kemkes.go.id/resources/download/pusdatin/infodatin/infodatin-tuberculosis-2018.pdf>.
 12. Guan, C. S. *et al.* (2020) 'CT findings of COVID-19 in follow-up: Comparison between progression and recovery', *Diagnostic and Interventional Radiology*, 26(4), pp. 301–307. doi: 10.5152/dir.2019.20176.
 13. Miyazato, Y. *et al.* (2020) 'Prolonged and Late-Onset Symptoms of Coronavirus Disease 2019', *Open Forum Infectious Diseases*, 7(11), pp. 4–6. doi: 10.1093/ofid/ofaa507.
 14. Biswas, M. *et al.* (2021) 'Association of Sex, Age, and Comorbidities with Mortality in COVID-19 Patients: A Systematic Review and Meta-Analysis', *Intervirology*, 64(1), pp. 36–47. doi: 10.1159/000512592.
 15. Zhang, H. *et al.* (2020) 'Risks and features of secondary infections in severe and critical ill COVID-19 patients', *Emerging Microbes and Infections*, 9(1), pp. 1958–1964. doi: 10.1080/22221751.2020.1812437.