

# 08. The Success of Elimination Diet in Indonesian

*by Azwin Mengindra Putera*

---

**Submission date:** 03-Apr-2023 12:06PM (UTC+0800)

**Submission ID:** 2054228375

**File name:** 08.\_The\_Success\_of\_Elimination\_Diet\_in\_Indonesian.pdf (1.26M)

**Word count:** 6393

**Character count:** 33537

# The Success of Elimination Diet in Indonesian Children with Food Allergy: The Role of Caregiver's Stress, Family Activities, and Coping

Azwin Mengindra Putera\*<sup>1</sup>, Irwanto<sup>1</sup>, Margarita Maria Maramis<sup>2</sup>

<sup>1</sup>Department of Child Health, Faculty of Medicine, Universitas Airlangga – Dr. Soetomo General Academic Hospital, Surabaya, Indonesia

<sup>2</sup>Department of Psychiatry, Faculty of Medicine, Universitas Airlangga – Dr. Soetomo General Academic Hospital, Surabaya, Indonesia

Corresponding Author: Azwin Mengindra Putera

Email: [azwinmengindraputera@gmail.com](mailto:azwinmengindraputera@gmail.com)

## ABSTRACT

**Background:** Elimination diet is the most effective treatment in treating children with food allergy. However, in its implementation, the success of elimination diet depends on a variety of factors, especially their caregiver.

**Objectives:** This study analyzed the correlation between caregiver's stress, coping strategy, and family's daily activities on the success of elimination diet in Indonesian children with food allergy.

**Methods:** Participants in this study were caregiver of Indonesian children with food allergy who received elimination diet for 4 weeks. Participants were measured for caregiver's stress, family's daily activities, and family's coping strategy after elimination diet. Caregiver's stress was measured using PSS, family's daily activities used FAIS, and while family's coping strategy used F-COPES. The data were analyzed using multiple linear regression, with  $p < 0.05$ .

**Results:** The success rate of elimination diet in Indonesian children with food allergy was 43.64%. There was no significant correlation between caregiver's stress and elimination diet in Indonesian children with food allergy ( $t = -0.145$ ; 95% CI  $-0.014 - 0.012$ ;  $p = 0.886$ ). A significant correlation was found between FAIS and elimination diet in Indonesian children with food allergy ( $t = -2.297$ ; 95% CI  $-0.179 - -0.012$ ;  $p = 0.026$ ). Moreover, caregiver's coping strategy significantly correlated with elimination diet in Indonesian children with food allergy ( $t = 16.154$ ; 95% CI  $0.034 - 0.043$ ;  $p < 0.001$ ).

**Conclusion:** The success of elimination diet in Indonesian children with food allergy is influenced by family's daily activities and coping strategy, while caregiver's stress has no significant influence.

**Keywords:** Caregiver's stress, family's coping strategy, family's daily activities, food allergy, elimination diet

## Correspondence:

Azwin Mengindra Putera

Department of Child Health, Faculty of Medicine, Universitas Airlangga – Dr. Soetomo General Academic Hospital, Surabaya, Indonesia

Email: [azwinmengindraputera@gmail.com](mailto:azwinmengindraputera@gmail.com)

## INTRODUCTION

Food allergy is the main cause of early atopic disease manifestation that can cause various other allergic manifestations [1]. The incidence of allergy in the world, particularly food allergy continues to increase, especially in children [2]. It was reported that in 2016, 2-10% of patients experienced food allergy, while as much as 5.9% of patients in Europe experienced food allergy. Moreover, 7.1% children in Canada had food allergy [3]. There is still no detailed data on the prevalence of food allergy patients in Indonesia in all age groups [4], but data on children with food allergy in Indonesia is reported to be 3-60% [5]. Food allergy disease is commonly found in children [3,6,7]. The prevalence of food allergy continues to increase by around 4-8% in children annually [8].

Elimination diet is the most effective treatment in handling patients with adverse reactions due to food allergy [1,9]. The method detects and removes specific antagonistic foods by formulating a nutritious hypoallergenic diet to ensure optimal health of people who are sensitive to food. However, this process is often tedious, time-consuming, and requires extraordinary knowledge, skills, and commitment [8]. In the case of children with food allergy, the success of food elimination diet depends on parents and children, how serious they are to follow and understand the elimination diet recommendation as much as possible. There are still many parents and children with food allergy who do not follow elimination diet properly due to various factors, both from parents and children [1,10].

Food allergy in children cannot be predicted and life threatening, thus increases caregiver's stress [11]. It was reported that 41% of caregivers experienced increased stress when their children were diagnosed with food allergy [11,12]. This condition causes the caregiver to limit children's activities in party and snacking at school. Daily activities such as shopping and eating out are frightening for children with food allergies and are even considered life threatening [12,13]. Adaptive coping is needed to solve these problems [14,15], so that the elimination diet given to children with food allergies can be successful [6,7].

Elimination diet is the best way to help patients cutting ties from foods that cause their daily allergic symptoms [1,9,16,17]. Data related to the success of elimination diet in Indonesian children with food allergy is very limited. Dr. Soetomo General Academic Hospital, Surabaya, as the main referral hospital in East Java, Indonesia, reported the number of Indonesian children with food allergy has increased annually, with 127 patients in 2017 and 143 patients in 2018. Based on the explanation above, it is necessary to conduct a analysis to examine correlation between caregiver's stress, coping strategy, and family's daily activities on the success of elimination diet in Indonesian children with food allergy.

## MATERIALS AND METHODS

### Participants

Participants in this study were caregiver of Indonesian children with food allergy, which consisted of biological

mother or father. The participant's inclusion criteria were caregiver with children aged 1 month to 18 years, children diagnosed with food allergy [4,18,19], and children receiving elimination diet. The exclusion criteria were caregiver or children who did not want to participate in the study and children aged <1 month. Participants first received explanation of their rights, obligations, and filled out an informed consent sheet.

**Design**

This study used a case control study design [20,21], by observing the administration of elimination diet in Indonesian children with food allergy. This research was conducted in Dr. Soetomo General Academic Hospital,

Surabaya, Indonesia, from May to August 2019. The participants were chosen using consecutive sampling, with a total of 55 participants (figure 1). Participants were not assessed prior to elimination diet. Assessment of allergy symptoms of Indonesian children with food allergy was carried out based on the response of emerging diseases such as allergic rhinitis [22], asthma [23], and atopic dermatitis [24]. The elimination diet was given for 4 weeks based on the Guideline for Diagnosis and Management of Food Allergy [18,25,26]. The participants were then evaluated for allergy symptoms and measured for characteristics, caregiver's stress, family's daily activities, and family's coping strategy.




Figure 1. Identification Flowchart of Number of Participants

**Ethical Approval**

This study has obtained a certificate of ethical clearance according to Declaration of Helsinki from ethical committee team in Dr. Soetomo General Academic Hospital and Faculty of Medicine Universitas Airlangga ethics Committee (1165/KEPK/V/2019).

**Elimination Diet**

Elimination diet was provided based on a history of response to food allergies and skin prick tests [16]. The consumption of allergy-causing food was stopped and replaced the nutritional intake of these foods in consultation with a nutritionist. Nutrients obtained from these foods should be replaced with foods containing similar nutrients [18,25,26]. The use of books in

elimination diet really helped caregivers and nutritionists make a diet plan for 4 weeks. Diet elimination was declared unsuccessful if the elimination diet was carried out >4 weeks and persistent signs of symptoms of food allergy.

**Measurement of Caregiver's Stress**

Caregiver's stress was measured using parental stress scale (PSS) with 18 questions. The scale is developed to measure stress level as parents [27]. The PSS used Likert scale of 1-5 (1 = strongly disagree, 2 = disagree, 3 = neutral, 4 = agree, 5 = strongly agree), with a minimum value of 18 and a maximum value of 90 [28]. This assessment tool was categorized into 3, namely mild (18 - 42), moderate (43 - 66), and severe (67-90) [29]. The

PSS was considered valid and reliable with Cronbach's alpha = 0.83 [27,28].

**Measurement of Family's Daily Activities**

Family's daily activities were measured using the food allergy impact scale (FAIS) to evaluate parents' perceptions about the impact of food allergy on eight aspects of family's daily activities, with the higher score indicating a greater impact. The scale consists of 32 questionnaire items, consisting of 8 activity aspects namely meal preparation (six items), family social activities (five items), caregiver-supervised child social activities (two items), family relations (three items), caregiver's stress and free time (three items), school or structured activities (six items), autonomous child social activities (three items), and employment and finances (three items). The FAIS reliability value was  $\geq 0.70$  [15,30].

**Measurement of Family's Coping Strategy**

The measurement of family coping was conducted using family crisis oriented personal evaluation scales (F-COPES) questionnaire, with a 5-point Likert's scale. This instrument displays 30 items of coping behavior designed to record problem solving, attitudes and behaviors developed by families to respond to problems or difficulties. This scale is divided into 5 subscales: obtaining social support, reframing, seeking spiritual

support, mobilizing family support, and passive assessment [31]. The Indonesian F-COPES was considered valid and reliable with Cronbach's alpha = 0.89 [32,33].

**Statistical Analysis**

The measured data were analyzed univariate and bivariate, where univariate data were displayed in the form of a frequency distribution or mean  $\pm$  standard deviation (SD). Meanwhile, the bivariate analysis used multiple linear regression tests. Statistical tests were significant if  $p < 0.05$ . Measurement data were analyzed using IBM SPSS Statistic software version 21.0 (IBM Corp., Armonk, NY, USA).

**RESULTS**

**Characteristics of Participants**

Most participants were male 40 (72.73;  $p = 0.349$ ), with the mean age of  $68.1 \pm 37.46$  years ( $p = 0.159$ ). Most participants came from low-income family (65.46%;  $p = 0.009$ ). Parents' income affects the process of treating children with food allergy. Most parents' education level was non-bachelor as much as 72.73% ( $p = 0.782$ ; Table 1). Most elimination diet was unsuccessful (56.36%). The mean age of participants who succeeded in implementing elimination diet was  $60.0 \pm 39.23$  years, and those who were unsuccessful were  $74.4 \pm 35.38$  ( $p = 0.692$ ).

Table 1. Characteristics of Participants

Characteristics of Participants	N (%)	p
Sex		
Male	40 (72.7)	0.349
Female	15 (27.3)	
Income		
Low	36 (65.5)	0.009*
High	19 (34.5)	
Educational Level		
Non-bachelor	40 (72.7)	0.782
Bachelor	15 (27.3)	
Elimination diet in Children		
Unsuccessful	31 (56.4)	-
Successful	24 (43.6)	

Abbreviations: \* significant  $p < 0.05$ ; \*\*significant  $p < 0.001$ .

**Correlation between Caregiver's Stress on Elimination Diet in Indonesian Children with Food Allergy**

The participant's stress value in this study ranged from 28 to 57, where the most participant had a stress value of 43 for 12 participants. The mean value of the participant's stress was  $45.29 \pm 4.67$ , with a median value of 45 (43 -

48). Most participants had stress in the moderate category as much as 89.09%. Participants who had mild PSS scores and successful elimination diet were as much as 83.3%, while participants who had moderate PSS scores and unsuccessful elimination diet were 61.2%. There was no significant correlation between

Table 2. Correlation between Caregiver's Stress, Family Activities, and Family Coping on the Success of Elimination Diet in Indonesian Children with Food Allergy

Caregiver's Condition	Successful Elimination Diet			Unsuccessful Elimination Diet			p
	Mean $\pm$ SD	Median (Q1-Q3)	Min-Max	Mean $\pm$ SD	Median (Q1-Q3)	Min-Max	
PSS	44.17 $\pm$ 5.65	45.00 (43.00 - 46.75)	28.00 - 54.00	46.16 $\pm$ 3.61	45.00 (43.00 - 49.00)	41.00 - 57.00	0.886
FAIS	5.01 $\pm$ 0.89	5.13 (4.48 - 5.47)	2.98 - 6.75	5.52 $\pm$ 0.53	5.33 (5.15 - 5.78)	4.31 - 6.52	0.026*
F-COPES	116.29 $\pm$ 3.91	116.50 (113.25 - 120.00)	109.00 - 122.00	95.45 $\pm$ 5.20	97.00 (94.00 - 99.00)	77.00 - 100.00	0.000**

Abbreviation: PSS, parental stress scale; FAIS, food allergy impact scale; F-COPES, family crisis oriented personal evaluation scales; \*significant  $p < 0.05$ ; \*\*significant  $p < 0.001$ .

Table 3. Comparison of Each FAIS Subscale to the Caregiver of Indonesian Children with Food Allergy

FAIS	2		3		4		5		6		7		8	
	95% CI	p	95% CI	p	95% CI	p	95% CI	p	95% CI	p	95% CI	p	95% CI	p
1	0.12-0.43	0.001*	0.48-1.31	0.000**	0.72-1.48	0.000**	0.96-1.68	0.000**	0.96-1.77	0.000**	0.26-0.97	0.001*	0.46-1.23	0.000**
2			0.17-0.94	0.005*	0.43-1.22	0.000**	0.70-1.38	0.000**	0.65-1.44	0.000**	-0.09-0.70	0.125	0.15-1.23	0.008*
3					-0.56-0.48	0.875	-0.12-0.90	0.135	0.97-1.00	0.019*	-0.69-0.24	0.329	-0.54-0.76	0.743
4							-2.71-0.70	0.380	0.04-1.01	0.067	-0.99-0.65	0.084	-0.83-0.32	0.372
5									-0.39-0.57	0.714	-1.34-0.24	0.005*	-0.90-0.04	0.031*
6											-1.34-0.23	0.007*	-1.11-0.01	0.053
7													-0.21-0.80	0.249

Abbreviation: FAIS, food allergy impact scale; 1, meal preparation; 2, family social activities; 3, caregiver-supervised child social activities; 4, family relations; 5, caregiver stress and free time; 6, school or structured

activities; 7, autonomous child social activities; 8, employment and finances; \*significant  $p < 0.05$ ; \*\*significant  $p < 0.001$ .

Table 4. Comparison of Each F-COPES Subscale to the Caregiver of Indonesian Children with Food Allergy

F-COPES	Reframing		Spiritual Support		Mobilizing Family Support		Passive Appraisal	
	95% CI	p	95% CI	p	95% CI	p	95% CI	p
Social Support	-6.14 -- 0.52	0.021*	11.81 -- 16.95	0.000**	14.62 -- 20.03	0.000**	16.76 -- 22.01	0.000**
Reframing			16.57 -- 18.85	0.000**	19.36 -- 21.95	0.000**	21.24 -- 24.18	0.000**
Spiritual Support					2.04 -- 3.85	0.000**	3.87 -- 6.13	0.000**
Mobilizing Family Support							0.96 -- 3.15	0.000**

Abbreviation: F-COPES, family crisis oriented personal evaluation scales; \*significant  $p < 0.05$ ; \*\*significant  $p < 0.001$ .

Caregiver's stress and elimination diet in Indonesian children with food allergy, where the PSS value of the participant with the successful elimination diet was  $44.17 \pm 5.65$ , while the PSS value of unsuccessful diet was  $46.16 \pm 3.61$  ( $t = -0.145$ ; 95 % CI -0.014 - 0.012;  $p = 0.886$ ; Table 2).

#### Correlation between Family's Daily Activities on Elimination Diet in Indonesian Children with Food Allergy

The FAIS value ranged from 2.98 – 6.75, where the mean value was  $5.30 \pm 0.75$  with a median value of 5.26 (5.08 – 5.66). The mean value of each FAIS subscale in the successful elimination diet group was as follows: meal preparation ( $5.86 \pm 0.85$ ), family social activities ( $5.58 \pm 1.05$ ), caregiver-supervised child social activities ( $3.64 \pm 2.77$ ), family relations ( $4.82 \pm 1.40$ ), caregiver's stress and free time ( $4.82 \pm 1.15$ ), school or structured activities ( $3.33 \pm 2.35$ ), autonomous child social activities ( $4.14 \pm 2.52$ ), and employment and finances ( $4.91 \pm 1.49$ ). Meanwhile, the FAIS value in the unsuccessful diet elimination group was as follows: meal preparation ( $6.37 \pm 0.76$ ), family social activities ( $6.10 \pm 0.82$ ), caregiver-supervised child social activities ( $4.92 \pm 2.12$ ), family relations ( $5.22 \pm 1.61$ ), caregiver's stress and free time

( $4.84 \pm 1.33$ ), school or structured activities ( $3.79 \pm 2.30$ ), autonomous child social activities ( $5.83 \pm 1.30$ ), and employment and finances ( $5.61 \pm 1.10$ ). There was a significant correlation between FAIS and elimination diet in Indonesian children with food allergy, where the caregiver's average FAIS value in the successful elimination diet group was  $5.01 \pm 0.89$ , while the value in the unsuccessful group was  $5.52 \pm 0.53$  ( $t = -2.297$ ; 95% CI -0.179 – -0.012;  $p = 0.026$ ; Table 2).

There was a significant correlation of each FAIS subscale value in caregiver of Indonesian children with food allergy. The correlation was as follows: meal preparation versus (vs) family social activities ( $0.28 \pm 0.57$ ;  $p = 0.001$ ); meal preparation vs caregiver-supervised child social activities ( $0.90 \pm 1.40$ ;  $p < 0.001$ ); meal preparation vs family relations ( $1.10 \pm 0.90$ ;  $p < 0.001$ ); meal preparation vs caregiver's stress and free time ( $1.32 \pm 1.33$ ;  $p < 0.001$ ); meal preparation vs school or structured activities ( $1.37 \pm 1.30$ ;  $p < 0.001$ ); meal preparation vs autonomous child social activities ( $0.62 \pm 1.25$ ;  $p = 0.001$ ); meal preparation vs employment and finances ( $0.84 \pm 1.43$ ;  $p < 0.001$ ); family social activities vs caregiver-supervised child social activities ( $0.56 \pm 1.26$ ;  $p = 0.005$ ); family social activities vs family relations ( $0.82 \pm 1.45$ ;  $p < 0.001$ ); family social activities vs caregiver's

stress and free time ( $1.04 \pm 1.28$ ;  $p < 0.001$ ); caregiver-supervised child social activities vs school or structured activities ( $0.55 \pm 1.38$ ;  $p = 0.019$ ); caregiver's stress and free time vs autonomous child social activities ( $-0.79 \pm 1.93$ ;  $p = 0.005$ ); caregiver's stress and free time vs employment and finances ( $-0.47 \pm 1.58$ ;  $p = 0.031$ ); and school or structured activities vs autonomous child social activities ( $-0.80 \pm 1.93$ ;  $p = 0.007$ ). The highest correlation value was obtained by meal preparation vs school or structured activities of  $1.37 \pm 1.30$  (95% CI 0.96 - 1.77;  $p < 0.001$ ). Meanwhile, the lowest comparison value was obtained in subscale school or structured activities vs autonomous child social activities of  $-0.80 \pm 1.93$  (95% CI -1.34 - -0.23;  $p = 0.007$ ; Table 3).

#### Family's Coping Strategy on the Success of Elimination Diet in Indonesian Children with Food Allergy

The participant's F-COPES value ranged from 77 to 122, where the mean F-COPES value was  $104.54 \pm 11.42$  with a median value of 100 (96 - 115). The mean value of each F-COPES subscale was as follows: social support of  $39.08 \pm 6.77$  (successful elimination diet) and  $23.29 \pm 5.22$  (unsuccessful diet elimination); reframing of  $34.08 \pm 3.34$  (successful elimination diet) and  $33.06 \pm 3.46$  (unsuccessful elimination diet); spiritual support of  $16.46 \pm 1.56$  (successful elimination diet) and  $15.29 \pm 2.22$  (unsuccessful elimination diet); mobilizing family support of  $13.38 \pm 2.75$  (successful elimination diet) and  $12.45 \pm 2.69$  (unsuccessful elimination diet); and passive assessment of  $11.54 \pm 3.58$  (successful elimination diet) and  $10.23 \pm 2.79$  (unsuccessful elimination diet). There was a significant correlation between caregiver's coping strategy and elimination diet in Indonesian children with food allergy, where the mean value of family coping with successful diet elimination was  $116.29 \pm 3.91$ , and  $95.45 \pm 5.20$  for unsuccessful group ( $t = 16.154$ ; 95% CI 0.034 - 0.043;  $p < 0.001$ ; Table 2).

There was a significant comparison of each F-COPES subscale value of caregiver of Indonesian children with food allergy. The highest comparison value was obtained in subscale reframing vs passive appraisal of  $22.71 \pm 5.44$  (95% CI 21.24 - 24.18;  $p < 0.001$ ). Meanwhile, the lowest comparison value was found on subscale social support vs spiritual support of  $-3.33 \pm 10.40$  (95% CI -6.14 - -0.52;  $p = 0.021$ ; Table 4).

#### DISCUSSION

The main treatment of allergy is eliminating the allergen causing the problem, so the main management of food allergy is elimination diet [34]. Some literatures stated that elimination diet is an effective treatment for patients with food allergies [6,7,16]. The success of elimination diet in children with food allergy can increase the nutritional intake of children in the stages of development and growth and eliminate the signs of food allergy symptoms [16,17]. The success of elimination diet is expected to prevent the occurrence of food allergy reactions, most of which are anaphylactic reactions [35]. The success of elimination diet in children is strongly influenced by the role of caregiver, as they are the closest people to the child. Any condition experienced by the caregiver greatly affects the child's condition, including the success of elimination diet in children with food allergy [11,36].

In this study, there was no significant correlation between caregiver's stress and the success of elimination diet in Indonesian children with food allergy. Previous

research pointed out that parents or caregivers of children with food allergy experienced increased stress [11] that eventually had negative effect on the knowledge, attitudes and behavior of the caregiver in assisting children in carrying out the elimination diet [37]. Caregivers with high stress level have difficulty to adapt to changes in lifestyle of elimination diet for children with food allergy, because the allergy-causing food must be stopped and replaced with those that have similar nutritional content but are hypoallergenic. These conditions result in increased costs and the need for dietary consultation with a nutritionist for the selection of foods that are good for consumption during elimination diet [11,38]. In addition, excessive stress leads to incorrect treatment chosen by caregivers [39]. Thus, caregiver's stress directly and indirectly affects the success of elimination diet in children with food allergy [11,37,38].

This study found that family's activities had a significant relationship with the success of elimination diet in Indonesian children with food allergy. Previous studies mentioned that food allergies greatly affect daily activities, in which the caregiver must pay attention to the labels of food items that children will consume [12]. In addition, caregivers of children with food allergy are forced to limit children's activities, especially in public activities (attending parties, eating together outside) [12,13,40]. Caregivers should pay extra attention to children at school because their risk of eating food allergens is very high [12,30]. Activities such as sleepovers and camps are also limited by the caregiver to minimize the risk of exposure to food allergens [15]. Daycare, birthday parties, field trips, and going to restaurants are also reported to be limited by the caregiver to prevent children with food allergies from consuming food allergens [12,41]. Activity limitation by the caregiver can increase the success of elimination diet by 70% [12,13].

Moreover, this study found a significant correlation between family's coping strategy and the success of elimination diet in Indonesian children with food allergy. Family's coping is important in overcoming the problem of food allergy in children, because family coping can determine the coping strategies used to seek treatment for children with food allergy [42]. The use of family coping in elimination diet in children with food allergy is very complex, which requires an effective strategy in its implementation. On the other hand, prolonged elimination diet causes psychological problems. Elimination diet in adults is effective to reduce food allergy load, but it requires further research in children patients regarding adaptive coping strategies [43]. Family coping affects the success of elimination diet in children with food allergies because family support is needed to play a crucial role in eliminating children's diets [6,7,11], because adapting children to eat hypoallergen foods to avoid allergic foods requires a long process [44,45]. However, some limitations should be noted. First, it is hoped that further research will investigate elimination diet in Indonesian children with food allergy by using a quasi / true experimental research design. Second, it is hoped the number of participants in future studies will be greater and use a randomized control trail technique so that the composition between the intervention group and control group is more balanced and minimizes bias. Third, it is hoped that further research classifies the ages

of children with food allergy in order to identify the most effective elimination diet given to which age group.

#### CONCLUSION

Elimination diet is an effective treatment for children with food allergy, and the role of caregiver greatly affects its success. Family activities greatly affect the success of elimination diet in children with food allergy because limiting social activity greatly supports its success. Likewise, family coping greatly affects elimination diet in Indonesian children with food allergy because the coping used by families in responding to food allergies is critical to success because children cannot make decisions. While caregiver's stress greatly influences the success of elimination diet in Indonesian children with food allergy, this study found no significant correlation between caregiver's stress and elimination diet in Indonesian children with food allergy.

#### ACKNOWLEDGEMENT

The authors would like to thank Dr. Windhu Purnomo, MS, Faculty of Public Health, Universitas Airlangga, Surabaya, Indonesia for his technical assistance for this study, particularly in the statistical analysis. We would also like to thank Fis Citra Ariyanto as our manuscript editor.

#### CONFLICT OF INTEREST

None.

#### Author's Contribution

All authors contributed toward data analysis, drafting and revising the paper, gave final approval of the version to be published and agree to be accountable for all aspects of the work.

#### FUNDING

None

#### REFERENCES

1. Walczak M, Grzelak T, Kramkowska M, Czyżewska K (2014) Food allergies in children—aspects of epidemiology and diet management. *Journal of Medical Science* 83 (3):255-259
2. Loh W, Tang MLK (2018) The Epidemiology of Food Allergy in the Global Context. *Int J Environ Res Public Health* 15 (9):2043. doi:10.3390/ijerph15092043
3. Savage J, Johns CB (2015) Food allergy: epidemiology and natural history. *Immunol Allergy Clin North Am* 35 (1):45-59. doi:10.1016/j.jiac.2014.09.004
4. Munasir Z, Muktiarti D (2013) The management of food allergy in Indonesia. *Asia Pac Allergy* 3 (1):23-28. doi:10.5415/apallergy.2013.3.1.23
5. Sumadiono, Muktiarti D, Setiabudiawan B, Irsa L, Wati KDK, Majangsari RGD (2015) Rekomendasi Ikatan Dokter Anak Indonesia Pencegahan Primer Alergi. 1 edn. Unit Kerja Koordinasi Alergi Imunologi: Ikatan Dokter Anak Indonesia,
6. Berry MJ, Adams J, Voutilainen H, Feustel PJ, Celestin J, Järvinen KM (2015) Impact of elimination diets on growth and nutritional status in children with multiple food allergies. *Pediatric allergy and immunology* : official publication of the European Society of Pediatric Allergy and Immunology 26 (2):133-138. doi:10.1111/pai.12348
7. Venter C, Mazzocchi A, Maslin K, Agostoni C (2017) Impact of elimination diets on nutrition and growth in children with multiple food allergies. *Current opinion in allergy and clinical immunology* 17 (3):220-226. doi:10.1097/aci.0000000000000358
8. Bora S, Rindfleisch JA (2018) Chapter 86 - The Elimination Diet. In: Raket D (ed) *Integrative Medicine* (Fourth Edition). Elsevier, pp 849-862.e846. doi:https://doi.org/10.1016/B978-0-323-35868-2.00086-4
9. Le T-M, Zijlstra WT, van Opstal EY, Knol MJ, L'Hoir MP, Knulst AC, Pasmans SGMA (2013) Food avoidance in children with adverse food reactions: Influence of anxiety and clinical parameters. *Pediatric Allergy and Immunology* 24 (7):650-655. doi:10.1111/pai.12114
10. Allen CW, Kemp AS, Campbell DE (2009) Dietary advice, dietary adherence and the acquisition of tolerance in egg-allergic children: a 5-yr follow-up. *Pediatric Allergy and Immunology* 20 (3):213-218. doi:10.1111/j.1399-3038.2008.00784.x
11. Birdi G, Cooke R, Knibb R (2016) Quality of Life, Stress, and Mental Health in Parents of Children with Parentally Diagnosed Food Allergy Compared to Medically Diagnosed and Healthy Controls. *J Allergy (Cairo)* 2016:1497375-1497375. doi:10.1155/2016/1497375
12. Bollinger ME, Dahlquist LM, Mudd K, Sonntag C, Dillinger L, McKenna K (2006) The impact of food allergy on the daily activities of children and their families. *Annals of Allergy, Asthma & Immunology* 96 (3):415-421. doi:10.1016/S1081-1206(10)60908-8
13. Cummings AJ, Knibb RC, King RM, Lucas JS (2010) The psychosocial impact of food allergy and food hypersensitivity in children, adolescents and their families: a review. *Allergy* 65 (8):933-945. doi:10.1111/j.1398-9995.2010.02342.x
14. Muñoz-Furlong A (2003) Daily coping strategies for patients and their families. *Pediatrics* 111 (6 Pt 3):1654-1661
15. Klinnert MD, McQuaid EL, Fedele DA, Faino A, Strand M, Robinson J, Atkins D, Fleischer DM, Hourihane JOB, Cohen S, Fransen H (2015) Children's Food Allergies: Development of the Food Allergy Management and Adaptation Scale. *J Pediatr Psychol* 40 (6):572-580. doi:10.1093/jpepsy/jsv009
16. Kim J, Kwon J, Noh G, Lee SS (2013) The effects of elimination diet on nutritional status in subjects with atopic dermatitis. *Nutr Res Pract* 7 (6):488-494. doi:10.4162/nrp.2013.7.6.488
17. Lozinsky AC, Meyer R, De Koker C, Dziubak R, Godwin H, Reeve K, Dominguez Ortega G, Shah N (2015) Time to symptom improvement using elimination diets in non-IgE-mediated gastrointestinal food allergies. *Pediatric allergy and immunology* : official publication of the European Society of Pediatric Allergy and Immunology 26 (5):403-408. doi:10.1111/pai.12404
18. Boyce JA, Assa'ad A, Burks AW, Jones SM, Sampson HA, Wood RA, Plaut M, Cooper SF, Fenton MJ, Arshad SH, Bahna SL, Beck LA, Byrd-Bredbenner C, Camargo CA, Jr, Eichenfield L, Furuta GT, Hanifin JM, Jones C, Kraft M, Levy BD, Lieberman P, Lucciolli S, McCall KM, Schneider LC, Simon RA, Simons FER, Teach SJ, Yawn BP, Schwanger JM (2011) Guidelines for the diagnosis and management of food allergy in the

- United States: summary of the NIAID-sponsored expert panel report. *Nutr Res* 31 (1):61-75. doi:10.1016/j.nutres.2011.01.001
19. Abrams EM, Sicherer SH (2016) Diagnosis and management of food allergy. *CMAJ* 188 (15):1087-1093. doi:10.1503/cmaj.160124
  20. Setia MS (2016) Methodology Series Module 2: Case-control studies--an efficient observational study design. *Nephron Clinical practice* 114 (1):c1-4. doi:10.1159/000242442
  22. Noronha L, Fox A, Du Toit G, Lack G (2017) Diagnosis and management of allergic rhinitis in children. *Prescriber* 28 (12):13-19. doi:10.1002/psb.1631
  23. Papadopoulos NG, Arakawa H, Carlsen KH, Custovic A, Gern J, Lemanske R, Le Souef P, Mäkelä M, Roberts G, Wong G, Zar H, Akdis CA, Bacharier LB, Baraldi E, van Bever HP, de Blic J, Boner A, Burks W, Casale TB, Castro-Rodriguez JA, Chen YZ, El-Gamal YM, Everard ML, Frischer T, Geller M, Gereda J, Goh DY, Guilbert TW, Hedlin G, Heymann PW, Hong SJ, Hossny EM, Huang JL, Jackson DJ, de Jongste JC, Kalayci O, Ait-Khaled N, Kling S, Kuna P, Lau S, Ledford DK, Lee SJ, Liu AH, Lockey RF, Lørdrup-Carlsen K, Lötvalld J, Morikawa A, Nieto A, Paramesh H, Pawankar R, Pohunek P, Pongracic J, Price D, Robertson C, Rosario N, Rossenwasser LJ, Sly PD, Stein R, Stick S, Szeffler S, Taussig LM, Valovirta E, Vichyanond P, Wallace D, Weinberg E, Wennergren G, Wildhaber J, Zeiger RS (2012) International consensus on (ICON) pediatric asthma. *Allergy* 67 (8):976-997. doi:10.1111/j.1398-9995.2012.02865.x
  24. Severity Scoring of Atopic Dermatitis: The SCORAD Index (1993). *Dermatology* 186 (1):23-31. doi:10.1159/000247298
  25. Rodriguez J, Crespo JF (2012) Chapter 6 - Food-induced Urticaria and Angioedema. In: James JM, Burks W, Eigenmann P (eds) *Food Allergy*. W.B. Saunders, Edinburgh, pp 75-82. doi:https://doi.org/10.1016/B978-1-4377-1992-5.00006-5
  26. Worm M, Reese I, Ballmer-Weber B, Beyer K, Bischoff SC, Classen M, Fischer PJ, Fuchs T, Huttegger I, Jappe U, Klimek L, Koletzko B, Lange L, Lepp U, Mahler V, Niggemann B, Rabe U, Raithe M, Saloga J, Schäfer C, Schnadt S, Schreiber J, Szépfalusi Z, Treudler R, Wagenmann M, Watzl B, Werfel T, Zuberbier T, Kleine-Tebbe J (2015) Guidelines on the management of IgE-mediated food allergies: S2k-Guidelines of the German Society for Allergology and Clinical Immunology (DGAKI) in collaboration with the German Medical Association of Allergologists (AeDA), the German Professional Association of Pediatricians (BVKJ), the German Allergy and Asthma Association (DAAB), German Dermatological Society (DDG), the German Society for Nutrition (DGE), the German Society for Gastroenterology, Digestive and Metabolic Diseases (DGVS), the German Society for Oto-Rhino-Laryngology, Head and Neck Surgery, the German Society for Pediatric and Adolescent Medicine (DGKJ), the German Society for Pediatric Allergology and Environmental Medicine (GPA), the German Society for Pneumology (DGP), the German Society for Pediatric Gastroenterology and Nutrition (GPGE), German Contact Allergy Group (DKG), the Austrian Society for Allergology and Immunology (Æ-GAI), German Professional Association of Nutritional Sciences (VDOE) and the Association of the Scientific Medical Societies Germany (AWMF). *Allergo J Int* 24:256-293. doi:10.1007/s40629-015-0074-0
  27. Pontoppidan M, Nielsen T, Kristensen IH (2018) Psychometric properties of the Danish Parental Stress Scale: Rasch analysis in a sample of mothers with infants. *PLoS One* 13 (11):e0205662-e0205662. doi:10.1371/journal.pone.0205662
  28. Berry JO, Jones WH (1995) The Parental Stress Scale: Initial Psychometric Evidence. *Journal of Social and Personal Relationships* 12 (3):463-472. doi:10.1177/0265407595123009
  29. Aggarwal A, Unni K, Prasad S (2018) A Comparative Study of Stress, Anxiety & Work Impairment in Parents of Children with Conduct Disorder. *Journal of Indian Association for Child & Adolescent Mental Health* 14 (4)
  30. Hoehn JL, Dahlquist LM, Hahn AL, Bollinger ME (2017) Parents of Children With Food Allergy: Gender Differences in Perceived Impact and Perceived Food Allergy Severity. *J Pediatr Psychol* 42 (2):186-197. doi:10.1093/jpepsy/jsw059
  31. Eaton PM, Davis BL, Hammond PV, Condon EH, McGee ZT (2011) Coping Strategies of Family Members of Hospitalized Psychiatric Patients. *Nursing Research and Practice* 2011. doi:10.1155/2011/392705
  32. Putera AM, Irwanto I, Maramis MM, Prasetyo RV, Soemyarso NA, Noer MS (2020) Effect of Mental Health Problems on the Quality of Life in Children with Lupus Nephritis. *Neuropsychiatr Dis Treat* 16:1583-1593. doi:10.2147/NDT.S250373
  33. Putera AM, Irwanto, Maramis MM (2020) Quality-of-Life (QoL) of Indonesian Children Living with HIV: The Role of Caregiver Stigma, Burden of Care, and Coping. *HIV/AIDS-Research and Palliative Care* 12:573-581. doi:10.2147/HIV.S269629
  34. Niggemann B, Heine RG (2006) Who should manage infants and young children with food induced symptoms? *Arch Dis Child* 91 (5):379-382. doi:10.1136/adc.2005.086603
  35. Staden U, Rolinck-Werninghaus C, Brewe F, Wahn U, Niggemann B, Beyer K (2007) Specific oral tolerance induction in food allergy in children: efficacy and clinical patterns of reaction. *Allergy* 62 (11):1261-1269. doi:10.1111/j.1398-9995.2007.01501.x
  36. Scherer E, Hagaman A, Chung E, Rahman A, O'Donnell K, Maselko J (2019) The relationship between responsive caregiving and child outcomes: evidence from direct observations of mother-child dyads in Pakistan. *BMC Public Health* 19 (1):252-252. doi:10.1186/s12889-019-6571-1
  37. Wright RJ, Berin MC (2014) Unlocking the stress-allergy puzzle: need for a more comprehensive stress model. *Annals of allergy, asthma & immunology : official publication of the American College of Allergy, Asthma, & Immunology* 113 (1):1-2. doi:10.1016/j.anaai.2014.05.007
  38. Lau G-Y, Patel N, Umasunthar T, Gore C, Warner JO, Hanna H, Phillips K, Mohd Zaki A, Hodes M, Boyle RJ (2014) Anxiety and stress in mothers of food-allergic children. *Pediatric Allergy and Immunology* 25 (3):236-242. doi:10.1111/pai.12203
  39. Begeen FM, Barnett J, Barber M, Payne R, Gowland MH, Lucas JS (2017) Parents' and caregivers' experiences



- and behaviours when eating out with children with a food hypersensitivity. *BMC Public Health* 18 (1):38-38. doi:10.1186/s12889-017-4594-z
40. Meyer R, Godwin H, Dziubak R, Panepinto JA, Foong R-XM, Bryon M, Lozinsky AC, Reeve K, Shah N (2017) The impact on quality of life on families of children on an elimination diet for Non-immunoglobulin E mediated gastrointestinal food allergies. *World Allergy Organ J* 10 (1):8-8. doi:10.1186/s40413-016-0139-7
41. Vargas PA, Sicherer SH, Christie L, Keaveny M, Noone S, Watkins D, Carlisle SK, Jones SM (2011) Developing a food allergy curriculum for parents. *Pediatric allergy and immunology : official publication of the European Society of Pediatric Allergy and Immunology* 22 (6):575-582. doi:10.1111/j.1399-3038.2011.01152.x
42. Aika S, Ito M, Yamamoto Y (2017) Food allergy response capabilities of mothers and related factors. *Nursing & health sciences* 19 (3):340-350. doi:10.1111/nhs.12351
43. Weiss D, Marsac ML (2016) Coping and posttraumatic stress symptoms in children with food allergies. *Annals of allergy, asthma & immunology : official publication of the American College of Allergy, Asthma, & Immunology* 117 (5):561-562. doi:10.1016/j.anai.2016.08.022
44. Polloni L, Toniolo A, Lazzarotto F, Baldi I, Foltran F, Gregori D, Muraro A (2013) Nutritional behavior and attitudes in food allergic children and their mothers. *Clin Transl Allergy* 3 (1):41-41. doi:10.1186/2045-7022-3-41
45. Shanahan L, Zucker N, Copeland WE, Costello EJ, Angold A (2014) Are children and adolescents with food allergies at increased risk for psychopathology? *J Psychosom Res* 77 (6):468-473. doi:10.1016/j.jpsychores.2014.10.005

## 08. The Success of Elimination Diet in Indonesian

---

### ORIGINALITY REPORT

---

17%

SIMILARITY INDEX

12%

INTERNET SOURCES

14%

PUBLICATIONS

0%

STUDENT PAPERS

---

### PRIMARY SOURCES

---

1	<a href="http://spiral.imperial.ac.uk">spiral.imperial.ac.uk</a> Internet Source	1%
2	Aliki Kalmpourtzidou, Ioannis Xinias, Charalampos Agakidis, Antigoni Mavroudi et al. "Diet Quality: A Neglected Parameter in Children With Food Allergies. A Cross-Sectional Study", <i>Frontiers in Pediatrics</i> , 2021 Publication	1%
3	<a href="http://www.medicopublication.com">www.medicopublication.com</a> Internet Source	1%
4	<a href="http://www.scienceopen.com">www.scienceopen.com</a> Internet Source	1%
5	Sylvia Nabila Azwa Ambad, Khairiah Mazdah Kalimin, Dayang Haryani Diana Ag Damit, Jasmine Vivienne Andrew. "The mediating effect of psychological empowerment on leadership styles and task performance of academic staff", <i>Leadership &amp; Organization Development Journal</i> , 2021 Publication	1%

---

6

[primarycare.imedpub.com](http://primarycare.imedpub.com)

Internet Source

1 %

7

Royke Tony Kalalo, Sasanti Yuniar, Fis Citra Ariyanto. "Effect of parental skills-based psychoeducation intervention on parental stress index and severity of children with autism spectrum disorders: A pilot study", *Annals of Medicine and Surgery*, 2021

Publication

1 %

8

Isnin Anang Marhana, Kadek Widianiti, Ety Hary Kusumastuti. "Conformity of Fine Needle Aspiration Biopsy (FNAB) and Core Needle Biopsy (CNB) in peripheral lung tumor patients: A cross-sectional study", *Annals of Medicine and Surgery*, 2022

Publication

1 %

9

Fedele, David A., Elizabeth L. McQuaid, Anna Faino, Matthew Strand, Sophia Cohen, Jane Robinson, Dan Atkins, Jonathan O'B Hourihane, and Mary D. Klinnert. "Patterns of Adaptation to Children's Food Allergies", *Allergy*, 2015.

Publication

1 %

10

Herley Windo Setiawan, Resti Yudhawati, Irmu Syafaah. "Association between serum PGE2 levels and degree of acid-fast bacilli positivity in sputum of pulmonary tuberculosis

1 %

patients", Annals of Medicine and Surgery,  
2021

Publication

---

11 [www.medscape.com](http://www.medscape.com) 1 %  
Internet Source

---

12 [assets.researchsquare.com](http://assets.researchsquare.com) <1 %  
Internet Source

---

13 [balimedicaljournal.org](http://balimedicaljournal.org) <1 %  
Internet Source

---

14 [pure.rug.nl](http://pure.rug.nl) <1 %  
Internet Source

---

15 Mizuho Konishi, Junko Kamoshida, Hisako Ogasawara, Mieko Takahashi, Yukihiro Ohya. "Tentative development of a psychoeducational program for alleviating psychological burden of mothers of young children with milk allergy", Journal of Health Psychology Research, 2019 <1 %  
Publication

---

16 [etd.lib.fsu.edu](http://etd.lib.fsu.edu) <1 %  
Internet Source

---

17 [www.researchgate.net](http://www.researchgate.net) <1 %  
Internet Source

---

18 [uir.unisa.ac.za](http://uir.unisa.ac.za) <1 %  
Internet Source

---

[dspace.nmmu.ac.za:8080](http://dspace.nmmu.ac.za:8080)

19

Internet Source

<1 %

20

[medicopublication.com](http://medicopublication.com)

Internet Source

<1 %

21

"Abstracts of the 18th Congress of the International Pediatric Nephrology Association, Venice (Italy), October 2019", *Pediatric Nephrology*, 2019

Publication

<1 %

22

[link.springer.com](http://link.springer.com)

Internet Source

<1 %

23

[e-space.mmu.ac.uk](http://e-space.mmu.ac.uk)

Internet Source

<1 %

24

[uilis.unsyiah.ac.id](http://uilis.unsyiah.ac.id)

Internet Source

<1 %

25

[www.pubfacts.com](http://www.pubfacts.com)

Internet Source

<1 %

26

"Allergy and Asthma", Springer Science and Business Media LLC, 2019

Publication

<1 %

27

Danielle Cole, Danielle Weiss, Kristen L. Kohser, Claudine Jones et al. "The Cellie Coping Kit for Children with Food Allergy: A Pilot Study", *Pediatric Allergy, Immunology, and Pulmonology*, 2017

Publication

<1 %

28

[clinisoft.org](http://clinisoft.org)

Internet Source

<1 %

---

29

[digital.library.unt.edu](http://digital.library.unt.edu)

Internet Source

<1 %

---

30

[myschool.ng](http://myschool.ng)

Internet Source

<1 %

---

31

[www.cambridge.org](http://www.cambridge.org)

Internet Source

<1 %

---

32

Cansin Sackesen, Derya Ufuk Altintas, Aysen Bingol, Gulbin Bingol et al. "Current Trends in Tolerance Induction in Cow's Milk Allergy: From Passive to Proactive Strategies", *Frontiers in Pediatrics*, 2019

Publication

<1 %

---

33

Christina A. Di Bartolo, Maureen K. Braun. "Pediatrician's Guide to Discussing Research with Patients", Springer Science and Business Media LLC, 2017

Publication

<1 %

---

34

[aacijournal.biomedcentral.com](http://aacijournal.biomedcentral.com)

Internet Source

<1 %

---

35

[blog.balimedicaljournal.org](http://blog.balimedicaljournal.org)

Internet Source

<1 %

---

36

[commons.emich.edu](http://commons.emich.edu)

Internet Source

<1 %

---

37	<a href="http://create.canterbury.ac.uk">create.canterbury.ac.uk</a> Internet Source	<1 %
38	<a href="http://iovs.arvojournals.org">iovs.arvojournals.org</a> Internet Source	<1 %
39	<a href="http://onlinelibrary.wiley.com">onlinelibrary.wiley.com</a> Internet Source	<1 %
40	<a href="http://ro.uow.edu.au">ro.uow.edu.au</a> Internet Source	<1 %
41	<a href="http://worldwidescience.org">worldwidescience.org</a> Internet Source	<1 %
42	<a href="http://www.foodprotect.org">www.foodprotect.org</a> Internet Source	<1 %
43	<a href="http://www.mdpi.com">www.mdpi.com</a> Internet Source	<1 %
44	A. J. Cummings. "The psychosocial impact of food allergy and food hypersensitivity in children, adolescents and their families: a review", <i>Allergy</i> , 02/2010 Publication	<1 %
45	"Pediatric Food Allergy", Springer Science and Business Media LLC, 2020 Publication	<1 %
46	Alina Kanikowska, Susanne Janisz, Dorota Mańkowska-Wierzbicka, Marcin Gabryel, Agnieszka Dobrowolska, Piotr Eder.	<1 %

"Management of Adult Patients with Gastrointestinal Symptoms from Food Hypersensitivity—Narrative Review", Journal of Clinical Medicine, 2022

Publication

---

47

C. Venter, M. Groetch, M. Netting, R. Meyer. "A patient-specific approach to develop an exclusion diet to manage food allergy in infants and children", Clinical & Experimental Allergy, 2018

Publication

---

<1 %

48

Emily M. Steiner, Danielle Weiss Byrne, Lynnda M. Dahlquist, Amy L. Hahn, Mary Elizabeth Bollinger. "Which foods should a child with food allergy avoid? The role of parental knowledge in food avoidance appraisals", Children's Health Care, 2020

Publication

---

<1 %

49

Linda Herbert, Eyal Shemesh, Bruce Bender. "Clinical Management of Psychosocial Concerns Related to Food Allergy", The Journal of Allergy and Clinical Immunology: In Practice, 2016

Publication

---

<1 %

50

Michael A. Golding, Nina V. Gunnarsson, Roelinde Middelveld, Staffan Ahlstedt, Jennifer LP Protudjer. "A scoping review of the caregiver burden of pediatric food allergy",

<1 %



# Annals of Allergy, Asthma & Immunology, 2021

Publication

51

[conservancy.umn.edu](http://conservancy.umn.edu)

Internet Source

<1 %

52

[hdl.handle.net](http://hdl.handle.net)

Internet Source

<1 %

53

[publications.aston.ac.uk](http://publications.aston.ac.uk)

Internet Source

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

# 08. The Success of Elimination Diet in Indonesian

---

GRADEMARK REPORT

---

FINAL GRADE

**/100**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---

PAGE 6

---

PAGE 7

---

PAGE 8

---