

# Source details

## Systematic Reviews in Pharmacy

Scopus coverage years: from 2010 to Present

(coverage discontinued in Scopus)

Publisher: Wolters Kluwer Health

ISSN: 0975-8453 E-ISSN: 0976-2779

Subject area: Pharmacology, Toxicology and Pharmaceutics: Pharmaceutical Science

Source type: Journal

CiteScore 2019

3.9

SJR 2019

0.424

SNIP 2019

3.180

[View all documents >](#) [Set document alert](#) [Save to source list](#)

[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)

### Improved CiteScore methodology

CiteScore 2019 counts the citations received in 2016-2019 to articles, reviews, conference papers, book chapters and data papers published in 2016-2019, and divides this by the number of publications published in 2016-2019. [Learn more >](#)

### CiteScore 2019

$$3.9 = \frac{463 \text{ Citations 2016 - 2019}}{119 \text{ Documents 2016 - 2019}}$$

Calculated on 06 May, 2020

### CiteScoreTracker 2020

$$1.3 = \frac{2,081 \text{ Citations to date}}{1,580 \text{ Documents to date}}$$

Last updated on 06 April, 2021 • Updated monthly

### CiteScore rank 2019

Category	Rank	Percentile
Pharmacology, Toxicology and Pharmaceutics	#53/169	68th
Pharmaceutical Science		

[View CiteScore methodology >](#) [CiteScore FAQ >](#) [Add CiteScore to your site >](#)

## About Scopus

- What is Scopus
- Content coverage
- Scopus blog
- Scopus API
- Privacy matters

## Language

- 日本語に切り替える
- 切换到简体中文
- 切换到繁體中文
- Русский язык

## Customer Service

- Help
- Contact us



## Systematic Reviews in Pharmacy

**COUNTRY**

India

Universities and  
research institutions in  
India**SUBJECT AREA AND  
CATEGORY**Pharmacology,  
Toxicology and  
Pharmaceutics  
Pharmaceutical  
Science**PUBLISHER**Wolters Kluwer Medknow  
Publications**H-INDEX****14****PUBLICATION TYPE**

Journals

**ISSN**

09758453, 09762779

**COVERAGE**

2010-2020

**INFORMATION**[Homepage](#)[How to publish in this  
journal](#)[editor@sysrevpharm.org](mailto:editor@sysrevpharm.org)**SCOPE**

The journal covers and publishes all articles related to Pharmacy subjects including some of the allied subjects. Articles with timely interest and newer research concepts will be given more preference. This journal also publishes manuscripts related to agriculture and agriculture sciences. Topics include all aspects of crop and animal physiology, modelling of crop and animal systems, the scientific underpinning of agronomy and husbandry, animal welfare and behaviour, soil science, plant and animal product quality, plant and animal nutrition, engineering solutions, decision support systems, land use, environmental impacts of agriculture and forestry, impacts of climate change, rural biodiversity, experimental design and statistical analysis, and the application of new analytical and study methods (including genetic diversity and molecular biology approaches).

 [Join the conversation about this journal](#)

FIND SIMILAR JOURNALS ?

1  
Tropical Journal of  
Pharmaceutical Research  
NGA

**37%**  
similarity

2  
Zhongguo Zhongyao Zazhi  
CHN


**35%**  
similarity

3  
Journal of Applied  
Pharmaceutical Science  
IND


**35%**  
similarity

4  
Internationa  
Pharmacolo  
PAK


SJR




Citations per document




Total Cites Self-Cites




External Cites per Doc Cites per Doc




% International Collaboration



Citable documents Non-citable documents



Cited documents Uncited documents







← Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scimagojr.com">
```

Metrics based on Scopus® data as of April 2020

M **MIK** 3 months ago

Dear Sir this Journal SRP is out of scoups please up date your data because the researchers is dependent on SJR in World and SJR is one of best.

reply

SCImago Team



**Melanie Ortiz** 2 months ago

Dear Sir/Madam,

Thank you very much for your comment.

All the metadata have been provided by Scopus /Elsevier in their last update sent to SCImago, including the Coverage's period data. The SJR for 2019 was released on 11 June 2020. We suggest you consult the Scopus database directly to see the current index status as SJR is a static image of Scopus, which is changing every day.

For further information, please contact Scopus support:

[https://service.elsevier.com/app/answers/detail/a\\_id/14883/kw/scimago/supporthub/scopus/](https://service.elsevier.com/app/answers/detail/a_id/14883/kw/scimago/supporthub/scopus/)

Best Regards, SCImago Team

J **Juliaans Marantika** 3 months ago

Dear SCImago Team,

Would you please write some reason why my Article Politeness in Foreign Language Teaching and Learning Process has been published in this Journal? I need it to provide an explanation regarding the relevance of the journal to my field of specialization, namely language

reply

SCImago Team



**Melanie Ortiz** 3 months ago

Dear Juliaans,

thank you for contacting us.

We are sorry to tell you that SCImago Journal & Country Rank is not a journal. SJR is a portal with scientometric indicators of journals indexed in Elsevier/Scopus.

Unfortunately, we cannot help you with your request, we suggest you contact the journal's

ISSN 1472-2148  
Vol. 17, No. 1, 2009

# Systematic Reviews in Pharmacy


[www.sysrevpharm.org](http://www.sysrevpharm.org)



The Society for Systematic Reviews in Pharmacy (SSRP) is an international organization of researchers, clinicians, and students who are interested in the development and practice of systematic reviews in pharmacy. The Society was founded in 1998 and has since then been working to promote the use of systematic reviews in pharmacy practice and research. The Society's main activities include the organization of international conferences, the publication of a journal, and the provision of training and support to researchers and clinicians.

Wiley-Blackwell  
Publishers





Online First

Current Issue

Archive

Aims and Scope

Abstracting & Indexing

Most Accessed Articles

Most Downloaded Articles

ORCID

Crossref

OPEN ACCESS

CC creative commons

## Editorial Board

### Editor-in-Chief

Dr. Ayad F. Alkaim  
University of Babylon, College of Science for Women, Babylon, Scopus Author ID: 55255310600, Iraq

### Editorial Board

Dr. Aygul Z. Ibatova  
Department of Natural Sciences  
Tyumen Industrial University, Scopus Author ID: 57191110632 <http://orcid.org/0000-0003-0565-8533>, Russia

Dr Ahmad Faisal Ismail  
Kulliyah of Dentistry  
International Islamic University Malaysia, Kuantan Campus, 25200 Kuantan, Pahang,  
Scopus Author ID: 35388596700, Malaysia

Dr. Huiliang ZHAO, Ph.D  
Guizhou Minzu University, Huaxi District, Guiyang, China

Dr. Mohd Armi Abu Samah  
International Islamic University Malaysia, (IIUM) 25200 Kuantan Pahang

Dr. Baded ramji  
Sri Lanka

Dr. Chris randea  
South Africa

Dr. Yingwen ZHAO  
Researcher of Guizhou Rural Economic and Social Development Research Institute,  
China

Dr. Li Zihan, Ph.D  
University of Glasgow, UK

Gabriela Cioca  
Faculty of Medicine, Pharmacology Department  
Lucian Blaga University of Sibiu, Romania, Lucian Blaga street, no 2A, Sibiu, Romania

Dariusz Nowak  
Municipal Hospital, Mickiewicz street no 12, 42-200 Czestochowa, Poland

Aleksandra Zyska  
Faculty of Medicine, Department of Physiology  
Opole University, Oleska street no 48, 45-052 Opole, Poland

Katarzyna Sznajder  
Faculty of Medicine, Clinical Department of Diagnostic Imaging  
Opole University, Oleska street no 48, 45-052 Opole, Poland

Jacek Jankowski  
Faculty of Medicine, Department of Family Medicine and Public Health  
Opole University, Oleska street no 48, 45-052 Opole, Poland

Luciano Benedini  
Universidad Nacional del Sur (National University of South-UNS), Bahía Blanca 8000,  
Argentina

Paula Messina  
Departamento de Biología  
Universidad Nacional del Sur (National University of South-UNS), Bioquímica y Farmacia,  
Bahía Blanca 8000, Argentina

Submit your article

Login

Register

### Most Viewed Articles

Dental Development between Assisted Reproductive Therapy (Art) and Natural Conceived Children: A Comparative Pilot Study Norzaiti Mohd Kenali, Naimah Hasanah Mohd Fathil, Norbasyirah Bohari, Ahmad Faisal Ismail, Roszaman Ramli SRP. 2020.

### Most Downloaded

Dental Development between Assisted Reproductive Therapy (Art) and Natural Conceived Children: A Comparative Pilot Study Norzaiti Mohd Kenali, Naimah Hasanah Mohd Fathil, Norbasyirah Bohari, Ahmad Faisal Ismail, Roszaman Ramli SRP. 2020.



According to new research from Rensselaer Polytechnic Institute (#RPI), the antiviral #remdesivir is 10 times more effective in treating cells infected with SARS-CoV-2 when combined with drugs currently used to treat hepatitis C virus (HCV).  
sysrevpharm.org

3h

Embed

View on Twitter

Michael Walsh  
Washington State University, College of Pharmacy and Pharmaceutical Sciences (CPPS),  
USA

Prof. Dr. Kittisak Jernsittiparsert  
Henan University, China

Amel Dawod Kamel Gudia, PhD  
Faculty of nursing  
Cairo University, Egypt

Arif Nur Muhammad Ansori  
Airlangga University, Scopus Author ID: 57195995342, <https://orcid.org/0000-0002-1279-3904>, Indonesia

Mohammed Nader Shalaby  
Suez Canal University, Associate Professor of Biological Sciences and Sports Health,  
Egypt

Dr. Faten Abo-Aziza Mohamed, PhD  
Associate Professor, Clinical Pathology and Stem Cell Research  
National Research Centre, Manager of Veterinary Division Central Lab (605), 33 El-  
Behoos St, Dokki, Cairo, Egypt

Professor Asim Ahmed Elnour Ahmed  
College of Pharmacy  
Al-Ain University of Science and Technology, UAE

S. Parasuraman, M.Pharm., Ph.D  
AIMST University, Malaysia

Copyright © 2021 Systematic Reviews in Pharmacy All Rights Reserved. Subject to change without notice from or liability to Systematic Reviews in Pharmacy



Copyright © 2021 Systematic Reviews in Pharmacy All Rights Reserved. Subject to change without notice from or liability to Systematic Reviews in Pharmacy. For best results, please use Internet Explorer or Google Chrome

#### POLICIES & JOURNAL LINKS

- Advertising Policy
- Author's Rights and Obligations
- Conflict of Interest Policy
- Digital Archiving & Preservation Policies
- Editorial Policies
- Peer Review Policy
- Editorial & Peer Review Process
- License Information
- Plagiarism Policy
- Privacy Policy
- Protection of Research Participants (Statement On Human Rights)
- Animal Rights
- Publishing Ethics
- Corrections, Retractions & Expressions of Concern
- Self-Archiving Policies
- Statement of Informed Consent
- Terms of Use







## 2020: Volume 11, Issue 11

### Research Article

#### Gadget Use, Pocket Money, and Snacking Habits of Children with and without Overweight/Obesity Problem in Surabaya, Indonesia

✎ *Trias Mahmudiono, Qonita Rachmah, Diah Indriani, Triska Susila Nindya, Calista Segalita, Susi Hidayah, Azizah Ajeng Pratiwi, Loh Su Peng*

SRP. 2020; 11(11): 1087 - 1090

» Abstract » PDF DOI: 10.31838/srp.2020.11.155

### Review Article

#### Ameliorative Effect Of Thiamine Pyrophosphate Against Cisplatin-Induced Reproductive System Damage Within Male Rats

✎ *Maitham Abd Ali Mnati, Bahir Abdul Razzaq Mshimesh, Mustafa Mohammed Ibraheem, Suzan Yousif Jasim*

SRP. 2020; 11(11): 396 - 403

» Abstract » PDF DOI: 10.31838/srp.2020.11.59

### Research Article

#### Neutrophil Extracellular Traps in Coronavirus Infection: Interaction Network Analysis

✎ *Amal Bouzid, Asmaa T Uthman, Noor N Al-Rawi, Natheer H Al-Rawi*

SRP. 2020; 11(11): 1091 - 1101

» Abstract » PDF DOI: 10.31838/srp.2020.11.156

### Review Article

#### An Observational Study Of Coronavirus (Covid-19) In Iraqi Patients At Al-Shifa Medical Center In Baghdad's Capital, Al-Rusafa

✎ *Saad Abdul Kareem Mohammed, Suha Hussein Ahmed, Ali I. Omran Al- Saadawi, Mohammed Mahmood Mohammed*

SRP. 2020; 11(11): 404 - 411

» Abstract » PDF DOI: 10.31838/srp.2020.11.60

### Review Article

#### Pathogenic Microorganisms, Toxigenic Fungi, Heavy Metal accumulation, and Toxic Materials Contamination in Natural Products

✎ *Huda S. Husni*

SRP. 2020; 11(11): 1772 - 1776

» Abstract » PDF DOI: 10.31838/srp.2020.11.247

### Review Article

» Abstract » PDF DOI: 10.31838/srp.2020.11.247

[Online First](#)

[Current Issue](#)

[Archive](#)

[Aims and Scope](#)

[Abstracting & Indexing](#)

[Most Accessed Articles](#)

[Most Downloaded Articles](#)

ORCID

Crossref

OPEN ACCESS

CC creative commons



## Tweets by @in\_systematic



Systematic Reviews in I  
@in\_systematic

According to new research from Rensselaer Polytechnic Institute (#RPI), the antiviral #remdesivir is 10 times more effective in treating cells infected with SARS-CoV-2 when combined with drugs currently used to treat hepatitis C virus (HCV).[sysrevpharm.org](http://sysrevpharm.org)

3h

Embed

View on Twitter

>> Abstract >> PDF DOI: 10.31838/srp.2020.11.61

### Review Article

#### The Therapeutic Effect Of Eucalyptus Microtheca Against The Giardiasis Induced Renal Damage In Male Rats

Abdulrazzaq Mohammed Hammood, Muhannad Shweash, and Hazim Ghazzay

SRP. 2020; 11(11): 418 - 422

>> Abstract >> PDF DOI: 10.31838/srp.2020.11.62

### Research Article

#### The Determinant of Exclusive Breastfeeding among Female Worker in Indonesia

Ina Kusriani, Mara Ipa, Agung Dwi Laksono, Noviati Fuada, Sri Supadmi

SRP. 2020; 11(11): 1102 - 1106

>> Abstract >> PDF DOI: 10.31838/srp.2020.11.157

### Review Article

#### Effect of COVID-19 on Bacterial Resistance

Hussam H. Tizkam, Osama Q. Fadhil, Esraa Ghazy

SRP. 2020; 11(11): 423 - 427

>> Abstract >> PDF DOI: 10.31838/srp.2020.11.63

### Research Article

#### Next-Generation Sequencing Technologies for Environmental DNA as an Efficient Bio Indicator for Bacterial Biodiversity in Tigris River, Iraq

Fikrat M Hassan, Warqaa Y. Salih and Halah H Al-Haideri

SRP. 2020; 11(11): 1107 - 1114

>> Abstract >> PDF DOI: 10.31838/srp.2020.11.158

### Review Article

#### Analysis of the Prophet Saleh in Al Quran through the food Science Approach to the Miracle of She-Camel

Endin Mujahidin, Nur Richana, Adian Husaini, Didin Hafidhuddin

SRP. 2020; 11(11): 771 - 775

>> Abstract >> PDF DOI: 10.31838/srp.2020.11.111

### Review Article

#### The Relationship of Family Support to the Success of the Treatment of Tuberculosis (TB) Patients in Kediri

Fauzan Adima, Chatarina U. Wahjuni, Hari Basuki Notobroto, Shrimarti Rukmini Devy

SRP. 2020; 11(11): 776 - 778

>> Abstract >> PDF DOI: 10.31838/srp.2020.11.112

SRP. 2020; 11(11): 1777 - 1782

» Abstract » PDF DOI: 10.31838/srp.2020.11.248

#### Research Article

##### Recent Development In Oxine Complexes And Their Medical Application: A Review

» Shatha M. H. Obaid, Waleed K. Mahdi, Falah H. Hussein, Yahya F. Al-Khafaji

SRP. 2020; 11(11): 428 - 437

» Abstract » PDF DOI: 10.31838/srp.2020.11.64

#### Review Article

##### Cytotoxic Activity of Ethanol Extract Legundi Leaf (Vitex trifolia L.) and N-Hexan, Ethyl Acetate and Ethanol-Water Fraction against MCF-7 Breast Cancer Cells

» Peni Indrayudha, Indah Gita Cahyani

SRP. 2020; 11(11): 779 - 785

» Abstract » PDF DOI: 10.31838/srp.2020.11.113

#### Review Article

##### Detecting Mycobacterium Tuberculosis through Bacterioferritin Gene Amplification by Real-Time PCR

» Raed Obaid Saleh, Nagham Maad Hamdi, Rana Kadhim Mohammed.

SRP. 2020; 11(11): 786 - 789

» Abstract » PDF DOI: 10.31838/srp.2020.11.114

#### Review Article

##### MIC of Some Plant Extracts against E. Coli Isolated from UTI

» Ibrahim Mohammed Saeed Hussein, Jalank Hameed Mahmoud, Halala Mohammed Abdulla

SRP. 2020; 11(11): 790 - 792

» Abstract » PDF DOI: 10.31838/srp.2020.11.115

#### Review Article

##### Acute Coronary Syndrome in Patients with Oncological Diseases

» N.A. Shanazarov, N.K. Seidalin, A.F. Khalirahmanov, G.D. Gatiyatullina, S.V. Zinchenko, K. F. Idrisova

SRP. 2020; 11(11): 1115 - 1120

» Abstract » PDF DOI: 10.31838/srp.2020.11.159

#### Review Article

##### Effects of a Cellular Camel Urinary Bladder Sub Mucosa on Chronic Full Thickness Skin Wound Healing in Equine Species

» Ahmed Kassem, Ayad N.D. Alhaki, Abd-Alhadi Jaithom Marzok.

SRP. 2020; 11(11): 793 - 797

» Abstract » PDF DOI: 10.31838/srp.2020.11.116

#### Mn+2 Precursors

The Success of Elimination Diet in Indonesian Children with Food Allergy: The Role of Caregiver's

Stress, Family Activities, and Coping

✎ Azwin Mengindra Putera, Irwanto, Margarita Maria Maramis

SRP. 2020; 11(11): 1604 - 1611

>> Abstract >> PDF DOI: 10.31838/srp.2020.11.225

Review Article

TELEVISION AND LANGUAGE OF COMMUNICATION

✎ Ridha Ahida.

SRP. 2020; 11(11): 55 - 62

>> Abstract >> PDF DOI: 10.31838/srp.2020.11.10

Research Article

Electronic Rumors of the Corona 19 Pandemic and its Impact on Psychological Reassurance in a Sample of the Society of Prince Sattam Bin Abdulaziz University.

✎ Mona AbdeLatif Al-Awad Khairallah, Asmaa Ghurmallah Al-Zahrani

SRP. 2020; 11(11): 1612 - 1625

>> Abstract >> PDF DOI: 10.31838/srp.2020.11.226

Research Article

The Effect of 8 Weeks National Academy of Sports Medicine Exercises Training on Posture, Shoulder Pain, and Functional Disability in Male with Upper Cross Syndrome

✎ Mohammed Challab Idan Almasoodi, Reza Mahdavinejad, Ghollamali Ghasmi

SRP. 2020; 11(11): 1826 - 1833

>> Abstract >> PDF DOI: 10.31838/srp.2020.11.256

Review Article

THE RESULTS OF THE RESEARCH OF THE INITIAL STATE OF READINESS OF FUTURE SOCIAL PEDAGOGUES TO WORK IN INCLUSIVE EDUCATION

✎ al-Farabi KazNU, al-Farabi KazNU, M. Auezov.

SRP. 2020; 11(11): 63 - 70

>> Abstract DOI: 10.31838/srp.2020.11.11

Review Article

Phytochemical Investigation Of Some Phenolic Compounds In Carthamus Tinctorius That Grown Naturally In Iraq

✎ Mohammed Abbas Fadhil, Enas Jawad Kadhim

SRP. 2020; 11(11): 1869 - 1874

>> Abstract >> PDF DOI: 10.31838/srp.2020.11.262

Review Article

Hydrothermal Synthesis Of  $\gamma$ -MnO<sub>2</sub> Nanostructures With Different Morphologies Using Different



# The Success of Elimination Diet in Indonesian Children with Food Allergy: The Role of Caregiver's Stress, Family Activities, and Coping

Azwin Mengindra Putera\*<sup>1</sup>, Irwanto<sup>1</sup>, Margarita Maria Maramis<sup>2</sup>

<sup>1</sup>Department of Child Health, Faculty of Medicine, Universitas Airlangga – Dr. Soetomo General Academic Hospital, Surabaya, Indonesia

<sup>2</sup>Department of Psychiatry, Faculty of Medicine, Universitas Airlangga – Dr. Soetomo General Academic Hospital, Surabaya, Indonesia

**Corresponding Author:** Azwin Mengindra Putera

**Email:** [azwinmengindraputera@gmail.com](mailto:azwinmengindraputera@gmail.com)

## ABSTRACT

**Background:** Elimination diet is the most effective treatment in treating children with food allergy. However, in its implementation, the success of elimination diet depends on a variety of factors, especially their caregiver.

**Objectives:** This study analyzed the correlation between caregiver's stress, coping strategy, and family's daily activities on the success of elimination diet in Indonesian children with food allergy.

**Methods:** Participants in this study were caregiver of Indonesian children with food allergy who received elimination diet for 4 weeks. Participants were measured for caregiver's stress, family's daily activities, and family's coping strategy after elimination diet. Caregiver's stress was measured using PSS, family's daily activities used FAIS, and while family's coping strategy used F-COPEs. The data were analyzed using multiple linear regression, with  $p < 0.05$ .

**Results:** The success rate of elimination diet in Indonesian children with food allergy was 43.64%. There was no significant correlation between caregiver's stress and elimination diet in Indonesian children with food allergy ( $t = -0.145$ ; 95% CI -0.014 – 0.012;  $p = 0.886$ ). A significant correlation was found between FAIS and elimination diet in Indonesian children with food allergy ( $t = -2.297$ ; 95% CI -0.179 – -0.012;  $p = 0.026$ ). Moreover, caregiver's coping strategy significantly correlated with elimination diet in Indonesian children with food allergy ( $t = 16.154$ ; 95% CI 0.034 – 0.043;  $p < 0.001$ ).

**Conclusion:** The success of elimination diet in Indonesian children with food allergy is influenced by family's daily activities and coping strategy, while caregiver's stress has no significant influence.

**Keywords:** Caregiver's stress, family's coping strategy, family's daily activities, food allergy, elimination diet

## Correspondence:

Azwin Mengindra Putera

Department of Child Health, Faculty of Medicine, Universitas Airlangga – Dr. Soetomo General Academic Hospital, Surabaya, Indonesia

**Email:** [azwinmengindraputera@gmail.com](mailto:azwinmengindraputera@gmail.com)

## INTRODUCTION

Food allergy is the main cause of early atopic disease manifestation that can cause various other allergic manifestations [1]. The incidence of allergy in the world, particularly food allergy continues to increase, especially in children [2]. It was reported that in 2016, 2-10% of patients experienced food allergy, while as much as 5.9% of patients in Europe experienced food allergy. Moreover, 7.1% children in Canada had food allergy [3]. There is still no detailed data on the prevalence of food allergy patients in Indonesia in all age groups [4], but data on children with food allergy in Indonesia is reported to be 3-60% [5]. Food allergy disease is commonly found in children [3,6,7]. The prevalence of food allergy continues to increase by around 4-8% in children annually [8].

Elimination diet is the most effective treatment in handling patients with adverse reactions due to food allergy [1,9]. The method detects and removes specific antagonistic foods by formulating a nutritious hypoallergenic diet to ensure optimal health of people who are sensitive to food. However, this process is often tedious, time-consuming, and requires extraordinary knowledge, skills, and commitment [8]. In the case of children with food allergy, the success of food elimination diet depends on parents and children, how serious they are to follow and understand the elimination diet recommendation as much as possible. There are still many parents and children with food allergy who do not follow elimination diet properly due to various factors, both from parents and children [1,10].

Food allergy in children cannot be predicted and life threatening, thus increases caregiver's stress [11]. It was reported that 41% of caregivers experienced increased stress when their children were diagnosed with food allergy [11,12]. This condition causes the caregiver to limit children's activities in party and snacking at school. Daily activities such as shopping and eating out are frightening for children with food allergies and are even considered life threatening [12,13]. Adaptive coping is needed to solve these problems [14,15], so that the elimination diet given to children with food allergies can be successful [6,7].

Elimination diet is the best way to help patients cutting ties from foods that cause their daily allergic symptoms [1,9,16,17]. Data related to the success of elimination diet in Indonesian children with food allergy is very limited. Dr. Soetomo General Academic Hospital, Surabaya, as the main referral hospital in East Java, Indonesia, reported the number of Indonesian children with food allergy has increased annually, with 127 patients in 2017 and 143 patients in 2018. Based on the explanation above, it is necessary to conduct an analysis to examine correlation between caregiver's stress, coping strategy, and family's daily activities on the success of elimination diet in Indonesian children with food allergy.

## MATERIALS AND METHODS

### Participants

Participants in this study were caregiver of Indonesian children with food allergy, which consisted of biological

mother or father. The participant's inclusion criteria were caregiver with children aged 1 month to 18 years, children diagnosed with food allergy [4,18,19], and children receiving elimination diet. The exclusion criteria were caregiver or children who did not want to participate in the study and children aged <1 month. Participants first received explanation of their rights, obligations, and filled out an informed consent sheet.

#### Design

This study used a case control study design [20,21], by observing the administration of elimination diet in Indonesian children with food allergy. This research was conducted in Dr. Soetomo General Academic Hospital,

Surabaya, Indonesia, from May to August 2019. The participants were chosen using consecutive sampling, with a total of 55 participants (figure 1). Participants were not assessed prior to elimination diet. Assessment of allergy symptoms of Indonesian children with food allergy was carried out based on the response of emerging diseases such as allergic rhinitis [22], asthma [23], and atopic dermatitis [24]. The elimination diet was given for 4 weeks based on the Guideline for Diagnosis and Management of Food Allergy [18,25,26]. The participants were then evaluated for allergy symptoms and measured for characteristics, caregiver's stress, family's daily activities, and family's coping strategy.




Figure 1. Identification Flowchart of Number of Participants

#### Ethical Approval

This study has obtained a certificate of ethical clearance according to Declaration of Helsinki from ethical committee team in Dr. Soetomo General Academic Hospital and Faculty of Medicine Universitas Airlangga ethics Committee (1165/KEPK/V/2019).

#### Elimination Diet

Elimination diet was provided based on a history of response to food allergies and skin prick tests [16]. The consumption of allergy-causing food was stopped and replaced the nutritional intake of these foods in consultation with a nutritionist. Nutrients obtained from these foods should be replaced with foods containing similar nutrients [18,25,26]. The use of books in

elimination diet really helped caregivers and nutritionists make a diet plan for 4 weeks. Diet elimination was declared unsuccessful if the elimination diet was carried out >4 weeks and persistent signs of symptoms of food allergy.

#### Measurement of Caregiver's Stress

Caregiver's stress was measured using parental stress scale (PSS) with 18 questions. The scale is developed to measure stress level as parents [27]. The PSS used Likert scale of 1-5 (1 = strongly disagree, 2 = disagree, 3 = neutral, 4 = agree, 5 = strongly agree), with a minimum value of 18 and a maximum value of 90 [28]. This assessment tool was categorized into 3, namely mild (18 - 42), moderate (43 - 66), and severe (67-90) [29]. The

PSS was considered valid and reliable with Cronbach's alpha = 0.83 [27,28].

**Measurement of Family's Daily Activities**

Family's daily activities were measured using the food allergy impact scale (FAIS) to evaluate parents' perceptions about the impact of food allergy on eight aspects of family's daily activities, with the higher score indicating a greater impact. The scale consists of 32 questionnaire items, consisting of 8 activity aspects namely meal preparation (six items), family social activities (five items), caregiver-supervised child social activities (two items), family relations (three items), caregiver's stress and free time (three items), school or structured activities (six items), autonomous child social activities (three items), and employment and finances (three items). The FAIS reliability value was  $\geq 0.70$  [15,30].

**Measurement of Family's Coping Strategy**

The measurement of family coping was conducted using family crisis oriented personal evaluation scales (F-COPES) questionnaire, with a 5-point Likert's scale. This instrument displays 30 items of coping behavior designed to record problem solving, attitudes and behaviors developed by families to respond to problems or difficulties. This scale is divided into 5 subscales: obtaining social support, reframing, seeking spiritual

support, mobilizing family support, and passive assessment [31]. The Indonesian F-COPES was considered valid and reliable with Cronbach's alpha = 0.89 [32,33].

**Statistical Analysis**

The measured data were analyzed univariate and bivariate, where univariate data were displayed in the form of a frequency distribution or mean  $\pm$  standard deviation (SD). Meanwhile, the bivariate analysis used multiple linear regression tests. Statistical tests were significant if  $p < 0.05$ . Measurement data were analyzed using IBM SPSS Statistic software version 21.0 (IBM Corp., Armonk, NY, USA).

**RESULTS**

**Characteristics of Participants**

Most participants were male 40 (72.73;  $p = 0.349$ ), with the mean age of  $68.1 \pm 37.46$  years ( $p = 0.159$ ). Most participants came from low-income family (65.46%;  $p = 0.009$ ). Parents' income affects the process of treating children with food allergy. Most parents' education level was non-bachelor as much as 72.73% ( $p = 0.782$ ; Table 1). Most elimination diet was unsuccessful (56.36%). The mean age of participants who succeeded in implementing elimination diet was  $60.0 \pm 39.23$  years, and those who were unsuccessful were  $74.4 \pm 35.38$  ( $p = 0.692$ ).

Table 1. Characteristics of Participants

Characteristics of Participants	N (%)	<i>p</i>
Sex		
Male	40 (72.7)	0.349
Female	15 (27.3)	
Income		0.009*
Low	36 (65.5)	
High	19 (34.5)	
Educational Level		0.782
Non-bachelor	40 (72.7)	
Bachelor	15 (27.3)	
Elimination diet in Children		-
Unsuccessful	31 (56.4)	
Successful	24 (43.6)	

Abbreviations: \* significant  $p < 0.05$ ; \*\*significant  $p < 0.001$ .

**Correlation between Caregiver's Stress on Elimination Diet in Indonesian Children with Food Allergy**

The participant's stress value in this study ranged from 28 to 57, where the most participant had a stress value of 43 for 12 participants. The mean value of the participant's stress was  $45.29 \pm 4.67$ , with a median value of 45 (43 -

48). Most participants had stress in the moderate category as much as 89.09%. Participants who had mild PSS scores and successful elimination diet were as much as 83.3%, while participants who had moderate PSS scores and unsuccessful elimination diet were 61.2%. There was no significant correlation between

Table 2. Correlation between Caregiver's Stress, Family Activities, and Family Coping on the Success of Elimination Diet in Indonesian Children with Food Allergy

Caregiver's Condition	Successful Elimination Diet			Unsuccessful Elimination Diet			<i>p</i>
	Mean $\pm$ SD	Median (Q1-Q3)	Min-Max	Mean $\pm$ SD	Median (Q1-Q3)	Min-Max	
PSS	44.17 $\pm$ 5.65	45.00 (43.00 - 46.75)	28.00 - 54.00	46.16 $\pm$ 3.61	45.00 (43.00 - 49.00)	41.00 - 57.00	0.886
FAIS	5.01 $\pm$ 0.89	5.13 (4.48 - 5.47)	2.98 - 6.75	5.52 $\pm$ 0.53	5.33 (5.15 - 5.78)	4.31 - 6.52	0.026*
F-COPES	116.29 $\pm$ 3.91	116.50 (113.25 - 120.00)	109.00 - 122.00	95.45 $\pm$ 5.20	97.00 (94.00 - 99.00)	77.00 - 100.00	0.000**

Abbreviation: PSS, parental stress scale; FAIS, food allergy impact scale; F-COPES, family crisis oriented personal evaluation scales; \*significant  $p < 0.05$ ; \*\*significant  $p < 0.001$ .



Table 3. Comparison of Each FAIS Subscale to the Caregiver of Indonesian Children with Food Allergy

FAIS	2		3		4		5		6		7		8	
	95% CI	p	95% CI	p	95% CI	p	95% CI	p	95% CI	P	95% CI	p	95% CI	p
1	0.12-0.43	0.001*	0.48-1.31	0.000**	0.72-1.48	0.000**	0.96-1.68	0.000**	0.96-1.77	0.000**	0.26-0.97	0.001*	0.46-1.23	0.000**
2			0.17-0.94	0.005*	0.43-1.22	0.000**	0.70-1.38	0.000**	0.65-1.44	0.000**	-0.09-0.70	0.125	0.15-1.23	0.008*
3					-0.56-0.48	0.875	-0.12-0.90	0.135	0.97-1.00	0.019*	-0.69-0.24	0.329	-0.54-0.76	0.743
4							-2.71-0.70	0.380	0.04-1.01	0.067	-0.99-0.65	0.084	-0.83-0.32	0.372
5									-0.39-0.57	0.714	-1.34-0.24	0.005*	-0.90-0.04	0.031*
6											-1.34-0.23	0.007*	-1.11-0.01	0.053
7													-0.21-0.80	0.249

Abbreviation: FAIS, food allergy impact scale; 1, meal preparation; 2, family social activities; 3, caregiver-supervised child social activities; 4, family relations; 5, caregiver stress and free time; 6, school or structured

activities; 7, autonomous child social activities; 8, employment and finances; \*significant  $p < 0.05$ ; \*\*significant  $p < 0.001$ .

Table 4. Comparison of Each F-COPES Subscale to the Caregiver of Indonesian Children with Food Allergy

F-COPES	Reframing		Spiritual Support		Mobilizing Family Support		Passive Appraisal	
	95% CI	p	95% CI	p	95% CI	p	95% CI	p
Social Support	-6.14 -- 0.52	0.021*	11.81 -- 16.95	0.000**	14.62 -- 20.03	0.000**	16.76 -- 22.01	0.000**
Reframing			16.57 -- 18.85	0.000**	19.36 -- 21.95	0.000**	21.24 -- 24.18	0.000**
Spiritual Support					2.04 -- 3.85	0.000**	3.87 -- 6.13	0.000**
Mobilizing Family Support							0.96 -- 3.15	0.000**

Abbreviation: F-COPES, family crisis oriented personal evaluation scales; \*significant  $p < 0.05$ ; \*\*significant  $p < 0.001$ .

Caregiver's stress and elimination diet in Indonesian children with food allergy, where the PSS value of the participant with the successful elimination diet was  $44.17 \pm 5.65$ , while the PSS value of unsuccessful diet was  $46.16 \pm 3.61$  ( $t = -0.145$ ; 95 % CI -0.014 - 0.012;  $p = 0.886$ ; Table 2).

**Correlation between Family's Daily Activities on Elimination Diet in Indonesian Children with Food Allergy**

The FAIS value ranged from 2.98 - 6.75, where the mean value was  $5.30 \pm 0.75$  with a median value of 5.26 (5.08 - 5.66). The mean value of each FAIS subscale in the successful elimination diet group was as follows: meal preparation ( $5.86 \pm 0.85$ ), family social activities ( $5.58 \pm 1.05$ ), caregiver-supervised child social activities ( $3.64 \pm 2.77$ ), family relations ( $4.82 \pm 1.40$ ), caregiver's stress and free time ( $4.82 \pm 1.15$ ), school or structured activities ( $3.33 \pm 2.35$ ), autonomous child social activities ( $4.14 \pm 2.52$ ), and employment and finances ( $4.91 \pm 1.49$ ). Meanwhile, the FAIS value in the unsuccessful diet elimination group was as follows: meal preparation ( $6.37 \pm 0.76$ ), family social activities ( $6.10 \pm 0.82$ ), caregiver-supervised child social activities ( $4.92 \pm 2.12$ ), family relations ( $5.22 \pm 1.61$ ), caregiver's stress and free time

( $4.84 \pm 1.33$ ), school or structured activities ( $3.79 \pm 2.30$ ), autonomous child social activities ( $5.83 \pm 1.30$ ), and employment and finances ( $5.61 \pm 1.10$ ). There was a significant correlation between FAIS and elimination diet in Indonesian children with food allergy, where the caregiver's average FAIS value in the successful elimination diet group was  $5.01 \pm 0.89$ , while the value in the unsuccessful group was  $5.52 \pm 0.53$  ( $t = -2.297$ ; 95% CI -0.179 - -0.012;  $p = 0.026$ ; Table 2).

There was a significant correlation of each FAIS subscale value in caregiver of Indonesian children with food allergy. The correlation was as follows: meal preparation versus (vs) family social activities ( $0.28 \pm 0.57$ ;  $p = 0.001$ ); meal preparation vs caregiver-supervised child social activities ( $0.90 \pm 1.40$ ;  $p < 0.001$ ); meal preparation vs family relations ( $1.10 \pm 0.90$ ;  $p < 0.001$ ); meal preparation vs caregiver's stress and free time ( $1.32 \pm 1.33$ ;  $p < 0.001$ ); meal preparation vs school or structured activities ( $1.37 \pm 1.30$ ;  $p < 0.001$ ); meal preparation vs autonomous child social activities ( $0.62 \pm 1.25$ ;  $p = 0.001$ ); meal preparation vs employment and finances ( $0.84 \pm 1.43$ ;  $p < 0.001$ ); family social activities vs caregiver-supervised child social activities ( $0.56 \pm 1.26$ ;  $p = 0.005$ ); family social activities vs family relations ( $0.82 \pm 1.45$ ;  $p < 0.001$ ); family social activities vs caregiver's

stress and free time ( $1.04 \pm 1.28$ ;  $p < 0.001$ ); caregiver-supervised child social activities vs school or structured activities ( $0.55 \pm 1.38$ ;  $p = 0.019$ ); caregiver's stress and free time vs autonomous child social activities ( $-0.79 \pm 1.93$ ;  $p = 0.005$ ); caregiver's stress and free time vs employment and finances ( $-0.47 \pm 1.58$ ;  $p = 0.031$ ); and school or structured activities vs autonomous child social activities ( $-0.80 \pm 1.93$ ;  $p = 0.007$ ). The highest correlation value was obtained by meal preparation vs school or structured activities of  $1.37 \pm 1.30$  (95% CI 0.96 - 1.77;  $p < 0.001$ ). Meanwhile, the lowest comparison value was obtained in subscale school or structured activities vs autonomous child social activities of  $-0.80 \pm 1.93$  (95% CI -1.34 - -0.23;  $p = 0.007$ ; Table 3).

#### Family's Coping Strategy on the Success of Elimination Diet in Indonesian Children with Food Allergy

The participant's F-COPES value ranged from 77 to 122, where the mean F-COPES value was  $104.54 \pm 11.42$  with a median value of 100 (96 - 115). The mean value of each F-COPES subscale was as follows: social support of  $39.08 \pm 6.77$  (successful elimination diet) and  $23.29 \pm 5.22$  (unsuccessful diet elimination); reframing of  $34.08 \pm 3.34$  (successful elimination diet) and  $33.06 \pm 3.46$  (unsuccessful elimination diet); spiritual support of  $16.46 \pm 1.56$  (successful elimination diet) and  $15.29 \pm 2.22$  (unsuccessful elimination diet); mobilizing family support of  $13.38 \pm 2.75$  (successful elimination diet) and  $12.45 \pm 2.69$  (unsuccessful elimination diet); and passive assessment of  $11.54 \pm 3.58$  (successful elimination diet) and  $10.23 \pm 2.79$  (unsuccessful elimination diet). There was a significant correlation between caregiver's coping strategy and elimination diet in Indonesian children with food allergy, where the mean value of family coping with successful diet elimination was  $116.29 \pm 3.91$ , and  $95.45 \pm 5.20$  for unsuccessful group ( $t = 16.154$ ; 95% CI 0.034 - 0.043;  $p < 0.001$ ; Table 2).

There was a significant comparison of each F-COPES subscale value of caregiver of Indonesian children with food allergy. The highest comparison value was obtained in subscale reframing vs passive appraisal of  $22.71 \pm 5.44$  (95% CI 21.24 - 24.18;  $p < 0.001$ ). Meanwhile, the lowest comparison value was found on subscale social support vs spiritual support of  $-3.33 \pm 10.40$  (95% CI -6.14 - -0.52;  $p = 0.021$ ; Table 4).

#### DISCUSSION

The main treatment of allergy is eliminating the allergen causing the problem, so the main management of food allergy is elimination diet [34]. Some literatures stated that elimination diet is an effective treatment for patients with food allergies [6,7,16]. The success of elimination diet in children with food allergy can increase the nutritional intake of children in the stages of development and growth and eliminate the signs of food allergy symptoms [16,17]. The success of elimination diet is expected to prevent the occurrence of food allergy reactions, most of which are anaphylactic reactions [35]. The success of elimination diet in children is strongly influenced by the role of caregiver, as they are the closest people to the child. Any condition experienced by the caregiver greatly affects the child's condition, including the success of elimination diet in children with food allergy [11,36].

In this study, there was no significant correlation between caregiver's stress and the success of elimination diet in Indonesian children with food allergy. Previous

research pointed out that parents or caregivers of children with food allergy experienced increased stress [11] that eventually had negative effect on the knowledge, attitudes and behavior of the caregiver in assisting children in carrying out the elimination diet [37]. Caregivers with high stress level have difficulty to adapt to changes in lifestyle of elimination diet for children with food allergy, because the allergy-causing food must be stopped and replaced with those that have similar nutritional content but are hypoallergenic. These conditions result in increased costs and the need for dietary consultation with a nutritionist for the selection of foods that are good for consumption during elimination diet [11,38]. In addition, excessive stress leads to incorrect treatment chosen by caregivers [39]. Thus, caregiver's stress directly and indirectly affects the success of elimination diet in children with food allergy [11,37,38].

This study found that family's activities had a significant relationship with the success of elimination diet in Indonesian children with food allergy. Previous studies mentioned that food allergies greatly affect daily activities, in which the caregiver must pay attention to the labels of food items that children will consume [12]. In addition, caregivers of children with food allergy are forced to limit children's activities, especially in public activities (attending parties, eating together outside) [12,13,40]. Caregivers should pay extra attention to children at school because their risk of eating food allergens is very high [12,30]. Activities such as sleepovers and camps are also limited by the caregiver to minimize the risk of exposure to food allergens [15]. Daycare, birthday parties, field trips, and going to restaurants are also reported to be limited by the caregiver to prevent children with food allergies from consuming food allergens [12,41]. Activity limitation by the caregiver can increase the success of elimination diet by 70% [12,13].

Moreover, this study found a significant correlation between family's coping strategy and the success of elimination diet in Indonesian children with food allergy. Family's coping is important in overcoming the problem of food allergy in children, because family coping can determine the coping strategies used to seek treatment for children with food allergy [42]. The use of family coping in elimination diet in children with food allergy is very complex, which requires an effective strategy in its implementation. On the other hand, prolonged elimination diet causes psychological problems. Elimination diet in adults is effective to reduce food allergy load, but it requires further research in children patients regarding adaptive coping strategies [43]. Family coping affects the success of elimination diet in children with food allergies because family support is needed to play a crucial role in eliminating children's diets [6,7,11], because adapting children to eat hypoallergen foods to avoid allergic foods requires a long process [44,45].

However, some limitations should be noted. First, is it is hoped that further research will investigate elimination diet in Indonesian children with food allergy by using a quasi / true experimental research design. Second, it is hoped the number of participants in future studies will be greater and use a randomized control trial technique so that the composition between the intervention group and control group is more balanced and minimizes bias. Third, it is hoped that further research classifies the ages

of children with food allergy in order to identify the most effective elimination diet given to which age group.

## CONCLUSION

Elimination diet is an effective treatment for children with food allergy, and the role of caregiver greatly affects its success. Family activities greatly affect the success of elimination diet in children with food allergy because limiting social activity greatly supports its success. Likewise, family coping greatly affects elimination diet in Indonesian children with food allergy because the coping used by families in responding to food allergies is critical to success because children cannot make decisions. While caregiver's stress greatly influences the success of elimination diet in Indonesian children with food allergy, this study found no significant correlation between caregiver's stress and elimination diet in Indonesian children with food allergy.

## ACKNOWLEDGEMENT

The authors would like to thank Dr. Windhu Purnomo, MS, Faculty of Public Health, Universitas Airlangga, Surabaya, Indonesia for his technical assistance for this study, particularly in the statistical analysis. We would also like to thank Fis Citra Ariyanto as our manuscript editor.

## CONFLICT OF INTEREST

None.

## Author's Contribution

All authors contributed toward data analysis, drafting and revising the paper, gave final approval of the version to be published and agree to be accountable for all aspects of the work.

## FUNDING

None

## REFERENCES

1. Walczak M, Grzelak T, Kramkowska M, Czyżewska K (2014) Food allergies in children—aspects of epidemiology and diet management. *Journal of Medical Science* 83 (3):255-259
2. Loh W, Tang MLK (2018) The Epidemiology of Food Allergy in the Global Context. *Int J Environ Res Public Health* 15 (9):2043. doi:10.3390/ijerph15092043
3. Savage J, Johns CB (2015) Food allergy: epidemiology and natural history. *Immunol Allergy Clin North Am* 35 (1):45-59. doi:10.1016/j.iac.2014.09.004
4. Munasir Z, Muktiarti D (2013) The management of food allergy in Indonesia. *Asia Pac Allergy* 3 (1):23-28. doi:10.5415/apallergy.2013.3.1.23
5. Sumadiono, Muktiarti D, Setiabudiawan B, Irsa L, Wati KDK, Majangsari RGD (2015) Rekomendasi Ikatan Dokter Anak Indonesia Pencegahan Primer Alergi. 1 edn. Unit Kerja Koordinasi Alergi Immunologi: Ikatan Dokter Anak Indonesia,
6. Berry MJ, Adams J, Voutilainen H, Feustel PJ, Celestin J, Järvinen KM (2015) Impact of elimination diets on growth and nutritional status in children with multiple food allergies. *Pediatric allergy and immunology : official publication of the European Society of Pediatric Allergy and Immunology* 26 (2):133-138. doi:10.1111/pai.12348
7. Venter C, Mazzocchi A, Maslin K, Agostoni C (2017) Impact of elimination diets on nutrition and growth in children with multiple food allergies. *Current opinion in allergy and clinical immunology* 17 (3):220-226. doi:10.1097/aci.0000000000000358
8. Bora S, Rindfleisch JA (2018) Chapter 86 - The Elimination Diet. In: Raketel D (ed) *Integrative Medicine (Fourth Edition)*. Elsevier, pp 849-862.e846. doi:https://doi.org/10.1016/B978-0-323-35868-2.00086-4
9. Le T-M, Zijlstra WT, van Opstal EY, Knol MJ, L'Hoir MP, Knulst AC, Pasmans SGMA (2013) Food avoidance in children with adverse food reactions: Influence of anxiety and clinical parameters. *Pediatric Allergy and Immunology* 24 (7):650-655. doi:10.1111/pai.12114
10. Allen CW, Kemp AS, Campbell DE (2009) Dietary advice, dietary adherence and the acquisition of tolerance in egg-allergic children: a 5-yr follow-up. *Pediatric Allergy and Immunology* 20 (3):213-218. doi:10.1111/j.1399-3038.2008.00784.x
11. Birdi G, Cooke R, Knibb R (2016) Quality of Life, Stress, and Mental Health in Parents of Children with Parentally Diagnosed Food Allergy Compared to Medically Diagnosed and Healthy Controls. *J Allergy (Cairo)* 2016:1497375-1497375. doi:10.1155/2016/1497375
12. Bollinger ME, Dahlquist LM, Mudd K, Sonntag C, Dillinger L, McKenna K (2006) The impact of food allergy on the daily activities of children and their families. *Annals of Allergy, Asthma & Immunology* 96 (3):415-421. doi:10.1016/S1081-1206(10)60908-8
13. Cummings AJ, Knibb RC, King RM, Lucas JS (2010) The psychosocial impact of food allergy and food hypersensitivity in children, adolescents and their families: a review. *Allergy* 65 (8):933-945. doi:10.1111/j.1398-9995.2010.02342.x
14. Muñoz-Furlong A (2003) Daily coping strategies for patients and their families. *Pediatrics* 111 (6 Pt 3):1654-1661
15. Klinnert MD, McQuaid EL, Fedele DA, Faino A, Strand M, Robinson J, Atkins D, Fleischer DM, Hourihane JOB, Cohen S, Fransen H (2015) Children's Food Allergies: Development of the Food Allergy Management and Adaptation Scale. *J Pediatr Psychol* 40 (6):572-580. doi:10.1093/jpepsy/jsv009
16. Kim J, Kwon J, Noh G, Lee SS (2013) The effects of elimination diet on nutritional status in subjects with atopic dermatitis. *Nutr Res Pract* 7 (6):488-494. doi:10.4162/nrp.2013.7.6.488
17. Lozinsky AC, Meyer R, De Koker C, Dziubak R, Godwin H, Reeve K, Dominguez Ortega G, Shah N (2015) Time to symptom improvement using elimination diets in non-IgE-mediated gastrointestinal food allergies. *Pediatric allergy and immunology : official publication of the European Society of Pediatric Allergy and Immunology* 26 (5):403-408. doi:10.1111/pai.12404
18. Boyce JA, Assa'ad A, Burks AW, Jones SM, Sampson HA, Wood RA, Plaut M, Cooper SF, Fenton MJ, Arshad SH, Bahna SL, Beck LA, Byrd-Bredbenner C, Camargo CA, Jr., Eichenfield L, Furuta GT, Hanifin JM, Jones C, Kraft M, Levy BD, Lieberman P, Lucciolli S, McCall KM, Schneider LC, Simon RA, Simons FER, Teach SJ, Yawn BP, Schwaninger JM (2011) Guidelines for the diagnosis and management of food allergy in the



- United States: summary of the NIAID-sponsored expert panel report. *Nutr Res* 31 (1):61-75. doi:10.1016/j.nutres.2011.01.001
19. Abrams EM, Sicherer SH (2016) Diagnosis and management of food allergy. *CMAJ* 188 (15):1087-1093. doi:10.1503/cmaj.160124
20. Setia MS (2016) Methodology Series Module 2: Case-control Studies. *Indian J Dermatol* 61 (2):146-151. doi:10.4103/0019-5154.177773
21. van Stralen KJ, Dekker FW, Zoccali C, Jager KJ (2010) Case-control studies--an efficient observational study design. *Nephron Clinical practice* 114 (1):c1-4. doi:10.1159/000242442
22. Noronha L, Fox A, Du Toit G, Lack G (2017) Diagnosis and management of allergic rhinitis in children. *Prescriber* 28 (12):13-19. doi:10.1002/psb.1631
23. Papadopoulos NG, Arakawa H, Carlsen KH, Custovic A, Gern J, Lemanske R, Le Souef P, Mäkelä M, Roberts G, Wong G, Zar H, Akdis CA, Bacharier LB, Baraldi E, van Bever HP, de Blic J, Boner A, Burks W, Casale TB, Castro-Rodriguez JA, Chen YZ, El-Gamal YM, Everard ML, Frischer T, Geller M, Gereda J, Goh DY, Guilbert TW, Hedlin G, Heymann PW, Hong SJ, Hossny EM, Huang JL, Jackson DJ, de Jongste JC, Kalayci O, Ait-Khaled N, Kling S, Kuna P, Lau S, Ledford DK, Lee SI, Liu AH, Lockey RF, Lødrup-Carlsen K, Løtvall J, Morikawa A, Nieto A, Paramesh H, Pawankar R, Pohunek P, Pongracic J, Price D, Robertson C, Rosario N, Rossenwasser LJ, Sly PD, Stein R, Stick S, Szefer S, Taussig LM, Valovirta E, Vichyanond P, Wallace D, Weinberg E, Wennergren G, Wildhaber J, Zeiger RS (2012) International consensus on (ICON) pediatric asthma. *Allergy* 67 (8):976-997. doi:10.1111/j.1398-9995.2012.02865.x
24. Severity Scoring of Atopic Dermatitis: The SCORAD Index (1993). *Dermatology* 186 (1):23-31. doi:10.1159/000247298
25. Rodriguez J, Crespo JF (2012) Chapter 6 - Food-induced Urticaria and Angioedema. In: James JM, Burks W, Eigenmann P (eds) *Food Allergy*. W.B. Saunders, Edinburgh, pp 75-82. doi:https://doi.org/10.1016/B978-1-4377-1992-5.00006-5
26. Worm M, Reese I, Ballmer-Weber B, Beyer K, Bischoff SC, Classen M, Fischer PJ, Fuchs T, Huttegger I, Jappe U, Klimek L, Koletzko B, Lange L, Lepp U, Mahler V, Niggemann B, Rabe U, Raithel M, Saloga J, Schäfer C, Schnadt S, Schreiber J, Szépfalusi Z, Treudler R, Wagenmann M, Watzl B, Werfel T, Zuberbier T, Kleine-Tebbe J (2015) Guidelines on the management of IgE-mediated food allergies: S2k-Guidelines of the German Society for Allergology and Clinical Immunology (DGAKI) in collaboration with the German Medical Association of Allergologists (AeDA), the German Professional Association of Pediatricians (BVKJ), the German Allergy and Asthma Association (DAAB), German Dermatological Society (DDG), the German Society for Nutrition (DGE), the German Society for Gastroenterology, Digestive and Metabolic Diseases (DGVS), the German Society for Oto-Rhino-Laryngology, Head and Neck Surgery, the German Society for Pediatric and Adolescent Medicine (DGKJ), the German Society for Pediatric Allergology and Environmental Medicine (GPA), the German Society for Pneumology (DGP), the German Society for Pediatric Gastroenterology and Nutrition (GPGE), German Contact Allergy Group (DKG), the Austrian Society for Allergology and Immunology (Æ-GAI), German Professional Association of Nutritional Sciences (VDOE) and the Association of the Scientific Medical Societies Germany (AWMF). *Allergo J Int* 24:256-293. doi:10.1007/s40629-015-0074-0
27. Pontoppidan M, Nielsen T, Kristensen IH (2018) Psychometric properties of the Danish Parental Stress Scale: Rasch analysis in a sample of mothers with infants. *PLoS One* 13 (11):e0205662-e0205662. doi:10.1371/journal.pone.0205662
28. Berry JO, Jones WH (1995) The Parental Stress Scale: Initial Psychometric Evidence. *Journal of Social and Personal Relationships* 12 (3):463-472. doi:10.1177/0265407595123009
29. Aggarwal A, Unni K, Prasad S (2018) A Comparative Study of Stress, Anxiety & Work Impairment in Parents of Children with Conduct Disorder. *Journal of Indian Association for Child & Adolescent Mental Health* 14 (4)
30. Hoehn JL, Dahlquist LM, Hahn AL, Bollinger ME (2017) Parents of Children With Food Allergy: Gender Differences in Perceived Impact and Perceived Food Allergy Severity. *J Pediatr Psychol* 42 (2):186-197. doi:10.1093/jpepsy/jsw059
31. Eaton PM, Davis BL, Hammond PV, Condon EH, McGee ZT (2011) Coping Strategies of Family Members of Hospitalized Psychiatric Patients. *Nursing Research and Practice* 2011. doi:10.1155/2011/392705
32. Putera AM, Irwanto I, Maramis MM, Prasetyo RV, Soemyarso NA, Noer MS (2020) Effect of Mental Health Problems on the Quality of Life in Children with Lupus Nephritis. *Neuropsychiatr Dis Treat* 16:1583-1593. doi:10.2147/NDT.S250373
33. Putera AM, Irwanto, Maramis MM (2020) Quality-of-Life (QoL) of Indonesian Children Living with HIV: The Role of Caregiver Stigma, Burden of Care, and Coping. *HIV/AIDS-Research and Palliative Care* 12:573-581. doi:10.2147/HIV.S269629
34. Niggemann B, Heine RG (2006) Who should manage infants and young children with food induced symptoms? *Arch Dis Child* 91 (5):379-382. doi:10.1136/adc.2005.086603
35. Staden U, Rolinck-Werninghaus C, Brewe F, Wahn U, Niggemann B, Beyer K (2007) Specific oral tolerance induction in food allergy in children: efficacy and clinical patterns of reaction. *Allergy* 62 (11):1261-1269. doi:10.1111/j.1398-9995.2007.01501.x
36. Scherer E, Hagaman A, Chung E, Rahman A, O'Donnell K, Maselko J (2019) The relationship between responsive caregiving and child outcomes: evidence from direct observations of mother-child dyads in Pakistan. *BMC Public Health* 19 (1):252-252. doi:10.1186/s12889-019-6571-1
37. Wright RJ, Berin MC (2014) Unlocking the stress-allergy puzzle: need for a more comprehensive stress model. *Annals of allergy, asthma & immunology : official publication of the American College of Allergy, Asthma, & Immunology* 113 (1):1-2. doi:10.1016/j.anai.2014.05.007
38. Lau G-Y, Patel N, Umasunthar T, Gore C, Warner JO, Hanna H, Phillips K, Mohd Zaki A, Hodes M, Boyle RJ (2014) Anxiety and stress in mothers of food-allergic children. *Pediatric Allergy and Immunology* 25 (3):236-242. doi:10.1111/pai.12203
39. Begen FM, Barnett J, Barber M, Payne R, Gowland MH, Lucas JS (2017) Parents' and caregivers' experiences

- and behaviours when eating out with children with a food hypersensitivity. *BMC Public Health* 18 (1):38-38. doi:10.1186/s12889-017-4594-z
40. Meyer R, Godwin H, Dziubak R, Panepinto JA, Foong R-XM, Bryon M, Lozinsky AC, Reeve K, Shah N (2017) The impact on quality of life on families of children on an elimination diet for Non-immunoglobulin E mediated gastrointestinal food allergies. *World Allergy Organ J* 10 (1):8-8. doi:10.1186/s40413-016-0139-7
41. Vargas PA, Sicherer SH, Christie L, Keaveny M, Noone S, Watkins D, Carlisle SK, Jones SM (2011) Developing a food allergy curriculum for parents. *Pediatric allergy and immunology : official publication of the European Society of Pediatric Allergy and Immunology* 22 (6):575-582. doi:10.1111/j.1399-3038.2011.01152.x
42. Aika S, Ito M, Yamamoto Y (2017) Food allergy response capabilities of mothers and related factors. *Nursing & health sciences* 19 (3):340-350. doi:10.1111/nhs.12351
43. Weiss D, Marsac ML (2016) Coping and posttraumatic stress symptoms in children with food allergies. *Annals of allergy, asthma & immunology : official publication of the American College of Allergy, Asthma, & Immunology* 117 (5):561-562. doi:10.1016/j.anai.2016.08.022
44. Polloni L, Toniolo A, Lazzarotto F, Baldi I, Foltran F, Gregori D, Muraro A (2013) Nutritional behavior and attitudes in food allergic children and their mothers. *Clin Transl Allergy* 3 (1):41-41. doi:10.1186/2045-7022-3-41
45. Shanahan L, Zucker N, Copeland WE, Costello EJ, Angold A (2014) Are children and adolescents with food allergies at increased risk for psychopathology? *J Psychosom Res* 77 (6):468-473. doi:10.1016/j.jpsychores.2014.10.005