

5

by Azhar Muhammad Helmi

Submission date: 09-Jun-2021 06:03PM (UTC+0800)

Submission ID: 1603376834

File name: tilapia-fish-oreochromis-niloticus--public-health-importance.pdf (1.14M)

Word count: 7469

Character count: 42698

A Review of *Salmonella sp.* in Tilapia fish (*Oreochromis niloticus*) : Public Health Importance

Azhar Muhammad Helmi¹, Akhmad Taufiq Mukti², Agoes Soegianto³, Ketut Mahardika⁴, Indah Mastuti⁴, Mustofa Helmi Effendi^{5*}, Hani Plumeriastuti⁶

¹Postgraduate Student on Faculty of Fisheries and Marine, Universitas Airlangga, Indonesia

²Department of Fish Health Management & Aquaculture, Faculty of Fisheries and Marine, Universitas Airlangga, Indonesia

³Department of Biology, Faculty of Sciences and Technology, Universitas Airlangga, Indonesia

⁴Center for Marine Cultivation Research and Fisheries Extension, Buleleng, Bali, Indonesia

⁵Department of Veterinary Public Health, Faculty of Veterinary Medicine, Universitas Airlangga, Indonesia

⁶Department of Veterinary Pathology, Faculty of Veterinary Medicine, Universitas Airlangga, Jl. Mulyorejo, FKH UNAIR, Kampus C Universitas Airlangga, Surabaya 60115, Indonesia

*Corresponding author: Mustofa Helmi Effendi

Email: mheffendi@yahoo.com

ABSTRACT

Tilapia fish or *Oreochromis niloticus* is a food security commodity from the fisheries sector and is traded internationally, but in Asian countries tilapia aquaculture is mostly maintained using traditional systems, besides that homemade feed is used to reduce production costs. Cultivation of tilapia with traditional systems and artificial feed that is integrated with other livestock has the potential to cause the danger of transmitting zoonotic pathogens from other livestock manure and feed residue. *Salmonella sp.* which is one of the zoonotic pathogens that can be transmitted from tilapia. The use of antibiotics in livestock and cultivation causes *Salmonella sp.* able to withstand some antibiotics. Salmonellosis in humans has become an important public health problem, incurring significant economic and medical costs worldwide. The incidence of salmonellosis due to fish consumption has become a concern of public health agencies in several countries, because increased consumption of fishery products, especially raw products, increases the risk. Pathogen exposure, especially in vulnerable groups, such as the elderly, has increased significantly. pregnant women and babies. The importance of this pathogen in fish can be assessed and evaluated, as records show that most *Salmonella* infections in humans are related to fish consumption.

Keywords: Tilapia fish, *Salmonella sp.*, Zoonotic pathogen, Public health

Correspondence:

Mustofa Helmi Effendi

Department of Veterinary Public Health, Faculty of Veterinary Medicine, Universitas Airlangga, Indonesia

Email: mheffendi@yahoo.com

INTRODUCTION

Integrated tilapia farming is a traditional practice by small-scale farmers in China as well as in other Asian countries. Fish are usually raised in ponds with livestock units, such as pigs, located on pond embankments, which allow drainage of livestock manure and excess feed into the pond as plankton and other fish feed [1, 2]. Although integrated agricultural systems are sustainable in many ways, they represent potential food safety hazards, for example transmission of zoonotic pathogens from faeces and accumulation of antimicrobials and others. residues originating from pig manure and feed [3, 4]. Tilapia is one of the easiest to cultivate and trade fishery food products internationally in the world, with an estimated 1.45 million tonnes produced in China in 2013 [5, 6].

In the Asia-Pacific region, farmed fish are fed commercial or homemade feed [7]. Homemade feed is used to reduce costs and usually consists of chicken offal and by-products produced during poultry processing, kitchen waste, and other by-products from the food industry [7, 8]. Homemade diets can be a potential source of foodborne pathogens, particularly *Salmonella* bacteria [9, 10], which can then be transmitted to farmed fish [11] and to humans. Salmonellosis in humans has become an important public health problem, creating significant economic and medical costs worldwide. In the United States, nontyphoidal *Salmonella* caused approximately 1 million foodborne illnesses, 13,000 hospitalizations, and 242 deaths in 2011 [12]. In Taiwan, *Salmonella* is one of the most frequently isolated foodborne pathogenic bacteria [13]. *Salmonella* is usually transmitted to humans through consumption of

contaminated food products. According to Bean et al. [14], approximately 7% of cases of human salmonellosis result from ingestion of contaminated food from fishery products, particularly ready-to-eat (RTE) products. In Hong Kong, detected *Salmonella* in smoked fish, salmon sushi, and salted fish products [15, 16, 17], which illustrates that *Salmonella* was continuously contaminated by these RTE products. In addition, human salmonellosis caused by contaminated fish products has been reported in European countries [18, 19], where processing is considered to be one of the important sources of contamination [20, 21]. Martinez-Urtaza and Liebana [21] and Morris et al. [22], who evaluated the incidence of *Salmonella* in seafood processing plants, surveillance of foodborne pathogens during processing is essential to a successful contamination control program. When monitoring *Salmonella* contamination, serotyping and molecular subtypes are often applied to trace the source of bacterial contamination [23, 24].

The incidence of salmonellosis due to fish consumption has become a concern of public health agencies in several countries, because the increased consumption of fish farming, especially raw products, which increases the risk of exposure to pathogens, especially in vulnerable groups, such as the elderly, increases significantly. pregnant women and infants [25, 26]. The importance of this pathogen in fish can be assessed and evaluated, as records show that fish is the source of transmission responsible for 7% of total foodborne outbreaks in Europe in 2016, with the majority of *Salmonella* infections in humans linked to

fish consumption caused by Salmonella Typhimurium and Enteritidis serovars [27, 28, 29, 30].

Overview of Salmonella sp.

Salmonella spp. are Gram-negative rod-shaped bacteria that cause salmonellosis. In humans, these pathogenic bacteria cause enteric fever and acute gastroenteritis [31]. Symptoms include mild to severe gastroenteritis, with an incubation period of 6–72 hours [31]. Salmonellosis outbreaks due to fish consumption have been reported in several countries. For example, salmonellosis caused by eel consumption, which is linked to fish farming in Italy and has been reported in Germany [32]. Several reports have linked the presence of Salmonella spp. in various fishes and shell fishes [33, 34]. Various hazards associated with natural fish farming come from the environment or from human or animal activities. Fish can be a vehicle for transmission of Salmonella which can be pathogenic for humans and has a high potential to transmit its antibiotic resistance gene to other pathogens via plasmids [35]. The potential for antibacterial agents to cause the development of resistance in fish pathogens is of worldwide concern [36].

Salmonella grows optimally at 37 °C, but these bacteria are able to grow in a temperature range of 5–47 °C [37]. However, dietary Salmonella exhibits varying survival behavior depending on the type of food matrix and storage conditions [37]. Studies show that Salmonella can survive for more than 16 weeks in frozen processed chicken products and for more than 9 months in frozen beef without much change in quantity [37]. Studies on the viability of Salmonella in chilled and frozen and fishery products are very limited. One study reported that Salmonella Enteritidis survived but did not grow at 3 °C, whereas bacteria thrived at 10 °C vacuum packed fish and poultry [38]. The amount of Salmonella in raw tuna was reduced by 1 to 2 logs after 12 days of storage at 5–7 °C, whereas in frozen storage Salmonella became undetectable after 42 days of storage. In frozen shrimp, populations of *S. Weltevreden* and *S. Senftenberg* decreased but were not eliminated during 12 weeks of storage [39, 40].

Zoonoses in humans

Infection can occur if fishery food contaminated with salmonella is consumed raw or undercooked. Salmonellosis transmitted through food from fisheries is a major cause of human morbidity worldwide [40, 41]. Salmonella spp. cause various diseases such as enteric fever and gastroenteritis. The majority of the estimated 1.3 million annual cases of salmonellosis result from consumption of contaminated food products especially pork, poultry, eggs and vegetables [41]. In comparison, fishery foods are less frequently associated with Salmonella infections although reports from Thailand, Malaysia and other Asian countries indicate that certain serovars, eg *S. Weltevreden*, may be primarily associated with aquatic environments and seafood [42]. Prevalence of Salmonella spp. may also be higher in seafood from Asia compared to seafood from other geographic regions [43, 44]. However, other studies argue that Salmonella spp. can last longer in tropical aquatic environments and should therefore be seen as part of the normal micro-flora of aquaculture products of the sea [45]. Salmonella can be disseminated as a result of water currents, underground springs and rainwater runoff carrying contaminated material [46–49]. Like *E. coli*, Salmonella is continuously released into the environment from infected humans, livestock, domestic animals, and wildlife [50]. Pathogenic and potentially pathogenic bacteria associated with fish

and shellfish include *Mycobacteria*, *Streptococcus iniae*, *Vibrio vulnificus*, *Vibrio* spp, *Aeromonads*, *Salmonella* spp, *Shigella* and others [51, 52, 53]. Human infection by these fish pathogens is usually through contact with infected fish during handling, water or other elements of the fish's living environment [54]. The initial microflora on the surface of the fish is directly related to the aquatic environment whereas the flora in the digestive tract is related to the type of food and condition of the fish [55]. Salmonella sp. is a causative agent of pathological diseases in humans, cattle, poultry and other livestock and Salmonella infection is a major cause of concern for humans, animals and the food industry. Characterization of more than 2,500 Salmonella serovars, the pathogen infects nearly all vertebrates but the severity of infection varies from one serovar to another depending on the specificity of the host [56]. Some Salmonella serovars are limited to one or several hosts while others have a wide spectrum of hosts. An understanding of the mechanisms involving host preference by one serovar over another is essential. This is used to increase our knowledge about host adaptability and will be instrumental in designing better prevention models. Methods involving identification of genetic markers for host specificity will prove to play a role in determining virulence factors for other bacterial pathogens that cause systemic infection [54, 56].

The low standard of health in the coastal areas increases the eutrophication process, thus creating a broad conducive environment for the survival of microbes that eventually infect fish. Analysis of fish tissue slurry showed that the fish harvested from the landing beaches along the coast that had been contaminated by Enterobacteriaceae were; *Salmonella*, *Shigella* and *E. coli* [57, 58, 59]. In the Asia-Pacific region, farmed fish are fed commercial or homemade feed [60]. Homemade feed is used to reduce costs and usually consists of chicken offal and byproducts produced during poultry processing, kitchen waste, and other by-products from the food industry [60, 61]. Homemade diets can be a potential source of foodborne pathogens, particularly *Salmonella* bacteria [9, 62], which can then be transmitted to farmed fish [63, 64] and, in turn, to humans. Detailed information on salmonella due to consumption of tilapia in Malaysia is lacking, as most cases of food poisoning are not reported to the authorities. However, the Malaysian National Public Health Laboratory reports that the five most common nontyphoidal *Salmonella* serovars are *Salmonella* Enteritidis, *Salmonella* Weltevreden, *Salmonella* Corvallis, *Salmonella* Typhimurium, and *Salmonella* Tshiongwé [65]. In Malaysia, freshwater fish are kept using aquaculture ponds, ex-mining ponds, freshwater cages, cement tanks, canvas tanks, and freshwater cage culture systems. Most of the freshwater cultivated fish are maintained using the aquaculture system (59.5%) and ex-mining ponds (25%). Catfish (58.1%) and tilapia (41.3%) were reared in ground ponds and ex-mining [65].

Transmission of bacteria from tilapia (*Oreochromis sp.*)

Fish is considered to be one of the most nutritious and highly desirable food ingredients due to its high nutritional value and is rich in protein, vitamins and unsaturated fatty acids. Fish contains n-3 polyunsaturated fatty acids which is a very important aspect for health-conscious people especially in affluent countries where the death rate from cardiovascular disease is high. Fish contains more easily digestible protein than protein found from other sources. Fish are of great concern for export

earnings because of their higher nutritional value, low cholesterol levels and presence of essential amino acids [64]. Bangladesh is the 5th in world aquaculture production, which accounted for half of the country's total fish production in 2015-2016 [65, 66]. This sector contributed 3.65% to GDP and 1.97% to foreign exchange earnings through exports [66]. It is the second largest export industry in Bangladesh and accounts for 2.5 percent of global shrimp production [66].

The main pathogens affecting cultivation include bacteria, fungi, viruses, and parasites [67-69]. Bacterial diseases have become a major concern for cultivation, especially with warm water temperatures [70]. Different bacterial species are reported to be pathogens against aquatic tilapia, including *Aeromonas hydrophila*, *Edwardsiella tarda*, *Flavobacterium columnare*, *Francisella* spp., *Yersinia ruckeri*, *Staphylococcus epidermidis*, *Vibrio vulnificus*, and *Streptococcus agalactiae* [71-79].

Fish was associated with 24% of foodborne disease outbreaks and 6% of all food poisoning, or foodborne illness [80]. Levels of pathogenic bacteria in tilapia are related to the environment and its handling before entering food markets and restaurants. The bacteria associated with tilapia fish can be transmitted by direct contact and cause foodborne illness. For example, handling of tilapia has been linked to an outbreak of *Vibrio vulnificus* in a Seattle supermarket [81]. Other foodborne pathogenic bacteria including *Salmonella enterica*, enteropathogenic *Escherichia coli*, *Listeria monocytogenes*, *Yersinia enterocolitica*, and *Klebsiella pneumoniae* were identified from fresh tilapia in the Kenyan freshwater fish chain [82]. Shigatoxigenic and enteropathogenic *Escherichia coli* was isolated from farmed tilapia (*Oreochromis niloticus*) in the northeast region of the state of Sao Paulo [83]. It was reported that the microbial quality of tilapia showed that all the tissues were contaminated by salmonella fecal coli. *Salmonella* can spread as a result of water currents, underground springs and rain runoff carrying contaminated material [84, 85]. Human infection by these fish pathogens is usually through contact with infected fish while handling them over, water or other elements of the fish's living environment [86]. These pathogenic organisms have been isolated from freshwater fish such as *Tilapia nilotica* Linn [87]. Tilapia is an important aquaculture production for food supply. Because on a global scale, fish and fish products are the most important sources of protein and it is estimated that more than 30% of the fish consumed by humans comes from aquaculture [88, 89].

Safe consumption of fish and fishery products requires adequate sanitary conditions from harvest to consumption [90, 91]. Consumption of fish and shellfish can cause illness due to infection or poisoning. Most of the foodborne diseases are caused by *Salmonella* spp., *Staphylococcus* spp. and *Escherichia coli*, usually associated with the consumption of fish infected with these bacterial species especially *Salmonella* and *E. coli* [92, 93]. *Salmonella* usually enters the human intestine with food, *Salmonella* spp. must overcome resistance to colonization mediated by the gut microbiota and the innate immune system. *Salmonella* by inducing inflammation and innate immune defense mechanisms. Many models have been developed to study *Salmonella* spp. interactions with the microbiota have helped to identify the factors needed to overcome colonization resistance and to mediate disease [10, 54, 93]. *Salmonella* infection in humans mainly includes typhoid fever and this infection, known as enteric fever, continues

to be one of the most serious public health problems worldwide. The presence of higher levels of *Salmonella* in fish causes several symptoms in the human body such as diarrhea, nausea, vomiting and abdominal pain. Human salmonellosis caused by contaminated fish products has been reported in European countries [18, 19], where processing is considered to be one of the important sources of contamination [20, 21].

ANTIBIOTIC RESISTANCE

Self-limiting gastroenteritis is the main clinical picture developed by *Salmonella*, which in severe cases may require fluid and electrolyte replacement. The use of antibiotics is reserved for patients with serious illness or a high risk of invasive disease [94]. The antibiotic therapy scheme for typhoid fever includes third generation cephalosporin antibiotics, quinolones and macrolides. However, recently it has developed between typhoid *Salmonellas* and non-typhoid strains with a high degree of resistance to quinolones and cephalosporins [95]. The emergence of several drugs resistant to *Salmonella* (MDR) is currently of worldwide concern, and the occurrence of *Salmonella* MDR in food is a risk condition, indicating an increase in the severity of foodborne disease, leading to increased hospitalization rates and the likelihood of death [96]. In contrast, the epidemiology of antimicrobial resistance of *Salmonella* spp. It is complex and can be influenced by factors such as: consumption of antibiotics, human travel, transmission between patients in hospitals, import and trade in food of animal origin or not, trade in live animals within the country or between countries and exposure through animals or the environment human [97].

The use of antibiotics for therapeutics and growth promoters in feed animals has shown that the main factor for the emergence of resistant isolates [98], has been reported from livestock [99-103], poultry [104-108], pets [109-112], fish [113-116], and from animal products. [117-120]. Several studies have reported that pet food, meat, milk and fishery products contaminated by a multi-resistant *S. aureus* strain which has been one of the common causes of severe nosocomial infection for a long time [121]. In contrast, a large number of studies have reported increased incidence of resistance among *Salmonella* spp. isolated from poultry, beef and fishery products [122].

Antibiotics are used for medicinal purposes and as growth promoters in livestock and aquaculture leading to the development of resistance [123]. Acquired resistance to tetracyclines and chloramphenicol has been associated with extensive use of antibiotics in aquaculture in several Asian countries [124]. Sapkota et al. [125] reported that of the top 13 aquaculture producing countries, 92% used oxytetracycline and 24% used chloramphenicol.

It was reported that antibiotic resistance in Gram-negative bacteria from pond culture was higher in ponds undergoing antimicrobial therapy or with a recent history of treatment than in ponds without antimicrobial treatment recently [126, 127]. In a previous study, it was reported that all *S. enteritidis* strains isolated from tilapia sold in a wet market in Selangor, Malaysia were susceptible to rifampin. The emergence of *Salmonella* serovars with high MAR index indicates that these serovars originate from environments where antimicrobials are often used as therapy as growth promoters in animal feed [128, 129]. Some drug-resistant *Salmonella* isolates have been suggested to be more virulent than non-multiple drug-resistant *Salmonella*

isolates [130]. Salmonella resistance to one or more antibiotics has been reported by many investigators [131]. Horizontal transfer of resistance genes in plasmids has been shown between bacteria in fish pond water [131] and in marine sediments [132]. Plasmids in *Salmonella* spp. has been reported to transfer antibiotic resistance and virulence properties [35].

CONCLUSION

The presence of higher levels of *Salmonella* sp in tilapia fish causes several symptoms in the human health such as diarrhea, nausea, vomiting and abdominal pain. *Salmonella* sp in tilapia obtained from the decay of animal waste and feed residue during the traditional cultivation process. *Salmonella* sp is often resistant to several antibiotics which allow it to be transferred to the surrounding environment, which is because in the process of tilapia pond cultivation it is mixed with disposal from livestock waste where there is a lot of antibiotic residue and while cultivation is still using antibiotics. This is what needs to be a public health concern about antibiotic resistance from isolates obtained in tilapia fish.

REFERENCES

- Little DC, Edwards P. Alternative strategies for livestockefish integration with emphasis on Asia. *Ambio* , 1999;28: 118-124.
- Petersen A, Andersen JS, Kaewmak T, Somsiri T, Dalsgaard A. Impact of integrated fish farming on antimicrobial resistance in a pond environment. *Appl. Environ. Microbiol.* 2002; 68: 6036-6042.
- Dhawan A, Kaur S. Effect of pig dung on water quality and polyculture of carp species during winter and summer. *Aquac. Int.* 2002;10: 297
- Shah SQ, Colquhoun DJ, Nikuli HL, Sorum H. Prevalence of antibiotic resistance genes in the bacterial flora of integrated fish farming environments of Pakistan and Tanzania. *Environ. Sci. Technol.* 2012;46: 8672-8679.
- FAO. The State of World Fisheries and Aquaculture, 2014; Rome.
- Zhang WB, Murray FJ, Liu LP, Little DC. A comparative analysis of four internationally traded farmed seafood commodities in China: domestic and international markets as key drivers. *Rev. Aquac.* 2015;0: 1-22.
- Food and Agriculture Organization of the United Nations. Report of the FAO Workshop on the On-Farm Feeding and Feed Management in Aquaculture, 2010; p. 4-5: FAO Fisheries and aquaculture report no. 949. Food and Agriculture Organization of the United Nations, Rome.
- Food and Agriculture Organization of the United Nations. Report of the Expert Workshop on the Application of Biosecurity Measures To Control *Salmonella* Contamination in Sustainable Aquaculture, 2010; p. 9-26: FAO Fisheries and aquaculture report no. 937. Food and Agriculture Organization of the United Nations, Rome.
- Burr WE and Helmboldt CF. *Salmonella* species contaminants in three animal by-products. *Avian Dis.* 1962; 6:441-443.
- Lunestad BT, Nesse L, Lassen J, Svihus B, Nesbakken T, Fossum K, Rosnes JT. *Salmonella* in fish feed; occurrence and implications for fish and human health in Norway. *Aquaculture* 2007; 265:1-8.
- Junior PG, Assunc,ão AW, Baldin JC, Amaral LA. Microbiological quality of whole and filleted shelf-tilapia. *Aquaculture* 2014; 433:196-200.
- Scallan E, Hoekstra RM, Angulo FJ, Tauxe RV, Widdowson MA, et al. Foodborne illness acquired in the United States—major pathogens. *Emerg. Infect. Dis.* 2011;17:7-15.
- Su HP, Chiu SI, Tsai JL, Lee CL, Pan TM. Bacterial food-borne illness outbreaks in northern Taiwan, 1995- 2001. *J. Infect. Chemother.* 2005; 11:146-151.
- Bean NH, Goulding JS, Lao C, Angulo FJ. Surveillance for foodborne-disease outbreaks—United States, 1988- 1992. *MMWR CDC Surveill. Summ.* 1996; 45:1-66.
- Cabedo L, Picart i Barrot L, Teixido i Canelles A. Prevalence of *Listeria monocytogenes* and *Salmonella* in ready-to-eat food in Catalonia, Spain. *J. Food Prot.* 2008; 71:855-859.
- Heinitz ML, Johnson JM. The incidence of *Listeria* spp, *Salmonella* spp, and *Clostridium botulinum* in smoked fish and shellfish. *J. Food Prot.* 1998; 61:318-323.
- Heinitz ML, Ruble RD, Wagner DE, Tatini SR. Incidence of *Salmonella* in fish and seafood. *J. Food Prot.* 2000; 63:579- 592.
- Fell G, Hamouda O, Lindner R, Rehmet S, Liesegang A, Prager R, Gericke B, Petersen L. An outbreak of *Salmonella* Blockley infections following smoked eel consumption in Germany. *Epidemiol. Infect.* 2000; 125:9-12.
- Guerin PJ, De Jong B, Heir E, Hasseltvedt V, Kapperud G, Styrmo K, et al. Outbreak of *Salmonella* Livingstone infection in Norway and Sweden due to contaminated processed fish products. *Epidemiol. Infect.* 2004; 132:889-895.
- Klaeboe H, Rosef O, Saebo M. Longitudinal studies on *Listeria monocytogenes* and other *Listeria* species in two salmon processing plants. *Int. J. Environ. Health Res.* 2005; 15:71-77.
- Martinez-Urtaza J, Liebana E. Use of pulsed-field gel electrophoresis to characterize the genetic diversity and clonal persistence of *Salmonella* Senftenberg in mussel processing facilities. *Int. J. Food Microbiol.* 2005; 105:153-163.
- Morris GK, Martin WT, Shelton WH, Wells JG, Brachman PS. *Salmonellae* in fish meal plants: relative amounts of contamination at various stages of processing and a method of control. *Appl. Microbiol.* 1970; 19:401-408.
- Kilic A, Demir O, Kocak N, Levent B, Eyigun CP, Tekbas OF, et al. Analysis of an outbreak of *Salmonella* Enteritidis by repetitive-sequence-based PCR and pulsed-field gel electrophoresis. *Intern. Med. (Tokyo)* 2010; 49:31-36.
- Weigel RM, Qiao B, Teferedegne B, Decker DA, Isaacson RE, et al. Comparison of pulsed field gel electrophoresis and repetitive sequence polymerase chain reaction as genotyping methods for detection of genetic diversity and inferring transmission of *Salmonella*. *Veter. Microbiol.* 2004; 100:205-217.
- Zhang J, et al. Prevalence of antimicrobial resistance of nontyphoidal *Salmonella* serovars in retail aquaculture products. *International Journal of Food Microbiology*, 2015;210:47-52.
- Paudyal NV. et al. Prevalence of foodborne pathogens in food from selected African countries—A meta-analysis. *International Journal of Food Microbiology*, 2017;249:35-43.

27. Santiago JDA. S. Bactérias patogênicas relacionadas à ingestão de pescados-revisão. *Arquivos de Ciências do Maret*, 2013;46:2.
28. SA. European Food Safety Authority Guidelines for reporting data on zoonoses, antimicrobial resistance and foodborne outbreaks using the EFSA data models for the data collection framework (DCF) to be used in 2017 for 2016 data. 2016; ISSN 2397-8325.
29. D. Characterization of extended-spectrum β -lactamase (ESBL) producing non-typhoidal *Salmonella* (NTS) from imported food products. *International Journal of Food Microbiology*. 2015; 214:12-17.
30. Center Diseases Control. Multistate Outbreak of *Salmonella* Paratyphi B variant L and tartrate and *Salmonella* Weltevreden infections linked to frozen raw tuna. 2015.
31. Hohmann EL. Nontyphoidal salmonellosis. *Clin. Infect. Dis.* 2001;32:263-269.
32. Fell G, Hamouda O, Lindner R, Rehm S, Liesegang A, Prager R, et al. An outbreak of *Salmonella* Blockley infections following smoked eel consumption in Germany. *Epidemiology and Infection* 2000;125: (1), 9-12.
33. Brands DA, Inman AE, Gerba CP, Mare J, Billington SJ, Saif LA. Prevalence of *Salmonella* spp. in oysters in the United States. *Applied and Environmental Microbiology* 2005;71: 893-897.
34. Duran GM, Marshall DL. Ready to eat shrimp as an international vehicle of antibiotic-resistant bacteria. *Journal of Food Protection* 2005;68 (11): 2395-2401.
35. Hradecka H, Karasova D, Rychlik I. Characterization of *Salmonella* enterica serovar Typhimurium conjugative plasmids transferring resistance to antibiotics and their interaction with the virulence plasmid. *Journal of Antimicrobial Chemotherapy* 2008;62 (5): 938-941.
36. Schnick RA. International harmonization of antimicrobial sensitivity determination for aquaculture drugs. *Aquaculture* 2001;196: 277-288.
37. Nychas GJE, Tassou CC. Growth/survival of *Salmonella* enteritidis on fresh poultry and fish stored under vacuum or modified atmosphere. *Letters in Applied Microbiology*, 1996;23(2): 115-119.
38. Liu C, Mou J, Su YC. Behavior of *Salmonella* and *Listeria monocytogenes* in raw yellowfin tuna during cold storage. *Foods (Basel, Switzerland)*, 2016;5(1): 16
39. Noda H, Chisuwa M, Kaneko M, Onoue Y, Takatori K, Hara-Kudo Y. Survival of *Salmonella* Weltevreden and *S. Senftenberg* in black tiger shrimp under frozen storage. *Shokuhin Eiseigaku Zasshi. Journal of the Food Hygienic Society of Japan*, 2009;50(2): 85-88.
40. Hassan R, Teclé S, Adcock B, Kellis M, Weiss J, Saupé et al. Multistate outbreak of *Salmonella* Paratyphi B variant L(+) tartrate(+) and *Salmonella* Weltevreden infections linked to imported frozen raw tuna: USA, March-July 2015. *Epidemiology & Infection*, 2018;146(11): 1461-1467.
41. Gong J, Zhang J, Xu M, Zhu C, Yu Y, Liu X, et al. Prevalence and fimbrial genotype distribution of poultry *Salmonella* isolates in China (2006 to 2012). *Appl. Environ. Microbiol.* 2014;80: 687-693.
42. Deekshit VK, Ballamoole KK, PraveenRai M, Karunasagar I, Karunasagar I. Draft genome sequence of multidrug resistant *Salmonella* enterica serovar Weltevreden isolated from seafood. *J. Genom.* 2015;3: 57-58.
43. Yan H, Li L, Alam MJ, Shinoda S, Miyoshi S, Shi L. Prevalence and antimicrobial resistance of *Salmonella* in retail foods in northern China. *Int. J. Food Microbiol.* 2010;143: 230-234.
44. Koonse B, Burkhardt W, Chirtel S, Hoskin GP. *Salmonella* and the sanitary quality of aquacultured shrimp. *J. Food Prot.* 2005;68: 2527-2532.
45. FAO. Report of FAO expert workshop on the application of biosecurity measures to control *Salmonella* contamination in sustainable aquaculture. In: FAO Fisheries and Aquaculture Report No. 937. FAO, Mangalore, India. 2010.
46. Fortes TP. et al. Ilhas de patogenicidade de *Salmonella* enterica: uma revisão. *Revista do Instituto Adolfo Lutz*, 2012;71:219227.
47. Wotzka SY. et al. *Salmonella* Typhimurium Diarrhea Reveals Basic Principles of Enteropathogen Infection and Disease Promoted DNA Exchange. *Cell Host & Microbe* 2017;21(12):443-454,
48. Chao W, Ding R, Chen R. Survival of pathogenic bacteria in environmental microcosms. *Chinese J. Microbial Immun.* 1987; Vol. 20: pp.339-348,
49. Abdel-Monem MH, Dowidar AA. Recoveries of *Salmonella* from soil in Eastern region of Saudi Arabia Kingdon, *J. Egypt. Public Health Assoc.* 1990; vol, 65: pp.61- 75,
50. Baudart J, Lemarchand K, Brisabois A, Lebaron P, Diversity of *Salmonella* strains isolated from the aquatic environment as determined by serotyping and amplification of ribosomal DNA spacer regions. *Appl. Environ. Microbiol.* 2000;66(4):1544-1552,
51. Lipp EK, Rose JB. The role of sea food in foodborne diseases in the United States of America. *Rev. Sci. Tech. OIE*, 1997;16:620640,
52. Zlotkin A, Eldar A, Ghifino C, Bercovier H. Identification of *Lactococcus garvieae* by PCR. *J. Clin. Microbiol.* 1989;36: 983985,
53. Bhaftopadhyay P. Fish - catching and handling. In: Robinson R.K. (ed.): Encyclopedia of Food Microbiology, Academic Press, London, 2000; vol.2, pp. 1547,
54. Acha PN, Szyfres B. Zoonoses and communicable diseases common to man and animals, Bacterioses and mycoses. 3rd ed. Scientific and Technical Publication, vol.1, No. 580, Pan American Health Organization, Regional Office of the WHO, Washington, USA, ISBN 92 75 31580 9. 2003; pp.384,
55. Liston J. Microbiology in fishery science, In Connell, JJ. (ed). Advances in Fish Science and Technology; Jubilee Conference of Torry Research Oxford: Fishing News Books Ltd, Farnham, UK, 1980.
56. Singh, V. *Salmonella* Serovars And Their Host Specificity. *J Vet Sci Anim Husb*, 2013; 1(3): 301.
57. Abila RO, Jansen EG. From Local to Global markets: The fish exporting and fishing meal industries of Lake Victoria- Structure, Strategies and Socio-economic Impacts in Kenya, IUCN Eastern Africa programme. Socioeconomics of Lake Victoria fisheries: Report No.2, The World Conservation Union, Nairobi, 1997.

58. Kayambo S, Sven EJ. Lake Victoria. Experience and lessons learnt. A case Study for Preliminary Risk Assessment Report. 2006; pp. 431 – 446.
59. Onyango D, Wandili S, Kakai R, Waindi EN, Isolation of *Salmonella* and *Shigella* from fish harvested along winam Gulf of Lake Victoria, Kenya, *J. infect. Dis. In Developing Countries*, 2008;2: 106111,
60. Food and Agriculture Organization of the United Nations. Report of the FAO Workshop on the On-Farm Feeding and Feed Management in Aquaculture, p. 4–5. FAO Fisheries and aquaculture report no. 949. Food and Agriculture Organization of the United Nations, Rome. 2010.
61. Food and Agriculture Organization of the United Nations. Report of the Expert Workshop on the Application of Biosecurity Measures to Control *Salmonella* Contamination in Sustainable Aquaculture, p. 9–26. FAO Fisheries and aquaculture report no. 937. Food and Agriculture Organization of the United Nations, Rome. 2010.
62. Lunestad BT, Nesse L, Lassen J, Svihus B, Nesbakken T, Fossum K, Rosnes JT. *Salmonella* in fish feed; occurrence and implications for fish and human health in Norway. *Aquaculture* 2007; 265:1–8.
63. Junior PG, Assunc,ão AW, Baldin JC, Amaral LA. Microbiological quality of whole and filleted shelf-tilapia. *Aquaculture* 2014; 433:196–200.
64. Torpdah M, Sorensen G, Lindstedt BA, Nielsen EM. Tandem repeat analysis for surveillance of human *Salmonella* Typhimurium infections. *Emerg. Infect. Dis.* 2007; 13:388–395.
65. Department of Fisheries Malaysia. Annual Fishery Statistic 2011. Department of Fisheries Malaysia, Putrajaya, Malaysia. 2011.
66. Department of Fisheries. National fish week, Ministry of Fisheries and Livestock, Government of the people's Republic of Bangladesh. 2016;19: 13-132.
67. Ramaiah N. A review on fungal diseases of algae, marine fishes, shrimps and corals. *Indian Journal of Marine Sciences*. 2006;35: 380-387.
68. Guo FC, Woo PT. Selected parasitosis in cultured and wild fish. *Veterinary Parasitology*. 2009;163: 207-216.
69. Vega-Heredia S, Mendoza-Cano F, Sanchez-Paz A. The infectious hypodermal and haematopoietic necrosis virus: a brief review of what we do and do not know. *Transboundary and Emerging Diseases*. 2012;59: 95-105.
70. Indiy TV, Ravindranath K, Sreeraman PK, et al. *Aeromonas salmonicida* associated with mass mortality of *Cyprinus carpio* and *Oreochromis mossambicus* in a freshwater reservoir in Andhra Pradesh, India. *Journal of Aquaculture in the Tropics*. 1994; 9: 259-268.
71. Huang S, Chen W, Shei M, et al. Studies on epizootiology and pathogenicity of *Staphylococcus epidermidis* in Tilapia *Oreochromis spp.* cultured in Taiwan. *Zoological Studies*. 1999; 38:178-188.
72. Chen CY, Wooster GA, Bowser PR. Comparative lipid chemistry and histopathology of tilapia infected with *Vibrio vulnificus* or *Streptococcus iniae* or exposed to carbon tetrachloride, gentamicin, or copper sulfate. *Aquaculture*. 2004; 239: 421-443.
73. Jimenez AP, Iregui CA, Figueroa J. In vitro/in vivo characterization and evaluation of *Aeromonas hydrophila* lipopolysaccharides (LPS). *Acta Biologica Colombiana*. 2008; 13: 147-162.
74. Zaza AE, Moustafa M, Abdelaziz M, et al. *Yersinia ruckeri* infection in cultured Nile tilapia, *Oreochromis niloticus*, at a semi-intensive fish farm in lower Egypt. *African Journal of Aquatic Science*. 2008; 33: 283-286.
75. Jeffery KR, Stone D, Feist SW, et al. An outbreak of disease caused by *Francisella sp.* in Nile tilapia *Oreochromis niloticus* at a recirculation fish farm in the UK. *Diseases of Aquatic Organisms*. 2010; 91:161-165.
76. Mohamed SG, Saleh WD. *Flavobacterium columnare* infection in cultured *Oreochromis niloticus*. *Assiut Veterinary Medical Journal*. 2010; 56: 15-30.
77. Ye X, Li J, Lu M, et al. Identification and molecular typing of *Streptococcus agalactiae* isolated from pond-cultured tilapia in China. *Fisheries Science*. 2011; 77: 623-632.
78. Iregui CA, Guarín M, Tibata VM, et al. Novel brain lesions caused by *Edwardsiella tarda* in a red tilapia *Oreochromis spp.* *Journal of Veterinary Diagnostic Investigation*. 2012; 24: 446-449.
79. Zhang Z, Lan J, Li Y, et al. The pathogenic and antimicrobial characteristics of an emerging *Streptococcus agalactiae* serotype IX in Tilapia. *Microbial Pathogenesis*. 2018; 122: 39-45.
80. CDC. Surveillance for Foodborne Disease Outbreaks United States Annual Report. GA, USA. 2013.
81. Food poisoning bulletin (FPB). *Vibrio* Outbreak Associated with Tilapia from Seattle Supermarket. 2017.
82. Long HA, Ngayo MO, Mwaniki M, et al. Microbiological Safety of Fresh Tilapia *Oreochromis niloticus* from Kenyan Fresh Water Fish Value Chains. *Journal of Food Protection*. 2018; 81: 1973-1981.
83. Dozo MV, Borges CA, Beraldo LG, et al. Shigatoxigenic and atypical enteropathogenic *Escherichia coli* in fish for human consumption. *Brazilian Journal of Microbiology*. 2018; 49: 936-941.
84. Thampuren N, Surendraraji A, Surendran P. Prevalence and characterization of typical and atypical *Escherichia coli* from fish sold at retail in Cochin, India *Journal Food Protection*, 2005; 68: 2208 – 2211.
85. Abdel-Monem MH. Dowidar AA. Recoveries of *Salmonella* from Soil in Eastern region of Saudi Arabic Kingdom, *Journal Egypt Public Health Association*. 1990;65: 61–75.
86. Acha PN, Szyfres BC. Zoonoses and communicable diseases common to man and animals. *Bacterioses and mycoses*. 3rd ed. Scientific and Technical Publication: 1: 580, Pan American Health Organization, Regional Office of the WHO, Washington, USA: ISBN 92 75 315809. 2003; pp 384.
87. D' Aoust J, Sewell A, Daley E, Greco P. Antibiotic resistance of agricultural and food borne, *Salmonella* isolates in Canada: 1986 – 1989. *Journal Food Protection*, 1992;55: 428 – 434.
88. Hastein T, Hjeltnes B, Lillehaug A, Utne Skare J, Berntssen M, Lundebye K. Food safety hazards that occur during the production stage: challenges for fish farming and the fishing industry Review Science. Technology, 2006;25: 607–625.

89. Morales G, Blanco L, Arias M, Chaves C. Bacteriological evaluation of fresh tilapia coming from the Northern Region of Costa Rica. *Archive Latinoam Nutrition*. 2004;54: 433 – 437.
90. 57. Adhi N. Prevalence and antimicrobial resistance of *Salmonella* spp. in raw retail frozen imported freshwater fish to Eastern Province of Saudi Arabia. *Asian Pacific Journal of Tropical Biomedicine*, 2014;4:234-238
91. 34. Habban AJM. Fish Farms as a source for parasites transport: Parasitological and developmental studies of *Prohemistomum vivax* with the ameliorating role of *Moringa oleifera* in the treatment. *J American Sci*. 2014;10: 6-14.
92. Yagoub SO. Isolation of Enterobacteriaceae and *Pseudomonas* spp. from raw fish sold in fish market in Khartoum state, *African J Bacterio Res*. 2009;1: 085-088.
93. Ahmer, B.M.M. and Gunn, J.S. Interaction of Salmonella spp. with the intestinal microbiota. *Frontier in Microbiology*, 2011; 2: 101
94. WHO, World Health Organization? Antimicrobial resistance: global report on surveillance. 2014, Publications of the World Health Organization are available on the WHO website (www.who.int) or can be purchased from WHO Press, World Health Organization, 20 Avenue Appia, 1211 Geneva 27, Switzerland.
95. Cosby DE. *Salmonella* and antimicrobial resistance in broilers: A review. *J. Appl. Poult. Res*. 2015; 24:408-426.
96. Crump JA, Heyderman RS. A Perspective on Invasive *Salmonella* Disease in Africa. *Clinical Infectious Diseases*. 2015;61(S4): S235-40,
97. 32. EC (European Centre for Disease Prevention and Control), EFSA (European Food Safety Authority), and EMA (European Medicines Agency), ECDC/EFSA/EMA second joint 23. report on the integrated analysis of the consumption of antimicrobial agents and occurrence of antimicrobial resistance in bacteria from humans and food-producing animals - Joint Interagency Antimicrobial Consumption and Resistance Analysis (JIACRA) Report. *EFSA Journal* 2017;15(7):4872
98. Barber DA, Miller GY, McNamara PE. Models of antimicrobial resistance and food-borne illness: examining assumptions and practical application. *J. Food Protect*. 2003;66: 700-709.
99. 65. Indri, M.H., Oktavianto, A and Hastutiek, P. Tetracycline Resistance Gene in *Streptococcus Agalactiae* Isolated from Bovine Subclinical Mastitis in Surabaya, Indonesia. *Philipp. Journal of Veterinary Medicine*. 2018; 55 (SI): 115-120.
100. Effendi M.H, Harijani N, Budiarto, Triningtya N.P, Tyasningsih W. and Plumeriastuti H. Prevalence of Pathogenic *Escherichia Coli* Isolated from Subclinical Mastitis in East Java Province, Indonesia. *Indian Vet. J*. 2019; 96 (03): 22 – 25.
101. Tyasningsih, W., Effendi, M. H., Budiarto, B., Syahputra, I. R. Antibiotic Resistance to *Staphylococcus aureus* and Methicillin Resistant *Staphylococcus aureus* (MRSA) Isolated from Dairy Farms in Surabaya, Indonesia. *Indian Vet. J*, 2019; 96(11), 27-31.
102. Putra, A.R. Effendi, M.H. Koesdarto, S. Suwarno, S. Tyasningsih, W. and Estoepangestie, A.T. Detection 42. the extended spectrum β -lactamase produced by *Escherichia coli* from dairy cows by using the Vitek-2 method in Tulungagung regency, Indonesia. *Iraqi Journal of Veterinary Sciences*, 2020; 34 (1): 203-207.
103. Putra ARS, Effendi MH, Koesdarto S, and Tyasningsih W. Molecular Identification of Extended Spectrum Beta-Lactamase (ESBL) Producing *Escherichia coli* Isolated from Dairy Cows in East Java Province, Indonesia. *Indian Vet. J*. 2019; 96 (10): 26 – 30.
104. Wibisono FJ, Sumiarto B, Untari T, Effendi MH, 21. matasari DA, Witaningrum AM. The Presence of Extended Spectrum Beta-Lactamase (ESBL) Producing *Escherichia coli* On Layer Chicken Farms In Blitar Area, Indonesia. *Biodiversitas*. 2020; 21 (6): 2667-2671.
105. Rahmahani J, Salamah, Mufasirin, Tyasningsih W, and Effendi MH. Antimicrobial Resistance Profile of *Escherichia coli* From Cloacal Swab of Domestic Chicken in Surabaya Traditional Market. *Biochem. Cell. Arch*. 2020; 20 (1): 2993-2997.
106. Wibisono, F.J., Sumiarto, B., Untari, T., Effendi, M.H., 21. matasari, D.A., Witaningrum, A.M. CTX Gene of Extended Spectrum Beta-Lactamase (ESBL) Producing *Escherichia coli* on Broilers in Blitar, Indonesia. *Sys Rev Pharm* 2020;11(7): 396-403.
107. Permatasari, D.A., Witaningrum, A.M., Wibisono, F.J., Effendi, M.H. Detection and prevalence of multidrug-resistant *Klebsiella pneumoniae* strains isolated from poultry farms in Blitar, Indonesia. *Biodiversitas*, 2020; 21 (10): 4642-4647.
108. Wibisono, F.J., Sumiarto, B., Untari, T., Effendi, M.H., Permatasari, D.A., Witaningrum, A.M. Short Communication: Pattern of antibiotic resistance on extended-spectrum beta-lactamases genes producing *Escherichia coli* on laying hens in Blitar, Indonesia. *Biodiversitas*, 2020; 21 (10): 4631- 4635.
109. Rahmaniar, R. P., Yunita, M. N., Effendi, M. H., Yanestria, S. M. Encoding Gene for Methicillin Resistant *Staphylococcus aureus* (MRSA) Isolated from Nasal Swab of Dogs. *Indian Vet. J*, 2020; 97(02), 37-40.
110. Kristianingtyas L, Effendi, MH, Tyasningsih W, Kurniawan F. Genetic 21. ntification of blaCTX-M Gene and blaTEM Gene on Extended Spectrum Beta Lactamase (ESBL) Producing *Escherichia Coli* from Dogs. *Indian Vet. J*. 2020; 97 (01): 17 – 21.
111. Decline, V., Effendi, M. H., Rahmaniar, R. P., Yanestria, S. M., Harijani, N. Profile of antibiotic-resistant and presence of methicillin-resistant *Staphylococcus aureus* from nasal swab of dogs from several animal clinics in Surabaya, Indonesia. *Intl J One Health*, 2020; 6(1), 90-94.
112. Yunita, M. N., Effendi, M. H., Rahmaniar, R. P., Arifah, S., Yanestria, S. M. Identification of Spa Gene for Strain Typing of Methicillin Resistant *Staphylococcus aureus* (MRSA) Isolated From Nasal Swab Of Dogs. *Biochem. Cell. Arch*. 2020; 20 (1), 2999-3004.
113. Helmi, AM, Mukti, AT, Soegianto, A and Effendi, MH. A Review of Vibriosis in Fisheries: Public Health Importance. *Sys Rev Pharm*, 2020;11(8):51-58.
114. Helmi, AM., Mukti, AT., Soegianto, A., Mahardika, K., Mastuti, I., Effendi, MH., Plumeriastuti, H. A Review of Bacterial Zoonoses and Antimicrobial Resistant (AMR) on Grouper fish (*Epinephelus sp.*). *Sys Rev Pharm* 2020;11(9):79-88.
115. Effendi MH, Bintari IG, Aksono EB, Hermawan IP. Detection of blaTEM Gene of *Klebsiella pneumoniae* Isolated from Swab of Food Producing Animals in

- East Java. *Tropical Animal Science Journal*. 2018;41(3):174-178.
116. Yanestria, S.M., Rahmaniar, R.P., Wibisono, F.J., Effendi, M.H. Detection of *invA* gene of *Salmonella* from milkfish (*Chanos chanos*) at Sidoarjo wet fish market, Indonesia, using polymerase chain reaction technique. *Veterinary World*, 2019; 12(1): 170-175.
 117. Effendi, MH, Harijani N, Yanestria SM, Hastutiek P. 2018. Identification of shiga toxin-producing *Escherichia coli* in raw milk samples from dairy cows in Surabaya, Indonesia. *Philippine J Vet Med*.55(S1):109-114.
 118. Harijani, N., Wandari, A., Effendi, M.H. and Tyasingsih. W. Molecular Detection of Encoding Enterotoxin C Gene and Profile of Antibiotic Resistant on *Staphylococcus Aureus* Isolated from Several Dairy Farms in East Java, Indonesia. *Biochem. Cell. Arch*. 2020; 20 (1): 3081-3085.
 119. Ramandinianto, S.C., Khairullah, A.R., Effendi, M.H., Tyasingsih, W. and Rahmahani, J. Detection of Enterotoxin type B gene on Methicillin Resistant *Staphylococcus aureus* (MRSA) isolated from raw milk in East Java, Indonesia. *Sys Rev Pharm*, 2020;11(7):290-298.
 120. Ramandinianto, S.C., Khairullah, A.R., Effendi, M.H. *MecA* gene and methicillin-resistant *Staphylococcus aureus* (MRSA) isolated from dairy farms in East Java, Indonesia. *Biodiversitas*, 2020; 21(8): 3562-3568.
 121. Cailhol J, Lailier R, Bouvet P, La Vieille S, Gauchard F, Sanders P, Brisabois, A. Trends in antimicrobial resistance phenotypes in non-typhoid *Salmonellae* from human and poultry origins in France. *Epidemiol. Infect.* 2006;134, 171–178.
 122. Serrano PH. Responsible use of antibiotics in aquaculture. FAO Fisheries Technical Paper. 2005; No. 469. FAO, Rome
 123. Mohamed S, Nagaraj G, Chua FHC, Wang YG. The use of chemicals in aquaculture in Malaysia and Singapore. In: Arthur, J.R, Lavilla-Pitogo, C.R, Subasinghe, R.P. (Eds.), Use of Chemicals in Aquaculture in Asia: Proceedings of the Meeting on the Use of Chemicals in Aquaculture in Asia. Aquaculture Department, Southeast Asian Fisheries Development Center.2000.
 124. Sapkota A, Sapkota AR, Kucharski M, Burke J, McKenzie S, Walker P, Lawrence R. Aquaculture practices and potential human health risks: current knowledge and future priorities. *Environment International* 2008;34 (8): 1215–1226.
 125. Doublet B, Weill FX, Fabre L, Chaslus-Dancla E, Cloeckaert A. Variant *Salmonella* genomic island 1 antibiotic resistance gene cluster containing a novel 3'-Naminoglycoside acetyltransferase gene cassette, *aac(3)-Id*, in *Salmonella enterica* serovar Newport. *Antimicrobial Agents and Chemotherapy* 2004;48 (10): 3806–3812.
 126. McPhearson RM, DePaolo A, Zywno SR, Motes ML, Guarino AM. Antibiotic resistance in Gram-negative bacteria from cultured catfish and aquaculture ponds. *Aquaculture* 1991;99: 203–211.
 127. Radu S, Sahilah AM, Rusul G, Samuel L, Zuraini MI, Nasreldin EH. Comparison of arbitrarily primed PCR, antibiotic resistance and plasmid profiling for differentiating *Salmonella Enteritidis* isolated from fish. *Asian Fisheries Science* 2000;13: 13–20.
 128. Singh S, Yadav AS, Singh SM, Bharti P. Prevalence of *Salmonella* in chicken eggs collected from poultry farms and marketing channels and their antimicrobial resistance. *Food Research International* 2010;43 (8): 2027–2030.
 129. Foley SL, Lynne AM. Food animal-associated *Salmonella* challenges: pathogenicity and antimicrobial resistance. *Journal of Animal Science* 2008;86: E173–E187.
 130. Pan ZM, Geng SZ, Zhou YQ, Liu ZY, Fang Q, Liu BB, Jiao XA. Prevalence and antimicrobial resistance in *Salmonella* spp. isolated from domestic animals in Eastern China. *Journal of Animal and Veterinary Advances* 2010;9 (17): 2290–2294.
 131. Aoki T. Resistance plasmids and the risk of transfer. In: Bernoth, E.-M, Ellis, A.E, Midtyling, P.J, Olivier, G, Smith, P. (Eds.), *Furunculosis: multidisciplinary fish disease research*. Academic Press, San Diego, 1997; pp. 433–440.
 132. Stewart GJ, Sinigalliano C.D. Detection of horizontal gene transfer by natural transformation in native and introduced species of bacteria in marine and syntetic sediments. *Applied and Environmental Microbiology* 1990;56 (6): 1818–1824.

ORIGINALITY REPORT

19%

SIMILARITY INDEX

13%

INTERNET SOURCES

15%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

-
- | | | |
|---|---|----|
| 1 | C. N. Groesbeck, J. M. DeRouchey, M. D. Tokach, R. D. Goodband, S. S. Dritz, J. L. Nelssen. "Effects of irradiation of feed ingredients added to meal or pelleted diets on growth performance of weanling pigs ¹ ", <i>Journal of Animal Science</i> , 2009
Publication | 1% |
| 2 | Www.arcjournals.org
Internet Source | 1% |
| 3 | www.e-repository.unsyiah.ac.id
Internet Source | 1% |
| 4 | David Foster, Adam Locker, Andrea Maldonado, Eileen O'Dea et al. "Publication of scientific data from EU - coordinated monitoring programmes and surveys", <i>EFSA Supporting Publications</i> , 2019
Publication | 1% |
| 5 | amif.org
Internet Source | 1% |
| 6 | environhealthprevmed.biomedcentral.com
Internet Source | 1% |

7	www.elsevier.es Internet Source	1 %
8	www.health.nt.gov.au Internet Source	<1 %
9	www.lvemp.org Internet Source	<1 %
10	G.O. Abakpa, V.J. Umoh, J.B. Ameh, S.E. Yakubu, J.K.P. Kwaga, S. Kamaruzaman. "Diversity and antimicrobial resistance of <i>Salmonella enterica</i> isolated from fresh produce and environmental samples", Environmental Nanotechnology, Monitoring & Management, 2015 Publication	<1 %
11	www.daff.gov.au Internet Source	<1 %
12	Www.mdpi.com Internet Source	<1 %
13	synapse.koreamed.org Internet Source	<1 %
14	academictree.org Internet Source	<1 %
15	mircenbiotech.in Internet Source	<1 %

www.nap.edu

16

Internet Source

<1 %

17

Bernard Lama Ngota. "Advancing the contextual factors impeding the development of aquapreneurship in Cameroon", *Entrepreneurial Business and Economics Review*, 2020

Publication

<1 %

18

Mariana Rodrigues Gomes da CRUZ, Ylka Jannielly Barbalho de Souza LEITE, Julyanna de Lima MARQUES, Sabrina Lunara Santos PAVELQUESI et al. "Microbiological quality of minimally processed vegetables commercialized in Brasilia, DF, Brazil", *Food Science and Technology*, 2019

Publication

<1 %

19

T Budiati, W Suryaningsih, S. Umaroh, B. Poerwanto, A Bakri, E Kurniawati. "Antimicrobial activity of essential oil from Indonesian medicinal plants against food-borne pathogens", *IOP Conference Series: Earth and Environmental Science*, 2018

Publication

<1 %

20

mro.massey.ac.nz

Internet Source

<1 %

21

worldwidescience.org

Internet Source

<1 %

22 www.ajol.info Internet Source <1 %

23 "Guide to Antimicrobial Use in Animals", Wiley, 2008 Publication <1 %

24 Amir Sapkota, Amy R. Sapkota, Margaret Kucharski, Janelle Burke, Shawn McKenzie, Polly Walker, Robert Lawrence. "Aquaculture practices and potential human health risks: Current knowledge and future priorities", Environment International, 2008 Publication <1 %

25 Abdel-Fattah M. El-Sayed. "Tilapia Co-culture in Egypt", Wiley, 2016 Publication <1 %

26 Anurag Jyoti, Rajesh Singh Tomar, Rishi Shanker. "Chapter 5 Nanosensors for the Detection of Pathogenic Bacteria", Springer Science and Business Media LLC, 2016 Publication <1 %

27 Lihong Wang, Long Li, Yan Lv, Qiaoling Chen, Junchang Feng, Xin Zhao. "Lactobacillus plantarum Restores Intestinal Permeability Disrupted by Salmonella Infection in Newly-hatched Chicks", Scientific Reports, 2018 Publication <1 %

28 biblio.ugent.be

<1 %

29

J Triastuti, D Kintani, E M Luqman, D Y Pujiastuti. " The motility and motion duration of jatimbunan tilapia () spermatozoa in different salinity ", IOP Conference Series: Earth and Environmental Science, 2018

Publication

<1 %

30

Rosalie A. Schnick. "International harmonization of antimicrobial sensitivity determination for aquaculture drugs", Aquaculture, 2001

Publication

<1 %

31

"Fishery Products", Wiley, 2009

Publication

<1 %

32

Nicole Rosenkötter, Timo Clemens, Kristine Sørensen, Helmut Brand. "Twentieth anniversary of the European Union health mandate: taking stock of perceived achievements, failures and missed opportunities – a qualitative study", BMC Public Health, 2013

Publication

<1 %

33

Todd R. Callaway. "Orange Peel Products Can Reduce *Salmonella* Populations in Ruminants", Foodborne Pathogens and Disease, 06/08/2011

Publication

<1 %

34

free-journal.umm.ac.id

Internet Source

<1 %

35

ynu.repo.nii.ac.jp

Internet Source

<1 %

36

Douglas E. Cosby, Nelson A. Cox, Mark A. Harrison, Jeanna L. Wilson, R. Jeff Buhr, Paula J. Fedorka-Cray. "Salmonella and antimicrobial resistance in broilers: A review", *Journal of Applied Poultry Research*, 2015

Publication

<1 %

37

Jie Yang, Zijing Ju, Yi Yang, Xiaonan Zhao, Zhiyu Jiang, Shuhong Sun. "Serotype, antimicrobial susceptibility and genotype profiles of Salmonella isolated from duck farms and a slaughterhouse in Shandong province, China", *BMC Microbiology*, 2019

Publication

<1 %

38

Ristori, Christiane Asturiano, Ruth Estela Gravato Rowlands, Cecília Geraldés Martins, Maria Luisa Barbosa, Júlia T.U. Yoshida, and Bernadette D.G. de Melo Franco. "Prevalence and Populations of *Listeria monocytogenes* in Meat Products Retailed in Sao Paulo, Brazil", *Foodborne Pathogens and Disease*, 2014.

Publication

<1 %

39

actavet.vfu.cz

Internet Source

<1 %

40

m.scirp.org

Internet Source

<1 %

41

qdoc.tips

Internet Source

<1 %

42

sinta.ristekbrin.go.id

Internet Source

<1 %

43

studentsrepo.um.edu.my

Internet Source

<1 %

44

Akoll, P, and WW Mwanja. "Fish health status, research and management in East Africa: past and present", African Journal of Aquatic Science, 2012.

Publication

<1 %

45

Beshada Haile Adanech, Kassa Getahun Temesgen. "Isolation and identification of Escherichia coli and Edwardsiella tarda from fish harvested for human consumption from Zeway Lake, Ethiopia", African Journal of Microbiology Research, 2018

Publication

<1 %

46

Chang-Ro Lee, Ill Cho, Byeong Jeong, Sang Lee. "Strategies to Minimize Antibiotic Resistance", International Journal of Environmental Research and Public Health, 2013

Publication

<1 %

47

Reem Y. Aljindan, Khaled R. Alkharsah.
"Pattern of increased antimicrobial resistance
of Salmonella isolates in the Eastern Province
of KSA", Journal of Taibah University Medical
Sciences, 2020

Publication

<1 %

48

Robert E. Levin. " The Use of Molecular
Methods for Detecting and Discriminating
Associated with Foods — A Review ", Food
Biotechnology, 2009

Publication

<1 %

49

Upadhyay, A., D.P. Karumathil, I. Upadhyaya,
V. Bhattaram, and K. Venkitanarayanan.
"Controlling Bacterial Antibiotic Resistance
Using Plant-Derived Antimicrobials", Antibiotic
Resistance, 2016.

Publication

<1 %

50

Zhang, Wen-Hui, Xiang-Yan Lin, Liang Xu, Xi-Xi
Gu, Ling Yang, Wan Li, Si-Qi Ren, Ya-Hong Liu,
Zhen-Ling Zeng, and Hong-Xia Jiang. "CTX-M-
27 Producing Salmonella enterica Serotypes
Typhimurium and Indiana Are Prevalent
among Food-Producing Animals in China",
Frontiers in Microbiology, 2016.

Publication

<1 %

51

irishvetjournal.biomedcentral.com

Internet Source

<1 %

52	publications.aston.ac.uk Internet Source	<1 %
53	vphcap.vet.cmu.ac.th Internet Source	<1 %
54	Akter, S, Z Ferdowshi, MN Islam, MAM Prodhan, and MYE Chowdhury. "Determination of Salmonella in egg shell and egg content in some selected areas of Bangladesh", International Journal of Natural Sciences, 2016. Publication	<1 %
55	Swathy J.S., Prabhakar Mishra, John Thomas, Amitava Mukherjee, Natarajan Chandrasekaran. "Antimicrobial potency of high-energy emulsified black pepper oil nanoemulsion against aquaculture pathogen", Aquaculture, 2018 Publication	<1 %
56	helda.helsinki.fi Internet Source	<1 %
57	nexusacademicpublishers.com Internet Source	<1 %
58	rjpbcs.com Internet Source	<1 %
59	www.seafdec.org.ph Internet Source	<1 %

60 D. Daffonchio, A. Cherif, S. Borin. <1 %
"Homoduplex and Heteroduplex
Polymorphisms of the Amplified Ribosomal
16S-23S Internal Transcribed Spacers
Describe Genetic Relationships in the "Bacillus
cereus Group"", Applied and Environmental
Microbiology, 2000
Publication

61 bmcevolbiol.biomedcentral.com <1 %
Internet Source

62 ehp.niehs.nih.gov <1 %
Internet Source

63 www.stir.ac.uk <1 %
Internet Source

64 www.zora.uzh.ch <1 %
Internet Source

65 ejournals.ph <1 %
Internet Source

66 www.samj.org.za <1 %
Internet Source

67 pesquisa.bvsalud.org <1 %
Internet Source

Exclude bibliography On