

Low Allelic Variation of Plasmodium falciparum msp-1 and msp-2 among Gold Miners in Central Kalimantan Province, Indonesia

by Heny Arwati

Submission date: 12-Mar-2023 02:19PM (UTC+0800)

Submission ID: 2035013620

File name: 52._Low_Allelic_Variation_of.pdf (420.7K)

Word count: 4640

Character count: 24022



1
Tehran University of Medical
Sciences Publication
<http://tums.ac.ir>

Iran J Parasitol

Open access Journal at
<http://ijpa.tums.ac.ir>



Iranian Society of Parasitology
<http://isp.tums.ac.ir>

Original Article

Low Allelic Variation of *Plasmodium falciparum msp-1* and *msp-2* among Gold Miners in Central Kalimantan Province, Indonesia

*Heny Arwati¹, Trylanti Lestarisa², Indria Augustina³, Etik Ainun Rohmah⁴, Sri Subekti⁵, Soedjajadi Keman⁶, Yoes Prijatna Dachlan¹

- 23
1. Department of Medical Parasitology, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia
2. Department of Public Health, Faculty of Medicine, Universitas Palangka Raya, Palangka Raya City, Indonesia
3. Department of Parasitology, Faculty of Medicine, Universitas Palangka Raya, Palangka Raya City, Indonesia
4. Entomology Study Group, Institute of Tropical Diseases, Universitas Airlangga, Surabaya, Indonesia
5. Department of Marine, Faculty of Marine and Fisheries, Universitas Airlangga, Surabaya, Indonesia
6. Department of Environmental Health, Faculty of Public Health, Universitas Airlangga, Surabaya, Indonesia

Received 16 Aug 2022
Accepted 25 Nov 2022

Abstract

Background: We aimed to find out the allelic variation of *Pfmsp-1* and *Pfmsp-2* among gold miners in Central Kalimantan Province, Indonesia using parasites' DNA isolated from archived RDT and GSBS

Methods: This study was done using the samples collected between 2017-2020 from health centers in Subdistrict of Mihing Raya, Danau Rawah, and Bukit Hindu as well as Kapuas District Health Laboratory in Central Kalimantan Province, Surabaya, Indonesia. Parasites DNA were isolated from RDT cartridges and GSBS of local and migrant gold miners. Species of *Plasmodium* were confirmed by single step PCR. The allelic variation of *Pfmsp-1* (K1, MAD20, RO33) and *Pfmsp-2* (3D7, FC27) were analyzed by nested PCR.

Results: *Pfmsp-1* gene was found in only two (22.22%) out of 9 local samples, and 3 (27.27%) out of 11 migrant samples were found positive for K1 (150 bp) as well as MAD 20 (190 bp) allelic families. *Pfmsp-2* gene were found in each one sample of 550 bp fragment in local (11.11%) and migrant samples (9.09%) for 3D7, and 2 samples of 300 bp fragments in local (22.22%) and 3 samples of 300 bp in migrant samples (27.27%). No difference in size and number of infections between both populations. The RO33 allelic family Alhamdulillah was not found in any sample.

Conclusion: Low allelic variation of *Pfmsp-1* and *Pfmsp-2* genes with monogenotype indicated the low intensity of malaria transmission among gold miners in the studied areas. Further, the transmission may occur locally in the mining sites.

Keywords:
Giemsa;
Blood smears;
Allelic variation;
Malaria;
Indonesia

***Correspondence**
Email:
heny-a@fk.unair.ac.id



Introduction

Malaria is a devastating disease that occurs mostly in tropical and subtropical countries around the world. Malaria is closely related to poverty and underdevelopment (1). Gold mining has played an important role in increasing national economic of mineral rich countries, but simultaneously can maintain the persistence of malaria in gold mining areas, especially illegal gold mining (2).

Malaria cases have been reported in gold mining areas in Indonesia (3, 4), French Guiana (5), Guyana, (6), and Colombia (7). In Indonesia, small-scale mining of the gold and coal, has increased since the onset of the economic crisis during about the year of 2000 (3). The new ponds ex illegal gold mining were left by the miners and became breeding places of *Anopheles* mosquito in West Kalimantan (8). Malaria patients were found in areas around illegal gold mines in South Kalimantan (9). The difficult access to the gold mines, which were located in the forests of the slopes of mountain contributed to the increased of malaria cases due to the increasing contact with *Anopheles* mosquito during their journey to and from gold mines (10). Gold miners in Central Kalimantan are five times more likely to be infected with malaria than non-miners (11). Malaria cases among gold miners in Kapuas and Gunung Mas Districts have been recorded by local health centers, however there was no formal publication.

Plasmodium falciparum merozoite surface protein-1 (*Pfmsp-1*) and *P. falciparum* merozoite surface protein-2 (*Pfmsp-2*) are the major protein that important for erythrocyte invasion (12), that can induce immune response in human (13), and are the promising candidate for an erythrocytic stage malaria vaccine (14). Three polymorphic allelic families in block two of *Pfmsp-1* gene has been identified as K1, MAD20, and RO33 (15). The *Pfmsp-2* has two allelic families in the central domain, and iden-

tified as 3D7 and FC27 (16). The genetic diversities of these two genes have been reported from different parts of the world (17, 18) to describe the population genetic structure of *P. falciparum* in the different malaria endemicity (19) as well as to distinguish treatment failure (20). Further, the genetic diversity of *P. falciparum* population is an important indicator of malaria transmission intensity in an area (21).

Parasite-based malaria diagnostic test either by microscopy examination or rapid diagnostic test (RDT) is recommended by the WHO for all patients suspected malaria before medication is provided (22). The used RDT and microscopy slides usually were then archived for years in the laboratory. Covid-19 pandemic has hampered malaria field survey such as active and passive case detection especially in collecting fresh blood from the patients for diagnosis, treatment, mapping as well as epidemiological and molecular studies. Alternatively, the archived RDT cartridges and microscopy slides of Giemsa-stained blood smears (GSBS) could be used as DNA sources for molecular study (23, 24). The RDT and GSBS have long been used to diagnose malaria among gold miners in Kapuas District.

Therefore, this study aimed to find out the allelic variation of *Pfmsp-1* and *Pfmsp-2* among gold miners in Kapuas District of Central Kalimantan Province, Indonesia using parasites' DNA isolated from archived RDT and GSBS.

Materials and Methods

Sample collection and ethics

This study was done during 2021 using the samples collected between 2017-2020 from health centers in Subdistrict of Mihing Raya, Danau Rawah, and Bukit Hindu as well as Kapuas District Health Laboratory in Central Kalimantan Province. The samples of positive *P. falciparum* and *P. vivax* those have been archived since the year of 2017 to 2020. The

collected RDT cartridges and microscopy slides have been used to diagnose malaria infection among gold miners who developed fever seeking medication to the health centers during that time. The origin of samples was further divided into local and migrant populations. Local population in the studied areas is called Dayak (25). Migrant population those came from various different ethnicities, from other cities, other provinces and even other islands who have lived in these districts for more than three months (9). The archived RDTs and GSBS microscopy slides were used as a source of DNA for molecular confirmation of *Plasmodium* species by single step PCR. Only DNA extracted from *P. falciparum* and mix infections were used for PCR-based genetic genotyping of *Pfmsp-1* and *Pfmsp-2* genes.

The research was done after receiving certificate of ethic for research from Health Research Ethics Committee, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia as stated on the certificate number 10/EC/KEPK/FKUA/2021 dated on January 18, 2021.

DNA extraction

The individual RDT cartridge was opened by using sterile scalpel and removed the nitrocellulose strip and stripped of any plastic covering the nitrocellulose (26). New scalpel was used for every single RDT cartridge. DNA extraction was done by using QIAamp DNA Blood Mini Kit (Qiagen, Hilden, Germany) following manufactures instruction.

The DNA from GSBS of microscopy slides was isolated by scratching off the smear and directly applied to the DNA isolation kit.

Species confirmation using single step PCR

31

The extracted DNA were then used as template for single step PCR amplification. Primers used for single step PCR based on the sequence of small sub unit ribosomal RNA (ssu-rRNA) of *P. falciparum* and *P. vivax*. Only three primers were used in this PCR. The first

primer amplifies sequence common to rRNA of both *P. falciparum* and *P. vivax*. Second primer is specific for *P. vivax* and third primer is specific for *P. falciparum*. First and second primers together produced a 266 bp product specific for *P. vivax*, whereas first and third primers produced a 346 bp product specific for *P. falciparum*. The sequences of primers and PCR condition applied for this PCR were as described (27, 28).

PCR amplification and allele detection

PCR-based detection of allelic variation in *Pfmsp-1* and *Pfmsp-2* was done according to Aubouy method including primers and PCR condition with slight modification (13). The PCR products of single step PCR were used as template for nested PCR. The PCR products were then electrophoresed on 2% agarose gel. Determination of allelic variation of *Pfmsp-1* was including K1, MAD20, and RO33, and *Pfmsp-2* was including 3D7 and FC27 allelic families. The allelic variation was detected by visualization of the gel under UV illuminator, the sizes of bands were confirmed based on the 100 bp DNA ladder (Invitrogen).

Results

Characteristic of samples

Based on the data obtained from the health center of Kapuas and Gunung Mas Districts, a total of 101 samples consisted of 55 (54.46%) from local population further consisted of 30 (29.70%) of RDT cartridge and 25 (24.75%) of microscopy slides of GSBS, and 46 (45.54%) from migrant population, consisted of 29 (28.71%) RDT cartridges and 17 (16.83%) microscopy slides. There was no data of the origin of the migrants, however the information from the health centers noted that migrant community came from outside both districts, outside the province, and even outside Kalimantan (Borneo) Island. All samples in this study were collected from gold miners who sought medication due to malaria

infection during 2017-2020. All miners were male aged between 15-65 years old. The result

of malaria diagnosis by RDT and microscopy examination are listed in Table 1.

Table 1: Malaria diagnosis by RDT and microscopy examination among gold miners originated from local and migrant population in Central Kalimantan Province during 2017-2020

	Number of sample (%)		
	LOCAL	MIGRANT	Total
<i>P. falciparum</i>	28(27.72)	23(22.77)	51(50.50)
<i>P. vivax</i>	27(26.73)	23(22.77)	50(49.50)
Total	55(54.45)	46(45.54)	101 (100)

Species identification by single step PCR

The purpose of this step was to confirm the species of *Plasmodium* based on DNA of parasites before further determination the allelic variation (Table 2). Twenty samples (19.80%) out of 101 samples were detected consisted of 7 samples (35%) of *P. falciparum* (346 bp), and

13 samples (65%) of mix infection of *P. falciparum* and *P. vivax* (346 bp and 226 bp) as seen in Fig. 1. The samples consisted of 9 (45%) local, and 11 (55%) migrant population origins. These 20 samples were used for nested PCR genotyping.

Table 2: RDT and microscopy diagnosis of malaria compared with those of single step PCR

Species	RDT and microscopy		Single-step PCR			
	Pf	Pv	Pf	Pv	Mix	Neg
Pf	51		6			45
Pv		50	1	3	13	33
Total		101			101	

Pf: *P. falciparum*; Pv: *P. vivax*; Mix: mix infection of Pf and Pv; Neg: negative




Fig. 1: Representative picture of single step PCR is showing single band of 346 bp for *P. falciparum* (a), 226 bp for *P. vivax* (b), and double band for mix infection of both species (c)

Allelic variation of *Pfmsp-1* and *Pfmsp-2*

Pfmsp-1 gene was found in only two (22.22%) out of 9 local samples, and 3 (27.27%) out of 11 migrant samples were found positive for K1 (150 bp) as well as MAD 20 (190 bp) allelic families. *Pfmsp-2* gene were found in each one sample of 550 bp fragment in local (11.11%) and migrant samples (9.09%) for 3D7, and 2 samples of 300 bp fragments in local (22.22%) and 3 samples

of 300 bp in migrant samples (27.27%). The RO33 allelic family Alhamdulillah was not found in any sample. The representative pictures of these allelic variation is shown in Fig. 2. Single band indicated a single clonal infection of *Plasmodium*. There was no different number of PCR fragments nor different size of the allelic family between indigenous and migrant population.




Fig. 2: Allelic variations of *Pmsp-1* and *Pfmsp-2* genes based on the DNA extracted from RDT cartridges and GSBS, are showing a 150 bp in K1, 190 bp in MAD20, and negative in RO33 of *Pfmsp-1* allelic family, a 550 bp of 3D7 and 300 bp of FC27 of *Pfmsp-2* allelic family

Discussion

19

Kalimantan (Borneo) island is one of the oldest artisanal and small-scale gold mining (ASGM) hotspot in Indonesia, where many people from outside the district even outside island areas came to work as illegal gold miners (4). The gold mining are often located in the forest or mountain slopes, where the miners have to walk along the areas where the contact with *Anopheles* mosquitoes were unavoidable without any kind of repellent to reach mining areas. This became an interesting field to study the genetic diversity of parasites among the malaria-infected gold miners. The results showed that the DNA could be extracted from RDT cartridges and GSBS samples archived during 2017-2020 at room temperature. Parasites were trapped in the nitro-cellulose of RDT for long time period at room temperature. The results of PCR proved that DNA of parasites in this condition remained

safe for molecular study. Further, the older RDT that were collected during 2014-2017 as a source of DNA have been reported for sequencing and studying the molecular marker of anti-malaria resistance in República da Guiné-Bissau, West Africa (29). The RDT stored at 19°-25°C were also good tool for malaria diagnosis and also useful for molecular epidemiology and *Plasmodium* genetic analysis (30).

The differences in host geographic origins with different immune status might cause the differences in the host immune response to parasites which ultimately create diversity in parasite genetics (30), which will complicate the identification of *P. vivax* and *P. falciparum* (31, 32). The antigenic diversity of *P. falciparum* and *P. vivax* populations in an area plays an important role in the formation of natural immunity against malaria. The rapid and severe destruction of erythrocytes corresponds to the number of parasites found in the blood

by microscopic examination of the blood smear. The parasitemia is also influenced by the characteristics of the patient, such as age, sex, and race. The factors such as meiotic recombination events during the fertilization process in *Anopheles* mosquito might cause genetic variation of malaria parasite strains (18, 33).

The allelic variation of *Pfmsp-1* and *Pfmsp-2* based on the DNA isolated from archived RDT and GSBS, proved the DNA preparation by this method could be used in the genetic study. The results showed that amplified product from the *Pfmsp-1*, *Pfmsp-2* genes were not varied in size, and there were low allelic variations between local and migrant gold miners. Besides, regarding the small number of samples, this condition might reflect the real condition of the circulating genes among the gold miners in the studied areas. Apparently, there was no new allelic variation were introduced by migrant gold miners. Malaria infection among migrant gold miners may occur in the location of gold mines. Therefore, the parasites that infect them probably are local parasites, and the genetic recombination in *Anopheles* mosquito occurred between local parasites, since the allelic variations were not different with those of local gold miners. This condition was also observed among *Pvmsp-1* and *Pvmsp-2* of *P. vivax*-infected patients in the same location (9).

Central Kalimantan Province is a malaria hypo endemic area, where the annual parasite incidence (API) is low in 2018 reached 0.24 (34). Only three districts of this province remained uncertified for malaria free (35). Low allelic variation of *Pfmsp-1* and *Pfmsp-2* genes with monogenotype indicated the low intensity of malaria transmission in the studied areas, since high rate of multigenotype infection indicated the high intensity of transmission(21). Genetic diversity of *Pfmsp-1* and *Pfmsp-2* in several districts in Indonesia are varies. Low genetic allelic variation found in this study is lower than that found in Lampung Province in

Sumatra Island, where six variations in length range from 200 to 1000 bp of *Pfmsp-1* gene (36). Genetic diversity of *Pfmsp-1* in malaria hyper endemic area of Papua Province showed that MAD20 allelic family was predominant (20.8%), RO33 was 8.2%, and K1 was only 4.2% (37). Interestingly, the RO33 allelic family was not found in Central Kalimantan Province. This result is different with that found in Pahang, Malaysia, where RO33 allelic family was predominant as monomorphic (38), in contrast with that found in Grande Comoro Island where RO33 allelic family were poorly represented (2.4-4.0%) (39), while in Myanmar-China border was higher (12.56%) (17). In Gabon, when the DNA was isolated from RDTs showed that K1 (93.9%) was found higher than MAD20 (73.3%) (40). The allelic prevalence of FC27 in *Pfmsp-2* gene (15,38%) in this current study was higher than that of 3D7 allelic family (7,69%), similar to that found in Pakistan (41) and China-Myanmar border region (17). On the other hand in India, FC27 (12-23.3%) was found lower than 3D7 (35.7-56%) (18).

The various genetic polymorphism reported from various countries around the world proved that different population with different geographic conditions has a wide variety of different genetic variations.

Conclusion

9
The low allelic diversity of *Pfmsp-1* and *Pfmsp-2* genes among gold miners in studied areas indicated the low malaria transmission intensity. Further, the transmission may only occur locally between local and migrants' populations in the gold mining sites. Therefore, gold mining plays an important role in malaria transmission, and plays a barrier to malaria elimination in these areas. The specific malaria control program must be implemented in these areas to control the transmission and to avoid the spread of malaria to other regions.

Acknowledgments

⁸ This study was supported by a grant from the Directorate General of Higher Education, Ministry of Education, Culture, Research and Technology of the Republic of Indonesia decree number 194/SP2H/LT/DRPM/2021. We would like to thank the heads and staffs of the following health centers: Puskesmas Miring Raya, Puskesmas Danau Rawah, and Puskesmas Bukit Hindu as well as the Kapuas District Health Laboratory for providing the samples.

Conflict of Interest

The authors declare that there is no conflict of interest.

References

1. Tusting LS, Rek J, Arinaitwe E, et al. Why is malaria associated with poverty? Findings from a cohort study in rural Uganda. Infect Dis Poverty. 2016; 5 (78).
2. Castellanos A, Chaparro-Narváez P, Morales-Plaza CD, et al. Malaria in gold-mining areas in Colombia. Mem Inst Oswaldo Cruz. 2016;111(1):59–66.
3. McMahon G, Subdibjo ER, Aden J, Bouzaher A, Dore G, Kunanayagam R. Mining and the environment in Indonesia: long-term trends and repercussions of the Asian economic crisis. EEASES Discuss Pap Ser. 2000; 21438: 1–46.
4. Kanakath P. Illegal gold mining operations in Kalimantan threaten lives and the environment. Indonesia Expat. 2015. Cited January 15, 2022 from <https://indonesiaexpat.id/business-property/illegal-gold-mining-operations-in-kalimantan-threaten-lives-and-the-environment/>
5. Douine M, Sanna A, Hiwat H, et al. Investigation of a possible malaria epidemic in an illegal gold mine in French Guiana: an original approach in the remote Amazonian Forest. Malar J. 2019; 18 (1):91.
6. Olapeju B, Adams C, Simpson J, et al. Malaria Care-Seeking and Treatment Ideation among Gold Miners in Guyana. PLoS One. 2020; 15(12): e0244454.
7. Endeshaw T, Gebre T, Ngondi J, et al. Evaluation of light microscopy and rapid diagnostic test for the detection of malaria under operational field conditions: a household survey in Ethiopia. Malar J. 2008, 7118. 2008; 7(118).
8. Bariyah K, Utomo B, Sulistyawati S, et al. Different Types of *Anopheles* Breeding Place in Low and High Malaria Case Areas. J Kesehat Masy. 2018;14(2):178–85.
9. Lestarisa T, Arwati H, Dachlan Y, Keman S, Safruddin D. The use of archived Giemsa-stained blood smears and RDT for PCR-based genotyping of *Plasmodium vivax*: merozoite surface protein-1 in Central Kalimantan Province, Indonesia. Afr J Infect Dis. 2022; 16(1):13–20.
10. Indriyati I, Rosanji A, Juhairiyah J, Yuana WT, Haryati E. Habitat Perkembangbiakan Spesifik *Anopheles* sp. di Tambang Emas Kura-Kura Banian (Perubahan Perilaku *Anopheles* sp.). Balaba J Litbang Pengendali Penyakit Ber-sumber Binatang Banjarnegara. 2018; (October):121–34.
11. Rahayu N, Suryatinah Y, Kusumaningtyas H. Discovery of Malaria Cases on Forest Workers in Public Health Center, Teluk Kepayang, Tanah Bumbu, South Kalimantan. BIO Web Conf. 2020; 20(01008).
12. Holder AA, Blackman MJ. What is the Function of MSP-I on the Malaria Merozoite? Parasitol Today. 1994; 10 (5):182–4.
13. Aubouy A, Migot-Nabias F, Deloron P. Polymorphism in Two Merozoite Surface Proteins of *Plasmodium falciparum* Isolates from Gabon. Malar J. 2003; 2(12).
14. Woehlbier U, Epp C, Kauth CW, et al. Analysis of Antibodies Directed against Merozoite Surface Protein 1 of the Human Malaria Parasite *Plasmodium falciparum*. Infect Immun. 2006; 74(2):1313–1322.
15. Contamin H, Fandeur T, Rogier C, et al. Different Genetic Characteristics of *Plasmodium falciparum* Isolates Collected during Successive Clinical Malaria Episodes in Senegalese Children. Am J Trop Med Hyg. 1996; 54(6):632–43.
16. Lê HG, Kang JM, Jun H, et al. Changing pattern of the genetic diversities of *Plasmodium falciparum* merozoite surface protein-1 and mero-

- zoite surface protein-2 in Myanmar isolates. Malar J. 2019; 18(1): 241.
17. Zhang C, Zhou H, Liu Q, Yang YM. Genetic polymorphism of merozoite surface proteins 1 and 2 of *Plasmodium falciparum* in the China–Myanmar border region. Malar J. 2019; 18:367.
18. Hussain HH, Sohail M, Kumar R, Branch OH, Adak T, Raziuddin M. Genetic diversity in merozoite surface protein-1 and 2 among *Plasmodium falciparum* isolates from malarious districts of tribal dominant state of Jharkhand, India. Ann Trop Med Parasitol. 2011; 105(8):579–92.
19. Mobegi VA, Loua KM, Ahoudi AD, et al. Population genetic structure of *Plasmodium falciparum* across a region of diverse endemicity in West Africa. Malar J. 2012; 11:223.
20. Collins WJ, Greenhouse B, Rosenthal PJ, Dorsey G. The Use of Genotyping in Antimalarial Clinical Trials: A Systematic Review of Published Studies from 1995–2005. Malar J. 2006; 5:122.
21. Soe T, Wu Y, Tun M, et al. Genetic diversity of *Plasmodium falciparum* populations in southeast and western Myanmar. Parasit Vectors. 2017; 10(1):322.
22. WHO. Diagnostic testing. 1990. Cited on February 20, 2020 from: <https://www.who.int/teams/global-malaria-programme/case-management/diagnosis>
23. Shavey CA, Morado J. DNA extraction from archived Giemsa-stained blood smears using polymerase chain reaction to detect host and parasitic DNA. J Histotechnol. 2012; 35(3).
24. Nguyen T, Mombo, BN Lalremruata A, et al. DNA recovery from archived RDTs for genetic characterization of *Plasmodium falciparum* in a routine setting in Lambaréne, Gabon. Malar J. 2019; 18 (1):336.
25. Sada C, Alas Y, Anshari M. Indigenous people of Borneo (Dayak): Development, social cultural perspective and its challenges. Cogent Arts & Humanities., 2019; 6(1):1665936.
26. WWARN. Preparation of Rapid Diagnostic Tests (RDTs) for DNA extraction. 11. 2014; 1–7. Cited on April 20, 2020 from: http://www.wwarn.org/sites/default/files/MOL06_RDTsForDNAExtraction.pdf
27. Patsoula E, Spanakos G, Sofianatou D, Parara M, Vakalis NC. A single-step, PCR-based method for the detection and differentiation of *Plasmodium vivax* and *P. falciparum*. Ann Trop Med Parasitol. 2003; 97(1):15–21.
28. Arwati H, Yotoprano S, Rohmah EA, Syafruddin D. Submicroscopic malaria cases play role in local transmission in Trenggalek District, East Java Province, Indonesia. Malar J. 2018; 17(1):2.
29. Nag S, Ursing J, Rodrigues A, et al. Proof of concept: used malaria rapid diagnostic tests applied for parallel sequencing for surveillance of molecular markers of anti-malarial resistance in Bissau, Guinea-Bissau during 2014–2017. Malar J. 2019; 18 (1):252.
30. Su X, Zhang C, Joy D. Host-Malaria Parasite Interactions and Impacts on Mutual Evolution. Front Cell Infect Microbiol. 2020; 10:587933.
31. Karunaweera N, Ferreira M, Munasinghe A, et al. Extensive microsatellite diversity in the human malaria parasite *Plasmodium vivax*. Gene. 2008; 410(1):105–112.
32. Imwong M, Nair S, Pukrittayakamee S, et al. Contrasting genetic structure in *Plasmodium vivax* populations from Asia and South America. Int J Parasitol. 2007; 37:1013–102.
33. Lê HG, Thái TL, Kang JM, et al. Genetic polymorphism of merozoite surface protein-1 and merozoite surface protein-2 in *Plasmodium falciparum* field isolates from Myanmar. Malar J. 2010; 9(1):131.
34. Sugihantono A. Bebas malaria prestasi bangsa. Media Indonesia. 2019. Cited on July 15, 2020 from: <https://mediaindonesia.com/read/detail/2355/64-bebas-malaria-prestasi-bangsa>
35. Trisnawati S. Tiga Kabupaten di Kalteng Belum Bebas Malaria. 2020. Cited on November 20, 2021 from: <https://rri.co.id/palangkaraya/tanggap-bencana/933034/tiga-kabupaten-di-kalteng-belum-bebas-malaria>
36. Triajayanti A, Utami N, Kurniawan B, Suwandi J. Identification of *Plasmodium falciparum* Merozoite Surface Protein-1 (Pfmsp-1) Gene from malaria patients In Hanura Area, Lampung, Indonesia. Bioscientia Medicina. 2019; 3(4):1–9.
37. Tanjung R, Sarungu Y, Imbiri M, et al. Genetic Diversity in Merozoite Surface Protein-1 of *Plasmodium falciparum* in Highlands: Wamena Papua Indonesia. Am J Infect Dis. 2018; 14(4):106–9.

38. Atroosh W, Al-Mekhlafi H, Mahdy M, Saif-Ali R, Al-Mekhlafi A, Surin J. Genetic diversity of *Plasmodium falciparum* isolates from Pahang, Malaysia based on *MSP-1* and *MSP-2* genes. *Parasites Vectors.* 2011; 4:233.
39. Mze N, Bogueau H, Diedhiou C, et al. Genetic diversity of *Plasmodium falciparum* in Grande Comore Island. *Malar J.* 2020; 19 (1):320.
40. Mawili-Mboumba D, Mbuyi M, M'bondoukwe N, Bouyou-Akoté M. *Plasmodium falciparum* Allelic Diversity: A Comparison of DNA Extraction from Isolates Collected on Rapid Diagnostic Tests (RDTs) and Filter Paper. *Iran J Parasitol.* 2021; 16(4):555–9.
41. Khan S, Ali R, Khan S, et al. Genetic Diversity of Polymorphic Marker Merozoite Surface Protein 1 (Msp-1) and 2 (Msp-2) Genes of *Plasmodium falciparum* isolates from Malaria Endemic Region of Pakistan. *Front Genet.* 2021;12: 751552.

Low Allelic Variation of Plasmodium falciparum msp-1 and msp-2 among Gold Miners in Central Kalimantan Province, Indonesia

ORIGINALITY REPORT



PRIMARY SOURCES

- | | | |
|---|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----|
| 1 | Submitted to Georgia Southern University
Student Paper | 2% |
| 2 | Jazia Sriti, Mouna Boulares, Youkabed Zarroug, Rym Essid, Nadia Fares, Selim Jallouli, Saida Djebbi, Nidhal Salem. "Potential application of essential oil of Lavandulastoechas in poultry meat during refrigerated storage", Journal of Food Safety and Hygiene, 2021
Publication | 1% |
| 3 | Soudabeh HEIDARI, Javad GHARECHAHI, Mehdi MOHEBALI, Behnaz AKHOUNDI, Shahab MIRSHAHVALADI, Bahareh AZARIAN, Homa HAJJARAN. "Western Blot Analysis of Leishmania infantum Antigens in Sera of Patients with Visceral Leishmaniasis", Iranian Journal of Parasitology, 2019
Publication | 1% |

- 4 Retno Desy Tri Lestari, Sri Subekti, Gunanti Mahasri. " The occurrence of Trematodes Infection in Cantang Grouper at Lamong Bay, Surabaya, Indonesia ", IOP Conference Series: Earth and Environmental Science, 2020
Publication
-
- 5 J Taylor. "Toward quantitative genetic analysis of host and parasite traits in the manifestations of Plasmodium falciparum malaria", Current Opinion in Genetics & Development, 2000
Publication
-
- 6 pubmed.ncbi.nlm.nih.gov
Internet Source
-
- 7 publikationen.uni-tuebingen.de
Internet Source
-
- 8 apip AMRULLAH, Obie FAROBIE, Lusi Ernawati. "Slow-pyrolysis of brown macroalgae Padina sp.: Product characterization and degradation kinetic mechanism", Research Square Platform LLC, 2023
Publication
-
- 9 Jiraporn Kuesap, Wanna Chaijaroenkul, Kanchanok Ketprathum, Puntanat Tattiayapong, Kesara Na-Bangchang. "Evolution of Genetic Polymorphisms of Plasmodium falciparum Merozoite Surface

Protein (PfMSP) in Thailand", The Korean Journal of Parasitology, 2014

Publication

10	onlinelibrary.wiley.com	1 %
11	www.researchsquare.com	1 %
12	www.frontiersin.org	1 %
13	www.scielo.br	1 %
14	Joseph Kirui, Josephine Malinga, Edna Sang, George Ambani et al. "Bottlenecks to Intervention Scale Up: Supply and Demand Side Perspectives From a Large, Community-based Trial of Malaria Testing", Research Square Platform LLC, 2021	<1 %
15	journal.um-surabaya.ac.id	<1 %
16	link.springer.com	<1 %
17	Submitted to St George's Hospital Medical School	<1 %
	bmcinfectdis.biomedcentral.com	

18

<1 %

19

Submitted to Universiti Teknologi MARA

<1 %

Student Paper

20

Aliehsan Heidari. "Genetic diversity in merozoite surface protein (MSP)-1 and MSP-2 genes of Plasmodium falciparum in a major endemic region of Iran", The Korean Journal of Parasitology, 2007

<1 %

Publication

21

Makhtar Niang, Laty G. Thiam, Cheikh Loucoubar, Abdourahmane Sow et al. "Spatio-temporal analysis of the genetic diversity and complexity of Plasmodium falciparum infections in Kedougou, southeastern Senegal", Parasites & Vectors, 2017

<1 %

Publication

22

Eduardo Massad, Gabriel Zorello Laporta, Jan Evelyn Conn, Leonardo Suveges Chaves et al. "The risk of malaria infection for travelers visiting the Brazilian Amazonian region: A mathematical modeling approach", Travel Medicine and Infectious Disease, 2020

<1 %

Publication

23

Heny Arwati, Ramadhani R. Bahalwan, Windya T. Hapsari, Kartika A. Wardhani, Kholida N. Aini, Putu I. B. Apsari, Puspa Wardhani.

<1 %

"Suppressive effect of goat bile in Plasmodium berghei ANKA infection in mice", Veterinary World, 2021

Publication

-
- 24 Mohd Ridzuan Mohd Abd Razak, Umi Rubiah Sastu, Nor Azrina Norahmad, Abass Abdul-Karim et al. "Genetic Diversity of Plasmodium falciparum Populations in Malaria Declining Areas of Sabah, East Malaysia", PLOS ONE, 2016 <1 %
- Publication
-
- 25 S. J. Patgiri, K. Sarma, N. Sarmah, N. Bhattacharyya et al. "Characterization of drug resistance and genetic diversity of Plasmodium falciparum parasites from Tripura, Northeast India", Scientific Reports, 2019 <1 %
- Publication
-
- 26 bmcmedicine.biomedcentral.com <1 %
- Internet Source
-
- 27 pure.rug.nl <1 %
- Internet Source
-
- 28 Amirah Amir, Fei-Wen Cheong, Jeremy R. De Silva, Yee-Ling Lau. "Diagnostic tools in childhood malaria", Parasites & Vectors, 2018 <1 %
- Publication
-

- 29 Adela Ida Jiram, Choo Huck Ooi, José Miguel Rubio, Shamilah Hisam et al. "Evidence of asymptomatic submicroscopic malaria in low transmission areas in Belaga district, Kapit division, Sarawak, Malaysia", Malaria Journal, 2019 <1 %
- Publication
-
- 30 Chenet, Stella M, Kristan A Schneider, Leopoldo Villegas, and Ananias A Escalante. "Local population structure of Plasmodium: impact on malaria control and elimination", Malaria Journal, 2012. <1 %
- Publication
-
- 31 G. F. Zhang. "Rapid diagnosis of the invasive species, *Frankliniella occidentalis* (Pergande): a species-specific COI marker : Molecular identification of *Frankliniella occidentalis*", Journal of Applied Entomology, 08/2011 <1 %
- Publication
-

Exclude quotes Off
Exclude bibliography On

Exclude matches Off

Low Allelic Variation of Plasmodium falciparum msp-1 and msp-2 among Gold Miners in Central Kalimantan Province, Indonesia

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9
