

# Antimalarial Activity of Mahagony Seed Ethanolic Extract in BALB/c Mice Infected with Plasmodium berghei ANKA and The Correlation of Parasitemia and Plasma Level of IFN- $\gamma$

*by Ni Ketut Yuliana Sari*

---

**Submission date:** 12-Mar-2023 02:21PM (UTC+0800)

**Submission ID:** 2035014393

**File name:** 53.\_Antimalarial\_Activity\_of\_Mahagony\_Seed\_Ethanolic.pdf (334.91K)

**Word count:** 4402

**Character count:** 23180

ORIGINAL ARTICLE

6  
**Antimalarial Activity of Mahogany Seed Ethanolic Extract in BALB/c Mice Infected with *Plasmodium berghei* ANKA and The Correlation of Parasitemia and Plasma Level of IFN- $\gamma$**

\*Ni Ketut Yuliana Sari<sup>1,2</sup>, Heny Arwati<sup>3,4</sup> and Indah Setyawati Tantular<sup>3,4,5</sup>

<sup>1</sup> Graduate student of Immunology, Department of Immunology, Postgraduate School, Universitas Airlangga, Surabaya 60285, Indonesia

<sup>2</sup> Department of Medical Laboratory Technology, Poltekkes Kemenkes Kupang, Kupang 85111, Indonesia

<sup>3</sup> Department of Medical Parasitology, Faculty of Medicine, Universitas Airlangga, Surabaya 60131, Indonesia

<sup>4</sup> Postgraduate School of Immunology, Universitas Airlangga, Surabaya 60285 Indonesia

<sup>5</sup> Institute of Tropical Diseases, Universitas Airlangga, Surabaya 60115, Indonesia

ABSTRACT

**Introduction:** The genus of *Plasmodium* is a parasite that can cause malaria. Interferon gamma (IFN- $\gamma$ ) is an inflammatory cytokine, that can control *Plasmodium* in the blood. World Health Organization (WHO) recommends Artemisinin-based Combination Therapies (ACTs) to treat malaria. However, some ethnic communities in Indonesia still consume herbal, one of which is mahogany seeds. Its effect on the immune system has not been much explained. **Materials and Methods:** Twenty-five male BALB/c mice were infected with *Plasmodium berghei* ANKA and divided into 5 groups. Groups I-III were treated with mahogany seeds ethanolic extract (MSEE) at the concentrations of 100%, 50% and 25% respectively. Group IV was a negative control (NC) which was treated with 0.5% sodium carboxymethyl cellulose (CMCNa). Group V was a positive control (PC) which was treated with dihydroartemisinin-piperazine (DHP) at 187.2 mg/kgBW. Treatments were given for four consecutive days. Parasitemia was observed every day from the first day to the fourth day post infection. Plasma level of IFN- $\gamma$  were measured by Sandwich ELISA. **Results:** Two way ANOVA analysis showed a significant difference in parasitemia based on the treatment day ( $p=0.000$ ) and the treatment group ( $p=0.000$ ) in infected mice. Treatment of MSEE showed a significant effect on plasma level of IFN- $\gamma$  ( $p=0.032$ ) in infected mice. Spearman's rho bivariate correlation test showed a significant positive correlation between parasitemia on the fourth day post treatment and plasma level of IFN- $\gamma$  ( $p=0.006$ ;  $r=0.589$ ). **Conclusion:** Antimalarial activity of MSEE was shown by the decreased of parasitemia along with the decreased of IFN- $\gamma$  level on the day 4 post infection in a concentration dependent manner.

**Keywords:** Malaria, mahogany seed, *Plasmodium berghei* ANKA, parasitemia, IFN- $\gamma$

Corresponding Author:

Ni Ketut Yuliana Sari, S.ST., M.Si., M.Imun.  
Email: niketutyuliana@poltekkeskupang.ac.id  
Tel: +6281246396440

*vivax*, *Plasmodium malariae*, *Plasmodium ovale*, and *Plasmodium knowlesi* are five *Plasmodium* species possess the ability to infect humans (3).

52  
INTRODUCTION

Malaria is a serious disease and remains the leading cause of death worldwide. Malaria is caused by a parasite of the genus *Plasmodium*, which is transmitted to humans through the bite of an infected female *Anopheles* mosquito (1). It was estimated that there were 241 million cases of malaria worldwide in 2020 in 85 malaria endemic countries, with 627.000 deaths (2). *Plasmodium falciparum*, *Plasmodium*

50  
*Plasmodium* infection induces the production of IFN- $\gamma$  from various innate and adaptive cells at various stages of the life cycle. IFN- $\gamma$  produced by CD4+ T cells optimally activates CD8+ T cells, B cells and macrophages. IFN- $\gamma$  affects isotype transfer in B cells which leads to the production of cytophilic antibodies capable of binding free parasites and blocking red blood cell invasion, mediating parasite clearance through opsonization, and binding to the surface of infected red blood cells by promoting antibody dependent phagocytosis. IFN- $\gamma$  production from CD4+ T cells also optimally activates

macrophages to phagocytose infected and parasite-free red blood cells (4).

The first-line treatment for malaria, currently recommended by WHO, is Artemisinin-based Combination Therapies (ACTs) (3). However, some ethnic communities in Indonesia still consume herbal plants that already exist and are obtained from generation to generation, one of which is mahogany seeds (*Swietenia mahagoni* (L.) Jacq.) (5). These plants have been reported to have antimalarial activity both *in vitro* and *in vivo* (6,7), but its effect on the immune system has not been much explained. This study aims to analyze the antimalarial activity of mahogany seeds ethanolic extract (MSEE) in BALB/c mice infected with *Plasmodium berghei* ANKA and the correlation between parasitemia and plasma level of IFN- $\gamma$ .

## MATERIALS AND METHODS

### Ethical clearance and research design

The proposal of this research has been reviewed by Ethical Committee of Universitas Airlangga Faculty of Dental Medicine as described on certificate number 779/HRECC.FODM/XII/2019. This research is a true experimental research with a post test only control group design. The treatment was given using the standard test method for curative activity (8).

### Experimental animals

The test animals in this study was BALB/c strain male mice aged 2-3 months, with a weight of  $\pm 20$ -30g, which were maintained and developed in the Animal Laboratory of the Department of Biochemistry, Airlangga University, Surabaya. The test animals were acclimatized for 7 days at a temperature of 25 °C with an exchange of dark and light every 12 hours and were given standard feed *ad libitum*. The mice were sacrificed after anesthetized by intraperitoneal injection of 87 mg/kg bodyweight ketamine. The dead animals were then burned in the incinerator.

### Plant material and MSEE preparation

The mahogany seeds used in this study were seeds from the mahogany plant species *Swietenia mahagoni* (L.) Jacq. They were obtained, determined and tested for phytochemical content at the Integrated Service Unit [Unit Pelayanan Terpadu, UPT] of the Herbal Materia Medika Laboratory, Batu City, East Java Province. *EEMS* was obtained from the extraction of mahogany seed powder by maceration technique using 96% ethanol. MSEE was made into three concentrations, 100%, 50% and 25% respectively. The 100% MSEE concentration was originally from maceration without is the result of maceration without dilution (crude extract). The extract was diluted using dilution using 0.5% CMCNa solution. MSEE concentration of 50% (v/v) was prepared in a ratio of 1:1 (3 mL crude extract : 3 mL CMCNa 0.5%) and MSEE concentration of 25% (v/v) was prepared in a ratio

of 1:3 (1.5 mL crude extract : 4.5mL CMCNa 0.5%). The results of the MSEE phytochemical test showed the presence of the flavonoids, tannins and triterpenoids. Infection of mice

Frozen isolates of *P. berghei* ANKA were thawed at 37 °C, then inoculated in 0.2 mL blood from donor mice intraperitoneally. Parasitemia was observed daily on Giemsa-stained thin smear. When parasitemia reached 10-20%, mice were sacrificed and blood were collected by cardiac puncture to infect the test mice. Twenty-five male BALB/c mice were infected with 0.2 mL of blood containing  $1 \times 10^6$  of infected red blood cells from donor mice. Test mice were then divided into 5 groups. Group I was treated with 100% MSEE, Group II was with 50% MSEE, Group III was with 25% MSEE, Group IV was negative control (NC) given 0.5% CMCNa and Group V was positive control (PC) treated with antimalarial drug of dihydroartemisinin-piperquin (DHP) at 187.2 mg/kgBW. The treatments were given when parasitemia reached  $\pm 1\%$ . The treatment was given for four consecutive days. Parasitemia was observed as described above. On the fourth day post treatment, the test mice were sacrificed to collect blood by cardiac puncture prior to the measurement of IFN- $\gamma$  level (10).

48

### Measurement of IFN- $\gamma$ level

Plasma level of IFN- $\gamma$  were measured on using the ELISA (Enzyme-Linked Immunosorbent Assay) Kit (The BioLegend, Singapore).

### Statistical analyses

The differences in mean of parasitemia were analyzed using two way analysis of variance (ANOVA), followed by Post Hoc (Tukey) analysis. The difference in mean of IFN- $\gamma$  levels were analyzed using Kruskal Wallis, followed by Mann-Whitney U analysis. The correlation between parasitemia and IFN- $\gamma$  level were analyzed using the Spearman correlation analysis. Values were statistically significant at  $p < 0.05$ .

16

## RESULTS

### Antimalarial activity of MSEE

The mean parasitemia of each treatment group is shown in Figure. The mean parasitemia in NC group has increased normally since the first day to the fourth day post treatment. Parasitemia in the PC group increased on the first day then continued to decrease until the fourth day post treatment. Parasitemia in the all group treated with MSEE on day four post treatment were lower than NC group, but higher than the PC group. This result indicated the antimalarial activity of MSEE.

Two Way ANOVA analysis to find out the the differences of parasitemia between day post treatment and between treatment groups resulted in significant differences, then was continued with the Post Hoc (Tukey) test. A summary of the results of the Post Hoc (Tukey) test is




Fig. 1: The mean of parasitemia from day 0 to day 4 post treatment in BALB/c mice infected with *P. berghei* ANKA which were given MSEE and the control group.

presented in Table I and II.

Based on the category of the treatment day, Post Hoc (Tukey) test showed a significance difference mean of parasitemia at day 1 and 4, day 2 and 4, day 3 and 4 post treatment and between treatment groups 100% MSEE, 50% MSEE and 25% MSEE with NC. Significant differences also occurred between the 50% MSEE treatment group with PC and between the NC and PC

Table I: Result of Post Hoc (Tukey) analysis of parasitemia based on treatment day category in BALB/c mice infected with *P. berghei* ANKA which were given MSEE and the control group

Treatment day	p-value	Information
Day 1 and day 2 post treatment	0.975	Not significant
Day 1 and day 3 post treatment	0.579	Not significant
Day 1 and day 4 post treatment	0.000*	Significant
Day 2 and day 3 post treatment	0.825	Not significant
Day 2 and day 4 post treatment	0.000*	Significant
Day 3 and day 4 post treatment	0.000*	Significant

\* p-value less than 0.05 was considered significant.

Table II: Result of Post Hoc (Tukey) analysis of parasitemia based on treatment group category in BALB/c mice infected with *P. berghei* ANKA which were given MSEE and the control group

Groups	p-value	Information
NC and PC	0.000*	Significant
NC and 100% MSEE	0.000*	Significant
NC and 50% MSEE	0.010*	Significant
NC and 25% MSEE	0.002*	Significant
PC and 100% MSEE	0.938	Not significant
PC and 50% MSEE	0.030*	Significant
PC and 25% MSEE	0.103	Not significant
100% MSEE and 50% MSEE	0.178	Not significant
100% MSEE and 25% MSEE	0.420	Not significant
50% MSEE and 25% MSEE	0.986	Not significant

\* p-value less than 0.05 was considered significant.

group, but not between the 100%, 50% and 25% MSEE concentrations.

Antimalarial activity of MSEE was also shown by the percentage of parasites' growth and its inhibition, where both percentages rising in a concentration-dependent manner (Table III). They were lower than those of PC group. The inhibition of parasites' growth by DHP in PC group was very high (137%) as parasitemia reached 0% at the end of experiment (Figure 1). Probit analysis based on the percentage of inhibition of MSEE resulted in an ED50 was 69.67%.

Table III: Percent growth and inhibition of parasites in BALB/c mice infected with *P. berghei* ANKA which were given MSEE and the control group

Groups	% of Growth	% of Inhibition
100% MSEE	1.21	62
50% MSEE	1.13	65
25% MSEE	0.18	75
NC	3.19	0
PC	-1.18	137

#### Plasma level of IFN- $\gamma$

The *P. berghei* ANKA-infected group which treated with 100% MSEE showed the highest mean of plasma level of IFN- $\gamma$  than other groups, while the lowest plasma level of IFN- $\gamma$  was shown by PC group. The plasma level of IFN- $\gamma$  in MSEE-treated groups increased along with the increased of MSEE concentration. The Kruskal Wallis analysis showed the differences in the mean of plasma levels of IFN- $\gamma$  among groups of mice treated with 100%, 50%, 25% MSEE, NC and PC treatment groups ( $p=0.032$ ). Mann-Whitney U analysis resulted in a significant difference in plasma level of IFN- $\gamma$  in all groups when compared with those in PC group ( $p=0.021$ ) (Table IV). However, the comparison among MSEE-treated groups and between MSEE-treated groups with NC were insignificantly different ( $p>0.05$ ). A summary of the mean of plasma levels and results of the Mann-Whitney U analysis is presented in Table IV. Bivariate correlation analysis was performed to determine the relationship between parasitemia on the fourth day post treatment with IFN- $\gamma$  levels. The results of the Spearman's rho correlation test showed a significant positive relationship between parasitemia on the fourth day post treatment and IFN- $\gamma$  levels (correlation

Table IV: A summary of the mean of plasma levels of IFN- $\gamma$  and their significant differences between MSEE-treated groups and PC group analyzed by Mann Whitney U test

Groups	Mean	SD	p*
100% MSEE	473.53	±550.82	0.021
50% MSEE	195.41	±42.27	0.021
25% MSEE	158.24	±89.08	0.021
NC	293.88	±154.51	0.021
PC	26.76	±15.27	

\*Significantly different when compared with PC.

coefficient ( $r$ ) = 0.589 and  $p$  = 0.006).

## DISCUSSION

This current study proved the antimalarial activity of MSEE as shown by the decreased of parasitemia and IFN- $\gamma$  level at the end of the experiment when compared with those in mice without any MSEE treatment (NC group), although the percentages inhibition of the parasites' growth were lower compared with that of PC group.

The antimalarial activity of MSEE in this study was probably caused by the bioactive compounds contained in MSEE. Based on the phytochemical test, the MSEE used in this study was positive for flavonoids, tannins and triterpenoids. Flavonoids inhibit the biosynthesis of fatty acids in the biochemistry of parasites and inhibit the entry of myoinositol into infected erythrocytes during the erythrocytic phase. These compound is required by *Plasmodium* for the development of the parasite stage during the erythrocytic phase (11,12). Flavonoids and their derivatives play a role in blocking the formation of haemozoin by forming free heme complexes which can be toxic to parasites so that parasites die (13).

Tannins can inhibit protease activity (14). Proteases are a group of enzymes that play an important role in the growth and invasion of *Plasmodium* parasites (15). In the erythrocytic stage, *Plasmodium* uses host hemoglobin as a food source for the development of the parasite stage (16). Aspartate and cysteine protease are used by *Plasmodium* to degrade hemoglobin into amino acids as a source of nutrition (17). The tannin content in MSEE may play a role in the inhibition of the *Plasmodium* protease enzyme, so that its growth and development are inhibited and prevent invasion of new erythrocytes. Oleanolic acid which is one of the triterpenoid compounds has antiplasmodium activity caused by its incorporation into the erythrocyte membrane and causes erythrocyte changes to form stomatocytes, thus interfering with the growth of *P. falciparum* strain 3D7 in vitro (18). The invasion and growth of *Plasmodium* in erythrocytes depends on the integrity and normal function of the erythrocyte membrane. Changes in the erythrocyte membrane will interfere with the growth of *Plasmodium* (19). The triterpenoid content in the MSEE in this study probably caused changes in the erythrocyte membrane, so that their growth was disturbed and the merozoites were unable to invade new erythrocytes.

The level of IFN- $\gamma$  in the NC group ( $293.88 \pm 154.51$ ) was higher than that in PC group ( $26.76 \pm 15.27$ ). The high parasitemia in NC group was responded by the host by secreting IFN- $\gamma$ . The higher parasitemia, the higher molecules which act as pathogen associated molecular pattern (PAMP) to activate more macrophage and dendritic cells to induce the IFN- $\gamma$  secretion by immune system tries to eliminate parasites by secreting IFN- $\gamma$ .

cytokines. IFN- $\gamma$  is a primary cytokine used to activate macrophages and promote the killing of intracellular pathogens (20). IFN- $\gamma$  is a Th1-derived pro-inflammatory cytokine that is important for the elimination of intracellular parasites (21).

In this study, the PC group had the lowest IFN- $\gamma$  levels compared to other groups. This is probably due to parasitemia in PC group is low, so it does not stimulate the secretion of IFN- $\gamma$  cytokines. The treatment of dihydroartemisinin therapy to BALB/c mice infected with *P. berghei* ANKA showed a decrease in IFN- $\gamma$  levels compared to untreated infected mice (22). In this study, the same thing happened, the levels of IFN- $\gamma$  in PC group that given with DHP were lower than those in NC. This results indicated that DHP is a potent antimalarial drug which was able to suppress the production of proinflammation cytokine such as IFN- $\gamma$ .

The mean of IFN- $\gamma$  level was increased along with the increased of MSEE concentration. However, this was not statistically significant. IFN- $\gamma$  levels in the 100% MSEE group had a standard deviation greater than the mean in that group. This happened because one sample of the study had a much higher IFN- $\gamma$  levels than the other sample. Although the mice as samples in this study have attempted to have small variations with uniformity of sex, age and body weight of mice, each mice may have different abilities in responding to a given infection. The immune system between individuals varies widely (23). Every individual has a specific immune system and is different from each other (24), that cause each individual responds differently to a given infection.

When compared to the PC group, the mean IFN- $\gamma$  levels in the MSEE group (100%, 50%, and 25%) were greater and showed a statistically significant difference. This is most likely relating to parasitemia as shown by the Spearman's correlation analysis. A potent antimalarial drug used as PC showed a sharp decrease in parasitemia, and reached 0% on day 4 post treatment along with the decreased of IFN- $\gamma$  level. Likewise, the decreased of IFN- $\gamma$  levels in MSEE-treated group of mice were followed by the decreases of parasitemia in a concentration dependent manner. This proved the role of IFN- $\gamma$  in parasite elimination. There was no statistically significant difference of the IFN- $\gamma$  levels between NC and all MSEE treatment groups. This is probably because parasitemia in NC can stimulate cell activation to secrete IFN- $\gamma$  in response to infection, whereas in the MSEE group besides parasitemia, the compounds contained can also activate cells to secrete IFN- $\gamma$ , so that in all these groups there are conditions that can trigger cell activation in secreting IFN- $\gamma$ .

The significant correlation between parasitemia on the fourth day post treatment with IFN- $\gamma$  levels showed that the immune cells in BALB/c mice tried to eliminate *P. berghei* ANKA infection by secreting IFN- $\gamma$ . In this study,

there was a tendency for IFN- $\gamma$  levels to be higher in the group with higher parasitemia. The same was true of studies assessing cytokine and antibody profiles in *P. falciparum* infected individuals in Southwest Nigeria. The results showed that IFN- $\gamma$  levels were higher in the group with symptomatic malaria, which had a higher percentage of parasitemia compared to asymptomatic malaria patients with lower parasitemia (25). An association between IFN- $\gamma$  levels and parasitemia were reported in *P. falciparum* infection which showed an increase in IFN- $\gamma$  levels at an increase in parasitemia levels (26). Another study showed that there was no relationship between IFN- $\gamma$  levels and parasite density, but descriptively, IFN- $\gamma$  levels were higher in the group with higher parasite density (27).

IFN- $\gamma$  provides important protection during the erythrocytic stage of *Plasmodium* infection (28). Erythrocytic stage triggers a strong IFN- $\gamma$  response during acute infection in *P. berghei*, *P. yoelii* and *P. chabaudi* infections in mice as well as in *P. falciparum* infections in humans (29). However, high IFN- $\gamma$  secretion during erythrocytic infection is associated with a predisposition to severe malaria, such as cerebral malaria. Proinflammatory cytokines such as IFN- $\gamma$  are illustrated as double swords. Under normal conditions, these cytokines are essential for controlling parasite growth and sustained protection against disease pathologies, but excessive and uncontrolled secretion can lead to multiple immunopathologies

## CONCLUSION

This study proved the antimalarial activity of MSEE in BALB/c mice infected with *P. berghei* ANKA. Treatment of MSEE caused a significant difference in the decreased of parasitemia and mean of IFN- $\gamma$  levels in a concentration dependent manner. The decreased of parasitemia correlated with the decreased of IFN- $\gamma$  levels at the day four post treatment .

## ACKNOWLEDGEMENTS

<sup>31</sup> The authors would like to thank Ministry of Health of the Republic of Indonesia for funding this research. We also thank the workers of the experimental animal laboratory, Department of Biochemistry, Universitas Alrlangga for the maintenance of mice during the research

## REFERENCES

1. Talapko J, Krlec I, Alebić T, Jukić M, Včev A. Malaria: The past and the present. *Microorganisms*. 2019;7(6):179. doi:10.3390/microorganisms7060179.
2. World Health Organization. World Malaria Report 2021. Geneva: World Health Organization; 2021. Licence: CC BY-NC-SA 3.0 IGO.
3. Buck E, Finnigan NA. Malaria. [Updated 2022 Apr 30]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2022 Jan-. [cited 2022 July 02]. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK551711/>.
4. King T, Lamb T. Interferon- $\gamma$ : The jekyll and hyde of malaria. *PLoS Pathog*. 2015;11(10):e1005118. doi:10.1371/journal.ppat.1005118.
5. Mufidah F, Zuhrotun A. Medicinal plants for malaria treatment in Indonesia based on ethnopharmacy. *Fitofarmaka : Jurnal Ilmiah Farmasi*. 2020; 10(2): 106-121.
6. Nugraha AS, Purnomo YD, Pratama ANW, et al. Isolation of antimalarial agents from Indonesian medicinal plants: *Swietenia mahagoni* and *Pluchea indica*. *Natural Product Communications*. 2022; Jan. doi:10.1177/1934578X211068926.
7. Astuti RUNW, Fathoni I, Rahayu T, Prasetyo AR. Bioactivities of mahoni's (*Swietenia mahagoni* (L.) jacq.) seed ethanolic extract on the parasitemic level of *plasmodium berghei* in mice mus musculus. *KnE Life Science*. 2015;2(1):422-426.
8. Attemene SDD, Beourou S, Tuo K, Gnonjdji AA, Konate A, Toure AO, et al. Antiplasmodial activity of two medicinal plants against clinical isolates of *Plasmodium falciparum* and *Plasmodium berghei* infected mice. *J Parasit Dis*. 2018 Mar;42(1):68-76. doi:10.1007/s12639-017-0966-7.
9. Riza H, Fahrurroji A, Supriyanto, Wicaksono A. The antimalarial activity of the water extract of simpur (*Dillenia indica* L.) leaves against *Plasmodium berghei* in mice. *Trad. Med*. 2019;24(1):28-32. doi:10.22146/mot.38626.
10. Rudrapal M, Chetia D. Plant flavonoids as potential source of future antimalarial leads. *Sys Rev Pharm*. 2018;8(1), pp.13-18.
11. MacRae JI, Lopaticki S, Maier AG, Rupasinghe T, Nahid A, Cowman AF, et al. *Plasmodium falciparum* is dependent on de novo myo-inositol biosynthesis for assembly of gpi glycolipids and infectivity. *Mol Microbiol*. 2014;91(4):762-76. doi:10.1111/mmi.12496.
12. Marlina E, Hairani R, Tjahjandarie TS, Tanjung M. Antiplasmodial activity of flavonoids from Macaranga tanarius leaves. *IOP Conf Series: Earth Environ Sci*. 2018;144: 012011.
13. Talukdar S, Ghosh K. Differential inhibition of digestive proteases by tannin in two size groups of rohu (*Labeo rohita*, Hamilton): a biochemical and zymography study. *Aquac Res*. 2018;50:449-56. doi:10.1111/are.13910.
14. Lilburn TG, Cai H, Zhou Z, Wang Y. Protease-associated cellular networks in malaria parasite *Plasmodium falciparum*. *BMC Genomics*. 2011;12(Suppl 5):S9. doi:10.1186/1471-2164-12-S5-S9.
15. Alam A. Serine proteases of malaria parasite *Plasmodium falciparum*: potential as antimalarial drug targets. *Interdiscip Perspect Infect Dis*. 2014:1-7. doi:10.1155/2014/453186.

16. Deu E. Proteases as antimalarial targets: strategies for genetic, chemical, and therapeutic validation. *FEBS J.* 2017;284(16): 2604-28. doi:10.1111/febs.14130.
17. Sairafianpour M, Bahreininejad B, Witt M, Ziegler HL, Jaroszewski JW, Staerk D. Terpenoids of *Salvia hydrangea*: two new, rearranged 20-norabietanes and the effect of oleanolic acid on erythrocyte membranes. *Planta Med.* 2003;69(9):846-50. doi:10.1055/s-2003-43212.
18. Wijayanti L, Yarso KY, Purwanto B, Mudigdo A, Suwito H, Dirgahayu P, et al. The antiplasmodial activity of chalcone derivative through the inhibition of haemozoin formation and the induction of stomatocytes formation. *Bali Med J.* 2018;8(1):365-70. doi:10.15562/bmj.v8i1.1196.
19. Actor J. *Introductory immunology, 2nd basic concepts for interdisciplinary applications.* 2nd ed. London: Elsevier, Academic Press; 2019.
20. Sylvester B, Gasarasi DB, Aboud S, Tarimo D, Masawe S, Mpembeni R, et al. Interferon- $\gamma$  and Interleukin-10 responses during clinical malaria episodes in infants aged 0-2 years prenatally exposed to *Plasmodium falciparum*: Tanzanian birth cohort. *J Trop Med.* 2018;2018:6847498. doi:10.1155/2018/6847498.
21. Zhang T, Zhang Y, Jiang N, Zhao X, Sang X, Yang N, et al. Dihydroartemisinin regulates the immune system by promotion of CD8<sup>+</sup> T lymphocytes and suppression of B cell responses. *Sci China Life Sci.* 2020;63(5):737-49. doi:10.1007/s11427-019-9550-4.
22. Brodin P, Davis MM. Human immune system variation. *Nat Rev Immunol.* 2017;17(1):21-9. doi:10.1038/nri.2016.125.
23. Afridi S, Azizullah. Individual immune system responds differently. *Immunol Case Reports.* 2017;1(1):5-10. doi:10.35841/immunology-case-reports.1.1.5-10.
24. Iriemenam NC, Okafor CM, Balogun HA, Ayede I, Omosun Y, Persson JO, et al. Cytokine profiles and antibody responses to *Plasmodium falciparum* malaria infection in individuals living in Ibadan, southwest Nigeria. *Afr Health Sci.* 2009;9(2):66-74.
25. Benjamin GY, Ajiboye T, Eneya RP, Dauda E, Whong CMZ, Jatau ED. Determination of interferon gamma associated with malaria parasitaemia among patients attending selected hospitals in Zaria, Kaduna State, Nigeria. *Bayero J Med Lab Sci.* 2016;1(1): 8-13.
26. Barkat H, Abd Alla AB, Galander A, Salah T, Elfaki T, Nasir A. Prevalence of malaria and quantification of cytokine levels during infection in East Nile locality, Khartoum State: a cross-sectional study. *F1000Res.* 2019;8:1529. doi:10.12688/f1000research.19217.2.
27. Gun SY, Claser C, Tan KS, Rñnia L. Interferons and interferon regulatory factors in malaria. *Mediators Inflamm.* 2014;2014:243713. doi:10.1155/2014/243713.
28. Perez-Mazliah D, Langhorne J. CD4 T-cell subsets in malaria: TH1/TH2 revisited. *Front Immunol.* 2015;5:671. doi:10.3389/fimmu.2014.00671.

# Antimalarial Activity of Mahagony Seed Ethanolic Extract in BALB/c Mice Infected with Plasmodium berghei ANKA and The Correlation of Parasitemia and Plasma Level of IFN- $\gamma$

## ORIGINALITY REPORT

19%

SIMILARITY INDEX

13%

INTERNET SOURCES

13%

PUBLICATIONS

4%

STUDENT PAPERS

## PRIMARY SOURCES

1	Submitted to Universitas Airlangga Student Paper	1%
2	<a href="http://spiral.imperial.ac.uk">spiral.imperial.ac.uk</a> Internet Source	1%
3	<a href="http://cyberleninka.org">cyberleninka.org</a> Internet Source	1%
4	Qiongming Xu, Lingchuan Guo, Xinhua Gu, Biao Zhang et al. "Prevention of colorectal cancer liver metastasis by exploiting liver immunity via chitosan-TPP/nanoparticles formulated with IL-12", Biomaterials, 2012 Publication	1%
5	Submitted to University of Newcastle Student Paper	1%
6	"Plant-derived Bioactives", Springer Science and Business Media LLC, 2020 Publication	1%



7	M Iigo, T Nakagawa, C Ishikawa, Y Iwahori, M Asamoto, K Yazawa, E Araki, H Tsuda. "Inhibitory effects of docosahexaenoic acid on colon carcinoma 26 metastasis to the lung", British Journal of Cancer, 1997 Publication	1 %
8	Maria A. F. Queiroz, Vânia N. Azevedo, Ednelza da S. G. Amoras, Tuane C. F. Moura et al. "IFNG +874A/T Polymorphism Among Asymptomatic HTLV-1-Infected Individuals Is Potentially Related to a Worse Prognosis", Frontiers in Microbiology, 2018 Publication	1 %
9	<a href="http://kubanni.abu.edu.ng">kubanni.abu.edu.ng</a> Internet Source	<1 %
10	Submitted to University of Queensland Student Paper	<1 %
11	<a href="http://journal.wima.ac.id">journal.wima.ac.id</a> Internet Source	<1 %
12	<a href="http://worldwidescience.org">worldwidescience.org</a> Internet Source	<1 %
13	<a href="http://etd.aau.edu.et">etd.aau.edu.et</a> Internet Source	<1 %
14	<a href="http://www.hindawi.com">www.hindawi.com</a> Internet Source	<1 %

15	Internet Source	<1 %
16	<a href="http://www.mdpi.com">www.mdpi.com</a> Internet Source	<1 %
17	"Comprehensive Analysis of Parasite Biology: From Metabolism to Drug Discovery", Wiley, 2016 Publication	<1 %
18	Submitted to University of College Cork Student Paper	<1 %
19	<a href="http://bmcbgenomics.biomedcentral.com">bmcbgenomics.biomedcentral.com</a> Internet Source	<1 %
20	Chao Wang, Arne Krüger, Xiaochen Du, Carsten Wrenger, Matthew R. Groves. "Novel Highlight in Malarial Drug Discovery: Aspartate Transcarbamoylase", Frontiers in Cellular and Infection Microbiology, 2022 Publication	<1 %
21	<a href="http://core.ac.uk">core.ac.uk</a> Internet Source	<1 %
22	<a href="http://medicineworld.org">medicineworld.org</a> Internet Source	<1 %
23	<a href="http://oamjms.eu">oamjms.eu</a> Internet Source	<1 %
24	<a href="http://coek.info">coek.info</a> Internet Source	<1 %

<1 %

25

Xindang Zhang, Hengzhi Wang, Jianwei Zhang, Beibei Lin, Lusi Chen, Qiumei Wang, Guangbin Li, Junming Deng. "Assessment of rapeseed meal as fish meal alternative in diets for juvenile Asian red-tailed catfish (*Hemibagrus wyckioides*)", *Aquaculture Reports*, 2020

Publication

<1 %

26

Yuelan Zhao, Lufeng Jiang, Teng Liu, Min Wang, Wenbo Cao, Yongzhan Bao, Jianhua Qin. "Construction and immunogenicity of the recombinant *Lactobacillus acidophilus* pMG36e-E0-LA-5 of bovine viral diarrhea virus", *Journal of Virological Methods*, 2015

Publication

<1 %

27

[academicjournals.org](http://academicjournals.org)

Internet Source

<1 %

28

[doi.org](http://doi.org)

Internet Source

<1 %

29

[e-nrp.org](http://e-nrp.org)

Internet Source

<1 %

30

[garuda.kemdikbud.go.id](http://garuda.kemdikbud.go.id)

Internet Source

<1 %

31

[psasir.upm.edu.my](http://psasir.upm.edu.my)

Internet Source

<1 %

32

[tropmedhealth.biomedcentral.com](https://tropmedhealth.biomedcentral.com)

Internet Source

&lt;1 %

33

"Melatonin and Melatonergic Drugs in Clinical Practice", Springer Nature, 2014

Publication

&lt;1 %

34

"Poster Session", Indian Journal of Gastroenterology, 2010

Publication

&lt;1 %

35

[1library.net](https://1library.net)

Internet Source

&lt;1 %

36

A. Hennessy, H. L. Pilmore, L. A. Simmons, D. M. Painter. "A Deficiency of Placental IL-10 in Preeclampsia", The Journal of Immunology, 1999

Publication

&lt;1 %

37

Evi Zahara, Iin Fitriani, Rahmi Rahmi, Emilda AS. "Effectiveness Of Combination Of Meditation And Positive Affirmations On Labor Duration And Apgar Score", Jurnal Kebidanan Malahayati, 2022

Publication

&lt;1 %

38

Lu Diao, Ming Li, Jin Tao, Xiaojun Xu, Yiqi Wang, Ying Hu. "Therapeutic effects of cationic liposomes on lupus-prone MRL/lpr mice are mediated via inhibition of TLR4-triggered B-cell activation", Nanomedicine: Nanotechnology, Biology and Medicine, 2021

&lt;1 %

39

Soilia Fertilita, Willy Sandhika, Desak Gede Agung Suprabawati. "The Cytotoxic Activity of Annona muricata Linn Leaves Ethanolic Extract (AMEE) on T47D Breast Cancer Cell Line", Medical Laboratory Technology Journal, 2020

Publication

---

<1 %

40

[dspace.hangtuah.ac.id](https://dspace.hangtuah.ac.id)

Internet Source

---

<1 %

41

[eprints.unm.ac.id](https://eprints.unm.ac.id)

Internet Source

---

<1 %

42

[repository.stikim.ac.id](https://repository.stikim.ac.id)

Internet Source

---

<1 %

43

[repository.um-surabaya.ac.id](https://repository.um-surabaya.ac.id)

Internet Source

---

<1 %

44

[researchonline.lshtm.ac.uk](https://researchonline.lshtm.ac.uk)

Internet Source

---

<1 %

45

[www.molvis.org](https://www.molvis.org)

Internet Source

---

<1 %

46

[www.researchsquare.com](https://www.researchsquare.com)

Internet Source

---

<1 %

47

[zjrms.com](https://zjrms.com)

Internet Source

---

<1 %

48

Francis Davoine, Sophie Lavigne, Jamila Chakir, Claudine Ferland, Marie-Ève Boulay, Michel Laviolette. "Expression of FcγRIII (CD16) on human peripheral blood eosinophils increases in allergic conditions", *Journal of Allergy and Clinical Immunology*, 2002

Publication

&lt;1 %

49

Sharma Arvind Virendra, Chandrakant Sahu, Ankur Kumar, Pooja Abrol Chawla. "Natural Antioxidants as Additional Weapons in the Fight against Malarial Parasite", *Current Topics in Medicinal Chemistry*, 2022

Publication

&lt;1 %

50

Daniel Osegi Okpokor, Olusola Ajibaye, Peter Mac Asaga, Ikechukwu Nwankwo, Anthony Danaan Dakul. "Pro- and anti-inflammatory response profile modulates Plasmodium falciparum malaria outcomes among subjects from Baiyeku, Ikorodu, Lagos, Nigeria.", *Research Square*, 2020

Publication

&lt;1 %

51

KEIZO SATO, MORITAKA SUGA, TAKAAKI AKAIKE, SHIGEMOTO FUJII, HIROYUKI MURANAKA, TOSHINORI DOI, HIROSHI MAEDA, MASAYUKI ANDO. "Therapeutic Effect of Erythromycin on Influenza Virus-induced

&lt;1 %

Lung Injury in Mice", American Journal of  
Respiratory and Critical Care Medicine, 1998

Publication

---

52

Samson Olaniyi, Olusegun A. Ajala, Sulaimon  
F. Abimbade. "Optimal Control Analysis of a  
Mathematical Model for Recurrent Malaria  
Dynamics", Operations Research Forum, 2023

Publication

---

<1 %

53

V.R. Gutlapalli, Aparna Sykam, Sandeep P.  
Tenali, Sujai Suneetha, Lavanya M. Suneetha.  
"High levels of plasma interferon gamma and  
+874T/A gene polymorphism is associated  
with HIV-TB co-infection", Human  
Immunology, 2016

Publication

---

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

# Antimalarial Activity of Mahagony Seed Ethanolic Extract in BALB/c Mice Infected with Plasmodium berghei ANKA and The Correlation of Parasitemia and Plasma Level of IFN- $\gamma$

---

GRADEMARK REPORT

---

FINAL GRADE

**/100**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---

PAGE 6

---