

# Toxicological Evaluation and Protective Effects of Ethanolic Leaf Extract of *Cassia spectabilis* DC on Liver and Kidney Function of *Plasmodium berghei*-Infected Mice

*by* Wiwied Ekasari

---

**Submission date:** 08-Dec-2022 04:03PM (UTC+0800)

**Submission ID:** 1975156003

**File name:** iver\_and\_Kidney\_Function\_of\_Plasmodium\_berghei-Infected\_Mice.pdf (8.27M)

**Word count:** 6018

**Character count:** 30975

## Research Article

# Toxicological Evaluation and Protective Effects of Ethanolic Leaf Extract of *Cassia spectabilis* DC on Liver and Kidney Function of *Plasmodium berghei*-Infected Mice

Wiwied Ekasari<sup>1</sup>,<sup>1</sup> Anisah Mahardiani<sup>1</sup>,<sup>1</sup> Nindya T. Putri<sup>1</sup>,<sup>1</sup> Tutik S. Wahyuni<sup>1</sup>,<sup>1</sup> and Heny Arwati<sup>2</sup>

<sup>1</sup>Department of Pharmaceutical Sciences, Faculty of Pharmacy, Universitas Airlangga, Surabaya 60115, Indonesia

<sup>2</sup>Department of Parasitology, Faculty of Medicine, Universitas Airlangga, Surabaya 60132, Indonesia

Correspondence should be addressed to Wiwied Ekasari; [wiwied-e@ff.unair.ac.id](mailto:wiwied-e@ff.unair.ac.id)

Received 21 November 2021; Accepted 13 January 2022; Published 15 February 2022

Academic Editor: Yoshiaki Hikasa

Copyright © 2022 Wiwied Ekasari et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Currently, the presence of antimalarial drug resistance has become a major obstacle in the treatment of malaria. To overcome the problem, a series of studies are needed to find new antimalarial drugs from plants. Previously, 90% ethanolic extract of *Cassia spectabilis* DC (EECS) leaves have been reported to have antimalarial activity *in vitro* against *Plasmodium falciparum* and *in vivo* against *Plasmodium berghei* ANKA. The research is conducted to find out the toxicity and protective effects of EECS on the liver and kidneys of mice infected with *P. berghei* ANKA. The acute and subacute toxicity tests were carried out on healthy mice that were given EECS at a dose of 150 mg/kg BW. An antimalarial activity test was carried out at doses of 150 and 200 mg/kg BW in *P. berghei*-infected mice. Regarding hepatomegaly, further plasma levels of hepatic enzyme were analyzed, as well as histopathological observation of the liver to determine the effect of the extract on liver. The kidney was observed histopathologically as well. The acute toxicity test of EECS showed that there was no mouse died at the highest dose, indicating safe for the mice. The subacute toxicity based on the histology data showed no significant difference in the liver and kidney of mice between the tested group and the healthy group. The histological and enzymatic effect of EECS in mice infected with *P. berghei* showed the histological and enzymatic effect that improved liver function and the histopathological effect on kidneys with the highest activity at a dose of 200 mg/kg BW compared with the negative control. The results showed the EECS was not toxic in mice and repaired the liver and kidney functions of *P. berghei* ANKA-infected mice, indicating a good candidate for antimalarial drug development.

## 1. Introduction

Until this day, the problem of *Plasmodium* parasitic resistance towards existing antimalarial drugs is still a major problem for the eradication of malaria [1–3]. Therefore, research on the discovery of new sources of antimalarial drugs, one of which is from medicinal plants, continues to be done [4]. *In vivo* and *in vitro* models have long been used in antimalarial testing. In the latter half of the 20th century, *Plasmodium berghei* or *Plasmodium yoelii* was used to infect rodents. Meanwhile, *P. berghei* became the most commonly used species for study at the liver stage, particularly the

formation of hypnozoite to investigate malarial recurrence [5–7].

One of the traditional plants in Indonesia, *Cassia spectabilis* DC of the Caesalpiniaceae family, has been proved experimentally *in vitro* against *P. falciparum* and in treating malaria *in vivo* against *P. berghei* [8, 9], indicating that *C. spectabilis* DC plant is very potential to be further developed as a candidate for antimalarial drugs. Previous works reported that an *in vivo* test on 90% ethanolic extract of *C. spectabilis* DC leaf against *P. berghei* ANKA in BALB/c mice showed that an ED<sub>50</sub> value was 131.5 mg/kg BW [9], and it is categorized as very good antimalarial activity [10].

Furthermore, the extract, fractions, subfractions, and isolated compound of *C. spectabilis* DC have been tested *in vitro* for their antimalarial activities. The active compound of this plant has been successfully identified as a compound that is identical to (-)-7-hydroxycassine, and its *in vitro* antimalarial activity test showed a very low  $IC_{50}$  of 0.016  $\mu\text{g/mL}$  [11], which is classified as a very strong antimalarial activity [12].

The research on this plant in relation to overcoming malarial drug resistance has been continued by examining the safety and effects of improving *C. spectabilis* DC leaf extract on the liver and kidney functions in *P. berghei* ANKA-infected mice. Research studies on the effect of medicinal plants that have antimalarial activity on the liver and kidney functions of mice infected with parasites have been widely carried out [13–16], as well as acute and subacute toxicity tests [17–20]. The liver has an important role in regulating physiological processes. This organ is involved in several vital functions such as metabolism, secretion, and storage. In addition, the liver has an important role in the detoxification and excretion of endogenous and exogenous compounds [21–24].

Malarial infection begins when sporozoites are injected by malarial parasite-infected female *Anopheles* mosquito bite. During blood feeding, infected female *Anopheles* injects the sporozoite stage of parasite [25, 26]. After about an hour traveling in human body, sporozoites then enter the liver, attack hepatocytes, and start the asexual cycle of exoerythrocytic schizogony. Inside the liver cells, the parasites multiply asexually, until they reach mature schizont, and finally, a large number of merozoites produce and enter the bloodstream after infecting hepatocyte burst. The infected hepatocytes causing liver damage due to the rupture of infected hepatocytes and merozoites enter the bloodstream and start the erythrocytic cycle in the red blood cells [27, 28]. Damage that occurs in the liver cells can increase the enzymes that work on liver function, especially transaminase enzyme and morphological changes in the appearance of liver [29, 30], such as hepatomegaly. Hepatosplenomegaly is a common feature of malarial infection in humans [31] and mice [32] caused by chronic exposure to malarial parasites. However, no enlargement of kidney in malaria infection.

The dysfunction of the liver can be detected by hepatocellular transamination of plasma glutamic oxaloacetic transaminase (SGOT) and plasma glutamic pyruvic transaminase (SGPT) released in plasma, or by histological examination of the tissue [33]. The most common damage is the activation of apoptotic cell death or hepatocyte necrosis [34–36]. Clinically significant kidney involvement is associated with the infection by *P. falciparum* and *P. malariae*. The infection of *P. falciparum* produces acute manifestations, ranging from asymptomatic, up to urinary disorders, and mild electrolyte disturbances for acute renal failure (ARF) or acute kidney injury (AKI) that require dialysis support [27, 37]. The cases of AKI in the complication of malaria are known to contribute to a high mortality rate, which is about 75% of cases. The histological study suggests

the presence of glomerulonephritis, acute tubular necrosis, and interstitial nephritis as the key hemodynamic factor in malaria-associated AKI [38]. Generally, the degree of kidney dysfunction can be detected by the presence of adequate amounts of protein in the urine and an increase in the plasma urea, creatinine, and plasma electrolyte levels [39].

The toxicity of 90% ethanolic extract of *C. spectabilis* DC (EECS) leaves in BALB/c mice, followed by the enzyme examinations, may affect histopathology and liver and kidney functions post-EECS administration.

## 2. Materials and Methods

**2.1. Plant Material and Preparation of EECS.** The *C. spectabilis* DC leaves were purchased and determined in LIPI (Indonesia Research Centre), Botanical Garden, Purwodadi, East Java, Indonesia (B-160/IPH.06/KS.02/III/2019). The specimen was deposited as the herbarium in the Department of Pharmacognosy and Phytochemistry, Faculty of Pharmacy, Universitas Airlangga. The leaves were rinsed thoroughly with tap water to remove extraneous contaminants, dried at 45°C, and ground into powder with a grinder. The extraction was carried out by macerating the powder plant materials (500 g) in a flask containing 2,500 mL of 90% ethanol (at 25–30°C) for 3 × 24 hours. The extraction solvent was separated, filtered through filter paper, and evaporated under reduced pressure by rotary evaporation. The ethanolic extract yield of 1000 g dried weight of *C. spectabilis* DC leaf powder was 10.20% (*w/w*) and was used in this experiment.

**2.2. Animals.** Male BALB/c mice aged 6–8 weeks, weight of 25–30 g, were used in the study. All mice were obtained from Farma Veterinary Center, Directorate General of Livestock and Animal Health Ministry of Agriculture, Surabaya, East Java, Indonesia. The animals were housed under standard conditions and fed with a stock diet and water *ad libitum*. The approval of the study protocol was obtained from the Ethics Committee for Animal Research, Universitas Airlangga, Indonesia, Number 2.KE.181.10.2018.

**2.3. Acute Toxicity Test.** EECS was weighted and resuspended with 0.5% sodium carboxymethyl cellulose (Na CMC) to obtain the desired doses. BALB/c mice were fasted for 24 hours before being fed with the extract. The animals were divided into three groups, each group containing five mice. The doses for each group were 1,250; 2,500; and 5,000 mg/kg BW, respectively. The general behavior of each mouse was observed continuously for one hour after each dose, intermittently every four hours, and thereafter over a period of 24 hours [40–42].

**2.4. Subacute Toxicity Test.** In the subacute toxicity test, mice were grouped into three groups and each group was treated with EECS at once, five times, and ten times of 150 mg/kg BW daily oral dose for 28 days; the normal control group was only given food and drink *ad libitum* [43]. The animals were observed for 28 days for any sign of toxicity. At the end

of observational period, all animals were sacrificed under ether anesthesia and vital organs such as livers and kidneys were removed from all animals for gross and histopathological examinations.

**2.5. Rodent's Parasite.** Chloroquine sensitivity of *Plasmodium berghei* ANKA was obtained from the Institute of Biomolecular Eijkman, Jakarta, Indonesia, and maintained by serial passaging that was used in this study.

**2.6. Effect of EECS on the Plasma Level of SGOT and SGPT.** This experiment was a suppressive test based on Peters [44]. This suppressive test was done prior to the evaluation of the EECS on the plasma level of SGOT and SGPT. The animals were infected with *P. berghei* ANKA and divided into five groups of A, B, C, D, and E. The animals were treated shortly after infection on day 0 ( $D_0$ ) and continued daily for three days ( $D_1$ – $D_3$ ). Group A as a negative control received 0.5% Na CMC orally. Groups B and C as the treated groups were given EECS at a single oral dose of 150 and 200 mg/kg BW, respectively. Group D was a positive control given chloroquine at a single oral dose of 100 mg/kg BW. Moreover, group E as a healthy control was not infected with parasites and was only given food and drink ad libitum. On the fourth day ( $D_4$ ), the blood smears of each mouse were prepared and examined microscopically. Further, plasma was collected prior to the measurement of SGOT and SGPT levels.

**2.7. Effect of EECS on Histopathology of Liver and Kidney.** At the end of the observational period, all animals were sacrificed under ether anesthesia prior to the liver and kidney removal. After weighing the organs, the tissues were then fixed in 10% formaldehyde solution and further processed for hematoxylin-eosin (H&E) staining. The histopathological observation was performed to find out the toxicological effect of EECS on the degenerative and necrotic cells of the tissues. The damages were graded based on Gibson-Corley et al. [45].

**2.8. Statistical Analysis.** The data were represented as mean  $\pm$  standard deviation (SD). One-way analysis of variance (ANOVA) was used to compare the normally distributed data among the treatments, followed by post hoc multiple comparison test when a different significance was obtained. When the data were not distributed normally, the Kruskal–Wallis and the Mann–Whitney U-test were used to assess the differences among treatments.

### 3. Results

**3.1. Acute Toxicity Test.** No mice died were observed within 24 hours postoral administration of the extract. The behavioral observation of the mice mediated no toxicity sign effects such as paw licking, hair erection, reduction in feeding activity, and increasing respiratory rate [46, 47].

TABLE 1: Number of animals' deaths in the acute toxicity test of EECS on BALB/c mice in 24 hours.

Doses (mg/kg BW)	Number of deaths (r)
1,250	0
2,500	0
5,000	0


Based on the number of animals' death from the three doses of extract (see Table 1), yielding  $r$  values were 0, 0, and 0, assuming there was no animal died at the highest dose (5,000 mg/kg BW).

**3.2. Subacute Toxicity Test.** The gross examination of the liver showed hepatomegaly, but there was no enlargement of the kidneys. The size and cell morphology of liver and kidney in both the treated and control groups were similar among the groups (see Figure 1). The results of histopathological changes in the liver and kidney tissues are summarized in Table 2. Statistically, there was no significant difference ( $p > 0.05$ ) and histopathological changes were observed in the liver and kidney tissues of the test groups when compared to the control groups.

**3.3. Effect of EECS on SGOT and SGPT.** The effect of EECS on biochemical parameters showed that the EECS intake induced a significant increase in plasma levels of SGOT at the dose of 150 mg/kg BW and was out of the normal range. The normal range of SGOT was 59–247 U/L and SGPT in mouse was 28–132 U/L [48], while the plasma level of SGPT was still in the normal range. At a dose of 200 mg/kg BW, the plasma levels of either SGOT or SGPT were in the normal range. There was a very sharp increase in the plasma levels of SGOT and SGPT in the negative control group, indicating liver damage (see Table 3).

The observations on the morphology of liver and kidneys of mice showed the enlargement of the liver (hepatomegaly) and showed differences in color, size, and texture in the liver. The morphology of kidneys showed no enlargement and was similar either in color, size, or texture among groups. Due to these reasons, the biochemistry examination of the kidney was not performed. Physically, there was an improvement in the liver of mice treated with 150 mg/kg BW of extract when compared to the control, although it was not as good as in mice treated with 200 mg/kg BW of extract. In addition, the appearance of the liver treated with 200 mg/kg BW extract was almost as same as the condition of experimental mouse liver treated with chloroquine and the healthy control group, where brownish-red and smooth texture of surface has appeared.

The results of histopathological changes in the liver and kidney tissues are summarized in Table 4. There were no significant histopathological changes observed in the liver and kidney tissues of the mice when compared to the control. The histopathological changes in liver and kidney are shown in Figures 2 and 3.



**FIGURE 1:** Histological section of the liver (a–d) and kidney (e–h) of mice (section stained with H&E,  $\times 400$ ). Green arrow: degenerative cell, yellow arrow: necrotic cell. (a and e) Single dose. (b and f) Five times of dose. (c and g) Ten times of dose. (d and h) Normal control groups.

TABLE 2: Lesion scores of liver and kidney of BALB/c mice in subacute oral toxicity of EECS.

Organ	Mean score of lesions	Groups			Normal control	Asymptotic significance ( $p < 0.05$ )
		EECS 1 × 150 mg/kg	EECS 5 × 150 mg/kg	EECS 10 × 150 mg/kg		
Liver	Necrosis	2.34 ± 0.10 <sup>a</sup>	2.29 ± 0.30 <sup>a</sup>	2.03 ± 0.92 <sup>a</sup>	2.37 ± 0.34 <sup>a</sup>	0.838
	Degeneration	2.20 ± 0.16 <sup>a</sup>	2.43 ± 0.21 <sup>a,b</sup>	2.77 ± 0.60 <sup>b,c</sup>	2.26 ± 0.25 <sup>a</sup>	0.029*
	Mean score	2.27 ± 0.13	2.36 ± 0.25	2.40 ± 0.76	2.31 ± 0.29	0.433
Kidney	Necrosis	1.94 ± 0.32 <sup>a</sup>	1.09 ± 0.60 <sup>b</sup>	1.54 ± 0.63 <sup>a,b</sup>	1.60 ± 0.48 <sup>a,b</sup>	0.060
	Degeneration	1.14 ± 0.49 <sup>a</sup>	1.74 ± 0.19 <sup>b</sup>	1.93 ± 0.71 <sup>a,b</sup>	1.86 ± 0.46 <sup>b,c</sup>	0.065
	Mean score	1.16 ± 0.40	1.41 ± 0.39	1.73 ± 0.67	1.73 ± 0.47	0.062

Data are expressed as mean ± standard deviation ( $n = 7$ ). The superscript (\*) denotes that the values were significantly different from control ( $p < 0.05$ ; Kruskal–Wallis test for global comparison of organ lesions among groups). The superscripts (a, b, c) indicate that the values in the same row with different superscript letters were significantly different ( $p < 0.05$ ; Mann–Whitney test). EECS: 90% ethanolic leaf extract of *Cassia spectabilis* DC.

TABLE 3: Differences in plasma level of SGOT and SGPT of *P. berghei* ANKA-infected mice treated with EECS compared with the control groups.

Groups	Parasitemia (%)		Suppression (%)	SGOT (U/L)	SGPT (U/L)
	$D_0$	$D_4$			
Negative control	1.24 ± 0.69	14.42 ± 7.86	—	658.75 ± 99.52	167.75 ± 32.37
EECS 150 mg/kg	2.13 ± 1.19	8.77 ± 4.42	51.85	384.75 ± 73.98	75.75 ± 35.52
EECS 200 mg/kg	0.98 ± 0.64	9.42 ± 3.27	58.93	216.00 ± 48.80	42.00 ± 4.50
Chloroquine	2.30 ± 1.44	0.14 ± 0.28	100.00	175.50 ± 45.59	41.25 ± 5.60
Healthy control	—	—	—	162.50 ± 59.18	97.25 ± 44.70

Data are expressed as mean ± standard deviation ( $n = 4$ ). No significant difference between the groups ( $p > 0.05$ ; one-way ANOVA followed by Tukey's multiple comparison test).  $D_0$ : day before the treatment;  $D_4$ : fourth day after treatment; SGOT: plasma glutamic oxaloacetic transaminase; SGPT: plasma glutamic pyruvic transaminase; EECS: 90% ethanolic leaf extract of *Cassia spectabilis* DC.

TABLE 4: Lesion scores of liver and kidney of *P. berghei* ANKA-infected mice treated with EECS.

Organ	Mean scores of lesions	Groups				
		Negative control	EECS 150 mg/kg	EECS 200 mg/kg	Chloroquine	Healthy control
Liver	Necrosis	1.30 ± 0.26 <sup>a</sup>	0.75 ± 0.34 <sup>a,b</sup>	0.60 ± 0.16 <sup>b</sup>	0.65 ± 0.10 <sup>b</sup>	0.50 ± 0.12 <sup>b</sup>
	Degeneration	2.45 ± 0.38 <sup>a</sup>	1.45 ± 0.19 <sup>b</sup>	1.10 ± 0.35 <sup>b</sup>	1.20 ± 0.28 <sup>b,c</sup>	0.65 ± 0.10 <sup>c</sup>
	Mean score	1.87 ± 0.32	1.10 ± 0.26	0.85 ± 0.25	0.92 ± 0.19	0.57 ± 0.11
Kidney	Necrosis	1.75 ± 0.50 <sup>a</sup>	1.55 ± 0.41 <sup>a,b</sup>	1.65 ± 0.19 <sup>a</sup>	1.40 ± 0.28 <sup>b</sup>	1.20 ± 0.36 <sup>c</sup>
	Degeneration	1.30 ± 0.74 <sup>a,b</sup>	1.20 ± 0.52 <sup>a,b</sup>	1.20 ± 0.16 <sup>a,b</sup>	1.10 ± 0.26 <sup>a</sup>	1.45 ± 0.68 <sup>b</sup>
	Mean score	1.52 ± 0.62	1.37 ± 0.46	1.42 ± 0.17	1.25 ± 0.27	1.32 ± 0.52

Data are expressed as mean ± standard deviation ( $n = 7$ ). No significant difference between the groups ( $p > 0.05$ ; Kruskal–Wallis test). The superscripts (a, b, c) indicate that the values in the same row with different superscript letters were significantly different ( $p < 0.05$ ; Mann–Whitney test). EECS: 90% ethanolic leaf extract of *Cassia spectabilis* DC.

#### 4. Discussion

Currently, the optimal use of available drugs and the development of a new approach to antimalarial chemotherapy are still needed. With serious consideration for safety issues, researches on phytochemistry, pharmacology, and biochemistry in medicinal plants traditionally used as antimalarial are urgently needed [49]. The plants that show good antimalarial activities that have been recorded until today come from 1,277 plants from 160 plant families [50, 51].

The acute toxicity test of the EECS showed that the EECS is not toxic to the BALB/c mice up to 5,000 mg/kg BW of dose, since there was no mouse died due to this highest dose of EECS. Following this test, the subacute toxicity test showed that there was no abnormality in mouse mobile activities and in water and food consumption.

In addition, the subacute toxicity test was based on the liver and kidney histopathology results (see Figure 1). The data obtained based on microscopic observations were analyzed statistically in the form using the Kruskal–Wallis statistical test to determine the differences in all population groups. Based on the results of the Kruskal–Wallis test, which evaluated liver necrosis damage, it was known that there were no significant differences between the groups. This could be seen from the value of  $p > 0.05$ . Then, the Mann–Whitney statistical test was also performed to determine the location of differences between the groups. Based on the results of the Mann–Whitney test on the level of liver necrosis damage, it was known that there were no significant differences between the values of  $p > 0.05$ . Likewise in liver degeneration damage, it could be seen that the four treatment groups showed no




FIGURE 2: Pathological lesions around the central vein of experimental mouse liver (section stained with H&E,  $\times 400$ ). (a) Normal hepatocyte cells around the central vein. (b) Degenerative nuclear hepatocyte. (c) Inflammatory cell infiltration. (d) Necrotic hepatocyte.

significant differences because there were no significant differences (not toxic to the liver).


The suppressive test of the EECS in this current research was performed to find out the effect of EECS on the function of the liver based on the plasma level of SGOT and SGPT. The doses of EECS used in this test were based on the previous suppressive test [9]. Furthermore, the plasma level of SGOT and SGPT in EECS-treated infected mice was maintained in the normal range compared with the controls and based on the reference of normal BALB/c mice [52], especially at the dose of 200 mg/kg BW. The statistical analysis showed no significant difference ( $p < 0.05$ ) in plasma level of SGOT and SGPT when compared with the healthy control group. These results indicated that the EECS was effective in reducing plasma levels of SGOT and SGPT to the normal levels. The excessive hepatic enzymes were released by the injured hepatocytes due to malarial infection [53]. In this study, the necrosis and degraded cells in the liver were repaired by the administration of EECS followed by the reduction in plasma level of SGOT and SGPT. These results indicated a very good liver functional improvement due to the EECS administration.

Apart from blood analysis, the histopathological analysis provides supportive evidence for biochemical

and hematological assessment [41]. Histopathological examination is the standard for evaluating treatment for pathological changes in tissues and organs [54]. Histopathological assessment of the effect of EECS on the liver samples treated in this study showed both necrosis and degeneration of hepatocytes and kidney cells (see Table 4). It is likely that these changes may reflect beneficial cellular adaptations of the extract to host tissues.

Based on the results, all doses showed significant differences in kidney lesion scores when compared to the controls (see Table 4). There was a significant difference between chloroquine and the healthy group in necrotic and degenerative cells of kidney. This significant difference shows that chloroquine provides a therapeutic effect on kidney repair in *P. berghei*-infected mice.

In the absence of significant differences between the treated group and chloroquine (see Table 4), it can be concluded that the therapeutic effect is not too different from being used as an alternative to antimalaria, which can reduce kidney damage. Furthermore, a dose of 200 mg/kg BW showed better liver repair activity than a dose of 150 mg/kg BW because there was no significant lesion score in the dose of 150 mg/kg BW compared with the negative control.



**FIGURE 3:** Histopathological appearances of kidney (section stained with H&E, ×400). Green arrow: degenerative cell, yellow arrow: necrotic cell, blue arrow: normal cell. (a) Negative control group. (b) 150 mg/kg BW of extract. (c) 200 mg/kg BW of extract. (d) Chloroquine. (e) Healthy control group.

The EECS has been proved to be not toxic in mice and was able to repair the liver and kidney functions in mice infected with *P. berghei* ANKA. The ability of EECS in heme detoxification

caused the parasite clearance [11] followed by further repairing of tissue in the main organ affected by the malarial infection such as liver and kidney as well as the plasma level of hepatic enzyme.



## 5. Conclusion

The EECS was not toxic in mice and repaired the liver and kidney functions of *P. berghei* ANKA-infected mice similarly to chloroquine, indicating a good candidate for antimalarial drug development.

## Data Availability

The data used to support the findings of this study are available from the corresponding author upon request.

## Conflicts of Interest

The authors declare that there are no conflicts of interest regarding the publication of this study.

## Acknowledgments

This research was supported by the Ministry of Education, Culture, Research and High Education of the Republic of Indonesia.

## References

- [1] M. Tanner, B. Greenwood, C. J. Whitty et al., "Malaria eradication and elimination: views on how to translate a vision into reality," *BMC Medicine*, vol. 13, no. 167, p. 167, 2015.
- [2] B. Hanboonkunupakarn and N. J. White, "The threat of antimalarial drug resistance," *Tropical Disease, Travel Medicine and Vaccines*, vol. 210 pages, 2016.
- [3] D. Menard and A. Dondorp, "Antimalarial drug resistance: a threat to malaria elimination," *Cold Spring Harbor Perspectives in Medicine*, vol. 7, no. 7, Article ID a025619, 2017.
- [4] H. Ginsburg and E. Deharo, "A call for using natural compounds in the development of new antimalarial treatments—an introduction," *Malaria Journal*, vol. 10, no. S1, 2011.
- [5] A. Rodriguez-acosta, H. J. Finol, M. Pulido-Mendez et al., "Liver ultrastructural pathology in mice infected with *Plasmodium berghei*," *Journal of Submicroscopic Cytology and Pathology*, vol. 30, no. 2, pp. 299–307, 1998.
- [6] E. Aidoo, F. K. Addai, J. Ahenkorah, B. Hottor, K. A. Bugyei, and B. A. Gyan, "Natural cocoa ingestion reduced liver damage in mice infected with *Plasmodium berghei* (NK65)," *Research and Reports in Tropical Medicine*, vol. 3, pp. 107–116, 2012.
- [7] R. Raphemot, D. Posfai, and E. R. Derbyshire, "Current therapies and future possibilities for drug development against liver-stage malaria," *Journal of Clinical Investigation*, vol. 126, no. 6, pp. 2013–2020, 2016.
- [8] W. Ekasari, T. S. Wahyuni, and R. A. S. Yudistira, "Potensi antimalaria dan pemeriksaan mikroskopik-fitokimia daun beberapa tanaman genus *Cassia*," *Jurnal Farmasi dan Ilmu Kefarmasian Indonesia*, vol. 2, no. 2, pp. 48–52, 2015.
- [9] W. Ekasari, T. S. Wahyuni, H. Arwati, and N. T. Putri, "Determination of effective dose of antimalarial from *Cassia spectabilis* leaf ethanol extract in *Plasmodium berghei*-infected mice," *African Journal of Infectious Diseases*, vol. 12, no. S1, pp. 110–115, 2018.
- [10] V. Muñoz, M. Sauvain, G. Bourdy et al., "A search for natural bioactive compounds in bolivia through a multidisciplinary approach: part I. evaluation of the antimalarial activity of plants used by the chacobo Indians," *Journal of Ethnopharmacology*, vol. 69, no. 2, pp. 127–137, 2000.
- [11] W. Ekasari, D. R. Basuki, H. Arwati, and T. S. Wahyuni, "Antiplasmodial activity of ethanolic extract of *Cassia spectabilis* DC leaf and its inhibition effect in heme detoxification," *BMC Complementary Medicine and Therapies*, vol. 21, no. 1, p. 71, 2021.
- [12] P. Rasoanaivo, E. Deharo, S. Ratsimamanga-Urverg, and F. Frappier, "Guidelines for the nonclinical evaluation of the efficacy of traditional antimalarials," in *Traditional Medicinal Plants and Malaria*, pp. 256–268, CRC Press, Boca Raton, FL, USA, 2004.
- [13] D. Jebasingh, D. D. Jackson, S. Venkataraman, and B. S. Emerald, "Physiochemical and toxicological studies of the medicinal plant *Cyperus rotundus* L. (cyperaceae)," *International Journal of Applied Research in Natural Products*, vol. 5, no. 4, pp. 1–8, 2012.
- [14] A. Rajan, U. Bagai, and S. Chandel, "Effect of artesunate based combination therapy with homeopathic medicine China on liver and kidney of *Plasmodium berghei* infected mice," *Journal of Parasitic Diseases: Official Organ of the Indian Society for Parasitology*, vol. 37, no. 1, pp. 62–67, 2013.
- [15] L. Kunlawong, N. Ponwiang, U. Jaihan, S. Srichairatanakool, C. Uthaipibull, and V. Somsak, "Anti-hemolytic effects of green and black tea aqueous extracts on *Plasmodium berghei*-infected mice," *Advanced Natural Products*, vol. 1, pp. 1–6, 2014.
- [16] S. Kalia, U. Bagai, and V. Gorki, "Effect of *Albizia lebbeck* on liver and kidney function of *Plasmodium berghei* infected mice," *International Journal of Pharma Sciences and Research*, vol. 6, no. 3, pp. 575–581, 2015.
- [17] L. E. Kuatsienu, C. Ansah, and M. B. Adinortey, "Toxicological evaluation and protective effect of ethanolic leaf extract of *Launaea taraxacifolia* on gentamicin induced rat kidney injury," *Asian Pacific Journal of Tropical Biomedicine*, vol. 7, no. 7, pp. 640–646, 2017.
- [18] S. A. Tadesse and Z. B. Wubneh, "Antimalarial activity of *syzygium guineense* during early and established plasmodium infection in rodent models," *BMC Complementary and Alternative Medicine*, vol. 17, no. 1, p. 21, 2017.
- [19] A. L. M. Kognou, A. D. Tchamgoue, L. R. Y. Tchokouaha et al., "Acute and sub-chronic toxicity studies of *Dichaetanthera africana* (hook. F.) jacq. fel. (melastomataceae) stem bark ethanol extract," *Journal of Applied Pharmaceutical Science*, vol. 8, no. 6, pp. 147–155, 2018.
- [20] P. Chaniad, T. Techarang, A. Phuwanjaroanpong, and C. Punsawad, "Antimalarial activity and toxicological assessment of *Betula alnoides* extract against *Plasmodium berghei* infections in mice," *Evidence-Based Complementary and Alternative Medicine*, vol. 2019, Article ID 2324679, 8 pages, 2019.
- [21] X. Gu and J. E. Manautou, "Molecular mechanisms underlying chemical liver injury," *Expert Reviews in Molecular Medicine*, vol. 14, no. e4, p. e4, 2012.
- [22] J. Chiang, "Liver physiology: metabolism and detoxification," in *Pathophysiology of Human Diseases*, pp. 1770–1782, Elsevier, San Diego, CA, USA, 2014.
- [23] R. Chatterjee and A. Mitra, "An overview of effective therapies and recent advances in biomarkers for chronic liver diseases and associated liver cancer," *International Immunopharmacology*, vol. 24, no. 2, pp. 335–345, 2015.

- [24] U. Ilyas, D. P. Katare, V. Aeri, and P. P. Naseef, "A review on hepatoprotective and immunomodulatory herbal plants," *Pharmacognosy Reviews*, vol. 10, no. 19, pp. 66–70, 2016.
- [25] G. A. Josling and M. Llinás, "Sexual development in plasmodium parasites: knowing when it's time to commit," *Nature Reviews Microbiology*, vol. 13, no. 9, pp. 573–587, 2015.
- [26] B. S. Laurent, T. A. Burton, S. Zubaidah et al., "Host attraction and biting behaviour of *Anopheles* mosquitoes in south Halmahera, Indonesia," *Malaria Journal*, vol. 16, no. 1, p. 310, 2017.
- [27] R. S. Barsoum, "Malarial acute renal failure," *Journal of the American Society of Nephrology*, vol. 11, no. 11, pp. 2147–2154, 2000.
- [28] N. Mohandas and X. An, "Malaria and human red blood cells," *Medical Microbiology and Immunology*, vol. 201, no. 4, pp. 593–598, 2012.
- [29] National Institute of Allergy and Infectious Disease (NIAID), *Understanding Malaria Fighting on Ancient Scourge*, NIH Publication No. 07–7139, Department of Health and Human Services, National Institutes of Health, Bethesda, MD, USA, 2007.
- [30] R. Bandyopadhyay, S. Bandyopadhyay, and A. Dutta, "Sickle cell hepatopathy," *Indian Journal of Pathology & Microbiology*, vol. 51, no. 2, pp. 284–285, 2008.
- [31] O. Prommano, U. Chairsi, G. D. Turner et al., "A quantitative ultrastructural study of the liver and the spleen in fatal *falciparum* malaria," *The Southeast Asian Journal of Tropical Medicine and Public Health*, vol. 36, no. 6, pp. 1359–1370, 2005.
- [32] A. A. Olayode, D. A. Ofusori, T. A. B. Ogunniyi, and O. S. Sakarya, "Histomorphological studies of the liver of *Plasmodium*-infected albino mice after administration of aqueous leaf extract of *Mangifera indica* (Linn.)," *Anatomy*, vol. 9, no. 3, pp. 168–176, 2015.
- [33] S. Biswas, N. D. Reddy, B. S. Jayashree, and C. M. Rao, "Evaluation of novel 3-hydroxyflavone analogues as HDAC inhibitors against colorectal cancer," *Advances in Pharmacological Sciences*, vol. 2016, Article ID 4751806, 14 pages, 2018.
- [34] N. C. Nayak, S. A. Sathar, S. Mughal, S. Duttagupta, M. Mathur, and P. Chopra, "The nature and significance of liver cell vacuolation following hepatocellular injury—an analysis based on observations on rats rendered tolerant to hepatotoxic damage," *Virchows Archiv*, vol. 428, no. 6, pp. 353–365, 1996.
- [35] M. E. Guicciardi, H. Malhi, J. L. Mott, and G. J. Gores, "Apoptosis and necrosis in the liver," *Comprehensive Physiology*, vol. 3, no. 2, pp. 977–1010, 2013.
- [36] M. Krishna, "Patterns of necrosis in liver disease," *Clinical Liver Disease*, vol. 10, no. 2, pp. 53–56, 2017.
- [37] R. Naqvi, E. Ahmad, F. Akhtar, A. Naqvi, and A. Rizvi, "Outcome in severe acute renal failure associated with malaria," *Nephrology Dialysis Transplantation*, vol. 18, no. 9, pp. 1820–1823, 2003.
- [38] L. C. Koopmans, M. E. van Wolfswinkel, D. A. Hesselink et al., "Acute kidney injury in imported *Plasmodium falciparum* malaria," *Malaria Journal*, vol. 14, no. 523, p. 523, 2015.
- [39] E. O. Ogbadoyi and B. Gabi, "Assessment of renal function in malaria patients in Minna, north central Nigeria," *African Journal of Infectious Diseases*, vol. 1, no. 1, pp. 57–64, 2007.
- [40] H. A. Twajj, A. Kery, and N. K. Al-Khazraji, "Some pharmacological, toxicological and phytochemical investigations on *Centaurea phyllocephala*," *Journal of Ethnopharmacology*, vol. 9, no. 2-3, pp. 299–314, 1983.
- [41] Organization for Economic Co-operation and Development (Oecd), "Test no. 423: acute oral toxicity—acute toxic class method," *OECD Guidelines for the Testing of Chemicals, Section 4*, OECD Publishing, Paris, France, 2002.
- [42] M. Firdaus, M. Astawan, D. Muchtadi, T. Wresdiyati, S. Waspadji, and S. S. Karyono, "Acute oral toxicity of methanol extract from brown algae *Sargassum echinocarpum*," *Jurnal Pengolahan Hasil Perikanan Indonesia*, vol. 15, no. 2, pp. 148–155, 2012.
- [43] P. Kupradinun, A. Tepsuwan, N. Tanthasri et al., "Toxicity testing of flowers on neem tree (*Azadirachta indica* A. juss)," *The Thai Journal of Veterinary Medicine*, vol. 40, no. 1, pp. 47–55, 2010.
- [44] W. Peters, J. H. Portus, and B. L. Robinson, "The chemotherapy of rodent malaria, XXII," *Annals of Tropical Medicine and Parasitology*, vol. 69, no. 2, pp. 155–171, 1975.
- [45] K. N. Gibson-Corley, A. K. Olivier, and D. K. Meyerholz, "Principles for valid histopathologic scoring in research," *Veterinary Pathology*, vol. 50, no. 6, pp. 1007–1015, 2013.
- [46] F. Monji, H. Hossein Tehrani, Z. Halvaei, and S. Arbabi Bidgoli, "Acute and subchronic toxicity assessment of the hydroalcoholic extract of *Stachys lavandulifolia* in mice," *Acta Medica Iranica*, vol. 49, no. 12, pp. 769–775, 2011.
- [47] K. Dondee, P. Bootprom, B. Saiphet, P. Borkaew, C. Klubsri, and V. Somsak, "Antimalarial activities of *Moringa oleifera* leaf extract against *Plasmodium berghei* ANKA infection in ICR mice," *International Journal of Innovative Research in Medical Science*, vol. 1, no. 5, pp. 194–201, 2016.
- [48] T.-H. Chow, Y.-Y. Lin, J.-J. Hwang et al., "Therapeutic efficacy evaluation of 111In-labeled PEGylated liposomal vinorelbine in murine colon carcinoma with multimodalities of molecular imaging," *Journal of Nuclear Medicine*, vol. 50, no. 12, pp. 2073–2081, 2009.
- [49] P. T. Reeves, C. Roesch, and M. N. Raghnaill, *Drug Action And Pharmacodynamics* MSD Veterinary Manual, Kenilworth, NJ, USA, 2016.
- [50] M. L. Willcox and G. Bodeker, "Traditional herbal medicines for malaria," *BMJ*, vol. 329, no. 7475, pp. 1156–1159, 2004.
- [51] K. Spelman, "'Silver bullet' drugs vs. traditional herbal remedies: perspectives on malaria," *HerbalGram*, vol. 84, pp. 44–55, 2009.
- [52] "Charles river research models: BALB/c mouse hematology," 2019, [https://www.animalab.eu/sites/%20all/pliki/produkty-dopobrania/balb\\_c\\_mouse\\_clinical\\_%20pathology\\_data.pdf](https://www.animalab.eu/sites/%20all/pliki/produkty-dopobrania/balb_c_mouse_clinical_%20pathology_data.pdf).
- [53] S. N. H. Akil, "The clinical and histopathological aspect of the liver, lung, and kidney in malaria," *Qanun Medika—Medical Journal Faculty of Medicine Muhammadiyah Surabaya*, vol. 3, no. 2, pp. 123–135, 2019.
- [54] D. Larrey, "Epidemiology and individual susceptibility to adverse drug reactions affecting the liver," *Seminars in Liver Disease*, vol. 22, no. 2, pp. 145–155, 2002.

# Toxicological Evaluation and Protective Effects of Ethanolic Leaf Extract of *Cassia spectabilis* DC on Liver and Kidney Function of *Plasmodium berghei*-Infected Mice

## ORIGINALITY REPORT

17%

SIMILARITY INDEX

9%

INTERNET SOURCES

16%

PUBLICATIONS

0%

STUDENT PAPERS

## PRIMARY SOURCES

- |   |   |     |
|---|---|-----|
| 1 | <a href="http://www.freepatentsonline.com">www.freepatentsonline.com</a><br>Internet Source   | 1%  |
| 2 | Huacheng Yan, Lin Sun, Yingmeng Ni, Juan Du, Dong Liu, Ping Wang, Jin Cao, Guofang Xu, Yi Tao, Ranran Dai, Wei Tang. "Influence of effective omalizumab treatment on eosinophils in allergic asthma with comorbidities", Research Square Platform LLC, 2022<br>Publication                  | <1% |
| 3 | <a href="http://www.sciencegate.app">www.sciencegate.app</a><br>Internet Source   | <1% |
| 4 | Akalu Mesfin, Mirutse Giday, Abebe Animut, Tilahun Teklehaymanot. "Ethnobotanical study of antimalarial plants in Shinile District, Somali Region, Ethiopia, and in vivo evaluation of selected ones against <i>Plasmodium berghei</i> ", Journal of Ethnopharmacology, 2012<br>Publication | <1% |

---

5	<a href="http://www.asjp.cerist.dz">www.asjp.cerist.dz</a> Internet Source	<1 %
6	Sakaewan Ounjaijean, Voravuth Somsak. "Effect of Allicin and Artesunate Combination Treatment on Experimental Mice Infected with <i>Plasmodium berghei</i> ", <i>Veterinary Medicine International</i> , 2022 Publication	<1 %
7	Wanna Chaijaroenkul, Murad A. Mubarak, Stephen A. Ward, Kesara Na-Bangchang. "Metabolite footprinting of <i>Plasmodium falciparum</i> following exposure to <i>Garcinia mangostana</i> Linn. crude extract", <i>Experimental Parasitology</i> , 2014 Publication	<1 %
8	<a href="http://ff.unair.ac.id">ff.unair.ac.id</a> Internet Source	<1 %
9	<a href="http://uhra.herts.ac.uk">uhra.herts.ac.uk</a> Internet Source	<1 %
10	B.Parimala Devi, R. Boominathan, Subhash C. Mandal. "Evaluation of antipyretic potential of <i>Cleome viscosa</i> Linn. (Capparidaceae) extract in rats", <i>Journal of Ethnopharmacology</i> , 2003 Publication	<1 %
11	Gedefaw Getnet Amare, Tadesse Awgichew, Solomon Ahmed, Zemene Demelash Kifle.	<1 %

---

"Evaluation of in vitro and in vivo Antiplasmodial Activity of the Leaf Latex of Aloe Weloensis (Aloaceae)", Research Square, 2020

Publication

---

12

Olusegun Akanbi, Jamiu Saliu, Aderotimi Adejuyigbe, Tobiloba Adefala, Abel Goodluck, Damilola Adedoyin. "Combination Therapy of Methanolic Leaf Extracts of Anogeissus leiocarpus and Terminalia avicennioides on Malaria Parasite Count and Its Effect on Some Biochemical Parameters in Mice Infected with Plasmodium berghei", International Journal of TROPICAL DISEASE & Health, 2017

Publication

---

<1 %

13

[lupinepublishers.com](http://lupinepublishers.com)

Internet Source

---

<1 %

14

[www.plantphysiol.org](http://www.plantphysiol.org)

Internet Source

---

<1 %

15

R. Naqvi. "Outcome in severe acute renal failure associated with malaria", Nephrology Dialysis Transplantation, 09/01/2003

Publication

---

<1 %

16

Sureshkumar, S.V.. "Hepatoprotective effect of extracts from Pergularia daemia Forsk.", Journal of Ethnopharmacology, 20060919

Publication

---

<1 %

17

Wiwied Ekasari, Dwi Widya Pratiwi, Zelmira Amanda, Suciati, Aty Widyawaruyanti, Heny Arwati. " Various Parts of Plants as New Sources of Antimalarial Drugs ", Evidence-Based Complementary and Alternative Medicine, 2019

Publication

---

&lt;1 %

18

H Rahmania, R Permana, D Latipudin, N Suwarno, T Puspitasari, N Nuryanthi, A Mushawwir. "Enhancement of the liver status of Sentul chickens from the starter phase induced by irradiated chitosan", IOP Conference Series: Earth and Environmental Science, 2022

Publication

---

&lt;1 %

19

Mohanambal Moorthy, Joon Joon Khoo, Uma D. Palanisamy. "Acute oral toxicity of the ellagitannin geraniin and a geraniin-enriched extract from Nephelium lappaceum L rind in Sprague Dawley rats", Heliyon, 2019

Publication

---

&lt;1 %

20

[revistabionatura.com](http://revistabionatura.com)

Internet Source

---

&lt;1 %

21

Heny Arwati, Windya T. Hapsari, Kartika A. Wardhani, Kholida N. Aini, Ramadhani R. Bahalwan, Puspa Wardhani, Willy Sandhika. "Acute and subacute toxicity tests of goat bile in BALB/c mice", Veterinary World, 2020

&lt;1 %

22

Ibrahim Halil Bahcecioglu, Nalan Kuzu, Kerem Metin, Ibrahim Hanifi Ozercan, Bilal Ustündağ, Kazim Sahin, Omer Kucuk. "Lycopene Prevents Development of Steatohepatitis in Experimental Nonalcoholic Steatohepatitis Model Induced by High-Fat Diet", *Veterinary Medicine International*, 2010

Publication

---

<1 %

23

Lisa M. Weatherly, Hillary L. Shane, Ewa Lukomska, Rachel Baur, Stacey E. Anderson. "Systemic toxicity induced by topical application of heptafluorobutyric acid (PFBA) in a murine model", *Food and Chemical Toxicology*, 2021

Publication

---

<1 %

24

Muharni Muharni, Julinar Julinar, Heni Yohandini, Fitrya Fitrya, Rima Melati. "Antihyperlipidemia activity of ethanol extract the stem bark of *Flacourtia rukam* on propylthiouracil-induced albino rats, *Rattus noverticus* (Wistar strain)", *IOP Conference Series: Materials Science and Engineering*, 2021

Publication

---

<1 %

25

[cyberleninka.org](http://cyberleninka.org)  
Internet Source

<1 %

---

26 de Mesquita, M.L.. "In vitro antiplasmodial activity of Brazilian Cerrado plants used as traditional remedies", Journal of Ethnopharmacology, 20070301  
Publication <1 %

---

27 haematologica.org  
Internet Source <1 %

---

28 life.dysona.org  
Internet Source <1 %

---

29 Diorge Jônatas Marmitt, Mohamad Hesam Shahrajabian. "Plant species used in Brazil and Asia regions with toxic properties", Phytotherapy Research, 2021  
Publication <1 %

---

30 M.I. Akpanabiatu. "Biochemical and Histological Effects of Eleophorbia drupifera Leaf Extract in Wistar Albino Rats", Pharmaceutical Biology (Formerly International Journal of Pharmacognosy), 4/1/2003  
Publication <1 %

---

31 Narain Nutham, Sakuna Sakulmettatham, Suwit Klongthalay, Palatip Chutoam, Voravuth Somsak. " Protective Effects of Stem Extract on Renal Damage and Hemolysis during Infection in Mice ", Journal of Pathogens, 2015  
Publication <1 %

---



32

Omoboyowa Damilola A, Nwodo Fred, Joshua Parker Eli. "Effects of Combined Treatment of *Tithonia diversifolia* and Chloroquine on Selected Organs of Chloroquine Resistant *Plasmodium yeolii* Infected Mice", *Asian Journal of Biological Sciences*, 2018

Publication

<1 %

33

Prapaporn Chaniad, Tachpon Techarang, Arisara Phuwajaroanpong, Prasit Na-ek, Parnpen Viriyavejakul, Chuchard Punsawad. "In Vivo Antimalarial Activity and Toxicity Study of Extracts of *Tagetes erecta* L. and *Synedrella nodiflora* (L.) Gaertn. from the Asteraceae Family", *Evidence-Based Complementary and Alternative Medicine*, 2021

Publication

<1 %

34

[sphinxsai.com](http://sphinxsai.com)

Internet Source

<1 %

35

Carmen R. Silva-Correa, Víctor E. Villarreal-La Torre, Anabel D. González-Siccha, José L. Cruzado-Razco et al. "Acute toxicity of aqueous extract of *Ambrosia arborescens* Mill. on biochemical and histopathological parameters in rats", *Toxicological Research*, 2021

Publication

<1 %

36

Julia Penna Coutinho, Anna Caroline Campos Aguiar, Pierre Alexandre dos Santos, Joaquim Corsino Lima et al. "Aspidosperma (Apocynaceae) plant cytotoxicity and activity towards malaria parasites. Part I: Aspidosperma nitidum (Benth) used as a remedy to treat fever and malaria in the Amazon", Memórias do Instituto Oswaldo Cruz, 2013

Publication

&lt;1 %

37

Kenneth Kurt Booth, Edward Cottington Webb. "Effect of Blockage of the Ducts of the Vomeronasal Organ on LH Plasma Levels during the "Whitten Effect" in Does", Veterinary Medicine International, 2011

Publication

&lt;1 %

38

OLAYODE, Ahmed A., OFUSORI, David A., OGUNNIYI, Titus A. B. and SAKA, Olusola S.. "Histomorphological studies of the liver of Plasmodium-infected albino mice after administration of aqueous leaf extract of Mangifera indica (Linn.)", Deomed Publishing, 2015.

Publication

&lt;1 %

39

Prapaporn Chaniad, Tachpon Techarang, Arisara Phuwajaroanpong, Chuchard Punsawad. " Antimalarial Activity and Toxicological Assessment of Extract against

&lt;1 %

Infections in Mice ", Evidence-Based  
Complementary and Alternative Medicine,  
2019

Publication

---

40

Upma Singh, Ravi S. Pandey. "Fertilizer industry effluent induced hematological, histopathological and biochemical alterations in a stinging catfish, *Heteropneustes fossilis* (Bloch, 1794)", *Environmental and Sustainability Indicators*, 2021

Publication

---

<1 %

41

Xinling Song, Weijun Cui, Zheng Gao, Jianjun Zhang, Le Jia. "Structural characterization and amelioration of sulfated polysaccharides from *Ganoderma applanatum* residue against CCl<sub>4</sub>-induced hepatotoxicity", *International Immunopharmacology*, 2021

Publication

---

<1 %

42

[japsonline.com](http://japsonline.com)

Internet Source

---

<1 %

43

[media.neliti.com](http://media.neliti.com)

Internet Source

---

<1 %

44

Alozieuwa Blessing, Mann Abdulahi, Kabiru Yusuf, Ogbadoyi Olofu. "Antimalarial Activity of Crude Extract and Fractions of *Phyllanthus amarus* in *Plasmodium berghei*-Infected Mice", *European Journal of Medicinal Plants*, 2018

<1 %

45

Ernest Small. "Evolution and Classification of Cannabis sativa (Marijuana, Hemp) in Relation to Human Utilization", The Botanical Review, 2015

Publication

---

<1 %

46

Liliane Laure Toukam, Bertrand Tatsinkou Fossi, Germain Sotoing Taiwe, Raymond Bess Bila et al. "In vivo antimalarial activity of a probiotic bacterium Lactobacillus sakei isolated from traditionally fermented milk in BALB/c mice infected with Plasmodium berghei ANKA", Journal of Ethnopharmacology, 2021

Publication

---

<1 %

47

Muluken Adela Alemu, Yared Andargie Ferede, Getu Tesfaw Addis, Sintayehu Asnakew Alemayehu et al. "Antiplasmodial Activity of 80% Methanolic Extract and Solvent Fractions of Stem Bark of Acacia tortilis in Swiss Albino Mice", Evidence-Based Complementary and Alternative Medicine, 2022

Publication

---

<1 %

48

Ogbadoyi, EO, and B Gabi. "Assessment Of Renal Function In Malaria Patients In Minna, North Central Nigeria", African Journal of Infectious Diseases, 2008.

Publication

<1 %

49

Pang-Kuei Hsu, Yueh-Ting Tsai, Yu-Cheng Lin, Chen-Meng Kuan. "Assessment of the acute and sub-acute toxicity of the ethanolic extract of the aerial parts of *Crassocephalum rabens* (Asteraceae) in rats", *Toxicology Reports*, 2022

Publication

&lt;1 %

50

Razina Rouf, Puja Ghosh, Md. Raihan Uzzaman, Dipto Kumer Sarker, Fatima Tuz Zahura, Shaikh Jamal Uddin, Ilias Muhammad. "Hepatoprotective Plants from Bangladesh: A Biophytochemical Review and Future Prospect", *Evidence-Based Complementary and Alternative Medicine*, 2021

Publication

&lt;1 %

51

Xiaorong Li, Yongjiang Luo, Lijuan Wang, Yuhang Li, Yanbin Shi, Yin Cui, Ming Xue. "Acute and subacute toxicity of ethanol extracts from *Salvia przewalskii* Maxim in rodents", *Journal of Ethnopharmacology*, 2010

Publication

&lt;1 %

52

Xin Guan, Meng Gao, Hong Xu, Chenghong Zhang, Hongyan Liu, Li Lv, Sa Deng, Dongyan Gao, Yan Tian. " Quercetin-loaded poly (lactic-glycolic acid)-  $\alpha$ -tocopheryl polyethylene glycol 1000 succinate nanoparticles for the targeted treatment of liver cancer ", *Drug Delivery*, 2016

Publication

&lt;1 %

53

[academicworks.cuny.edu](http://academicworks.cuny.edu)

Internet Source

&lt;1 %

54

[journals.plos.org](http://journals.plos.org)

Internet Source

&lt;1 %

55

[www.scielo.br](http://www.scielo.br)

Internet Source

&lt;1 %

56

[www.scienceopen.com](http://www.scienceopen.com)

Internet Source

&lt;1 %

57

[www.tandfonline.com](http://www.tandfonline.com)

Internet Source

&lt;1 %

58

[www.wwarn.org](http://www.wwarn.org)

Internet Source

&lt;1 %

59

Evelin Fabjan, Etje Hulzebos, W. Mennes, A. H. Piersma. "A Category Approach for Reproductive Effects of Phthalates", *Critical Reviews in Toxicology*, 2008

Publication

&lt;1 %

60

John O. Olanlokun, Wisdom Oshireku Abiodun, Oluwakemi Ebenezer, Neil A. Koorbanally, Olufunso Olabode Olorunsogo. "Curcumin modulates multiple cell death, matrix metalloproteinase activation and cardiac protein release in susceptible and resistant *Plasmodium berghei*-infected mice", *Biomedicine & Pharmacotherapy*, 2022

Publication

&lt;1 %

61

Sherin Ramadan Hamad. "Possible ameliorative effect of nano-chocolate fortified with zinc on the hepatic and testicular toxicity induced by cadmium on male Swiss albino mice: histopathological and histomorphometrical studies", *The Journal of Basic and Applied Zoology*, 2019

Publication

&lt;1 %

62

Abrham Belachew Muluye, Eshetie Melese, Getnet Mequanint Adinew. "Antimalarial activity of 80% methanolic extract of *Brassica nigra* (L.) Koch. (Brassicaceae) seeds against *Plasmodium berghei* infection in mice", *BMC Complementary and Alternative Medicine*, 2015

Publication

&lt;1 %

63

Loubna Kharchoufa, Mohamed Bouhrim, Nouredine Bencheikh, Soufiane El Assri et al. "Acute and Subacute Toxicity Studies of the Aqueous Extract from *Haloxylon scoparium* Pomel (*Hammada scoparia* (Pomel)) by Oral Administration in Rodents", *BioMed Research International*, 2020

Publication

&lt;1 %

64

Muhydeen Abiodun Abdulraheem, Medard Ernest, Ifeoma Ugwuany, Saori Nishikawa et al. " High prevalence of and in co-infections with in asymptomatic malaria parasite

&lt;1 %

# carriers in Southwest Nigeria ", Cold Spring Harbor Laboratory, 2021

Publication

---

65

Sakaewan Ounjaijean, Chonticha Romyasamit, Voravuth Somsak. "Evaluation of Antimalarial Potential of Aqueous Crude *Gymnema Inodorum* Leaf Extract against *Plasmodium berghei* Infection in Mice", Evidence-Based Complementary and Alternative Medicine, 2021

<1 %

Publication

---

Exclude quotes On

Exclude matches Off

Exclude bibliography On



# Toxicological Evaluation and Protective Effects of Ethanolic Leaf Extract of *Cassia spectabilis* DC on Liver and Kidney Function of *Plasmodium berghei*-Infected Mice

---

GRADEMARK REPORT

---

FINAL GRADE

**/100**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---

PAGE 6

---

PAGE 7

---

PAGE 8

---

PAGE 9

---