

Nigella sativa toothpaste promotes anti-inflammatory and anti-destructive effects in a rat model of periodontitis

by Eka Fitria Augustina

Submission date: 04-Apr-2023 09:50AM (UTC+0800)

Submission ID: 2055173348

File name: in_a_rat_model_of_periodontitis_ARCHIVE_OF_ORAL_BIOLOGY_2022.pdf (2.62M)

Word count: 6416

Character count: 36565



Nigella sativa toothpaste promotes anti-inflammatory and anti-destructive effects in a rat model of periodontitis

Ernie Maduratna Setiawatie^{a,*}, Maria Apriliani Gani^b, Retno Puji Rahayu^c, Noer Ulfah^a, Shafira Kurnia^a, Eka Fitria Augustina^a, Desi Sandra Sari^d

^a Department of Periodontics, Faculty of Dental Medicine, Universitas Airlangga, Surabaya, East Java 60132, Indonesia

^b Doctoral Programme of Pharmaceutical Sciences, Faculty of Pharmacy, Universitas Airlangga, Surabaya, East Java 60115, Indonesia

^c Department of Oral and Maxillofacial Pathology, Faculty of Dental Medicine, Universitas Airlangga, Surabaya, East Java 60132, Indonesia

^d Department of Periodontics, Faculty of Dentistry, University of Jember, Jember, East Java 68121, Indonesia

ARTICLE INFO

Keywords:

Black cumin
Antibacterial toothpaste
Dental plaque
Porphyromonas gingivalis
Infectious disease

ABSTRACT

Objective: Periodontitis is an infectious disease that results in gingiva tissue damage. This study aimed to evaluate the effects of *Nigella sativa* (*N. sativa*) toothpaste in a periodontitis tissue repair based on inflammation and periodontal extracellular matrix in vivo.

Design: The periodontitis disease model was developed using Wistar rats infected with *Porphyromonas gingivalis* (*P. gingivalis*). The rats were divided into three main groups as follows: those that did not receive any toothpaste treatment; those that were treated with *N. sativa* toothpaste twice a day (simultaneously with *P. gingivalis* induction); and normal healthy rats. The rats were sacrificed after 1 and 7 days of animal modeling. The number of inflammatory cells, matrix metalloproteinase (MMP)1 + and MMP8 + cells, levels of cytokines (interleukin-1 β (IL-1 β) and prostaglandin E2 (PGE2)) and density of collagen type 1 were determined in the gingival tissues of the rats.

Results: The rats treated with *N. sativa* toothpaste had significantly lower numbers of neutrophils, macrophages and lymphocytes than the non-treated rats after 1 and 7 days of treatment; likewise, the levels of IL-1 β and PGE2 were lower in the treated experimental rats. In addition, the group treated with *N. sativa* toothpaste had fewer numbers of MMP1 + and MMP8 + cells and higher collagen density after 1 and 7 days of administration.

Conclusions: *N. sativa* toothpaste exhibited anti-inflammatory effects by reducing both inflammatory cell count and activity. Additionally, *N. sativa* toothpaste demonstrated anti-destructive effects on the periodontal extracellular matrix. Thus, *N. sativa* toothpaste might be potentially used for the management of periodontitis.

1. Introduction

Periodontitis is a multifactorial chronic illness induced by oral bacteria, characterized by a host-mediated inflammation in the periodontal tissues accompanied by dysbiotic plaque biofilms, which leads to the gradual degradation of the tooth-supporting apparatus and loss of periodontal attachment. It is one of the most common causes of tooth loss and can affect the masticatory efficiency, appearance, self-esteem and quality of life of the individual (Muñoz-Carrillo et al., 2019). Periodontal disease affects 20–50% of the population worldwide (Sanz,

D'aiuto, Deanfield, & Fernandez-Avilés, 2010). The average proportion increase in the global burden of periodontitis, oral cancer and dental caries from 1990 to 2010 was reported to be 45.6% (Jin et al., 2016). In developing countries such as Indonesia, the prevalence of periodontitis was expected to be higher than 70% (Susanto et al., 2011).

A variety of salivary and serum pro-inflammatory biomarkers were found in periodontitis patients. Isola, Polizzi, Santonocito, Alibrandi, and Williams (2021) found that an elevated serum and salivary Nod-like receptor family pyrin domain-containing protein-3 (NLRP3), a caspase recruitment domain (ASC), caspase-1, and interleukin (IL)-1 β were

Abbreviations: COL1, collagen type 1; IgA, immunoglobulin A; IHC, immunohistochemistry; IL-1 β , interleukin-1 β ; MMP1, matrix metalloproteinases 1; MMP8, matrix metalloproteinases 8; *N. sativa*, *Nigella sativa*; *P. gingivalis*, *Porphyromonas gingivalis*; PGE2, prostaglandin E2; TNF- α , tumor necrosis factor- α .

* Correspondence to: Faculty of Dental Medicine, Airlangga University, Prof. Dr. Moestopo Street, Surabaya, East Java 60132, Indonesia.

E-mail addresses: ernie-m-s@fkg.unair.ac.id (E.M. Setiawatie), maria.apriliiani.gani-2019@ff.unair.ac.id (M.A. Gani), a-retno-p-r@fkg.unair.ac.id (R.P. Rahayu), noer-u@fkg.unair.ac.id (N. Ulfah), shafira-k-s@fkg.unair.ac.id (S. Kurnia), eka-f-a@fkg.unair.ac.id (E.F. Augustina), desi_sari.fkg@unej.ac.id (D.S. Sari).

<https://doi.org/10.1016/j.archoralbio.2022.105396>

Received 7 July 2021; Received in revised form 3 February 2022; Accepted 6 March 2022

Available online 9 March 2022

0003-9969/© 2022 Elsevier Ltd. All rights reserved.

found in periodontitis patients. This was also similar in the IL-6 (Isola, Lo Giudice et al., 2021) and other pro-inflammatory cytokines levels (Liao, Fei, Shen, Yin, & Lu, 2014; Rangbulla, Nirola, Gupta, & Batra, 2017). It has been reported that recruitment in NLRP3-caspase 1 leads to the recruitment of inflammatory cells such as neutrophils (Strowig, Henao-Mejia, Elinav, & Flavell, 2012). The inflammatory cells then will release pro-inflammatory cytokines and mediators, such as tumor necrosis factor- α (TNF- α), IL1 β , and prostaglandin E2 (PGE2). Moreover, macrophages and lymphocytes contribute to the production of collagenases, including matrix metalloproteinases 1 (MMP1) and 8 (MMP8), which degrade the organic matrix in periodontal tissues (Kinney et al., 2014; Rangbulla et al., 2017). Other proteins such as Galectin-3 and asymmetric dimethylarginine (ADMA) also involved in periodontitis progression and its association with cardiovascular diseases (Isola, Polizzi, Alibrandi, Williams, & Lo Giudice, 2021; Isola et al., 2020). Considering their importance in periodontitis pathophysiology, inflammation factors are required as biomarkers for periodontitis.

The management of periodontitis varies from a lifestyle change, local and systemic pharmacotherapy, and surgery. A meta-analysis reported that scaling, root planing, and systemic antibiotic therapy led to beneficial clinical effects in aggressive periodontitis (Gonc, 2015). Even so, there are various side effects of antibiotics therapy in periodontitis, mostly related to microbiologically adverse effects and increase in bacterial resistance (Jepsen & Jepsen, 2016). Besides, regenerative surgical treatment such as bone graft is essential to replace the bone destroyed (Khotib et al., 2021). Despite its success in dental implantation, biomaterials used as bone grafts are also reported to cause plaque accumulation, gingival inflammation, and the potentially toxic effects caused by released components (Isola, 2021).

Periodontitis is caused by microbial biofilms formed by oral bacteria such as *Porphyromonas gingivalis* (*P. gingivalis*) that occur due to improper oral health (Mekhemar, Hassan, & Dörfer, 2020). Maintenance of oral hygiene by methods such as tooth brushing is an inexpensive but effective way to prevent and treat periodontitis (AlAttas, Zahran, & Turkistany, 2016). Nahak, Tedjasulaksana, and Raiyanti (2018) reported that tooth brushing with antibacterial toothpaste reduced the number of periodontal pathogens in the dental plaque. *Nigella sativa* (*N. sativa*) is a medicinal plant with a variety of pharmacological effects (Ahmad et al., 2013; Nurhan, Gani, Budiati, Siswodiardjo, & Khotib, 2021). A study by Senthilnathan, Ilango, Abiram, Vummidi, and Mahalingam (2020) showed that *N. sativa* had an antibacterial effect against periodontitis-causing bacteria, *P. gingivalis* and *Prevotella intermedia*. In addition, we have recently shown the non-toxic effects of toothpaste containing *N. sativa* and 2% sodium lauryl sulfate and non-sodium lauryl sulfate on fibroblasts (Setiawatie et al., 2021). These findings indicate that *N. sativa* can be used as the main ingredient in toothpastes for adjuvant therapy in periodontitis.

Considering the incidence of periodontitis, there should be an effective and inexpensive way to prevent and treat this disease. The antibacterial effect of *N. sativa* has been widely reported (Bakathir & Abbas, 2011; Salem et al., 2010; Senthilnathan et al., 2020). However, to the best of our knowledge, there is no information regarding the anti-inflammatory and anti-destructive effects of toothpaste containing *N. sativa*. Thus, this study aimed to evaluate the effects of toothpaste containing *N. sativa* in the repair periodontitis tissue based on the neutrophil, macrophage, and lymphocyte cell counts, collagen type 1 fibers (COL1) density, (IL)-1 β and PGE2 levels, as well as MMP1 + and MMP8 + cells in vivo. The results of this study will contribute in finding an effective and inexpensive ways in periodontitis management.

2. Materials and methods

2.1. Toothpaste fabrication

Na CMC (also known as sodium carboxymethyl cellulose salt) was placed in a beaker containing hot distilled water (80 °C) and stirred.

Next, cold distilled water was added to the Na CMC mixture and stirred until homogenous. Titanium dioxide was crushed in a mortar, mixed with CaCO₃ until homogeneous and placed in a bottle glass containing saccharine. This mixture was then combined with *N. sativa* extract and stirred until homogeneous. The Na CMC mucilage was added to the mixture and homogenized. Finally, sorbitol, aqua, silica, sodium monofluorophosphate, sodium benzoate, flavor, allantoin, methylparaben and CL 42090 (colorant) were added to the mixture and stirred to form a paste.

2.2. In vivo study

Thirty-six male Wistar rats (*Rattus norvegicus*) aged 5–6 months (bodyweight, 250–350 g) were adapted to the laboratory environment for at least 1 week and housed under standard laboratory conditions. All procedures conducted on animals were approved by the Health Research Ethical Clearance Commission at the Faculty of Dental Medicine, Universitas Airlangga (approval number: 564/HRECC.FODM/XII/2020). The periodontitis model was developed as described by Budhy, Arundina, Surboyo, and Halimah (2021). Briefly, *P. gingivalis* (Pg ATCC 33277 PK/5, Thermo Scientific; 1 × 10 colony-forming units in 20 μ l of PBS) was injected into the gingival sulcus between the two lower mandibular incisors using a 0.5-mL syringe. The injection was performed every 3 days for 14 days. The initial signs of periodontitis, which included a reddish pigmentation of the gingiva, swelling of the interdental incisive central mandibular area and interdental resorption (during histopathological investigation) were observed from the 14th day.

One milligram of *N. sativa* toothpaste was administered twice a day simultaneously with the induction of *P. gingivalis* in the gingival sulcus of the mandibular incisors. The animals were randomly divided into three main groups. The normal group were healthy controls, and the two remaining groups were used as periodontitis model groups. The periodontitis model groups were further subdivided into four groups as follows: PG day 1 group (the rats were sacrificed after day 1); PG + toothpaste day 1 group, which comprised rats that received *N. sativa* toothpaste and were sacrificed after day 1; PG day 7 group (the rats were sacrificed after day 7); and PG + toothpaste day 7 group, which comprised rats that received *N. sativa* toothpaste treatment and were sacrificed after day 7. All subgroups were contained five rats. The rats were euthanized using chloroform. The gingival tissues from the incisors to the molars were collected and fixed in 10% neutral buffered formalin for 3 days.

2.3. Histological staining

Histological staining was performed to determine the number of inflammatory cells and the collagen density. The tissue samples were implanted in a paraffin block. The paraffin blocks were made by dehydrating the tissue in various grades of ethanol (70–100%, 60 min each). The tissues were then washed with xylol (3 times, 15 min each), followed by liquid paraffin infiltration and transfer (3 times, 60 min each in a 60 °C incubator). A rotary microtome was used to cut the paraffin blocks (thickness, 4–6 μ m). The slides were stained with haematoxylin-eosin (Budiati et al., 2021) and Masson trichrome, according to the standard procedures.

2.4. Immunohistochemistry

The expression of MMP1 and MMP8 was investigated via immunohistochemistry (IHC). The paraffin blocks were proceed the same as in the histological staining. Briefly, after being implanted in paraffin blocks, the gingiva tissue was dehydrated and washed with xylol. Moreover, the samples were infiltrated, transferred, and cut into paraffin blocks. The IHC procedure was conducted as described by Budiati et al. (2021). Briefly, the histological slide was deparaffinized

using xylol (3 times, 5 min), followed by rehydration using alcohol (absolute to 70%, 4 min each) and rinsed under running water (5 min). The slide was blocked using endogenous peroxidase 0.5% (5 min) and rinsed with running water (5 min). Antigen retrieval was performed in a de-cloaking chamber, following which, the specimen was cooled for

approximately 20 min, washed with PBS (3 min) and snipper-blocked for 15 min. The slide was then incubated with rabbit anti-rat MMP1 (BS-0463R, Bioss, Massachusetts, USA, 1:250 dilution) or rabbit anti-rat MMP8 (orb214259, Biorbyt, Cambridge, UK, 1:250 dilution) for 60 min and washed with PBS for 3 min. Moreover, the Trekkie Universal link




Fig. 1. Representative histological images of gingiva tissues stained with haematoxylin-eosin (total magnification, 100× and 400×), lymphocyte (red arrow), macrophage (yellow arrow), and neutrophil (black arrow) (a). Histophotometric quantification of neutrophils, macrophages and lymphocytes. Each bar represents the mean cell count ± standard deviation (SD) of five different visual fields in five rats. **p < 0.01, ***p < 0.001 based on the one-way ANOVA test (b).

was carried out on the slide (20 min), and the slide was rewashed with PBS for 3 min. Trekavidin-HRP Label was applied to the slide (10 min); following which, the specimen was washed with PBS (3 min). The slide was reacted with chromogen DAB + buffer substrate (2–5 min) and washed with running water (5 min). After counterstaining with haematoxylin (1–2 min) and washing under running water (2 times, 5 min each), the slide was dehydrated with alcohol (70% to alcohol absolute, 5 min each), cleared in xylol (3 times, 5 min each), mounted (Ecomount) and covered with a coverslip.

2.5. Histophotometry

Quantitative assessment of inflammatory cells in the gingival connective tissue on the lingual side was performed by an investigator blinded to the animal groups as described by Sha, Garib, Azeez, and Gul (2020). A light microscope with an image analyzing system (Motic, ToupTek, ToupView) was used to perform the quantification (magnification, 400 ×). A grid of 16 squares was used to split each collected image. The mean ± standard deviation (SD) values of the neutrophils, macrophages and lymphocytes were determined. The density of the collagen fibers (%) was determined by calculating the area formed by the collagen fibers at the healing center and dividing it by the total area of measurement (Budi & Astuti, 2019). On the basis of the IHC results, the mean percentage ± SD values of immunopositive cells expressing MMP1 and MMP8 were determined.

2.6. Enzyme-linked immunosorbent assay

Blood (3 mL) was collected from each rat via intracardiac puncture before sacrifice, transferred to empty tubes and centrifuged for 10 min at 3000 rpm. The obtained serum sample was used to measure the levels of PGE2 and IL-1β. The procedures were conducted using the Prostaglandin E2 Assay Kit (PKGE004B, R&D Systems, Minneapolis, MN, USA) and Rat IL-1β/IL-1F2 Immunoassay Kit (RLB00, R&D Systems, Minneapolis, MN, USA), according to the manufacturers' instructions.

2.7. Statistical analysis

One-way analysis of variance was used to analyze the data; the significance level was set at 0.05. The statistical tests were performed using SPSS version 24.0 (IBM Corporation, Armonk, NY, USA).

3. Results

The numbers of neutrophils, macrophages and lymphocytes were qualitatively (Fig. 1a) and quantitatively ($p < 0.001$; Normal vs. PG Day 1, Normal vs. PG Day 7; Fig. 1b) increased after induction with *P. gingivalis*. Furthermore, induction with *P. gingivalis* decreased the COL1 density ($p < 0.001$, Normal vs. PG Day 1, Normal vs. PG Day 7; Fig. 2). The number of neutrophils was decreased after days 1 and 7 of treatment with *N. sativa* toothpaste ($p < 0.001$; Fig. 1b). Likewise, the administration of *N. sativa* toothpaste resulted in a decrease in macrophage cell counts after the first and seventh days of treatment ($p < 0.001$; Fig. 1b). In the case of the lymphocytes, a decrease in number was noted after the seventh day of treatment compared to the group without treatment ($p < 0.001$; Fig. 1b); this value was not significantly different from that in the normal group ($p = 0.125$; Normal vs. PG Day 1; Fig. 1b). Significant increases in the density of COL1 were observed after days 1 and 7 of using *N. sativa* toothpaste ($p < 0.001$; Fig. 2).

Furthermore, there was a significant increase in IL-1β serum concentration in the PG Day 1 and PG Day 7 groups compared with those in the normal group ($p < 0.001$; Fig. 3a). This indicated that the periodontitis model was successful based on the pro-inflammatory cytokine levels. However, treatment with the *N. sativa* toothpaste decreased the IL-1β serum concentration at both time points ($p < 0.01$, Normal vs. PG + Toothpaste Day 1, Normal vs. Toothpaste Day 7; Fig. 3a). Similarly, P.




Fig. 2. Density of COL1 (%) based on Masson trichrome staining. Each bar represents the mean ± SD of five rats. *** $p < 0.001$, one-way ANOVA.

gingivalis induction increased PGE2 serum concentration in the PG Day 1 and PG Day 7 groups compared with those in the normal group ($p < 0.001$; Fig. 3b), whereas treatment with *N. sativa* toothpaste, decreased PGE2 serum concentration at both time points ($p < 0.01$, Normal vs. PG + Toothpaste Day 1, Normal vs. Toothpaste Day 7; Fig. 3b).

The administration of *P. gingivalis* increased the number of MMP1 + and MMP8 + cells in the PG Day 1 and PG Day 7 groups compared with that in the normal group, both qualitatively (Fig. 4a) and quantitatively ($p < 0.001$; Fig. 4b), whereas treatment with *N. sativa* toothpaste decreased the cell percentage at both time points ($p < 0.01$, Normal vs. PG + Toothpaste Day 1, Normal vs. Toothpaste Day 7; Fig. 4a).

4. Discussion

The current study evaluated the effects of *N. sativa* toothpaste in rat model of periodontitis induced by *P. gingivalis*. The current study found that *N. sativa* decreased the number of neutrophils, macrophages, and lymphocytes in periodontal tissue, as well as IL-1β and PGE2 serum concentration. Likewise, the *N. sativa* toothpaste decreased MMP1 + and MMP8 + cells and increased COL1 density in the affected tissue.

N. sativa belongs to the Ranunculaceae family and is a plant with a long history of medicinal use. It is also known as black seed and is often found in Southern Europe, North Africa and Southwest Asia. This plant has a wide range of pharmacological effects, including antihypertensive, analgesic, anti-inflammatory, antibacterial and antioxidant properties (Ahmad et al., 2013). It is widely known for its antibacterial properties. One study reported that 300 mg/mL of *N. sativa* inhibited the growth of *Staphylococcus aureus*, as demonstrated by the presence of a clear zone from the paper disc diffusion method (Bakathir & Abbas, 2011). This antibacterial effect was observed in a clinical study, where the combination of 2 g of *N. sativa* seeds with 40 mg of omeprazole in patients with non-ulcer dyspepsia beneficially eliminated *Helicobacter pylori* when compared with a triple-drug therapy using clarithromycin, amoxicillin and omeprazole (Salem et al., 2010). The amount of thymoquinone, the active component, ranges from 30% to 48% in the *N. sativa* seeds (Ahmad et al., 2013; Mekhemar et al., 2020). Chaieb, Kouidhi, Jrah, Mahdouani, and Bakhrouf (2011) reported that thymoquinone inhibited the formation of biofilms by various bacteria with a minimum inhibitory




Fig. 3. Levels of IL-1 β (a) and PGE2 (b) in rat serum. Each bar represents the mean \pm SD in five rats.

concentration ranging from 8 to 32 g/mL. Because of its potent antibacterial activity, *N. sativa* is potentially used as a therapeutic agent or adjuvant in bacterial infections.

Periodontal disease, the most common cause of tooth loss, is a chronic inflammatory disease of the periodontium characterized by the loss of periodontal ligaments and the destruction of surrounding alveolar bone in its advanced stages. Periodontal disease activates inflammatory and immune mechanisms that play a role in the development of systemic disease (Wada & Kamisaki, 2010). Moreover, periodontitis is defined as an inflammatory disease caused by bacteria that destroy the gingiva around the teeth (AlAttas et al., 2016). Inflammation in periodontitis is triggered by organisms in oral microbial biofilms, particularly gram-negative anaerobes such as *P. gingivalis* (Mekhemar et al., 2020). The imbalance between microbial communities resulted in the host's chronic inflammatory response. Induction by *P. gingivalis* linked to the activation of the inflammasome, mainly the NLRP3. Activation of NLRP3 cleaves the pro-form of inflammatory cytokines, such as IL-1 β , to the mature form with the help of caspase-1. Activation of caspase-1 also led to the death of osteoprogenitors and osteoclastic bone resorption. *P. gingivalis* activates the NLRP3 inflammasome, which may cause long-term inflammation and promote atherosclerosis and other periodontal diseases (Ding et al., 2020; Rocha et al., 2020).

Other proteins in periodontitis also have been suggested as independent risk factors of cardiovascular diseases (Isola, Polizzi, Alibrandi et al., 2021; Isola et al., 2020). Higher serum and salivary levels of Gal-3 and ADMA were found in periodontitis patients with coronary heart disease (Isola, Polizzi, Alibrandi et al., 2021; Isola et al., 2020). Gal-3 belongs to the β -galactoside-binding lectin family released from inflammatory cells (Velickovic et al., 2021). Moreover, ADMA is an endogenous inhibitor of NO synthase linked to endothelial dysfunction (Okada et al., 2021). This pathological mechanism elevated serum and salivary NLRP in periodontitis patients (Isola, Polizzi, Santonocito et al., 2021), which also positively correlated with caspase-1, IL-1 β , IL-18, Gal-3, and ADMA levels (Bostanci et al., 2009; Isola, Polizzi, Alibrandi et al., 2021; Isola, Polizzi, Santonocito et al., 2021; Okada et al., 2021). Because of this, patients with periodontitis have a higher risk of systemic inflammation (Hasturk & Kantarci, 2015).

The periodontitis disease model was successfully conducted in this study. The presence of some humoral and cellular makers could be considered as a primary sign of a successful periodontitis model. Li, Zhang, and Wang (2020) reported an increase in the number of inflammatory cells, particularly neutrophils, in chronic periodontitis tissues compared with that in healthy periodontal tissues. Increases in the levels of cytokines and chemokines, such as immunoglobulin A (IgA), IL-1 β , TNF- α , and PGE2, have also been reported in tissues affected by

periodontitis (Liao et al., 2014; Rangbulla et al., 2017). In the current study, an increase in the number of neutrophils, macrophages and lymphocytes was observed in the gingival tissue specimens of the rats. The induction of periodontitis following the application of bacteria increased the levels of the inflammatory factors IL-1 β and PGE2 in the peripheral blood of the experimental animals, thus indicating the development of innate inflammation in the experimental animals.

Periodontitis is characterized by the presence of gingival tissue destruction (Kawai et al., 2006), which is marked by a decrease in the density of abundant proteins, such as COL1, in the periodontal extracellular matrix (Gursoy et al., 2013). A decrease in COL1 is a result of an increase in the levels of collagenase enzymes, such as MMP8 and MMP1, in the tissue (Kinney et al., 2014; Rangbulla et al., 2017). In the current study, we showed that rats with periodontitis presented with increased numbers of MMP1 + and MMP8 + cells. MMP1 is synthesized by macrophages (Romanelli et al., 1999), whereas MMP8 is a neutrophil-driven collagenase (Gursoy et al., 2013). The increase in MMP1 and MMP8 levels in the current study was the result of an increase in the number of neutrophils and macrophages, which led to the damage of the periodontal extracellular matrix resulting in a reduction in the density of the collagen fibers.

The current study showed that *N. sativa* toothpaste treatment can reduce the number of neutrophils, macrophages, lymphocytes, IL-1 β and PGE2 levels. Neutrophils, the most abundant white blood cells, are important in the innate immune system. They function as phagocytes to kill extracellular pathogens; similarly, macrophages are phagocytic cells that help promote tissue healing (Scott & Krauss, 2011). Moreover, lymphocytes are responsible for the adaptive immune response (Hajishengallis & Korostoff, 2017). They secrete cytokines and mediators of inflammation, including TNF- α , IL-1 β and PGE2, which initiate the inflammation cascade and result in chronic inflammation (Kinney et al., 2014; Liao et al., 2014; Rangbulla et al., 2017). The decreased number of inflammatory cells and mediators in the current study proved that the administration of *N. sativa* toothpaste can treat acute inflammation and prevent chronic inflammation in the animal model of periodontitis.

COL1 is the most abundant protein in the periodontal extracellular matrix (Gursoy et al., 2013; Zhang, Li, Yan, & Huang, 2018). MMP1 and MMP8 are collagenases and members of the MMP family. They are reported to degrade the extracellular matrix in periodontal disease (Romanelli et al., 1999). The current study showed that the administration of *N. sativa* toothpaste increased COL1 density and decreased MMP1 + and MMP8 + cells in rats with periodontitis, thus exhibiting the anti-destructive effects of *N. sativa*.

The anti-inflammatory and anti-destructive activity of *N. sativa* may




Fig. 4. MMP1 and MMP8 immunohistochemical staining in the gingival tissues (magnification, 400× and 1000×), MMP1 + cells (red arrow), MMP8 + cells (yellow arrow) (a). Percentage of MMP1 and MMP8 immunopositive cells. Each bar represents the mean percentage of immunopositive cells ± SD observed in 10 different visual fields in five rats. ***p < 0.001, one-way ANOVA (b).

be due to the presence of the active compound thymoquinone. Thymoquinone shows the major pharmacological actions of *N. sativa*, including anti-inflammatory, antioxidant, antibacterial, analgesic, as well as hypoglycemic, and anticarcinogenic (Ahmad et al., 2013; Chaieb et al., 2011; Mekhemar et al., 2020). Moreover, thymoquinone was reported to reduce several pro-inflammatory cytokines, such as ILs, TNF-α (Shaterzadeh-Yazdi, Noorbakhsh, Hayati, Samarghandian, & Far-khondeh, 2018) and MMP8 (Elamrousy, 2021). Furthermore, thymoquinone demonstrated its antibacterial effect by inhibiting biofilm formation by *Fusobacterium nucleatum* and *P. gingivalis* (Tantivitayakul, Kaypetch, & Muadchiengka, 2020). Thus, taken together, these findings

indicate that *N. sativa* possesses antibacterial, anti-inflammatory and anti-destructive properties, particularly in tissues with periodontitis.

5. Conclusions

The effects of using toothpaste containing *N. sativa* on an animal model of periodontitis were examined in this study. The *N. sativa* toothpaste exhibited anti-inflammatory effects by reducing the numbers of neutrophils, macrophages and lymphocytes in the rat tissues. *N. sativa* also reduced inflammatory cell activity by decreasing the levels of IL-1β and PGE2. Furthermore, *N. sativa* toothpaste demonstrated anti-

destructive effects on the periodontal extracellular matrix by decreasing the levels of MMP1 and MMP1 and increasing COL1 density. Thus, *N. sativa* toothpaste can be potentially used as a therapeutic agent or adjuvant for the management of periodontitis. However, further studies in humans are required to confirm these findings.

Funding

This work was supported by the Ministry of Finance of the Republic of Indonesia, through RISPRO research schemes [grant number PRJ-49/LPDP/2019].

CRediT authorship contribution statement

Ernie Maduratna Setiawatie: Conceptualization, Data curation, Funding acquisition, Investigation, Methodology, Resources, Supervision, Writing – review & editing. **Maria Apriliani Gani:** Conceptualization, Formal analysis, Software, Validation, Visualization, Roles/ Writing – original draft. **Retno Puji Rahayu:** Conceptualization, Formal analysis, Investigation, Methodology. **Noer Ulfah:** Resources, Supervision, Validation. **Shafira Kurnia:** Investigation, Project administration, Validation, Writing – review & editing. **Eka Fitria Augustina:** Investigation, Project administration, Writing – review & editing. **Desi Sandra Sari:** Investigation, Project administration, Writing – review & editing.

Acknowledgements

The authors thank the Faculty of Dental Medicine, Universitas Airlangga, Surabaya, Indonesia, for the support during the study.

Declaration of Competing Interest

None.

References

- Ahmad, A., Husain, A., Mujeeb, M., Khan, S. A., Najmi, A. K., Siddique, N. A., ... Anwar, F. (2013). A review on therapeutic potential of *Nigella sativa*: A miracle herb. *Asian Pacific Journal of Tropical Biomedicine*, 3(5), 337–352. [https://doi.org/10.1016/S2221-1691\(13\)60075-1](https://doi.org/10.1016/S2221-1691(13)60075-1)
- AlAttas, S. A., Zahran, F. M., & Turkistany, S. A. (2016). *Nigella sativa* and its active constituent thymoquinone in oral health. *Saudi Medical Journal*, 37, 235–244. <https://doi.org/10.15537/smj.2016.3.13006>
- Bakathir, H. A., & Abbas, N. A. (2011). Detection of the antibacterial effect of *Nigella sativa* ground seeds with water. *African Journal of Traditional, Complementary and Alternative Medicines*, 8, 159–164. <https://doi.org/10.4314/ajtcam.v8i2.63203>
- Bostanci, N., Emingil, G., Saygan, B., Turkoglu, O., Atilla, G., Curtis, M. A., & Belibasakis, G. N. (2009). Expression and regulation of the NALP3 inflammasome complex in periodontal diseases. *Clinical and Experimental Immunology*, 157(3), 415–422. <https://doi.org/10.1111/j.1365-2249.2009.03972.x>
- Budhy, T. I., Arundina, I., Surboyo, M. D. C., & Halimah, A. N. (2021). The effects of rice husk liquid smoke in porphyromonas gingivalis-induced periodontitis. *European Journal of Dentistry*. <https://doi.org/10.1055/s-0041-1727554>
- Budi, H. S., & Astuti, E. R. (2019). The MMP-2, MMP-9 expression and collagen density of the ambonese banana stem sap administration on wound healing. *Journal of International Dental and Medical Research*, 12, 492–497.
- Budiati, A. S., Gani, M. A., Samirah, Ardianto, C., Raharjanti, A. M., Septiani, I., ... Khotib, J. (2021). Bovine hydroxyapatite-based bone scaffold with gentamicin accelerates vascularization and remodeling of bone defect. *International Journal of Biomaterials*, 2021, 1–7. <https://doi.org/10.1155/2021/5560891>
- Chaieb, K., Kouidhi, B., Jrah, H., Mahdouani, K., & Bakhrouf, A. (2011). Antibacterial activity of Thymoquinone, an active principle of *Nigella sativa* and its potency to prevent bacterial biofilm formation. *BMC Complementary and Alternative Medicine*, 11, 29. <https://doi.org/10.1186/1472-6882-11-29>
- Ding, P.-H., Yang, M.-X., Wang, N.-N., Jin, L.-J., Dong, Y., Cai, X., & Chen, L.-L. (2020). Porphyromonas gingivalis-induced NLRP3 inflammasome activation and its downstream interleukin-1 β release depend on caspase-4. *Frontiers in Microbiology*, 11. <https://doi.org/10.3389/fmicb.2020.01881>
- Elamrousy, W. (2021). Effect of locally delivered 0.2% Thymoquinone gel in subjects with grade C periodontitis: Clinical, immunological and microbiological assessment. *European Journal of Molecular & Clinical Medicine*, 05, 89–101. <https://doi.org/10.31838/ejmcm.05.01.11>
- Gonc, C. (2015). Systemic antibiotics in the treatment of aggressive periodontitis. A systematic review and a Bayesian Network. *Journal of Clinical Periodontology*, 42, 647–657. <https://doi.org/10.1111/jcpe.12427>
- Gursoy, U. K., Könönen, E., Huuononen, S., Tervahartala, T., Pussinen, P. J., Suominen, A. L., & Sorsa, T. (2013). Salivary type I collagen degradation end-products and related matrix metalloproteinases in periodontitis. *Journal of Clinical Periodontology*, 40, 18–25. <https://doi.org/10.1111/jcpe.12020>
- Hajishengallis, G., & Korostoff, J. M. (2017). Revisiting the Page & Schroeder model: The good, the bad and the unknowns in the periodontal host response 40 years later. *Periodontology* 2000, 75, 116–151. <https://doi.org/10.1111/prd.12181>
- Hasnurk, H., & Kantarci, A. (2015). Activation and resolution of periodontal inflammation and its systemic impact. *Periodontology* 2000, 69(1), 255–273. <https://doi.org/10.1111/prd.12105>
- Isola, G. (2021d). Interface between periodontal tissues and dental materials: Dynamic changes and challenges. *Coatings*, 11(5). <https://doi.org/10.3390/coatings11050485>
- Isola, G., Alibrandi, A., Currò, M., Matarese, M., Ricca, S., Matarese, G., & Kocher, T. (2020). Evaluation of salivary and serum asymmetric dimethylarginine (ADMA) levels in patients with periodontal and cardiovascular disease as subclinical marker of cardiovascular risk. *Journal of Periodontology*, 91(8), 1076–1084. <https://doi.org/10.1002/JPER.19-0446>
- Isola, G., Lo Giudice, A., Polizzi, A., Alibrandi, A., Murabito, P., & Indelicato, F. (2021b). Identification of the different salivary interleukin-6 profiles in patients with periodontitis: A cross-sectional study. *Archives of Oral Biology*, 122(November 2020), Article 104997. <https://doi.org/10.1016/j.archoralbio.2020.104997>
- Isola, G., Polizzi, A., Alibrandi, A., Williams, R. C., & Lo Giudice, A. (2021c). Analysis of galectin-3 levels as a source of coronary heart disease risk during periodontitis. *Journal of Periodontal Research*, 56(3), 597–605. <https://doi.org/10.1111/jre.12860>
- Isola, G., Polizzi, A., Santonocito, S., Alibrandi, A., & Williams, R. C. (2021a). Periodontitis activates the NLRP3 inflammasome in serum and saliva. *Journal of Periodontology*. <https://doi.org/10.1002/JPER.21-0049>
- Jepsen, K., & Jepsen, S. (2016). Antibiotics/antimicrobials: Systemic and local administration in the therapy of mild to moderately advanced periodontitis. *Periodontology* 2000, 71(1), 82–112. <https://doi.org/10.1111/prd.12121>
- Jin, L. J., Lamster, I. B., Greenspan, J. S., Pitts, N. B., Scully, C., & Wamakulasuriya, S. (2016). Global burden of oral diseases: Emerging concepts, management and interplay with systemic health. *Oral Diseases*, 22, 609–619. <https://doi.org/10.1111/odi.12428>
- Kawai, T., Matsuyama, T., Hosokawa, Y., Makihira, S., Seki, M., Karimbux, N. Y., ... Taubman, M. A. (2006). B and T lymphocytes are the primary sources of RANKL in the bone resorptive lesion of periodontal disease. *American Journal of Pathology*, 169, 987–998. <https://doi.org/10.2353/ajpath.2006.060180>
- Khotib, J., Gani, M. A., Budiati, A. S., Lestari, M. L. A. D., Rahadiansyah, E., & Ardianto, C. (2021). Signaling pathway and transcriptional regulation in osteoblasts during bone healing: Direct involvement of hydroxyapatite as a biomaterial. *Pharmaceuticals*, 14(7), 615. <https://doi.org/10.3390/ph14070615>
- Kinney, J. S., Morelli, T., Oh, M., Braun, T. M., Ramseier, C. A., Sugai, J. V., & Giannobile, W. V. (2014). Crevicular fluid biomarkers and periodontal disease progression. *Journal of Clinical Periodontology*, 41, 113–120. <https://doi.org/10.1111/jcpe.12194>
- Li, W., Zhang, Z., & Wang, Z. M. (2020). Differential immune cell infiltrations between healthy periodontal and chronic periodontitis tissues. *BMC Oral Health*, 20, 1–10. <https://doi.org/10.1186/s12903-020-01287-0>
- Liao, C.-H., Fei, W., Shen, Z.-H., Yin, M.-P., & Lu, C. (2014). Expression and distribution of TNF- α and PGE2 of periodontal tissues in rat periodontitis model. *Asian Pacific Journal of Tropical Medicine*, 7, 412–416. [https://doi.org/10.1016/S1995-7645\(14\)60067-5](https://doi.org/10.1016/S1995-7645(14)60067-5)
- Mekhemar, M., Hassan, Y., & Dörfer, C. (2020). *Nigella sativa* and thymoquinone: A natural blessing for periodontal therapy. *Antioxidants*, 9, 1–19. <https://doi.org/10.3390/antiox9121260>
- Muñoz-Carrillo, J. L., Hernández-Reyes, V. E., García-Huerta, O. E., Chávez-Ruvalcaba, F., Chávez-Ruvalcaba, M. I., Chávez-Ruvalcaba, K. M., & Díaz-Alfaro, L. (2019). Pathogenesis of periodontal disease. *Periodontal diseases - A clinician's guide*. Intech Open. <https://doi.org/10.5772/1274>
- Nahak, M. M., Tedjasulaksana, R., & Raiyanti, I. G. A. (2018). Effectiveness of using toothpaste and mouthwash of beluntas leaf ethanol extract in reducing *Streptococcus mutans* bacteria number in tooth plaque. *Journal of International Dental and Medical Research*, 11, 830–833.
- Nurhan, A. D., Gani, M. A., Budiati, A. S., Siswodiwardjo, S., & Khotib, J. (2021). Molecular docking studies of *Nigella sativa* L and Curcuma xanthorrhiza Roxb secondary metabolites against histamine N-methyltransferase with their ADMET prediction. *Journal of Basic and Clinical Pharmacy and Pharmacology*, 32(4), 795–802. <https://doi.org/10.1515/jbc>
- Okada, A., Murata, T., Matin, K., Ariyoshi, M., Otsuka, R., Yamashita, M., & Hanada, N. (2021). Effect of advanced periodontal self-care in patients with early-stage periodontal diseases on endothelial function: An open-label, randomized controlled trial. *PLoS One*, 16(9 September), 1–13. <https://doi.org/10.1371/journal.pone.0257247>
- Rangbulla, V., Nirola, A., Gupta, M., & Batra, P. (2017). Salivary IgA, interleukin-1 β and MMP-8 as salivary biomarkers in chronic periodontitis patients. *The Chinese Journal of Dental Research*, 20, 43–51. <https://doi.org/10.3290/j.cjdr.a37741>
- Rocha, F. R. G., Delitto, A. E., de Souza, J. A. C., González-Maldonado, L. A., Wallet, S. M., & Rossa Junior, C. (2020). Relevance of caspase-1 and Nlrp3 inflammasome on inflammatory bone resorption in a murine model of periodontitis. *Scientific Reports*, 10(1), 1–11. <https://doi.org/10.1038/s41598-020-64685-y>
- Romanelli, R., Mancini, S., Laschinger, C., Overall, C. M., Sodek, J., & McCulloch, C. A. G. (1999). Activation of neutrophil collagenase in periodontitis. *Infection and Immunity*, 67, 2319–2326. <https://doi.org/10.1128/iai.67.5.2319-2326.1999>

- Salem, E. M., Yar, T., Bamosa, A. O., Al-Quorain, A., Yasawy, M. I., Alsulaiman, R. M., & Randhawa, M. A. (2010). Comparative study of *Nigella sativa* and triple therapy in eradication of *Helicobacter Pylori* in patients with non-ulcer dyspepsia. *Saudi Journal of Gastroenterology*, *16*, 207–214. <https://doi.org/10.4103/1319-3767.65201>
- Sanz, M., D'aiuto, F., Deanfield, J., & Fernandez-Avilés, F. (2010). European workshop in periodontal health and cardiovascular disease - Scientific evidence on the association between periodontal and cardiovascular diseases: A review of the literature. *European Heart Journal, Supplement*, *12*, B3–B12. <https://doi.org/10.1093/eurheartj/suq003>
- Scott, D. A., & Krauss, J. L. (2011). Neutrophils in periodontal inflammation. *Periodontal Disease*, *15*, 56–83. <https://doi.org/10.1159/000329672>
- Senthilnathan, K., Ilango, P., Abiram, T., Vunmudi, V. A., & Mahalingam, A. P. (2020). Evaluation of antibacterial activity of *Nigella sativa* seed extract against *Porphyromonas gingivalis* and *Prevotella intermedia*. *Journal of Interdisciplinary Dentistry*, *10*, 51–55. <https://doi.org/10.4103/jid.jid>
- Setiawatie, E. M., Sari, D. S., Wahyudadi, B. S., Fitriya, E., Kurnia, S., Bargowo, L., & Gani, M. A. (2021). Viability of *Nigella sativa* toothpaste with SLS compared non-SLS on fibroblast cell culture. *Journal of International Dental and Medical Research*, *14*, 525–528.
- Sha, A. M., Garib, B. T., Azeez, S. H., & Gul, S. S. (2020). Effects of curcumin gel on osteoclastogenic bone markers in experimental periodontitis and alveolar bone loss in wistar rats. *Journal of Dental Sciences*, *16*, 905–914. <https://doi.org/10.1016/j.jds.2020.09.015>
- Shaterzadeh-Yazdi, H., Noorbakhsh, M.-F., Hayati, F., Samarghandian, S., & Farkhondeh, T. (2018). Immunomodulatory and anti-inflammatory effects of thymoquinone. *Cardiovascular & Hematological Disorders-Drug Targets*, *18*, 52–60. <https://doi.org/10.2174/1871529x18666180212114816>
- Strowig, T., Henao-Mejia, J., Elinav, E., & Flavell, R. (2012). Inflammasomes in health and disease. *Nature*, *481*(7381), 278–286. <https://doi.org/10.1038/nature10759>
- Susanto, H., Nesse, W., Dijkstra, P. U., Agustina, D., Vissink, A., & Abbas, F. (2011). Periodontitis prevalence and severity in Indonesians with type 2 diabetes. *Journal of Periodontology*, *82*, 550–557. <https://doi.org/10.1902/jop.2010.100285>
- Tantivitayakul, P., Kaypetch, R., & Muadchiengka, T. (2020). Thymoquinone inhibits biofilm formation and virulence properties of periodontal bacteria. *Archives of Oral Biology*, *115*, Article 104744. <https://doi.org/10.1016/j.archoralbio.2020.104744>
- Velickovic, M., Arsenijevic, A., Acovic, A., Arsenijevic, D., Milovanovic, J., Dimitrijevic, J., ... Arsenijevic, N. (2021). Galectin-3, possible role in pathogenesis of periodontal diseases and potential therapeutic target. *Frontiers in Pharmacology*, *12* (March), 1–14. <https://doi.org/10.3389/fphar.2021.638258>
- Wada, K., & Kamisaki, Y. (2010). Roles of oral bacteria in cardiovascular diseases - From molecular mechanisms to clinical cases: Involvement of *porphyromonas gingivalis* in the development of Human Aortic Aneurysm. *Journal of Pharmacological Sciences*, *113*(2), 115–119. <https://doi.org/10.1254/jphs.09R22FM>
- Zhang, L., Li, X., Yan, H., & Huang, L. (2018). Salivary matrix metalloproteinase (MMP)-8 as a biomarker for periodontitis. *Medicine*, *97*, 1–6. <https://doi.org/10.1097/MD.00000000000009642>

Nigella sativa toothpaste promotes anti-inflammatory and anti-destructive effects in a rat model of periodontitis

ORIGINALITY REPORT

19%

SIMILARITY INDEX

11%

INTERNET SOURCES

17%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1	link.springer.com Internet Source	1%
2	j-ips.org Internet Source	1%
3	Christopher Igwe Idumah. "Emerging trends in Poly(lactic-co-glycolic) acid bionanoarchitectures and applications", Cleaner Materials, 2022 Publication	1%
4	theses.gla.ac.uk Internet Source	1%
5	Aram Mohammed Sha, Balkees Taha Garib, Shokhan Hamaali Azeez, Sarhang Sarwat Gul. "Effects of curcumin gel on osteoclastogenic bone markers in experimental periodontitis and alveolar bone loss in wistar rats", Journal of Dental Sciences, 2020 Publication	<1%
6	cirst2.openum.ca Internet Source	<1%

7 Md Faruque Ahmad, Fakhruddin Ali Ahmad, Syed Amir Ashraf, Hisham H. Saad et al. "An updated knowledge of Black seed (*Nigella sativa* Linn.): Review of phytochemical constituents and pharmacological properties", *Journal of Herbal Medicine*, 2021
Publication

<1 %

8 assets.researchsquare.com
Internet Source

<1 %

9 kmbase.medic.or.kr
Internet Source

<1 %

10 Asmaa Abou-Bakr, Radwa R. Hussein, Eman Khalil, Enji Ahmed. "The frequency of periodontitis in end-stage renal disease on hemodialysis in a sample of Egyptian population: multi-center clinical cross-sectional study", *BMC Oral Health*, 2022
Publication

<1 %

11 C Jessica, T T Alwadris, S R Prasetyo, R Puspitawati, E I Auerkari. "Association of interleukin 8 -251 A/T gene polymorphism with periodontitis in Indonesia", *Journal of Physics: Conference Series*, 2018
Publication

<1 %

12 mdpi-res.com
Internet Source

<1 %

13 Moeko Momokita, Atsushi Abe, Kanae Shibata, Hiroki Hayashi et al. "Association between oral function and nutrition in gastric cancer patients undergoing chemotherapy", *Oral Diseases*, 2022
Publication <1 %

14 scholar.unair.ac.id
Internet Source <1 %

15 Asti Meizarini, Aryati, Wibi Riawan, Astari Puteri. "Anti-inflammatory properties of a wound dressing combination of zinc oxide and turmeric extract", *Veterinary World*, 2018
Publication <1 %

16 www.researchgate.net
Internet Source <1 %

17 Natalia Calva-Candelaria, María Estela Meléndez-Camargo, Hortensia Montellano-Rosales, Alan R. Estrada-Pérez et al. "Oenothera rosea L´Hér. ex Ait attenuates acute colonic inflammation in TNBS-induced colitis model in rats: in vivo and in silico myeloperoxidase role", *Biomedicine & Pharmacotherapy*, 2018
Publication <1 %

18 Karen L. Syrjala, Steven K. Sutton, Heather S. L. Jim, Jennifer M. Knight et al. "Cancer and treatment distress psychometric evaluation <1 %

over time: A BMT CTN 0902 secondary analysis", Cancer, 2017

Publication

19

[bmcgenomics.biomedcentral.com](https://www.bmcgenomics.biomedcentral.com)

Internet Source

<1 %

20

repositorio.uchile.cl

Internet Source

<1 %

21

tessera.spandidos-publications.com

Internet Source

<1 %

22

N.S. Tamashiro, P.M. Duarte, T.S. Miranda, S.S. Maciel, L.C. Figueiredo, M. Faveri, M. Feres. "Amoxicillin Plus Metronidazole Therapy for Patients with Periodontitis and Type 2 Diabetes", Journal of Dental Research, 2016

Publication

<1 %

23

Nurina Febriyanti Ayuningtyas, Fatma Yasmin Mahdani, Togu Andrie Simon Pasaribu, Muhammad Chalim et al. "Role of Candida albicans in Oral Carcinogenesis", Pathophysiology, 2022

Publication

<1 %

24

mts.intechopen.com

Internet Source

<1 %

25

topsecretapiaccess.dovepress.com

Internet Source

<1 %

26

Ali NE, Abd El kareem HM, Elnoury HA, Mahmoud AR. "EFFECT OF OXYMATRINE ON TRICLOSAN-INDUCED OVARIAN TOXICITY IN FEMALE RATS.", Egyptian Journal of Occupational Medicine, 2021

Publication

<1 %

27

Aniek S. Budiatin, Maria A. Gani, Samirah, Chrismawan Ardianto et al. "Bovine Hydroxyapatite-Based Bone Scaffold with Gentamicin Accelerates Vascularization and Remodeling of Bone Defect", International Journal of Biomaterials, 2021

Publication

<1 %

28

George A. Burdock. "Assessment of black cumin (Nigella sativa L.) as a food ingredient and putative therapeutic agent", Regulatory Toxicology and Pharmacology, 2022

Publication

<1 %

29

Jacqueline R. Hwang, Alexander M. Cartron, Amor Khachemoune. " A review of plant - based therapy in dermatology ", International Journal of Dermatology, 2021

Publication

<1 %

30

Retno Indrawati Roestamadji, Nahdah Izdihar Nastiti, Meircurius Dwi Condro Surboyo, Anis Irmawati. "The Risk of Night Shift Workers to the Glucose Blood Levels, Saliva, and Dental Caries", European Journal of Dentistry, 2019

<1 %

31	bora.uib.no Internet Source	<1 %
32	helda.helsinki.fi Internet Source	<1 %
33	ijrr.com Internet Source	<1 %
34	kubanni.abu.edu.ng Internet Source	<1 %
35	library.umac.mo Internet Source	<1 %
36	onlinelibrary.wiley.com Internet Source	<1 %
37	openarchive.ki.se Internet Source	<1 %
38	scholarshare.temple.edu Internet Source	<1 %
39	www.scilit.net Internet Source	<1 %
40	www.spandidos-publications.com Internet Source	<1 %
41	Äzlem Yilmaz. "ATP-dependent activation of an inflammasome in primary gingival	<1 %

epithelial cells infected by *Porphyromonas gingivalis*", Cellular Microbiology, 02/2010

Publication

42

A.H. REBAR. "Comparative Acute Toxicity of Four Nickel Compounds to F344 Rat Lung", Toxicological Sciences, 1986

Publication

<1 %

43

Ahmad M. Eid, Nidal A. Jaradat, Nagib A. Elmarzughi, Raed Alkowni, Fatima Hussien, Laila A. Ayyash, Maher Sawafta, Hadeel Danaa. "Anti-Microbial and Free Radical Scavenging Activities of Nigella Sativa Colloidal-Emulgel", Letters in Drug Design & Discovery, 2019

Publication

<1 %

44

Daniela Stupphann. "Intracellular and surface RANKL are differentially regulated in patients with ankylosing spondylitis", Rheumatology International, 08/2008

Publication

<1 %

45

Gaetano Isola, Alessandro Polizzi, Angela Alibrandi, Ray C. Williams, Antonino Lo Giudice. "Analysis of galectin - 3 levels as a source of coronary heart disease risk during periodontitis", Journal of Periodontal Research, 2021

Publication

<1 %

46

Hokamura, Kazuya, and Kazuo Umemura. "Roles of Oral Bacteria in Cardiovascular

<1 %

Diseases " From Molecular Mechanisms to Clinical Cases: Porphyromonas gingivalis Is the Important Role of Intimal Hyperplasia in the Aorta", Journal of Pharmacological Sciences, 2010.

Publication

47

Hoo-Seob Park, Hae-Sung Nam, Hyung-Seok Seo, Soo-Jeong Hwang. "Change of periodontal inflammatory indicators through a 4-week weight control intervention including caloric restriction and exercise training in young Koreans: a pilot study", BMC Oral Health, 2015

Publication

48

Jepsen, Karin, and Søren Jepsen. "Antibiotics/antimicrobials: systemic and local administration in the therapy of mild to moderately advanced periodontitis", Periodontology 2000, 2016.

Publication

49

Yamaguchi, Y., T. Kurita-Ochiai, R. Kobayashi, T. Suzuki, and T. Ando. "Activation of the NLRP3 inflammasome in Porphyromonas gingivalis-accelerated atherosclerosis", Pathogens and Disease, 2015.

Publication

50

Yang Li, Cheng Wen, Jialin Zhong, Junqi Ling, Qianzhou Jiang. " OG1RF induces apoptosis in

<1 %

<1 %

<1 %

<1 %

MG63 cells via caspase - 3/ - 8/ - 9 without activation of caspase - 1/GSDMD ", Oral Diseases, 2021

Publication

51 Yang Li, Junqi Ling, Qianzhou Jiang. "Inflammasomes in Alveolar Bone Loss", Frontiers in Immunology, 2021 <1 %

Publication

52 Zhi Wang, Xi Wang, Yiqun Jia, Liling Wen et al. "Porphyromonas gingivalis promotes colorectal carcinoma by activating the hematopoietic NLRP3 inflammasome", Cancer Research, 2021 <1 %

Publication

53 archive-ouverte.unige.ch <1 %

Internet Source

54 espace.library.uq.edu.au <1 %

Internet Source

55 pdfs.semanticscholar.org <1 %

Internet Source

56 portlandpress.com <1 %

Internet Source

57 scirp.org <1 %

Internet Source

58 www.karger.com <1 %

Internet Source

59	www.nature.com Internet Source	<1 %
60	www.tandfonline.com Internet Source	<1 %
61	www.thieme-connect.com Internet Source	<1 %
62	www.trjfas.org Internet Source	<1 %
63	"Clinical Cases in Dental Hygiene", Wiley, 2018 Publication	<1 %
64	ERBOGA, Mustafa, ERBOGA F?DANOL, Zeynep, DONMEZ BOZDEM?R, Yeliz, AKTAS, Cevat and KANTER, Mehmet. "Nigella sativa attenuates bleomycin-induced pulmonary fibrosis in rats by inhibition of inflammation, fibrosis, and inducible nitric oxide synthase", Ondokuz may?s ?niversitesi T?p Fak?ltesi, 2015. Publication	<1 %
65	Fernanda R G. Rocha, Andrea E. Delitto, Joao A. Chaves de Souza, Laura A. González-Maldonado et al. "Relevance of Caspase-1 and Nlrp3 Inflammasome on Inflammatory Bone Resorption in A Murine Model of Periodontitis", Scientific Reports, 2020 Publication	<1 %

66

Prathiba Gnanasekaran, Anitha Roy, Nagabhishek Sirpu Natesh, Varshini Raman, Priyanka Ganapathy, Madan Kumar Arumugam. "Removal of microbial pathogens and anticancer activity of synthesized nano-thymoquinone from Nigella sativa seeds", Environmental Technology & Innovation, 2021

Publication

<1 %

67

etheses.whiterose.ac.uk

Internet Source

<1 %

68

hdl.handle.net

Internet Source

<1 %

69

Alrieta H. Teixeira, Jordânia M. de Oliveira Freire, Luzia H. T. de Sousa, Antônia T. Parente et al. "Stemodia maritima L. Extract Decreases Inflammation, Oxidative Stress, and Alveolar Bone Loss in an Experimental Periodontitis Rat Model", Frontiers in Physiology, 2017

Publication

<1 %

70

Edible Medicinal And Non-Medicinal Plants, 2013.

Publication

<1 %

71

Gang Chen, Joao Pedra. "The Inflammasome in Host Defense", Sensors, 2009

Publication

<1 %

72

Javier González-Ramírez, Nicolás Serafín-Higuera, Marina Concepción Silva Mancilla, Gustavo Martínez-Coronilla et al. "Chapter 2 Use of Biomarkers for the Diagnosis of Periodontitis", IntechOpen, 2020

Publication

<1 %

73

Kaitlyn A. Stanton, Barbara A. McCracken. "An oral rinse reduces cytokine secretion and promotes proliferation in gingival tissues – A pilot study ", Clinical and Experimental Dental Research, 2021

Publication

<1 %

74

Kanako Higuchi, Ziauddin SM, Yasunori Yamashita, Yukio Ozaki, Atsutoshi Yoshimura. "Initial periodontal treatment affects nucleotide-binding domain leucine-rich repeat-containing protein 3 inflammasome priming in peripheral blood mononuclear cells", Archives of Oral Biology, 2020

Publication

<1 %

75

Kejia Lv, Guohua Wang, Chenlu Shen, Xia Zhang, Hua Yao. "Role and mechanism of the nod-like receptor family pyrin domain-containing 3 inflammasome in oral disease", Archives of Oral Biology, 2019

Publication

<1 %

76

Kübra Aral, Michael R. Milward, Paul R. Cooper. "Inflammasome dysregulation in

<1 %

human gingival fibroblasts in response to periodontal pathogens", Oral Diseases, 2021

Publication

77

Muhammad Torequl Islam, Md. Roich Khan, Siddhartha Kumar Mishra. "An updated literature-based review: phytochemistry, pharmacology and therapeutic promises of *Nigella sativa* L.", Oriental Pharmacy and Experimental Medicine, 2019

Publication

<1 %

78

Shabina Ishtiaq Ahmed, Sehrish Jamil, Humaira Ismatullah, Rashid Hussain et al. "A comprehensive perspective of traditional Arabic or Islamic medicinal plants as an adjuvant therapy against COVID-19", Saudi Journal of Biological Sciences, 2023

Publication

<1 %

79

Volkan Şengül, Zeliha Güney, Şivge Kurgan, Canan Önder, Muhittin A. Serdar, Meral Günhan. "Evaluation of salivary and serum methylated arginine metabolites and nitric oxide synthase in advanced periodontitis patients", Clinical Oral Investigations, 2022

Publication

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

Nigella sativa toothpaste promotes anti-inflammatory and anti-destructive effects in a rat model of periodontitis

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8
