



Source details

International Journal of Public Health Science

Scopus coverage years: from 2020 to Present

Publisher: Intelektual Pustaka Media Utama

ISSN: 2252-8806 E-ISSN: 2620-4126

Subject area: [Medicine: Medicine \(miscellaneous\)](#) [Medicine: Health Policy](#) [Social Sciences: Health \(social science\)](#)
[Medicine: Public Health, Environmental and Occupational Health](#) [Nursing: Nutrition and Dietetics](#)

Source type: Journal

[View all documents >](#)

[Set document alert](#)

[Save to source list](#) [Source Homepage](#)

CiteScore 2021

0.4



SJR 2021

0.142



SNIP 2021

0.201



[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)

i Improved CiteScore methodology



CiteScore 2021 counts the citations received in 2018-2021 to articles, reviews, conference papers, book chapters and data papers published in 2018-2021, and divides this by the number of publications published in 2018-2021. [Learn more >](#)

CiteScore **2021**

$$0.4 = \frac{80 \text{ Citations 2018 - 2021}}{180 \text{ Documents 2018 - 2021}}$$

Calculated on 05 May, 2022

CiteScoreTracker 2022

$$0.7 = \frac{207 \text{ Citations to date}}{314 \text{ Documents to date}}$$

Last updated on 05 August, 2022 • Updated monthly

CiteScore rank 2021

Category	Rank	Percentile
Medicine		
Medicine (miscellaneous)	#231/276	16th
Medicine		
Health Policy	#226/265	14th
Social Sciences		
Health (social)	#281/323	13th

[View CiteScore methodology >](#) [CiteScore FAQ >](#) [Add CiteScore to your site](#)

About Scopus

[What is Scopus](#)
[Content coverage](#)
[Scopus blog](#)
[Scopus API](#)
[Privacy matters](#)

Language

[日本語版を表示する](#)
[查看简体中文版本](#)
[查看繁體中文版本](#)
[Просмотр версии на русском языке](#)

Customer Service

[Help](#)
[Tutorials](#)
[Contact us](#)


ELSEVIER

[Terms and conditions](#) ↗ [Privacy policy](#) ↗

Copyright © [Elsevier B.V](#) ↗. All rights reserved. Scopus® is a registered trademark of Elsevier B.V.

We use cookies to help provide and enhance our service and tailor content. By continuing, you agree to the [use of cookies](#) ↗.



also developed by scimago:  SCIMAGO INSTITUTIONS RANKINGS

SJR Scimago Journal & Country Rank

[Home](#) [Journal Rankings](#) [Country Rankings](#) [Viz Tools](#) [Help](#) [About Us](#)

Latest Research Findings


Nature Research Journals

Groundbreaking Global Science Research Delivered to You & Online Archive Access From 1997.


nature.com [OPEN](#)

[Close](#)

International Journal of Public Health Science

COUNTRY	SUBJECT AREA AND CATEGORY	PUBLISHER	H-INDEX
<p>Indonesia</p> <p> Universities and research institutions in Indonesia</p>	<p>Medicine</p> <ul style="list-style-type: none"> Health Policy Medicine (miscellaneous) Public Health, Environmental and Occupational Health <p>Nursing</p> <ul style="list-style-type: none"> Nutrition and Dietetics <p>Social Sciences</p> <ul style="list-style-type: none"> Health (social science) 	<p>Intelektual Pustaka Media Utama</p>	<p>4</p>
PUBLICATION TYPE	ISSN	COVERAGE	
Journals	22528806, 26204126	2020-2021	



 Join the conversation about this journal


Reviews, News, & Analysis

Online & Print Weekly Journal


Groundbreaking Global Science Research Journal. Get Your Online & Print Subscription Now!








nature.com

OPEN

 Quartiles

B





International Journal of Public Health Science

Health Policy

Q4

best quartile

SJR 2021

0.14

powered by scimagojr.com

← Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scimi
```

SCImago Graphica

Explore, visually communicate and make sense of data with our **new data visualization tool.**

HOME ABOUT LOGIN REGISTER SEARCH CURRENT ARCHIVES ANNOUNCEMENTS

Home > Archives > Vol 5, No 4

Vol 5, No 4

December 2016

DOI: <http://doi.org/10.11591/ijphs.v5i4>

Table of Contents

Rapid-Acting Antidepressants and Underlying Mechanisms Samuel E Wilson, Meta Chen, Hiren Darji	PDF 347-353
Sensitivity and Specificity of Rapid Diagnostic Test with Microscopic Gold Standard to Identify Plasmodium Species Sahrir Sillehu, Heny Arwati, Yoes Prijatna Dachlan, Sudjadi Keman	PDF 354-358
The Support to Improve Self Efficacy and Healing of Drugs Addict Rosdiana Rosdiana, Bhisma Murti, Mahendra Wijaya, Suwanto Suwanto	PDF 359-366
Responses of Diabetes Mellitus Patients Who Used Complementary Medicine Anita Joeliantina, Mangestuti Agil, M. Bagus Qomaruddin, Arijanto Jonosewojo, Kusnanto Kusnanto	PDF 367-374
Prevalence and Associated Factors of Hypertension among Civil Servants Working in Arba Minch Town, South Ethiopia Aschenaki Kalssa, Gistane Ayele, Alemu Tamiso, Tadele Girum	PDF 375-383
Knowledge and Attitude of Students on Antimicrobial Resistance at Debre Markos University, Ethiopia Desalegne Amare Zelellw, Habtamu Mellie Bizuayehu	PDF 384-391
Availability of Adequately Iodized Salt at Household Level and Associated Factors in Dire Dawa, Eastern Ethiopia Anteneh Berhane Yaye, Negga Baraki, Birhanu Seyum Endale	PDF 392-399
Relaxation Progressive Muscle Program on Exercise Behavior and Clinical Outcomes among Hypertension Patients Rian Adi Pamungkas, Wahyu Kirana, Florensa Florensa	PDF 400-405
Ecological Social Development Model of Health Behavior of Conduct Achievement MDGs 5 Hilmi Yumni, Fendy Suhariadi, Oedojo Sudirham, Baiq Dewi Harmani R, Hasyim As'ari, Noer Saudah	PDF 406-414
Dealing with the Daily Emergency Care: A Case among the Waste Pickers in Surabaya Loetfia Dwi Rahariyani, Nikmatul Fadilah, Yohanes Kambaru Windi, Nursalam Nursalam	PDF 415-421
Determinants Levels of Urinary 8-Hydroxydeoxyguanosine among Chromium Electroplating Workers Yuliani Setyaningsih, Indwiani Astuti, Adi Heru Husodo	PDF 422-426
Adolescent's Self-Efficacy for Early Marriage in South Bangka Regency, Indonesia Antarini Antarini, Putri Rhadiyah, Tesza Permata, Rizka Marcelly, Dini Montovani	PDF 427-432
The Role of Village Surveillance Officer to Prevent Dengue Hemorrhagic Fever Sunarsih Sunarsih, Kuntoro Kuntoro, Chatrin U.W, Nugroho Susanto	PDF 433-436
Arsenic Exposure to Fish and Shellfish Consumption among Community in Makassar, Indonesia Musfirah Musfirah, Sri Damayanty	PDF 437-443
Association of Physical Activity with Co-morbid Conditions in Geriatric Population Aliya Hisam, Mahmood Ur Rahman, Rimsha Khan, Sundus Ilyas, Anosha Jabbar, Syed Awais Kazmi, Umer Mushtaq	PDF 444-449
Job Satisfaction among the Nurses of Makueni District Hospital, Kenya Anthony wambua Mathulu, Benard Wambua Mbithi	PDF 450-457
Ageing and Secular Social Involvement: Exploring to Subjective Wellbeing Rachmah Indawati, Kuntoro Kuntoro, Hari Basuki Notobroto, Mochammad Bagus Qomaruddin	PDF 458-464
Environmental Health Risk Analysis of Paraquat Exposure in Palm Oil Plantations Maksuk Maksuk, Tan Malaka, Suheryanto Suheryanto, Abu Umayah	PDF 465-469
The Influence of Lavender Aroma Therapy on Decreasing Blood Pressure in Hypertension Patients	PDF 470-478

USER

Username

Password

Remember me

[Login](#)

CITATION ANALYSIS

- Scopus
- Google Scholar
- Scholar Metrics
- Scinapse
- Dimensions

SPECIAL LINKS

- Author Guideline
- Editorial Boards
- Online Submissions
- Abstracting and Indexing
- Publication Ethics
- Visitor Statistics
- Contact Us

LINK PER ISSUE

2022: [Mar](#), [Jun](#), [Sep](#), [Dec](#)
 2021: [Mar](#), [Jun](#), [Sep](#), [Dec](#)
 2020: [Mar](#), [Jun](#), [Sep](#), [Dec](#)
 2019: [Mar](#), [Jun](#), [Sep](#), [Dec](#)
 2018: [Mar](#), [Jun](#), [Sep](#), [Dec](#)
 2017: [Mar](#), [Jun](#), [Sep](#), [Dec](#)
 2016: [Mar](#), [Jun](#), [Sep](#), [Dec](#)
 2015: [Mar](#), [Jun](#), [Sep](#), [Dec](#)
 2014: [Mar](#), [Jun](#), [Sep](#), [Dec](#)
 2013: [Mar](#), [Jun](#), [Sep](#), [Dec](#)
 2012: [Jul](#), [Dec](#)

JOURNAL CONTENT

Search

Search Scope [All](#)

[Search](#)

Browse

- By Issue
- By Author
- By Title

INFORMATION

- For Readers
- For Authors
- For Librarians

Agustina Boru Gultom, Surita Ginting, Elni Lorensi Silalahi

[Pain and Suffering: The Reality of Being Positive with HIV/AIDS](#)
Retno Lestari, Wenny Nugrahati Carsita

[PDF](#)
479-483

[Subchronic Toxicity of Green Algae \(*Spyrogyra* sp.\) Ethanolic Extract on Hematologic Parameters](#)
Nina Salamah, Wahyu Widyarningsih, Hari Susanti, Anggita Devi, Anita Wening Sejati, Zahra Alya Putri

[PDF](#)
484-488

International Journal of Public Health Science (IJPHS)
p-ISSN: 2252-8806, e-ISSN: 2620-4126



[View IJPHS Stats](#)



This work is licensed under a [Creative Commons Attribution-ShareAlike 4.0 International License](#).

Home > About the Journal > **Editorial Team**

Editorial Team

Advisory Board

[Prof. Hans-Olov Adami](#), Harvard School of Public Health, United States
[Assoc. Prof. Dr. Luoping Zhang](#), University of California, Berkeley, United States
[Dr. Khalid M. Al Aboud](#), King Faisal Specialist Hospital and Research Centre, Saudi Arabia

Editor-in-Chief

[Prof. Dr. Jay G. Silverman](#), University of California, San Diego School of Medicine, United States

Managing Editor

[Dr. Lina Handayani](#), Universitas Ahmad Dahlan, Indonesia

Associate Editors

[Assoc. Prof. Dr. Mohd Hasni Jaafar](#), Universiti Kebangsaan Malaysia, Malaysia
[Assoc. Prof. Dr. Henry Odhianoson Imhonde](#), Ambrose Alli University, Nigeria
[Prof. Dr. Mane Abhay Babruwahan](#), Smt. Kashibai Navale Medical College, India
[Dr. Fazal Shirazi](#), MD Anderson Cancer Center, United States
[Dr. Miguel A. Mayer MD, PhD, MPH, MSc, Dipl.](#), Universitat Pompeu Fabra, Spain
[Dr. Muhiuddin Haider](#), University of Maryland, United States
[Dr. Tassanee Rawiworrakul](#), Mahidol University, Thailand
[Dr. Xiaoli Gao](#), University of Texas Health Science Center at San Antonio, United States

Editorial Board Members

[Prof. Dr. George Williams Rutherford](#), Global Health Sciences, University of California, San Francisco, United States
[Prof. Dr. Jens Aagaard-Hansen](#), Danish Bilharziasis Laboratory, Denmark
[Prof. Dr. Louise H. Taylor](#), Global Alliance for Rabies Control, Manhattan, United States
[Prof. Dr. Mogens Vyberg](#), Aalborg Universitet, Denmark
[Dr. Abhinand Thavalappil](#), Ryerson University, Canada
[Dr. Stephanie S. Rothenberg](#), University of Pittsburgh Medical Center, United States
[Dr. Aletha Yvette S. Akers](#), Magee-Womens Hospital, Pittsburgh, United States
[Dr. Andrew Winokur](#), University of Connecticut School of Medicine, United States
[Dr. Barbara Abrams](#), University of California, United States
[Dr. Barbara K. Campbell](#), University of California, San Francisco, United States
[Dr. Christopher Bowie](#), University of Canterbury, New Zealand
[Dr. Erika Villavicencio-Ayub](#), Universidad Nacional Autónoma de México, Mexico
[Dr. Fhumulani Mavis Mulaudzi](#), University of Pretoria, South Africa
[Dr. Jennifer A. Fish](#), The University of South Australia, Australia
[Dr. Li-Ping Zou](#), Department of Paediatrics, Chinese PLA General Hospital, China
[Dr. Phayong Thepakorn](#), Sirirhorn College of Public Health, Thailand
[Dr. Rob M.G. van Bommel](#), Catharina Hospital, Netherlands
[Dr. Tarik Bereket](#), University of Toronto, Canada
[Dr. Vera Fernandes](#), Hospital de Braga, Portugal

International Journal of Public Health Science (IJPHS)
 p-ISSN: 2252-8806, e-ISSN: 2620-4126



[View IJPHS Stats](#)



This work is licensed under a [Creative Commons Attribution-ShareAlike 4.0 International License](#).

USER

Username
 Password
 Remember me
[Login](#)

CITATION ANALYSIS

- Scopus
- Google Scholar
- Scholar Metrics
- Scinapse
- Dimensions

SPECIAL LINKS

- Author Guideline
- Editorial Boards
- Online Submissions
- Abstracting and Indexing
- Publication Ethics
- Visitor Statistics
- Contact Us

LINK PER ISSUE

2022: [Mar](#), [Jun](#), [Sep](#), [Dec](#)
 2021: [Mar](#), [Jun](#), [Sep](#), [Dec](#)
 2020: [Mar](#), [Jun](#), [Sep](#), [Dec](#)
 2019: [Mar](#), [Jun](#), [Sep](#), [Dec](#)
 2018: [Mar](#), [Jun](#), [Sep](#), [Dec](#)
 2017: [Mar](#), [Jun](#), [Sep](#), [Dec](#)
 2016: [Mar](#), [Jun](#), [Sep](#), [Dec](#)
 2015: [Mar](#), [Jun](#), [Sep](#), [Dec](#)
 2014: [Mar](#), [Jun](#), [Sep](#), [Dec](#)
 2013: [Mar](#), [Jun](#), [Sep](#), [Dec](#)
 2012: [Jul](#), [Dec](#)

JOURNAL CONTENT

Search
 Search Scope
[Search](#)

Browse

- By Issue
- By Author
- By Title

INFORMATION

- For Readers
- For Authors
- For Librarians

Sensitivity and Specificity of Rapid Diagnostic Test with Microscopic Gold Standard to Identify Plasmodium Species

Sahrir Sillehu¹, Heny Arwati², Yoes Prijatna Dachlan³, Sudjajadi Keman⁴

^{1,4}Faculty of Public Health, Airlangga University, Surabaya, Indonesia

^{2,3}Faculty of Medicine, Airlangga University, Surabaya Indonesia

Article Info

Article history:

Received Feb 25, 2016

Revised May 20, 2016

Accepted May 26, 2016

Keyword:

Microscopic malaria
Rapid Diagnostic Test
Sensitivity
Specificity

ABSTRACT

Malaria is a main health problem in islands area which is under developed and isolated. Nation-wide, in 2014 Maluku province was recorded to have Annual Malaria Incident (AMI) value of 30.4%, positive incidents of 13.30%, ABER 3.76%, SPR 21.50%, and Annual Parasite Incident (API) 8.10%, while South Buru Regency has a value of Annual Malaria Incident (AMI) of 14.49%, 494 positive incidents, ABER 1.12%, SPR 60.91%, and Annual Parasite Incident (API) 6.86%. The purpose of this study was to identify Plasmodium species in malaria incidents in Namrole Subdistrict, South Buru Regency, Maluku Province. Observational research with a sample of 64 respondents for symptomatic and asymptomatic malaria. The instrument for the research was Rapid Diagnostic Test (RDT) and microscopic Gold Standard. Result: Malaria examination by using RDT suggested 3 kinds of parasites, i.e., *P. falciparum*, *P. Vivax*, and a mix between *P. falciparum* and *P. vivax*. Most parasites found were *P. falciparum* 56.3%. The accuracy of RDT examination was proven with microscopic test and the result suggested that the RDT sensitivity was 100% and the specificity was 63.3%. Positive predictive value was 92.9% and negative predictive value was 100%, both were for positive likelihood ration of 2.75%. While for negative likelihood ration of 0%, the value of degree of conformity (Kappa) between RDT and microscopic is 0%. RDT has one benefit that it can be used to conduct malaria diagnosis rapidly, particularly in isolated areas. The benefit of Rapid Diagnostic Test (RDT) was that it could be used in remote and isolated areas to conduct diagnosis. RDT is highly effective and efficient.

Copyright © 2016 Institute of Advanced Engineering and Science.
All rights reserved.

Corresponding Author:

Sahrir Sillehu,
Doctoral Student at Faculty of Public Health,
Airlangga University,
Campus C UNAIR, Mulyorejo, Surabaya, Indonesia.
Email: umay_01@yahoo.com

1. INTRODUCTION

Malaria is a type of infection caused by protozoa from Plasmodium genus. Malaria is an infectious disease which is very dominant in tropical and sub tropical areas. It can cause millions of death annually. Based on information and profiles of 99 countries worldwide, the controlling and elimination of Malaria incidents has progressing globally. As many as 80 countries are currently in supervision phase, while 19 others are in pre-elimination phase of malaria. It is estimated that 3.3 billion people around the globe are in a risk of getting infected by malaria. In Africa, the risk of getting infected by malaria is around 81% case, it is estimated that 91% mortalities occur on children aged under 5 years and pregnant woman [1].

In Indonesia, Malaria is one of the indicators of the targets of Millennium Development Goals (MDGs), i.e., stopping the spread and reducing malaria incidents in 2015. This can be indicated from the

decrease of morbidity and mortality caused by malaria. Global Malaria Programme (GMP) stated that malaria is an illness that must be continuously observed, monitored, and evaluated and it takes a formulation of appropriate policies and strategies [2].

Staid that the increasing incidents of malaria from year to year and the occurrence of Malaria Extraordinary Incidents during the period of 1997-2000 were caused by some factors including: 1) environmental change leading to wider area for malaria breedings, 2) high people mobility, 3) rain season changes, it has been much longer than dry season in the recent years, 4) lingering economic crisis bringing about malnutrition in some areas that they become adverse to malaria exposure, 5) low level of concern from the society to contribute in the efforts of preventing and mitigating malaria [3].

Indonesia is a country still facing the risk of malaria, some 80% of regencies/towns in Indonesia can be included into malaria endemic category. Malaria attacks people domiciling in remote areas with poor environmental condition, hard to reach transportation and communication, and limited access to health facilities. The number of reported clinical incidents in 2009 were as many as 1,143,024 people with positive case, based in laboratory examination were 199,577 people. Regarding mortality of infectious and non infectious disease, malaria ranks number 6 (4.6%) [4]. According to the data of Ministry of Health of Republic of Indonesia, generally malaria sufferers in Indonesia are inhabitants of remote and hard to reach areas, and live in poverty. On top of that, from 1998 to 2003, the occurrences of Extraordinary Incidents (KLB) included 14 provinces and 26 regencies in 162 villages with the number of sufferers of 27,956 people and 328 mortalities) [5].

The development of malaria program in Maluku Province from 2011-2014 suggested the following; in 2011 the number of malaria declared as positively suffering malaria per 1,000 citizen was as many as 13.857%, ABER 2.37%, SPR 34.48%, API 8.10% In 2012; positive incidents 16.69%, ABER 2.62%, SPR 37.64%, API 9.86%. In 2013; positive incidents 16.508%, ABER 3.32%, SPR 29.34%, API 9.76%. In 2014; positive incidents 13.307%, ABER 3.76%, SPR 21.50%, API 8.10% (Health Profile, 2014) [6].

South Buru Regency in 011, the number of positive incidents of malaria was as many as 477, ABER 0.91%, SPR 72.49%, API 6.63%. In 2012; positive incidents 208, ABER 0.53%, SPR 54.02%, API 2.89%. In 2013; positive incidents 361, ABER 0.69%, SPR 71.91%, API 5.01%. In 2014; positive incidents 494, ABER 1.12%, SPR 60.91%, API 6.86% [7].

In South Buru regency, there are some isolated people inhabiting mountainous or inland areas with limited infrastructures. These people still live primitively and do not do many interactions with modern people life. While their daily life depends on the surrounding natures or they live vegetatively so to say. If any member of their families suffers from malaria, they will attempt to cure them with traditional ingredients taken from natural herbs around them [8].

Clinical manifestations which are specifically found in malaria endemic areas, which can be used as initial step in diagnosing clinical malaria, are very helpful in rapid, appropriate, and rational diagnosis of malaria, however, clinical malaria diagnosis must be confirmed with the result of microscopic test, which remains a gold standard in diagnosing malaria. Lacking of workforce in the field of laboratory technicians skilled enough in using microscope to properly diagnose malaria is one of the factors leading to slow medication and inappropriate malaria diagnosis [9].

Mobility or movement of inhabitants from malaria endemic areas to non endemic areas, inhabitants traveling and coming home for certain period of time (incubation period) of Plasmodium, potentially lead to the transmission or infection of malaria. People's migration bring about the movement of population infected by *P. falciparum*, especially in children. Some reports mentioned that Multiplicity of Infection (MOI) will reflect individuals infected by having immunity against malaria [10].

According to the WHO, RDT is alternative method based dipstick major clinical manifestations of malaria, especially in places that do not have a qualified technician and microscopic means. In addition, helpful RDT unit emergency medical services, when extraordinary event malaria, as well as in areas that are very difficult and disadvantaged are not available clinical laboratory facilities [11].

This research aimed to know the sensitivity and specificity of Rapid Diagnostic Test with Microscopic as gold standard in conducting diagnosis of malaria incidents in Namrole Subdistrict, South Buru Regency, Maluku Province.

2. RESEARCH METHOD

Based on the type, this research is an observational research which was conducted in Namrole subdistrict, South Buru Regency, Maluku Province. The samples used in this research were as many as 64 respondents with symptomatic and asymptomatic malaria. The tool used in this research was Rapid Diagnostic Test (RDT) and Microscopic as Gold Standard [12-16]. Data gathering technique was conducted by taking blood samples in all age groups, which is then continued with blood checking to detect the

presence of parasitic species of Plasmodium by using Rapid Diagnostic Tool (RDT) and Microscopic as Gold Standard for sensitivity and specificity test.

3. RESULTS AND ANALYSIS

Table 1 shows that most of malaria infection was occurred toward children in the age of 0-10 years old in 39 respondents (60.9%), meanwhile, the age group of 21-30 years old and 31-40 years old were 9 respondents (14.1%).

Table 1. Distribution of Frequency Based on Age Group (n = 64)

Age Group	N	%
0-10 years old	39	60.9
11-20 years old	2	3.1
21-30 years old	9	14.1
31-40 years old	9	14.1
>40 years old	5	7.8

Table 2 shows that according to the identification result of *P.falciparum* species was in 36 respondents (56.3%) more than *P.vivax* species, meanwhile, mix infection was only 10.9%.

Table 2. Distribution of Identification Result of Plasmodium species by Using RDT

Type of Parasite	N	%
<i>P. Falcifarum</i>	36	56.3
<i>P. Vivax</i>	14	21.9
Mix	7	10.9
No detected	7	10.9
Total	64	100

Table 3 shows that according to the identification result of Plasmodium species by using giemsa painting removal thick and thin blood drops was found Plasmodium stage in microscopic, which was *P.falciparum* species with gametocytes in 27 respondents (42%).

Table 3. Distribution of Identification Result of Plasmodium according to Microscopic Test as a Gold Standard

Type of Parasite	N	%
Pf. StdGametosit	27	42.2
Pf. StdCincin	7	10.9
Pv. StdTropozoid	10	15.7
Pv. StdCincin	2	3.1
Pf. StdGametosit + ring	7	10.9
No detected	11	17.2
Total	64	100

Table 4 shows that parasitemia distribution according to age group was mostly on the age of 0-10 years old with *P. falciparum* species, *P.vivax* species, and Mix Infection.

Table 4. Distribution of Parasitemia according to Age Group

Age Group (Year)	P. Falcifarum	P. Vivax	Mix
0 - 10	23	7	5
11 - 20	0	0	0
21 - 30	3	5	0
31 - 40	5	1	1
> 40 years old	5	1	1
Total	36	14	7

According to sensitivity and specificity test (Table 5) that RDT was a tool that was used to identify the type of parasite species in the diagnosis of malaria in remote and isolated area.

Table 5. Accuracy Rapid Diagnostic Test (RDT)

	Microscopic (+)	Microscopic (-)	Total
RDT (+)	53	4	57
RDT (-)	0	7	7
Total	53	11	64

Sensitivity	: 100%
Specificity	: 63%
Positive Predictive Value	: 92.9%
Negative Predictive Value	: 100%
Likelihood ratio of a Positive Test	: 2.75
Likelihood ratio of a Negative Test	: 0
Cohen's kappa (Unweighted)	: 0

4. DISCUSSION

This research was a Cross Sectional diagnostic test, which was conducted in Namrole subdistrict, South Buru Regency, Maluku Province. The samples of this research were 64 people indicated as suffering from malaria, of 64 respondents 7 people were declared negative from malaria after laboratory test. Most sufferers were in age under 1 year old to 10 years old, as many as 39 sufferers or 60.9%. Children aged under 1 year to 10 years old were the most sufferers in this research, this was due that children of that age were still adverse to the exposure of environmental change, and their immunity were relatively weaker than those of adults. Plasmodium falciparum was the most found parasites on children, with the number of plasmodium falciparum as many as 23 sufferers. While 7 others were positively having plasmodium vivax, while 5 of whom were mixed.

Based on the result of malaria checking by using RDT instrument, it was known that there were 3 types i.e., P. falciparum, P. vivax, and Mixed. Most detected parasites is P. falciparum 56.3%, then P. Vivax 21.9%, and mixed 10.9%. After proven with advance test i.e. Microscopic as Gold Standard, it was suggested that 4 results of RDT were not confirmed with microscopic test. Besides, there were 2 samples which was identified with RDT as P. Vivax, however microscopic test suggested that the parasites were P. falciparum std gametocyte.

In Table 5, the accuracy of test with Rapid Diagnostic Test (RDT) instrument proven with microscopic suggested a result that by using 2x2 table, the sensitivity of RDT instrument was 100%, while the specificity of RDT instrument was 63.3% for early detection of malaria. Positive predictive value was 92.9%, while negative predictive value was 100% with positive likelihood ratio of 2.75, negative likelihood ratio was 0, degree of conformity (Kappa) between RDT and microscopic was 0.

This research was supported by the research in Keruak Community health Centre, East Lombok Regency, suggesting sensitivity value of 100%, specificity of 96.99%, positive likelihood value of 83.2%, and negative predictive value of 100%. It can be concluded that RDT has good reliability validity to be used as diagnostic instrument [17].

Results of research conducted it is known that sensitivity value and specificity value of microscopic were better than RDT in malaria test for pregnant woman [4]. The sensitivity of rapid diagnostic test in this research is very affected by parasite density. Generally, parasite density in rapid diagnostic test (+) and microscopic (+) were relatively higher compared to that of rapid diagnostic test (-) and microscopic [9], [18].

Thus, RDT can be recommended for early detection of malaria, especially in remote, inland, and isolated area, compared to that of microscopic test. Rapid Diagnostic Test instrument (RDT) has benefit in detecting plasmodium rapidly and easily. The lacking of RDT is that it can not determine the amount of parasites in one's blood.

5. CONCLUSION

The conclusion based on the results are: 1) The benefit of microscopic as Gold standard is that this instrument has the highest accuracy value for identifying plasmodium species and parasitaemia level calculation, however it must be performed by skilled technicians in conducting diagnosis, 2) Microscopic test

suggested better result than that of Rapid Diagnostic Test, particularly for low parasites density. Rapid Diagnostic test suggested shortage that can be used as alternative malaria test method, but Rapid Diagnostic test is still very necessary, particularly in areas lacking of laboratory technicians and for surveillance purpose, 3) The benefit of Rapid Diagnostic Test (RDT) was that it could be used in remote and isolated areas to early detect malaria in the effort of conducting malaria diagnosis. Besides, RDT can be recommended and was proven very effective and efficient to be performed in case of Extraordinary Incidents.

ACKNOWLEDGEMENTS

We would like to thank the malaria patients in Namrole Sub District for their willingness to participate in this study. We gratefully acknowledge the Head and Staffs of Department of Health of South Buru District, Namrole Sub District as well as the Head and Staffs of Namrole Community Health Center (*Puskesmas*).

REFERENCES

- [1] World Health Organization, "The World Malaria Report," Geneva, World Health Organization, 2013.
- [2] Health Ministry of Indonesia, "Technical Guidelines for Examination of Malaria Parasites. Directorate of controls animal-borne diseases," Jakarta, Health Ministry of Indonesia, 2010.
- [3] Achmadi, "Management of Area-Based Diseases Eradication," Jakarta, Publisher Book Kompas, 2005.
- [4] Health Ministry of Indonesia, "Regional Health Research (Riskesdas 2007)," Jakarta, Health Ministry of Indonesia, 2007.
- [5] World Health Organization, "Malaria elimination. A field manual for low and moderate endemic countries," Geneva, World Health Organization, 2007.
- [6] Health Department of Maluku Province, "Health Profile of Maluku Province," 2014.
- [7] Health Department of South Buru District, "Health Profile of South Buru District," 2014.
- [8] Latuconsina, "Pataheri and Pasuno Initiation Ritual," Naulu tribe, Center Maluku District, 2007.
- [9] Siahaan, "Comparative of Rapid Diagnostic Test with Microscopy to Diagnose Malaria Diseases," *Jurnal Kesehatan Masyarakat Nasional*, vol/issue: 5(6), 2011.
- [10] B. M. Holder, "What is the Function of MSP-1 on the Malaria Merozoit?" *Focus*, pp. 1-3, 2005.
- [11] World Health Organization, "Malaria Rapid Diagnostic Test Performance Results of WHO Product testing of Malaria RDT: Round 1," Geneva, World Health Organization, 2008.
- [12] World Health Organization, "Guidelines for The Treatment of Malaria," Geneva, World Health Organization, 2006.
- [13] Q. A. Acton, "Issue in Global Pacific Community and Institutional Health," Georgia, Scholarly Edition, 2012.
- [14] Q. A. Acton, "Malaria New Insights for The Health Care Professional," Georgia, Scholarly Edition, 2013.
- [15] Brunett G.W., "CDC Health Information Institutional Travel," Oxford, The Yellow Book, 2016.
- [16] Soedarto, "Malaria, Global Epidemiology of Plasmodium Anopheles on the Management of Malaria Patient," Jakarta, Sagung Seto, 2011.
- [17] Arum I. L., *et al.*, "Diagnostic Test Method Using Imunokromatografi compared with microscopic examination Malaria Plasmodium Falciparum," *Indonesian Journal Of Clinical Pathology And Medical Laboratory*, 2006.
- [18] R. Sinaga, "The Sensitivity and Specificity Assessment of the Rapid Diagnostic Test (RDT) to The Gold Standard of Microscope Slides of Blood for Early Detection of Malaria in Pregnancy," 2006.