

39. Genetic Diversity of *Plasmodium falciparum* Glutamate Rich Protein Patients Attending the Merauke Hospital in Papua Province, Indonesia

by Thomas Tandi Manu

Submission date: 21-Oct-2022 02:51PM (UTC+0800)

Submission ID: 1931342700

File name: Attending_the_Merauke_Hospital_in_Papua_Province,_Indonesia..pdf (666.14K)

Word count: 3651

Character count: 18037

Genetic Diversity of *Plasmodium falciparum* Glutamate Rich Protein in Patients Attending the Merauke Hospital in Papua Province, Indonesia

Thomas Tandi Manu¹, Puspa Wardhani², Heny Arwati³, Aryati²

¹ Master Program of Basic Medical Science, Faculty of Medicine, Airlangga University, Surabaya, Indonesia

² Department of Clinical Pathology, Faculty of Medicine, Airlangga University/Dr. Soetomo Hospital, Surabaya, Indonesia. E-mail: puspa_pk@yahoo.co.id

³ Department of Medical Parasitology, Faculty of Medicine, Airlangga University, Surabaya, Indonesia

ABSTRACT

Malaria remains an important health problem in Indonesia with the highest transmission in Papua Province, an eastern part of this country. The genetic diversity of malaria parasites is the main problem in understanding several aspects of malaria infections and the dynamics of their transmission, which also play a role in the development of a vaccine. *Plasmodium falciparum* is the deadliest of the human malaria parasites. *Plasmodium falciparum* glutamate-rich protein (Pfglurp) is one of the many erythrocytic stages antigens currently under development for a vaccine. The Pfglurp gene has been extensively used as a marker to investigate the genetic diversity, Multiplicity of Infection (MOI), the level of malaria transmission, immunity against malaria, as well as a discriminatory instrument to distinguish new from recrudescence infections of the field parasite population. Thus, this genotyping study aimed to find out the genetic population of *P.falciparum* at the Merauke District, Province of Papua, Indonesia. DNA samples were isolated from Dried Blood Spots (DBS) obtained from *P.falciparum* infected patients in the Regional Public Hospital of Merauke, Province of Papua, Indonesia during May 2019-July 2019. The isolated DNAs were then amplified for nested Polymerase Chain Reaction (PCR) prior to Pfglurp genotyping. The glurp gene was identified in all 51 DBS samples of *P.falciparum*-infected patients, and 18 variants of allele were found. Among them, 45.10% were found to bear multigenotype infections. The size of the dominant allele (12.5%) was 701-750 bp. The MOI was 1.58. The genetic population of *P.falciparum* in Merauke Hospital has contained a higher percentage of multigenotypes compared with monogenotypes indicating the high transmission of malaria in the studied area.

Keywords: *Plasmodium falciparum*, Pfglurp, genotyping, polymerase chain reaction

Malaria remains an important health problem in Indonesia. Papua province has the highest burden of malaria in Indonesia. Merauke is a district easternmost in Indonesia and still has the problem of malaria, the district orders directly with the state of Papua New Guinea. Annual Parasite Incidence (API) in the district Merauke in 2010 with 19/1000 inhabitants.¹ Although all *Plasmodium* species can be found in Papua, including *Plasmodium knowlesi* formerly found in Kalimantan, the most common type of infection in Papua is by *Plasmodium falciparum* and *Plasmodium vivax*. However, *P.falciparum* is the main cause of morbidity and mortality of malaria in Papua.^{2,3}

Genetic diversity of *P.falciparum* is the major characteristic and a factor by, which the parasites survive the immune response of hosts. It results from allelic polymorphism, recombination, chromosome rearrangements, and antigenic variation.⁴ Genetic diversity of malaria parasites represents a major

issue in understanding several aspects of malaria infection and disease transmission dynamics and has hampered the malaria vaccine development.⁵ The genetic diversity was influenced by several factors such as the irregular use of anti-malaria drugs, which cause parasite resistance to the drugs, people mobilization from malaria-endemic area to non-endemic ones.⁶

Plasmodium falciparum glutamate-rich protein (Pfglurp) is a protein expressed on both pre-erythrocytic and erythrocytic stages of the parasite as well as newly emerging merozoites has high immunogenicity and acts as an antibody target involved in cellular inhibitor in monocytes, has been extensively studied as a vaccine candidate.⁷⁻⁹ The Pfglurp gene has been extensively used as a marker to investigate the genetic diversity, Multiplicity of Infection (MOI), the level of malaria transmission, immunity against malaria, as well as a discriminatory instrument to distinguish new from recrudescence

infections of the field parasite population.¹⁰

The genotyping of Pfglurp in the Merauke district, Papua Province has not yet been reported. This study was aimed to evaluate the genetic population of *P.falciparum* by nested PCR among malaria-infected patients attending the Merauke Hospital in Papua Province, Indonesia.

METHODS

Ethical clearance was obtained from the Ethics Committee for Health Research, Faculty of Medicine, Airlangga University with number 169/EC/KEPK/FKUA/2019. Dried Blood Spots (DBS) were acquired from patients suffering from malaria who visited the Emergency Department, Specialist Outpatient Clinic, and Inpatient Department of the Regional Public Hospital of Merauke, Papua during May-July 2019. The DNA isolation was performed according to the protocols of Norgen's Dried Blood Spot DNA Isolation Kit (Norgen, Thorold, Canada). The DNA was then stored at -20°C until used.

Two kinds of nested PCR were performed to identify Plasmodium species and to identify Pfglurp genotype variants. The PCR to genotype the Pfglurp used primers specific for glurp gene locus as shown in Table 1.¹¹

The first PCR mixture consisted of 12.5 µL of 2x Go Taq green master mix (Promega, Madison, USA.), 2.5 µL of each primer forward and reverse (10 µM) and 1 µL of DNA template. The total volume was 25 µL. The PCR condition for glurp genotype can be seen in Table 2. Electrophoresis of PCR products was

performed in a 2% agarose gel along with the DNA marker of 100 bp interval (Promega, Madison, USA). The DNA products were then visualized and documented by GeldocEZ imager (Biorad, USA). Furthermore, MOI was determined by dividing the total number of fragments by the number of samples positive containing glurp.¹⁰

The data were descriptively analyzed by counting the number of DNA fragments. The size of fragments was then differentiated every 50 bp of band sizes to identify the number of allelic variants and their frequency distribution. The allelic frequency was calculated by dividing the number of a particular allele by the total number of samples positive for that allelic family of the gene.⁷ Multiplicity of infection was defined as the number of parasite genotypes per infection, which was obtained by dividing the total number of fragments by the number of samples positive containing glurp.^{7,9,10,12,13} Twenty samples with more than one allelic variant were considered as a multigenotype infection while the presence of a single allelic variant was considered as monogenotype infection.¹²

RESULT AND DISCUSSIONS

The age of patients was grouped based on Depkes RI 2009 to categorize the genotype based on age group. The grouping of age can be seen in Table 3.¹⁴

Molecular diagnosis by nested PCR resulted in 51 samples, which were positive *P.falciparum* consisting of 34 (66.7%) males and 17 (33.3%) females. The

Table 1. Primer sequences for genotyping of Pfglurp

Amplification	Primer	Sequence (5' → 3')
Primary	GF3	ACATGCAAGTGTGATCCTGAA
	GF4	TGTAGGTACCACGGGTTCTTGTGG
Secondary	GF4	TGTAGGTACCACGGGTTCTTGTGG
	GNF	TGTTCACTGAACAATTAGATTAGATCA

Table 2. The PCR condition for glurp genotype

PCR Step	Nested 1			Nested 2		
	T (°C)	Time (minutes)	Number of Cycle	T (°C)	Time (minutes)	Number of Cycle
Initial denaturation	95	5	1	94	5	1
Denaturation	94	1	30	94	1	35
Annealing	58	2		59	2	
Extension	72	2		72	2	
Final extension	72	10		72	10	1

T: Temperature

31

average age of patients was 10.08 years old. The representative picture of PCR-based diagnosis can be seen in Figure 1.

Genotyping by nested PCR resulted in all 51 *P.falciparum*-infected samples containing the glurp gene, and the second nest PCR showed the allelic variation of glurp genotype with the size of bands ranging from 250 bp–1315 bp. The representative picture of PCR-based glurp genotyping can be seen in Figure 2. Eighteen variants of the Pfglurp gene were found consisted of the PCR fragments with sizes ranging from 201 to 1250 bp. Correlation of the variants of Pfglurp genotype with a group of age showed that age group of 17-25 years old (late teen) contained the highest number of variants (19 variants) and the lowest number found in 0-5 years old age group (toddler), which contained 3 variants. The increased age of the number of variants tended to increase, however, decreased in 56-65 years old of age (late elderly) as shown in Table 4.

The Pfglurp gene locus has plenty of allele variations, enabling this genotyping to identify *P.falciparum* genetic variation in a population.¹⁵ The Pfglurp gene allele variation is directly associated with endemicity and the number of evaluated

samples.¹⁶ The glurp genotype variation found in this study ranged from 250 bp to 1315 bp in size similar to in Pesawaran District, Province of Lampung, Indonesia, where 200-1200 bp were found.¹⁷ A different result was inferred from a study in Central Sumba, East Nusa Tenggara, with a glurp gene of 700-1100 bp, and a study in Sulawesi and Kalimantan identified 580-119 bp.^{18,19} The finding of 18 variants of the allele in this current study was the same as found in West Cambodia, while in southwestern Nigeria was 12, in moderate malaria-endemic areas in East Malaysia showed 5 allele variants.^{12,20,21} This finding proved the different locations have a different population of parasite genetics as well. These diversities indicated the level of malaria endemicity worldwide. The three dominant allele variants were 701-750 bp, 651-700 bp, and 951-100 bp, respectively, which showed more than a 10% frequency (Figure 3). These allelic variant profiles are valuable baseline data for continued monitoring of polymorphisms associated with antimalarial drug resistance in these areas.

Based on the number of genotypes in glurp gene locus analysis resulted in the MOI of 1.57. The number of Pfglurp variant genotypes per 51 *P.falciparum*-infected samples found in the Merauke Hospital was 80. The MOI rate usually reflects the rate of transmission intensity, although its correlation is not linear.²² The low MOI in the isolates with a higher parasite density shows the presence of dominant monogenotype infection without significant rivalry.⁵ The monogenotype infection found in this study was 54.90% and multigenotype infection was 45.10%, as shown in Table 5. A monogenotype infection is an infection caused by a single genotype, whereas a multigenotype infection is caused by two or more genotypes in one sample within the same locus.¹⁸ The multigenotype infection found in this study was higher than that was found in Thailand-Myanmar

Table 3. Grouping of the age of *P.falciparum*-infected patients at the Regional Public Merauke Hospital

Category	Age (years)
Toddler	0–5
Children	6–11
Early teens	12–16
Late teens	17–25
Young adults	26–35
Adults	36–45
Early elderly	46–55
Late elderly	56–65



Figure 1. Representative picture of *P.falciparum* species M: marker. The positive *P.falciparum* samples were at 206 bp (white arrow)




Figure 2. Representative picture of PCR-based Pfglurp genotyping. Genotype containing one PCR fragment is monogenotype and two or more fragments are multigenotype

Table 4. The frequency of genotype based on age group

Variant Allele	Size (bp)	Age (years)								Frequency (%)
		0-5	6-11	12-16	17-25	26-35	36-45	46-55	56-65	
1	201-250	-	1	-	1	-	2	1	1	6 ((7.5)
2	251-300	-	-	-	-	1	-	1	-	2 (2.5)
3	351-400	-	-	-	1	-	-	-	-	1 (1.25)
4	401-450	-	1	1	2	2	-	1	-	7 (8.75)
5	451-500	-	-	1	1	-	-	-	-	2 (2.5)
6	501-550	-	-	-	1	-	-	-	1	2 (2.5)
7	601-650	-	-	-	-	-	-	1	-	1 (1.25)
8	651-700	-	-	2	2	2	-	2	1	9(11.25)
9	701-750	-	1	-	4	1	1	2	1	10 (12.5)
10	751-800	1	-	2	-	1	1	-	2	7 (8.75)
11	801-850	-	-	-	1	1	2	-	-	4 (5)
12	851-900	-	-	3	1	1	1	1	-	7 (8.75)
13	901-950	1	-	-	1	-	1	1	-	4 (5)
14	951-1000	-	2	1	2	1	1	-	1	8 (10)
15	1001-1050	-	-	1	-	-	-	-	-	1 (1.25)
16	1050-1100	1	-	-	-	1	-	-	-	2 (2.5)
17	1151-1200	-	-	2	1	-	1	2	-	6 (7.5)
18	1301-1350	-	-	-	1	-	-	-	-	1 (1.25)
Number (%)		3 (3.75)	5 (6.25)	13 (16.25)	19 (23.75)	11(13. 75)	10 (12.5)	12 (15)	7 (8.75)	80 (100)




Figure 3. Pie chart of the frequency distribution (%) of PfgIurp gene in *P.falciparum* patients at the Regional Public Hospital of Merauke. The three dominant sizes of allelic variant genotype were 701-750 bp (12.5%), 651-700 bp (11.25%), and 951-1000 bp (10%)

Table 5. Summary of glurp genotypes in *P.falciparum*-infected patients attending the Merauke Hospital, Papua

Kind of Genotypes	Number of Samples (%)	Number of Genotypes (%)
Monogenotype	28 (54.90)	28 (35)
Multigenotype		
2 Genotype	18 (35.30)	36 (45)
3 Genotype	4 (7.84)	12 (15)
4 Genotype	1 (1.96)	4 (5)
Total number of multigenotype	23 (45.10)	52 (65)
Total number of genotype	51 (100)	80 (100)
MOI = 1.57		

border, which was 15.3%, while in Western Cambodia the multigenotype infection was 8%.^{21,23} The high rate of multigenotype infection found in this study showed that the intensity of transmission at the study site was still high.¹³ A study in Gabon that evaluated the correlation between multigenotype infection and cytoadherence (the adherence of parasite-infected erythrocytes) showed that multigenotype infection, as it turned out, had no significant effect on cytoadherence or the severity of malaria. The severity of malaria was also not affected by the type of certain genotype and the amount of infecting *P.falciparum* genotype.²⁴

The use of glurp gene locus has its benefits in epidemiology tracking and to evaluate the transmission rate and the correlation of its factors to the immune system. An immunology study in a high and low malaria contagious area showed a high prevalence of glurp antibodies in adults, as well as a significant correlation between a specific high level of glurp antibody with low parasite density and clinical protection against malaria. Furthermore, *P.falciparum* infection often triggers antibody production as a response to glurp, which naturally inhibits in-vitro growth of *P.falciparum*, with or without contribution from monocytes, making glurp eligible for parasitemia control.²⁵

CONCLUSION AND SUGGESTION

The genetic population of *P.falciparum* in the Merauke Hospital containing a higher percentage of multigenotypes compared with monogenotypes indicated the high transmission of malaria in the study area. As long as the population mobility is still high, the genetic transformation will be high and so will the transmission of malaria.

ACKNOWLEDGEMENTS

This study was supported by the Directorate of Research and Public Service of the Ministry of

Research and Technology, Republic of Indonesia contract number: 544/UN3.14/LT/2019. Researchers thank the Laboratory Technologists who helped in blood collection and microscopic examination.

REFERENCES

- Shinta S, Marjana P. Distribusi dan perilaku vektor malaria di Kabupaten Merauke, Papua. Buletin Penelitian Kesehatan, 2016; 43(4): 219-230.
- Surjadjaja C, Surya A, Baird JK. Epidemiology of *Plasmodium vivax* in Indonesia. American Journal of Tropical Medicine and Hygiene, 2016; 95(69): 121-132.
- WHO. Indonesia South-East Asia region-malaria profile. Published 2018. Available from: https://www.who.int/malaria/publications/country-profiles/profile_idn_en.pdf?ua=1 (accessed 20 August, 2020).
- Kemp DJ, Cowman AF, Walliker D. Genetic Diversity in *Plasmodium falciparum*. Advances in Parasitology, 1990; 29: 75-149.
- Kidima W, Nkwengulila G. Multiplicity of infections among children under five years with uncomplicated malaria in Kibaha, Tanzania. Journal of Parasitology Research, 2015; 2015: 1-6.
- Sillehu S, Arwati H, Dachlan YP, Keman S. Genetic polymorphism of *Plasmodium falciparum* merozoite surface protein-1 (Pfmsp-1) in closed and opened community at South Buru District, Maluku Province. DamaInt J Res, 2016; 1(9): 1-4.
- Kumar D, Dhiman S, Rabha B, Goswami D, Deka M, et al. Genetic polymorphism and amino acid sequence variation in *Plasmodium falciparum* glurp R2 repeat region in Assam, India, at an interval of five years. Malar J, 2014; 13(1): 1-8.
- Patel P, Bharti PK, Bansal D, Raman RK, Mohapatra PK, et al. Genetic diversity and antibody responses against *Plasmodium falciparum* vaccine candidate genes from Chhattisgarh, Central India: Implication for vaccine development. PLoS One, 2017; 12(8): 1-17.
- Kaur H, Sehgal R, Goyal K, Makkar N, Yadav R, et al. Genetic diversity of *Plasmodium falciparum* merozoite surface protein-1 (block 2), glutamate-rich protein and sexual stage antigen Pfs25 from Chandigarh, North India. Trop Med Int Heal, 2017; 22(12): 1590-1598.

10. Abamecha A, El-Abid H, Yilma D, Addisu D, Ibenthal A, *et al.* Genetic diversity and genotype multiplicity of *Plasmodium falciparum* infection in patients with uncomplicated malaria in Chewaka district, Ethiopia. *Malar J*, 2020; 19(1):1-9.
11. Zhou X, Huang JL, Njuabe MT, Li SG, Chen JH, *et al.* A molecular survey of febrile cases in malaria-endemic areas along China-Myanmar border in Yunnan Province, People's Republic of China. *Parasite*, 2014; 21(27): 1-8.
12. Razak MRMA, Sastu UR, Norahmad NA, Abdu-Karim A, Muhammad A, *et al.* Genetic diversity of *Plasmodium falciparum* populations in malaria declining areas of Sabah, East Malaysia. *PLoS One*, 2016; 11(3):1-22.
13. Soe TN, Wu Y, Tun MW, Xu X, Hu Y, *et al.* Genetic diversity of *Plasmodium falciparum* populations in Southeast and Western Myanmar. *Parasit Vectors*, 2017; 10(1): 322.
14. Amin M Al, Juniati D. Klasifikasi kelompok umur manusia berdasarkan analisis dimensi fraktal box counting dari citra wajah dengan deteksi Tepi Canny. *J Ilm Mat*, 2017; 2(6): 1-10.
15. Mwingira F, Nkwengulila G, Schoepflin S, Sumari D, Beck H, *et al.* *Plasmodium falciparum* msp1, msp2 and glurp allele frequency and diversity in sub-Saharan Africa. *Malar J*, 2011; 10(1): 1-10.
16. Maestre A, Arango E, Carmona-Fonseca J. Status of allele frequency and diversity of *Plasmodium falciparum* msp1, msp2 and glurp before implementation of an Artemisinin-based combined therapy in Northwestern Colombia. *Colomb Med*, 2013; 44(4): 208-212.
17. Aziz R, Kumiawan B, Mutiara H, Suwandi JF. Identifikasi gen *Plasmodium falciparum* glutamate rich protein (Pfglurp) dari penderita malaria di wilayah kerja Puskesmas Hanura Kabupaten Pesawaran Provinsi Lampung. *J Major*, 2018; 7(2): 108-111.
18. Mau F, Murhandarwati EEH. Keragaman genetik dari msp 1, msp 2, dan glurp pada *Plasmodium falciparum* di Kabupaten Sumba Tengah, Nusa Tenggara Timur. *Bul Penelit Kesehat*, 2016; 44(2): 77-84.
19. Handayani S, Salwati E, Tjitra E. Keragaman genetik petanda *P.falciparum* dari specimen subyek penelitian monitoring Dihydroartemisinin-Piperrakuin di Kalimantan dan Sulawesi. *Media Penelit dan Pengemb Kesehat*, 2012; 22(3): 120-130.
20. Funwei RI, Thomas BN, Falade CO, Ojurongbe O. Extensive diversity in the allelic frequency of *Plasmodium falciparum* merozoite surface proteins and glutamate-rich protein in rural and urban settings of Southwestern Nigeria. *Malar J*, 2018; 17(1):1-8.
21. Gosi P, Lanteri CA, Tyner SD, Se Y, Lon C, *et al.* Evaluation of parasite sub-populations and genetic diversity of the msp1, msp2 and glurp genes during and following artesunate monotherapy treatment of *Plasmodium falciparum* malaria in Western Cambodia. *Malar J*, 2013; 12(1):403.
22. Paul REL, Hackford I, Brockman A, Muller-Graf C, Price R, *et al.* Transmission intensity and *Plasmodium falciparum* diversity on the Northwestern border of Thailand. *Am J Trop Med Hyg*, 1998; 58(2): 195-203.
23. Congpuong K, Sukaram R, Prompan Y, Dornae A. diversity of the msp-1, msp-2, and glurp genes of *Plasmodium falciparum* isolates along the Thai-Myanmar borders genetic. *Asian Pac J Trop Biomed*, 2014; 4(8): 598-602.
24. Touré FS, Ouwe-Missi-Oukem-Boyer O, Mezui-Me-Ndong J, Ndong-Atome GR, Bisvigou U, *et al.* Cytoadherence and genotype of *Plasmodium falciparum* strains from symptomatic children in Franceville, Southeastern Gabon. *Clin Med Res*, 2007; 5(2): 106-113.
25. Pratt-Riccio LR, Perce-da-Silva D de S, Lima-Junior J da C, Theisen M, Santos F, *et al.* Genetic polymorphisms in the glutamate-rich protein of *Plasmodium falciparum* field isolates from a malaria-endemic area of Brazil. *Mem Inst Oswaldo Cruz*, 2013; 108(4): 523-528.

39. Genetic Diversity of Plasmodium falciparum Glutamate Rich Protein Patients Attending the Merauke Hospital in Papua Province, Indonesia

ORIGINALITY REPORT

15%

SIMILARITY INDEX

12%

INTERNET SOURCES

13%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- | | | |
|---|--|----|
| 1 | journal2.unusa.ac.id
Internet Source | 1% |
| 2 | Tolla NDIAYE, Mouhamad Sy, Amy Gaye, Katherine J. Siddle et al. "Molecular epidemiology of Plasmodium falciparum by multiplexed amplicon deep sequencing in Senegal", Research Square Platform LLC, 2020
Publication | 1% |
| 3 | Fanomezantsoa Ralinoro, Tovonahary Angelo Rakotomanga, Rianasoambolanoro Rakotosaona, Danielle A. Doll Rakoto et al. "Genetic diversity of Plasmodium falciparum populations in three malaria transmission settings in Madagascar", Malaria Journal, 2021
Publication | 1% |
| 4 | www.tandfonline.com
Internet Source | 1% |
| 5 | downloads.hindawi.com
Internet Source | 1% |

6	"Poster Presentations P001 - P142", Pediatric Nephrology, 2004 Publication	1 %
7	www.researchsquare.com Internet Source	1 %
8	Voahangy Hanitriniaina Isabelle ANDRIANARANJAKA, Elisabeth Ravaoarisoa, Tovonanahary Angelo Rakotomanga, Fanomezantsoa Ralinoro et al. "DNA recovery from used malaria RDT to detect Plasmodium species and to assess Plasmodium falciparum genetic diversity: A pilot study in Madagascar", Research Square Platform LLC, 2022 Publication	1 %
9	www.amedeo.com Internet Source	1 %
10	f1000research.com Internet Source	1 %
11	d.lib.msu.edu Internet Source	<1 %
12	repository.unair.ac.id Internet Source	<1 %
13	www.nature.com Internet Source	<1 %

keperawatan.unri.ac.id

14	Internet Source	<1 %
15	www.mdpi.com Internet Source	<1 %
16	damaacademia.com Internet Source	<1 %
17	kipdf.com Internet Source	<1 %
18	www.bioflux.com.ro Internet Source	<1 %
19	themimu.info Internet Source	<1 %
20	Cang-Lin Zhang, Hong-Ning Zhou, Quan Liu, Ya-Ming Yang. "Genetic polymorphism of merozoite surface proteins 1 and 2 of Plasmodium falciparum in the China-Myanmar border region", Malaria Journal, 2019 Publication	<1 %
21	Otchere Addai-Mensah, Bismarck Dinko, Mark Noagbe, Selassie Louis Ameke et al. "Plasmodium falciparum histidine-rich protein 2 (PfHRP2) diversity in Ghana", Research Square Platform LLC, 2020 Publication	<1 %

22 S. Jordan. "Population structure of *Plasmodium falciparum* isolates during an epidemic in southern Mauritania", *Tropical Medicine and International Health*, 10/2001
Publication <1 %

23 journals.plos.org
Internet Source <1 %

24 norgenbiotek.com
Internet Source <1 %

25 Anyirékun Fabrice Somé, Thomas Bazié, Issaka Zongo, R. Serge Yerbanga et al. "Plasmodium falciparum msp1 and msp2 genetic diversity and allele frequencies in parasites isolated from symptomatic malaria patients in Bobo-Dioulasso, Burkina Faso", *Parasites & Vectors*, 2018
Publication <1 %

26 Bharti, Praveen K, Man M Shukla, Yagya D Sharma, and Neeru Singh. "Genetic diversity in the block 2 region of the merozoite surface protein-1 of *Plasmodium falciparum* in central India", *Malaria Journal*, 2012.
Publication <1 %

27 Wahyu Yuniati, Kuswantoro Rusca Putra, Dhelya Widasmara. "Brøset Violence Checklist (BVC) versus Dynamic Appraisal of Situational Aggression-Inpatient Version (DASA-IV): their

sensitivity, specificity and accuracy as a predictor of violence in 24 hours among schizophrenia patients", MEDISAINS, 2020

Publication

28

mafiadoc.com

Internet Source

<1 %

29

publisher.uthm.edu.my

Internet Source

<1 %

30

Puspa Wardhani, Trieva Verawaty Butarbutar, Christophorus Oetama Adiatmaja, Amarensi Milka Betaubun, Nur Hamidah, Aryati.

"Performance comparison of two malaria rapid diagnostic test with real time polymerase chain reaction and gold standard of microscopy detection method", Infectious Disease Reports, 2020

Publication

<1 %

31

"Poster Abstracts", Nephrology, 5/2008

Publication

<1 %

32

Nancy O. Duah, Sena A. Matrevi, Neils B. Quashie, Benjamin Abuaku, Kwadwo A. Koram. "Genetic diversity of Plasmodium falciparum isolates from uncomplicated malaria cases in Ghana over a decade", Parasites & Vectors, 2016

Publication

<1 %

33

eprints.undip.ac.id

Internet Source

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

39. Genetic Diversity of Plasmodium falciparum Glutamate Rich Protein Patients Attending the Merauke Hospital in Papua Province, Indonesia

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6
