


# Journal of Public Health in Africa

Volume 17

Number 1

Blackwell  
Publishing

blackwell  
publishing





### Death risk among COVID-19 patients with diabetes mellitus

Dahlia Dahlia, Kurnia Dwi Artanti, **Arief Hargono**, Santi Martini, Nayla Mohamed Gomaa Nasr, Chung-Yi Li

<https://doi.org/10.4081/jphia.2022.2399>

150 PDF: 176



### Proximate analysis and fiber content of smoothies fortified with Chia seeds

Farah Paramita, Septa Katmawanti, Agung Kurniawan

<https://doi.org/10.4081/jphia.2022.2407>

146 PDF: 173



### We are facing some barriers: A qualitative study on the implementation of kangaroo mother care from the perspectives of healthcare providers

Ory Tifani Rahmatika, Ronal Surya Aditya, Ah. Yusuf, Reem lafi Almutairi, Daifallah M. Al Razeeni, Siti Kotijah, Anita Sulistyorini

<https://doi.org/10.4081/jphia.2022.2407>





Journal of  
Public Health in  
Africa

[HOME](#) / Editorial Board

# Editorial Board

## Editors-in-Chief

### **Prof. Nicaise Ndembi**

Institute of Human Virology, *Nigeria*

### **Prof. Vittorio Colizzi**

University of Rome Tor Vergata, *Italy*

Faculty of Science & Technology, Evangelic University of Cameroon, *Cameroon*

[vittorio.colizzi@publichealthinafrica.org](mailto:vittorio.colizzi@publichealthinafrica.org)

## Managing Editor

### **Mrs. Emanuela Fusinato**

PAGEPress, *Italy*

## Board Members

### **Dr. John Nkengasong**

Africa Centres for Disease Control and Prevention, *Cameroon*

### **Dr. Chikwe Ihekweazu**

Nigeria Centres for Disease Control and Prevention, *Nigeria*

### **Dr. Alex Riolexus Ario**

Uganda National Institute of Public Health, Ministry of Health, *Uganda*

### **Dr. Elvira Singh**

National Cancer Registry, National Health Laboratory Service, *South Africa*

### **Prof. Abderrahmane Maaroufi**

Institut Pasteur du Maroc [National Public Health Institute], *Morocco*

### **Dr. Ileshi Jani**

National Institute of Health, *Mozambique*

### **Dr. Ebba Abate**

Ethiopian Public Health Institute, Addis Ababa, *Ethiopia*

### **Dr. Mazyanga Lucy Mazaba Liwewe**

Zambia National Public Health Institute, *Zambia*

**Prof. Vincent Batwala**

Mbarara University of Science & Technology, *Uganda*

**Dr. Giacomo Paganotti**

Botswana-University of Pennsylvania, *Botswana*

**Dr. Georges Etoundi**

Direction de la Lutte Contra la Maladie, les Epidemies et les Pandemies, *Cameroon*

**Prof. Jacques Simpore**

Joseph Ki-Zerbo University, *Burkina Faso*

**Dr. Tolbert Nyenswah**

Johns Hopkins University Bloomberg School of Public Health, *United States*

**Dr. Souha Bougatef**

National Observatory of New and Emerging Diseases, *Tunisia*

**Dr. Raji Tajudeen**

Africa Centres for Disease Control and Prevention, *Ethiopia*

**Dr. Mohammed Abdulaziz**

Africa Centres for Disease Control and Prevention, *Ethiopia*

**Prof. Epée Emilienne**

University of Yaoundé I, *Cameroon*

**Dr. Nafisah Chotun**

Africa Centres for Disease Control and Prevention, *Ethiopia*

**FOR AUTHORS****SUBMIT YOUR PAPER**

---

Guide for Authors

---

Benefits for Authors

---

How to write a scientific paper

---

How to write a Review article

---

Article Processing Charge

---

**FOR REVIEWERS**

---

Benefits for Reviewers

---

## Death risk among COVID-19 patients with diabetes mellitus

Dahlia Dahlia,<sup>1</sup> Kurnia Dwi Artanti,<sup>2</sup>  
Arief Hargono,<sup>2</sup> Santi Martini,<sup>2</sup>  
Nayla Mohamed Gomaa Nasr,<sup>3</sup>  
Chung-Yi Li,<sup>2,4</sup>

<sup>1</sup>Student of Bachelor Program of Public Health, Universitas Airlangga, Indonesia; <sup>2</sup>Division of Epidemiology, Faculty of Public Health, Universitas Airlangga, Indonesia; <sup>3</sup>Doctoral candidate at the University of Debrecen, Faculty of Medicine, Department of Public Health and Epidemiology, Debrecen, Hungary; <sup>4</sup>Department of Public Health, National Cheng Kung University, Taiwan

### Abstract

Deaths from COVID-19 are increasing in patients with comorbidities. One of the most common comorbidities is diabetes mellitus. The researchers wanted to see how having diabetes affected the mortality rate of COVID-19 participants. This investigation is a case control observational analytical study. Different types of people, called “cases,” and “controls,” complete the research sample. Each group had 68 responders, for a grand total of 136. Medical records from COVID-19 patients treated at Airlangga University Hospital, Surabaya, between March 2020 and September 2021 serve as the study’s secondary data source. The purpose of this study’s data analysis is to calculate an odds ratio. Patients with COVID-19 with concomitant diabetes mellitus had an increased risk of death, and this risk increased with age, gender, and COVID-19 symptoms. In contrast, education, occupation, and laboratory results were not significantly related to mortality among COVID-19 individuals with concomitant diabetes mellitus (GDA status). The results of this study show that COVID-19 patients with concomitant diabetes mellitus are at a higher risk of death if they are over the age of 65, if they are male, and if they have severe symptoms.

### Introduction

COVID-19 or commonly called Coronavirus disease 2019 by the World Health Organization is a new viral disease that is currently making people around the world anxious and has a significant impact

on human life, especially in the health sector. The corona virus is likely to evolve further, quicker, causing more severe infection organ failure and, ultimately, death. This state of emergency is observed even more frequently in patients with previous health problems or a history of illness.<sup>1</sup> It was found that the mortality rate due to COVID-19 is increasing in patients with comorbidities. Based on a report from the Central of Disease Control (CDC), COVID-19 is twelve times more deadly for patients with comorbidities than patients without comorbid diseases. The mortality rate (CFR) for patients with cardiovascular disease is 10.5%, 7.3% for people with diabetes mellitus, 6.3% for patients with chronic respiratory disease, 6% for patients with hypertension, and 5.6% for cancer patients.<sup>2</sup>

Comorbidity is a risk factor for the severity of COVID-19. One of the comorbidities is diabetes mellitus. The problem with diabetes is primarily a poorer prognosis, not a greater chance of contracting the virus. Diabetes mellitus is characterized by high glucose levels. High glucose levels tend to worsen the disease a patient has, including COVID-19 itself. This is because high glucose levels can affect the ability of the virus to infect humans, increase the risk of inflammation and compromise the body’s immune system. Therefore, the morbidity and mortality rates of COVID-19 in patients with diabetes mellitus are significantly higher than in non-diabetic patients. This study aimed to analyze the risk of death in COVID-19 patients with comorbid diabetes mellitus.

### Materials and Methods

The research method used in this study was a cross-sectional analytical observational study. The time frame for this study was between October 2021 and April 2022. The study relied on secondary sources of information, specifically the medical records of inpatients at the Universitas Airlangga Hospital in Surabaya between March 2020 and September 2021. All 257 people enrolled in COVID-19 who also had diabetes mellitus were analyzed in this study. Patients with COVID-19 and concomitant diabetes mellitus who did not survive were considered the case population, while those who did survive were considered the control group. A sample was selected from the study population that met the inclusion criteria during the period March 2020 to September 2021 with a ratio of the case group to the control group 1:1. The total sample in this study was 136 people.

Data analysis was performed univari-

Correspondence: Santi Martini, Faculty of Public Health, Universitas Airlangga, Jl. Mulyorejo, Surabaya, 60115, East Java, Indonesia  
Tel.: +62 315920948, +72 5924618.  
E-mail: santi-m@fkm.unair.ac.id

Key words: Age, COVID-19, Death, Diabetes mellitus, Men

Contributions: Design the work: DD. Data collection: KDA. Data analysis: DD. Interpretation of data: DD, SM. Drafting: DD. Review & editing: DD, KDA, AH, SM, NMG, CYL.

Conflict of interest: The authors declare that there is no conflict of interest.

Acknowledgments: We would like to thank Kurnia Dwi Artanti, Arief Hargono, Santi Martini, Division of Epidemiology Faculty of Public Health, Airlangga University, Nayla Mohamed Gomaa Nasr, Doctoral candidate at the University of Debrecen (Hungary), Faculty of Medicine, Department of Public Health and Epidemiology and Chung Yi-Li, Department of Public Health, College of Medicine, National Cheng Kung University, Taiwan as mentors who supported this journal.

Conference presentation: This article was presented at the 4<sup>th</sup> International Scientific Meeting on Public Health and Sports (ISMOPHS 2022).

Received for publication: 6 October 2022.  
Accepted for publication: 14 November 2022.

This work is licensed under a Creative Commons Attribution NonCommercial 4.0 License (CC BY-NC 4.0).

©Copyright: the Author(s), 2022  
Licensee PAGEPress, Italy  
Journal of Public Health in Africa 2022; 13(s2):2399  
doi:10.4081/jphia.2022.2399

*Publisher's note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.*

ately and bivariate. In this study, univariate analysis was intended to identify the mortality rate in COVID-19 patients with comorbid diabetes mellitus based on patient characteristics including age, gender, education and occupation and the mortality rate in COVID-19 patients based on patient clinical characteristics which include symptoms of COVID-19 and laboratory results (RBS

status). Bivariate analysis was conducted to determine the relationship between each independent variable in the form of risk factors including age, gender, education, occupation, COVID-19 symptoms and laboratory results (RBS status) with the dependent variable, namely death in COVID-19 patients.

## Results

### Respondent characteristics description

The description of respondent characteristics is presented in Table 1 which shows that the sample size studied was 136 people. As a percentage, the elderly made up 38.2% of the COVID-19 cohort with concomitant diabetes mellitus. Fifty-seven percent were male. With 41.2%, high school diploma or equivalent is the most prevalent degree of education. Of the COVID-19 patients who also have diabetes mellitus, 62.5% are employed. COVID-19 patients who also had diabetes mellitus had a 50% higher chance of having moderate symptoms. A majority (55.7%) of COVID-19 patients with concomitant diabetes mellitus had uncontrolled blood sugar levels as determined by a Random Blood Sugar Status (RBA).

### Risk factors for COVID-19 patient mortality with comorbid diabetes mellitus

Table 2 displays the causes of death among COVID-19 patients. If we look at the OR value, we see that respondents who are 65 and older have a 2.10 times higher mortality rate than those who are 20-35. The 95% CI ranged from (1.82 to OR 5.37) with a mean of 2.88. It means that COVID-19 patients with concomitant diabetes mellitus have a higher death rate as they get older. According to the OR value, men have a 2.16 times higher mortality rate than women. As a result, COVID-19 patients who also had diabetes mellitus died at a higher rate if they were male. The 95% CI for the correlation between education and likelihood of having a health problem was (0.72 to OR 8.59). This finding suggests that among COVID-19 patients who also had diabetes mellitus, there was no correlation between educational attainment and mortality. 95% confidence interval for the occupation was (0.68, OR, 2.74). Patients with COVID-19 and concomitant diabetes mellitus showed no statistically significant association between occupation and mortality. People with severe symptoms are 10.07 times more likely to die than those with no symptoms, as indicated by the OR value. The 95% CI ranged from (OR 2.21) to (OR

45.91). What this means is that COVID-19 individuals who also have diabetes mellitus have a far higher risk of dying from their condition. The odds ratio (95% CI) for the status of random blood sugar was (0.92–3.61). Patients with COVID-19 with concomitant diabetes mellitus showed no statistically significant correlation between Random Blood Sugar status and mortality.

## Discussion

### COVID-19 patients with concurrent diabetes mellitus: does older age increase the risk of death

Old age may worsen an individual's health problem due to the risk factors that influence it. In the aging process, the body undergoes physiological changes, resulting in the body being more susceptible to disease. Therefore, the elderly often develops health problems. This happens due to body cell degradation, thus, the body's resistance and functions decrease and the risk factors for disease increase.<sup>3</sup> Age can also affect the severity of disease and death in patients. This also applies to COVID-19 patients. Patients with diabetes mellitus are at an increased risk of dying. A combination of advanced age and diabetes mellitus predicts a lower chance of success.<sup>4</sup> Individuals older than 65 years old accounted for 80% of COVID-19 fatalities. Therefore, old age

might be viewed as a risk factor for mortality caused by COVID-19.<sup>5</sup> This study's findings are consistent with those of Raden (2020), who found that patients older than 64 faced a 2.097-fold increased probability of dying.<sup>6</sup> It was also found in the study by Clement Drew (2021) that COVID-19 patients older than 60 had a 6.71-fold increased risk of mortality.<sup>7</sup> Research by Zhang (2020) using the logistic regression method shows that age affects patient mortality, and these findings are important to our investigation.<sup>8</sup>

### Comparative analysis of gender and mortality in COVID-19 patients with concurrent diabetes mellitus

One of the risk factors for dying from COVID-19 is one's gender. This occurs because men and women's immune systems are fundamentally different, as are their lifestyles.<sup>9</sup> Hormonal differences contribute to the higher mortality rate in men. Through immunological effects, the hormone estrogen on the female immune system will have a positive effect on fighting infection. In addition, Sexual hormones like progesterone, which is only found in females, play a significant impact in both the innate and adaptive immune systems.<sup>10</sup> Similar findings were found by Clement Drew (2021), who found that males had a mortality rate 2.65 times that of females.<sup>7</sup> Biswas's study also found that men had a 1.86 times higher mortality rate than women.<sup>11</sup> Also, Siagian's (2020) research revealed that

**Table 1. Respondent characteristics description.**

Variables	Frequency (n)	Percentage (%)
Age		
20-35	6	4.4
36-55	50	36.8
56-65	52	38.2
>65	28	20.6
Gender		
Male	69	50.7
Female	67	49.3
Education		
Elementary	18	13.2
Junior High	28	20.6
Senior/Vocational High	56	41.2
College	34	25.0
Occupation		
Employed	85	62.5
Unemployed	51	37.5
COVID-19 Symptoms		
Asymptomatic	11	8.1
Mild	14	10.3
Moderate	68	50.0
Severe	43	31.6
RBS Status		
Uncontrolled	76	55.9
Controlled	60	44.1

male COVID-19 patients had a much greater mortality rate than female COVID-19 patients.<sup>12</sup>

### Death in COVID-19 patients with concurrent diabetes mellitus and low levels of education

The results of the study showed that among COVID-19 patients who also had diabetes mellitus, higher levels of education were not associated with a lower risk of death. In keeping with the findings of Margareth's (2020) study, which found no association between educational attainment and COVID-19 mortality, we find the following.<sup>13</sup> In addition, Linda's (2021) research found no correlation between patients' levels of education and their risk of dying from COVID-19.<sup>14</sup> A person's motivation to take preventative measures against COVID-19 may be influenced by their degree of education.<sup>15</sup> However, education is not a risk factor for COVID-19 individuals who also have type 2 diabetes mellitus. Comorbid diabetes mellitus is a major contributor to mortality in COVID-19 patients, as these patients have a poor prognosis overall.

### Death in COVID-19 patients with concurrent diabetes mellitus and their occupation

COVID-19 patients with concomitant diabetes mellitus did not have a different

death rate based on their career, according to the study's findings. This finding is consistent with the findings of Shahir's (2020) study, which found no correlation between patients' occupations and their risk of dying from COVID-19.<sup>17</sup> However, a different study found a correlation between patients' occupations and their risk of dying from COVID-19. The researchers noted that the workplace is a key setting for the spread of COVID-19. The type of one's work greatly alters the likelihood that one will be exposed to COVID-19.<sup>18</sup> The occupations mentioned in the study were those with a high potential for exposure to COVID-19, such as those in the medical field.<sup>19</sup> In addition, occupations that are included in high risk and comorbid diseases groups are more likely to cause COVID-19-related deaths.<sup>4</sup>

### The relationship between COVID-19 symptoms and the mortality of COVID-19 patients with comorbid diabetes mellitus

Based on the severity of the symptoms, COVID-19 is classified as asymptomatic, mild, moderate, severe, or critical. When the body is infected with COVID-19, it reacts by displaying symptoms. The body's defenses, in response to the cell death brought on by the virus's replication process, will set off an inflammatory cascade, giving birth to a wide range of symptoms. The development of symptoms is evi-

dence that the virus has successfully infected cells and is actively replicating and spreading to new targets. Accordingly, symptoms may increase a patient's likelihood of dying from COVID-19. Similar findings were found by Du (2020), who found that individuals with severe symptoms have a 7.35-fold higher probability of death.<sup>21</sup> This finding is consistent with that of Zheng and Santos (2020) in Brazil, who found that COVID-19 patients with severe symptoms had a higher mortality rate.<sup>22,23</sup> Patients with COVID-19 who also have diabetes mellitus are at increased risk for both severe COVID-19 symptoms and death, as reported by Khumar (2020).<sup>24</sup>

### The association between retinol-based sympathetic neurotransmitters and death in COVID-19 patients with concurrent diabetes mellitus

The study found no association between Random Blood Sugar (RBS) status and mortality among COVID-19 participants with concomitant diabetes mellitus. It is in line with the study by Linda (2021) stating that there was no relationship between Random Blood Sugar (RBS) status and the mortality of COVID-19 patients.<sup>5</sup> Blood sugar level is a clinical indicator for diagnosing diabetes mellitus. High blood sugar levels that interact with other factors such as old age, unhealthy diet, low physical

**Table 2. Mortality risk factors for COVID-19 patients with comorbid diabetes mellitus.**

Variables	Died		Recovered		Total		OR	CI 95%		
	n	%	n	%	n	%		Lower	Upper	
<b>Age</b>										
20-35	3	4.4	3	4.4	6	4.4	1.36	0.25	7.40	
36-55	26	38.3	24	35.3	50	36.8	1.47	0.67	3.22	
56-65	22	32.3	30	44.1	53	38.2	1			
>65	17	25.0	11	16.2	28	20.6	2.10	1.82	5.37	
<b>Gender</b>										
Male	41	60.3	28	41.1	69	50.7	2.16	1.09	4.30	
Female	27	39.7	40	58.9	69	49.3				
<b>Education</b>										
Elementary	9	13.2	9	13.2	18	13.2	2.50	0.72	8.59	
Junior High	8	11.7	20	29.5	28	20.6	1			
Senior/Vocational High	30	44.1	26	38.2	56	41.2	2.88	1.08	7.63	
College	21	31.0	13	19.1	34	25.0	4.03	1.38	11.80	
<b>Occupation</b>										
Employed	45	66.2	40	58.9	85	62.5	1.37	0.68	2.74	
Unemployed	23	33.8	28	41.1	51	37.5				
<b>COVID-19 Symptoms</b>										
Asymptomatic	3	4.4	8	11.8	11	8.1	1			
Mild	2	3.0	12	17.7	14	10.3	0.44	0.60	3.28	
Moderate	29	42.6	39	57.3	68	50.0	1.98	0.48	8.13	
Severe	34	50.0	9	13.2	43	31.6	10.07	2.21	45.91	
<b>RBS Status</b>										
Uncontrolled	43	63.2	33	48.6	76	55.9	1.82	0.92	3.61	
Controlled	25	36.8	35	51.4	60	44.1				

activity can modulate immune and inflammatory responses. Thus, patients with diabetes mellitus are susceptible to diseases including exposure to COVID-19.<sup>6</sup> In contrast to the HbA1C test, patients with high HbA1C showed higher inflammatory parameters, increased renal function and blood viscosity. Increased HbA1C will be associated with clinical COVID-19 patients. Elevated HbA1C can serve as an indicator of long-term glycemic status. This is because COVID-19 is associated with worse outcomes in patients with poorly controlled diabetes mellitus and higher mortality in those who already have the disease. Improper glycemic management contributes to poor clinical outcomes and an increased risk of death in COVID-19 patients.<sup>26</sup>

## Conclusions

This study found that COVID-19 patients with severe symptoms, those who were male, and those who were older than 65 and who had diabetes mellitus had the highest mortality rates. Patients who are >65 years old, male and have severe COVID-19 symptoms are included in the high-risk group. High risk groups should receive appropriate treatment, supervision and care.

## References

- Asfahan, S., Kunal, D., Naveen, D., Ram, N., Priyank, J. and Mehul, A., 2020. Extrapolation of mortality in COVID-19: Exploring the role of age, sex, co-morbidities and health-care related occupation. *Monaldi Archives for Chest Disease*, 90(2), pp. 313–317. Available at: 10.4081/monaldi.2020.1325
- Baker, M. G., Peckham, T. K. and Seixas, N. S., 2020. Estimating the burden of United States workers exposed to infection or disease: A key factor in containing risk of COVID-19 infection. *PLoS ONE*, 15(4), pp. 4–11. Available at: 10.1371/journal.pone.0232452
- Biswas, M., Rahaman, S., Biswas, T.K., Haque, Z. and Ibrahim, B., 2021. Association of Sex, Age, and Comorbidities with Mortality in COVID-19 Patients: A Systematic Review and Meta-Analysis. *Intervirology*, 64(1), pp. 36–47. Available at: 10.1159/000512592
- Chan, J. F. W., Yuan, S., Kok, K.H., Chu, H., Yang, J., Xing, F., Liu, J., Yip, C.C., Poon R.W., Tsoi, H.W., Lo, S.K., Chan, K.H., Poon, V.K., Chan, W.M., Ip J.D., Cai J.P., Cheng, V.C., Chen, H., Hui, C.K. and Yuen, K.A., 2020. A familial cluster of pneumonia associated with the 2019 novel coronavirus indicating person-to-person transmission: a study of a family cluster. *The Lancet*, 395(10223), pp. 514–523 Available at: 10.1016/S0140-6736(20)30154-9
- Devan Hawkins, Letitia Davis, D. K., 2021. COVID-19 deaths by occupation Massachusetts March 1 July 31 2020. Available at: <https://publons.com/publon/10.1002/ajim.23227>
- Drew, C. and Adisasmita, A. C., 2021. Gejala dan komorbid yang memengaruhi mortalitas pasien positif COVID-19 di Jakarta Timur, Maret-September 2020. *Tarumanagara Medical Journal*, 3(3), pp. 274–283. Available at: <http://journal.untar.ac.id/index.php/tmj/article/view/11742>
- Du, R. H., Yang, C.Q., Wang, W., Cao, T.Z., Li, M., Guo, G.Y., Du, J., Zheng, C.L., Zhu, Q., Hu, M., Li, X.Y., Peng, P. and Shi, H.Z., 2020. Predictors of mortality for patients with COVID-19 pneumonia caused by SARS-CoV-2: A prospective cohort study. *European Respiratory Journal*, 55(5). Available at: 10.1183/13993003.00524-2020
- Hussain, A., Bhowmik, B. and Cristina, N., 2020. COVID-19 and diabetes: Knowledge in progress. *Diabetes Research and Clinical Practice*, (January). Available at: 10.1016/j.diabres.2020.108142
- Imam, Z., Odish, F., Gill, I., O'Connor, D., Armstrong, J., Vanood, A., Ibrionke, O., Hanna, A., Ranski, A. and Halalau, A., 2020. Older age and comorbidity are independent mortality predictors in a large cohort of 1305 COVID-19 patients in Michigan, United States. *Journal of Internal Medicine*, 288(4), pp. 469–476. Available at: 10.1111/joim.13119
- Kumar, Ashish. Arora, Anil. Sharma, Praveen. Anil, S., 2020. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information. *Diabetes & Metabolic Syndrome: Clinical Research & Reviews*, 14(January), pp. 535–545. Available at: <https://www.sciencedirect.com/science/article/abs/pii/S1871402120301090?via%3Dihub>
- Li, P., Chen, L., Liu, Z., Pan, J., Zhou, D., Wang, H., Gong, H., Fu, Z., Song, Q., Min, Q., Ruan, S., Xu, T., Cheng, F. and Li, X., 2020. Clinical features and short-term outcomes of elderly patients with COVID-19. *Int J Infect Dis*, 97:245-250. Available at: 10.1016/j.ijid.2020.05.107
- Lim, S., Bae, J.H., Kwon, H.S. and Nauck, M.A., 2021. COVID-19 and diabetes mellitus: from pathophysiology to clinical management. *Nature Reviews Endocrinology*, 17(1), pp. 11–30. Available at: 10.1038/s41574-020-00435-4
- Luh, N. P. P. and Wisnu, K. S. K., 2021. Peran HbA1c Dalam Prognosis dan Prognosis COVID-19 pada Pasien Diabetes Mellitus. 12(3), pp. 682–688. Available at: 10.15562/ism.v12i3.1113
- Miller, C., 2012. *Nursing for wellness in older adult: Theory and practice* (6th Ed.). Philadelphia: Lippincott Williams & Wilkins. Available at: <https://ocw.ui.ac.id>
- No, V. and Mona, N., 2020. Konsep Isolasi Dalam Jaringan Sosial Untuk Meminimalisasi Efek Contagious (Kasus Penyebaran Virus Corona Di Indonesia). *Jurnal Sosial Humaniora Terapan*, 2(2), pp. 117–125. Available at: 10.7454/jsht.v2i2.86
- Noor, F. M. and Islam, M. M., 2020. Prevalence and Associated Risk Factors of Mortality Among COVID-19 Patients: A Meta-Analysis. *Journal of Community Health*, 45(6), pp. 1270–1282. Available at: 10.1007/s10900-020-00920-x
- Raden Muhammad Ali, S., Resty Varia, T. and Djazuly, C., 2020. Analisis Faktor Risiko Kematian Dengan Penyakit Komorbid Covid-19. *Jurnal Keperawatan Silampari*, 4(1), pp. 1689–1699. Available at: 10.31539/jks.v4i1.1587
- Santos, M. M., Lucena, E.E.S., Lima, K.C., Brito, A.A.C., Bay, M.B. and Bonfada, D., 2020. Survival and predictors of deaths of patients hospitalized due to COVID-19 from a retrospective and multicenter cohort study in Brazil. *Epidemiology and Infection*. Available at: 10.1017/S0950268820002034
- Siagian, T. H., 2020. MENCARI KELOMPOK BERISIKO TINGGI TERINFEKSI VIRUS CORONA DENGAN DISCOURSE NETWORK ANALYSIS. *Jurnal Kebijakan Kesehatan Indonesia*, 09(02), pp. 98–106. Available at: 10.22146/jkki.55475
- Silver, S. R., Jia, L., Winifred, L. B., Taylor, L.S. and Matthew, R.G., 2020. Prevalence of Underlying Medical Conditions Among Selected Essential Critical Infrastructure Workers —



- Behavioral Risk Factor Surveillance System, 31 States, 2017–2018. *MMWR. Morbidity and Mortality Weekly Report*, 69(36), pp. 1244–1249. Available at: 10.15585/mmwr.mm6936a3
21. Simatupang, M. D. and Arcana, I. M., 2020. Risiko Kematian Pasien Covid-19 dan Faktor yang Memengaruhinya. 2020, pp. 889–898. Available at: 10.34123/semnasoffstat.v2021i1.1085
22. Suprayitno, E., Sylvina, R., Adivtiani, R. and Muhti, Y. P., 2020. Pengetahuan dan Sikap Masyarakat dalam Pencegahan COVID-19. *Journal Of Health Science (Jurnal Ilmu Kesehatan)*, 5(2), pp. 68–73. Available at: 10.24929/jik.v5i2.1123
23. Wenham, C., Smith, J. and Morgan, R., 2020. COVID-19: the gendered impacts of the outbreak. (January). Available at: 10.1089/AID.2019.0118.1
24. Widiastuti, L., 2021. Perbedaan Kualitas Hidup Pasien COVID-19 Dengan Comorbid. 5, pp. 233–239. Available at: 10.31539/jks.v5i1.2833
25. Zhang, J., Wang, X., Jia, X., Li, J., Hu, K., Chen, G., Wei, J., Gong, Z., Zhou, C., Yu, M., Lei, H., Cheng, F., Zhangm B., Xu, Y., Wang, G. and Dong, W., 2020. Risk factors for disease severity, unimprovement, and mortality in COVID-19 patients in Wuhan, China. *Clin Microbiol Infect*, 26(6): 767-772. Available at: 10.1016/j.cmi.2020.04.012
26. Zheng, Z., Anwar, S., Raymond, P., Makhyani, J.A., Ian, H. and Budhi, A., 2020. The COVID-19 resource centre is hosted on Elsevier Connect , the company ' s public news and information. *Journal of Infection*. Available at: <https://persi.or.id>