

TRAP-positive osteoclast precursors mediate ROS/NO-dependent bactericidal activity via TLR4

by Irma Josefina Savitri

Submission date: 11-Apr-2023 10:45AM (UTC+0800)

Submission ID: 2061150167

File name: cursors_mediate_ROS_NO_dependent_bacterial_activity_via_TLR4.pdf (3.64M)

Word count: 9152

Character count: 48794



ELSEVIER

Contents lists available at ScienceDirect

33

Free Radical Biology and Medicine

journal homepage: www.elsevier.com/locate/freeradbiomed

Original article

TRAP-positive osteoclast precursors mediate ROS/NO-dependent bactericidal activity via TLR4



Kazuaki Nishimura^{a,b}, Satoru Shindo^{a,c}, Alexandru Movila^a, Rayyan Kayal^d,
Albassam Abdullah^{a,d}, Irma Josefina Savitri^e, Atsushi Ikeda^a, Tsuguno Yamaguchi^{a,f},
Mohammed Howait^d, Ayman Al-dharrab^d, Abdulghani Mira^d, Xiaozhe Han^{a,g,*},
Toshihisa Kawai^{a,g,*}

^a The Forsyth Institute, Department of Immunology and Infectious Diseases, Cambridge, MA, USA

^b Tohoku University Hospital, Maxillo-oral Disorders, Sendai, Japan

^c The University of Tokushima Graduate School, Department of Conservative Dentistry, Institute of Health Biosciences, Tokushima, Japan

^d King Abdulaziz University, Faculty of Dentistry, Jeddah, Saudi Arabia

^e Airlangga University, Department of Periodontics, Jawa Timur, Indonesia

^f Research and Development Headquarters, LION Corporation, 100 Tajima Odawara, Kanagawa, Japan

^g Harvard School of Dental Medicine, Department of Oral Medicine, Infection and Immunity, Boston, MA, USA

ARTICLE INFO

Article history:

Received 3 May 2016

Received in revised form

17 June 2016

Accepted 20 June 2016

Available online 22 June 2016

Keywords:

Osteoclast

Bacteria

Phagocytosis

Toll-like receptor

Tartrate-resistant acid phosphatase

Reactive oxygen species

Nitric oxide

RANKL

ABSTRACT

Osteoclastogenesis was induced by RANKL stimulation in mouse monocytes to examine the possible bactericidal function of osteoclast precursors (Ocp) and mature osteoclasts (OCm) relative to their production of NO and ROS. Tartrate-resistant acid phosphatase (TRAP)-positive Ocp, but few or no OCm, phagocytized and killed *Escherichia coli* in association with the production of reactive oxygen species (ROS) and nitric oxide (NO). Phagocytosis of *E. coli* and production of ROS and NO were significantly lower in TRAP+ Ocp derived from Toll-like receptor (TLR)-4 KO mice than that derived from wild-type (WT) or TLR₂-KO mice. Interestingly, after phagocytosis, TRAP+ Ocp derived from wild-type and TLR₂-KO mice did not differentiate into OCm, even with continuous exposure to RANKL. In contrast, *E. coli*-phagocytized TRAP+ Ocp from TLR4-KO mice could differentiate into OCm. Importantly, neither NO nor ROS produced by TRAP+ Ocp appeared to be engaged in phagocytosis-induced suppression of osteoclastogenesis. These results suggested that TLR4 signaling not only induces ROS and NO production to kill phagocytized bacteria, but also interrupts OCm differentiation. Thus, it can be concluded that TRAP+ Ocp, but not OCm, can mediate bactericidal activity via phagocytosis accompanied by the production of ROS and NO via TLR4-associated reprogramming toward phagocytic cell type.

© 2016 Elsevier Inc. All rights reserved.

1. Introduction

Osteoclast precursors (Ocp) are originated from macrophage/monocyte-lineage hematopoietic precursor cells [1,2]. Differentiation from hematopoietic precursor cells to mature osteoclasts (OCm) capable of resorbing bone is elicited by stimulation with

16

receptor activator nuclear factor- κ B ligand (RANKL) and macrophage-colony stimulating factor (M-CSF) [3]. Especially, M-CSF plays a key role in the induction of RANK, a receptor for RANKL, on Ocp upstream of RANKL-mediated activation of osteoclastogenesis [4]. Indeed, the critical role of M-CSF in the course of osteoclastogenesis was demonstrated by M-CSF-gene knockout (KO) mice that are completely absent of OCm and show osteopetrotic phenotypes [5–8]. It is known that M-CSF-primed monocytes can also differentiate into macrophages in the absence of RANKL and perform antibacterial function [9]. In other words, M-CSF-primed monocytes can differentiate into either macrophages or osteoclasts. Nonetheless, it is largely unknown if monocytes stimulated with both M-CSF and RANKL can mediate antibacterial activity. Both Ocp and OCm have been found in infectious bone loss lesion of periodontitis [10–12] and septic prosthesis loosening [13], but it

* Corresponding author at: Forsyth Institute Department of Immunology and Infectious Diseases, 245 First Street, Cambridge, MA 02142, USA.

E-mail addresses: knishimura@dent.tohoku.ac.jp (K. Nishimura), c301351004@tokushima-u.ac.jp (S. Shindo), amovila@forsyth.org (A. Movila), rkayal@kau.edu.sa (R. Kayal), albassam.a@hotmail.com (A. Abdullah), irmaudijanto@gmail.com (I.J. Savitri), aikeda@forsyth.org (A. Ikeda), tsuguno@gmail.com (T. Yamaguchi), mhowait@icloud.com (M. Howait), aaldharab@kau.edu.sa (A. Al-dharrab), amira@kau.edu.sa (A. Mira), XHan@forsyth.org (X. Han), tkawai@forsyth.org (T. Kawai).

<http://dx.doi.org/10.1016/j.freeradbiomed.2016.06.021>
0891-5849/© 2016 Elsevier Inc. All rights reserved.

is also largely unknown if OCp and/or OCm can mediate anti-bacterial activity.

TLR signaling plays a crucial role in the induction of bacterial phagocytosis by macrophages [14]. Activation of TLRs induces macrophages to produce proinflammatory mediators, such as nitric oxide (NO), reactive oxygen species (ROS; e.g., peroxides, superoxide and hydroxyl radical), TNF α and IL-1 β [15,16]. Among them, NO and ROS accumulated in phagolysosomes are significantly engaged in bactericidal effects on bacteria phagocytosed by macrophages [17,18]. Interestingly, it is reported that ROS, including superoxide and hydrogen peroxide, are crucial components that upregulate osteoclast differentiation [19–21]. Although the effects of NO on OCp cells are controversial, NO is indeed produced by osteoclasts [22].

In summary, these reported findings have demonstrated 1) the effects of TLRs on NO/ROS production by macrophages that phagocytize bacteria and 2) the engagement of NO/ROS in RANKL-induced osteoclastogenesis. However, it has not been established that TLR signaling can trigger the expressions of NO/ROS and thus facilitate the phagocytosis and killing of bacteria by OCp and/or OCm. Based on these lines of evidence, we hypothesized that TRAP+ OCp, but not OCm, can mediate bactericidal activity via phagocytosis utilizing ROS and NO produced in response to bacteria-derived TLR activation. To test this hypothesis, we performed bacterial phagocytosis assays using OCp and OCm induced from RAW264.7 cells, and primary culture of OCp and OCm derived from bone marrow of TLR2-KO and TLR4-KO as well as their wild type (WT) mice, employing *Escherichia coli* as a model bacterium.

2. Material and methods

2.1. Osteoclastogenesis in vitro

Bone marrow-derived monocyte (BMM) cells were generated as described previously [23,24]. In brief, bone marrow cells isolated from TLR2-KO mice, TLR4-KO mice and their WT (C57BL/6) mice were cultured (1×10^5 cells/well; 96-well plate) in α -modified minimal essential medium (α -MEM) (Life Technologies, Beverly, MA) supplemented with 10% fetal bovine serum (FBS) (Atlanta Biologicals, Lawrenceville, GA). After initial incubation of bone marrow cells with 20 ng/ml M-CSF (BioLegend, San Diego, CA) for 2 days, adherent cells were used as bone marrow-derived monocyte (BMM). Those BMM cells were further stimulated with sRANKL (50 ng/ml) in the presence of M-CSF (20 ng/ml) for various times to induce OCp and OCm. The BMM cells that were incubated in the presence of M-CSF (20 ng/ml) without RANKL for the same periods were used as control macrophages. For the assay using RAW264.7 cells, single cell suspension (2×10^3 cells/well; 96-well plate) in α -MEM with 10% FBS was stimulated with sRANKL (50 ng/ml). Osteoclasts induced in the culture were stained for their expression of TRAP using a commercially available kit (Sigma-Aldrich, St. Louis, MO).

2.2. Bacterial strains and culture

E. coli strain (ATCC27325, Manassas, VA) was cultured overnight at 37 °C in Luria-Bertani (LB) broth medium under constant agitation at 240 rpm. The bacterial number was measured by spectrophotometer at OD 580 nm.

2.3. Assay to determine bacterial phagocytosis and killing by osteoclasts and macrophages

The method described previously [25–27] was performed with minor changes. In brief, the confluent cells in a 96-well plate with

or without RANKL were cultured in antibiotic-free medium overnight prior to the bacterial phagocytosis assay. Live bacteria harvested at mid-log growth curve were co-cultured with the cells for 1 h in antibiotic-free medium at an MOI of 10. After 1 h of co-culture, extracellular bacteria that did not enter cells were killed by treatment with gentamicin (500 μ g/ml; Life Technologies) applied over the course of 1 h. After additional incubation for 1 h, 2 h and 3 h, live bacteria remaining inside cells were released by osmotic shock-dependent disruption using distilled water (100 ml/well), followed by extensive pipetting (20 strokes/well). Bacterial suspension was spread to LB agar (Sigma-Aldrich) plates, and the number of colonies was counted. The Colony Forming Unit (CFU) corresponds to the number of colonies observed in an agar plate. The CFU was further normalized based on the OCps present in the well, and expressed as CFU/1000 OCps. The protocol used for this assay is illustrated in Fig. 1A.

In some experiments, after the antibiotics treatment, BMM cells or RAW264.7 cells were cultured in fresh medium in the presence or absence of NAC (1 mM), L-NAME (1 mM) or D-NAME (1 mM). Subsequently, the CFU of survived *E. coli* was also monitored, while the culture supernatant was subject to the measurements of NO $_2^-$ and TOS (see below).

2.4. Flow cytometry analysis of pHrodo™ uptake by RAW264.7 cells and BMM cells

Phagocytosis of bacteria by osteoclasts and macrophages was also evaluated using flow cytometry. This was performed using pHrodo™ BioParticles™ [28]. Phagocytized pHrodo-labeled *E. coli* fluoresced in an acidic pH-dependent manner. Briefly, pHrodo (fixed *E. coli*) was incubated with live RAW264.7 cells or BMM cells for 1 h at 37 °C. Cells incubated with pHrodo on ice were used as negative control. Cells were harvested by Falcon™ cell scraper (Thermo Fisher), washed three times, and then subjected to flow cytometry or confocal fluorescent microscopy. Data were analyzed using FlowJo software (Treestar, Ashland, OR).

2.5. Determination of nitrite (NO $_2^-$) concentrations






The levels of nitric oxide (NO) in the macrophage/osteoclast medium were measured by Griess assay [29,30]. Briefly, 50 μ L of culture supernatants were gently mixed with an equal volume of 2% sulfanilamide (Sigma) solution and incubated in the dark at room temperature for 10 min. After incubation, 50 μ L of naphthyl ethylenediamine dihydrochloride (Sigma) in 5% phosphoric acid solution were added to the reaction and incubated in the dark at room temperature for another 10 min. The absorbance at 540 nm was measured in a microplate reader. Nitrite concentration, an indicator of NO production, was calculated from a NaNO $_2$ standard curve.

2.6. Total oxidant status (TOS) assay


TOS in cells was measured following the method previously reported by Erel [31]. The oxidants in the sample oxidize the ferrous ion-o-dianisidine complex to ferric ion. The ferric ion produced a colored complex with xylenol orange in an acidic medium. The color intensity, as measured spectrophotometrically, was related to the total amount of oxidant molecules present in the sample. The assay was calibrated with hydrogen peroxide (H $_2$ O $_2$), and the results were expressed in terms of micromolar hydrogen peroxide equivalent per liter (μ mol H $_2$ O $_2$ equivalent/L).

2.7. Immunofluorescence microscopy


Macrophages and osteoclasts imaged with confocal microscopy

A Protocol to examine the bacteria phagocytized and killed by OC**B MF: 48h Medium alone****D MF: 96h Medium alone****C OC: 48h w RANKL****E OC: 96h w RANKL****F TRAP Staining**


a) MF: 48h Medium alone






c) MF: 96h Medium alone



b) OC: 48h w RANKL



d) OC: 96h w RANKL

**G****H**

were plated on eight-well chamber slides (Lab-Tek Chamber Slide, Thermo Fisher). Images were acquired on a confocal microscope (Zeiss LSM780, Carl Zeiss Microscope, Thornwood, NY). To observe phagocytized pHrodo™ *E. coli* BioParticles™, cells were fixed in 4% paraformaldehyde for 30 min and stained with Alexa Fluor® 488-conjugated phalloidin (Life Technologies) and counterstained with DAPI (Life Technologies).

2.8. Statistical analysis

All assays were conducted at least in triplicate, and each study was repeated 3 times. The collected data were analyzed using the Student's *t*-test for the comparison between two groups, or a one-way ANOVA with post hoc Tukey's test for the comparisons among different groups. *P* values of < 0.05 were considered statistically significant.

3. Results

3.1. RANKL-stimulated osteoclasts can phagocytize bacteria

In order to examine if RANKL-stimulated osteoclast precursors (OCp) and/or mature osteoclasts (OCm) could phagocytize bacteria, *E. coli* was incubated with RANKL-stimulated or non-stimulated RAW264.7 cells for various times (Fig. 1). More specifically, RAW264.7 cells were preincubated with or without RANKL for 2 days or 4 days prior to the addition of *E. coli*. The phagocytized bacterial number (CFU/1000 mononuclear cells) by the cytoplasm of RAW264.7 cells detected at baseline (0 h) was higher in nonstimulated RAW264.7 cells than that in the cytoplasm of RAW264.7 cells prestimulated with RANKL for 48 h (non-stimulated, 159 ± 65 CFU/1000 mononuclear cells; RANKL-prestimulated, 43 ± 7 CFU/1000 mononuclear cells; $p < 0.01$) (Fig. 1B and C). Even with RANKL stimulation for 96 h, the bacteria phagocytized by RAW264.7 cells at baseline (0 h) showed no significant difference from those phagocytized by RAW264.7 cells prestimulated with RANKL for 48 h (Fig. 1D and E). According to TRAP staining (Fig. 1F), all RAW264.7 cells were TRAP-positive after stimulation with RANKL for more than 48 h, while the number of multinuclear OCm significantly increased in those RAW264.7 cells prestimulated with RANKL for 96 h (TRAP+ multinuclear OCm cell number: 48 h, 15 ± 7 cells/well; 96 h, 256 ± 65 cells/well; data not shown in Fig. 1). Both nonstimulated and RANKL-prestimulated RAW264.7 cells appeared to kill phagocytized bacteria in a time-dependent fashion. Such observations were more evident when the % of killing effects was calculated (Fig. 1G and H). RAW264.7 cells were prestimulated with RANKL for 2 days and 4 days, and after 3 h of incubation with *E. coli* at each time point, $62.1 \pm 1.0\%$ and $50.8 \pm 4.9\%$ of bacteria were killed, respectively ($p < 0.01$), suggesting that osteoclast differentiation impairs bactericidal effects by RANKL-prestimulated RAW264.7 cells. No statistical difference was observed in bactericidal effects by unstimulated RAW264.7 cells cultured in medium alone for 2 and 4 days (Fig. 1G and H).

3.2. Mononuclear, but not multinuclear, osteoclasts can phagocytize

bacteria

Fixed *E. coli* labeled with a pH-sensitive fluorescent dye (pHrodo) was used to examine the phagocytosis of bacteria by RANKL-stimulated mononuclear and multinuclear cells. Upon phagocytization of pHrodo-labeled *E. coli* in the acidic environment of host cell cytoplasm, red fluorescence is emitted. After incubation of RAW264.7 cells with pHrodo-labeled *E. coli*, the number of bacteria positive for red pHrodo was counted under the fluorescent microscope (Fig. 2A, B and C). After 1 h incubation with bacteria, $91.7 \pm 3.2\%$ of unstimulated RAW264.7 cells showed bacterial entry into cytoplasm. In contrast, $28.8 \pm 5.2\%$ of RANKL-prestimulated RAW264.7 mononuclear cells showed cytoplasmic entry of pHrodo-labeled *E. coli*. On the other hand, only $2.3 \pm 0.4\%$ multinuclear cells showed phagocytosis of pHrodo-labeled *E. coli* ($p < 0.01$) (Fig. 2F). It is important to note that all RANKL-prestimulated RAW264.7 cells were TRAP-positive, whereas non-stimulated RAW264.7 cells showed no TRAP staining pattern in separately prepared culture (Fig. 2D and E). To summarize, nearly 80% of non-stimulated RAW264.7 cells showed cytoplasmic entry of pHrodo-labeled *E. coli*, whereas about 30% of RANKL-prestimulated mononuclear, but not multinuclear, RAW264.7 cells internalized pHrodo-labeled *E. coli*. Therefore, these results support the phagocytosis of *E. coli* by mononuclear OCp. According to flow cytometry, the mononuclear cell population in RANKL-prestimulated RAW264.7 cells showed a smaller percentage (49.9%) of pHrodo-positive cells compared to unstimulated RAW264.7 cells (75.6%), corresponding to the results obtained using fluorescent microscopy.

3.3. Osteoclasts produce ROS and NO in response to bacterial stimulation

Since the production of NO and ROS is significantly promoted in macrophage phagocytosis [17,18], we hypothesized that NO and ROS produced by RANKL-stimulated RAW264.7 cells could kill the phagocytized bacteria. To test this idea, RAW264.7 cells, both unstimulated and prestimulated with RANKL for 48 h and 96 h, incubated with *E. coli* induced a significantly elevated production of ROS (Fig. 3A) and NO (Fig. 3D). However, no temporal difference was noted in the production of either NO or ROS from RAW264.7 cells in response to *E. coli* stimulation after two or four days of preincubation with or without RANKL, suggesting that similar levels of NO and ROS are produced by OCp at early stage (48 h) and mid-stage (96 h) of differentiation. The addition of N-acetyl-L-cysteine (NAC), an ROS inhibitor, or NG-nitro-L-arginine methyl ester (L-NAME), an NO inhibitor, but not the addition of control NG-nitro-D-arginine methyl ester (D-NAME), significantly suppressed the *E. coli*-induced production of ROS or NO in RAW264.7 cells prestimulated with or without RANKL for 48 h (Fig. 3B and C, ROS production; E and F, NO production) and 96 h (not shown). The possible bactericidal roles of ROS and NO on *E. coli* phagocytized by RANKL-stimulated OC were further evaluated in the following experiments using primary culture of OCp isolated from mouse bone marrow (Fig. 6).

Fig. 1. Bacteria killing effects by RAW264.7 cells stimulated with/without RANKL. (A) After preincubation of RAW264.7 cells with or without RANKL (50 ng/ml) for 48 or 96 h, *E. coli* entered into the cytoplasm of RAW264.7 cells were evaluated following the protocol described in Materials and Methods. The time line for each specific procedure performed in this experiment is illustrated in [A]. (B–E) The numbers of bacteria which phagocytized by RAW264.7 cells and survived inside of RAW264.7 cells were expressed as a colony forming unit (CFU)/1000 mononuclear cells. RAW264.7 cells were preincubated in medium alone for 48-h [B] and 96-h [D]. RAW264.7 cells were pre-incubated for 48-h [C] and 96-h [E] with RANKL. (F) TRAP-staining of RAW264.7 cells incubated with or without RANKL for 48-h (a: medium alone, b: with RANKL) and for 96-h (c: medium alone, d: with RANKL) are shown. (G and H) The rates of killed bacteria (%) after each incubation time (1, 2 and 3 h) were calculated based on the comparison with the phagocytized live bacteria counted at base line (0 h). The results of RAW264.7 cells preincubated with or without RANKL for 48-h [G] and for 96-h [H] are shown. Abbreviation in Figure: MF; macrophage (no-stimulated RAW264.7), OC; osteoclasts (RAW264.7 cells stimulated with RANKL). Each column and bar represents mean \pm SD. * $p < 0.05$, ** $p < 0.01$, and *** $p < 0.001$, n.s. = not significant.




Fig. 2. The uptake of pHrodo™ Red *E. coli* BioParticles[®] by RAW 264.7 cells. Fluorescent images of phagocytized *E. coli* BioParticles[®] in the cytoplasm of RAW264.7 cells that were prestimulated with or without RANKL are shown. (A–C) RAW264.7 cells were preincubated without RANKL [A] or with RANKL [B and C] for 96-h and co-cultured with pHrodo Red *E. coli* BioParticles[®] for 1 h, then, counter-stained with AlexaFluor 488 conjugated phalloidin (green) and DAPI (blue). Only the phagocytized pHrodo *E. coli* by RAW264.7 cells show the red fluorescent. In the large osteoclast, little or no red fluorescent was detected [B]. In contrast, small osteoclasts showed the red-fluorescent + pHrodo *E. coli* in the cytoplasm [C]. Images were captured using a fluorescent microscope at 600 × magnification. (D and E): TRAP staining patterns of RAW264.7 cells that were preincubated with or without RANKL for 96 h is shown [D: no RANKL, E: with RANKL]. (F): The % of pHrodo positive RAW264.7 cells was counted from the captured A–C images. ** $p < 0.01$ [G and H] In order to quantify the phagocytized pHrodo™ *E. coli* by RAW cells, after 1 h of co-culture with pHrodo *E. coli*, respective RAW264.7 cells, either prestimulated without (G) or with RANKL (H) for 96 h, and then were removed from tissue culture flask using a cell scraper. The resulting single cell suspension of respective RAW cells were subject to FACS analysis. The cells received pHrodo *E. coli* are indicated in blue and negative controls are in red samples, respectively.

3.4. TLR4 is engaged in the induction of bacterial phagocytosis

To investigate whether TLRs are involved in phagocytosis of RANKL-stimulated monocytes (OCp and/or OCm), bone marrow-derived monocytes (BMM) isolated from WT, TLR2-KO and TLR4-KO mice were used (Fig. 4A–E). BMM isolated from different strains of mice were stimulated with M-CSF and RANKL, or M-CSF alone, for 48 h or 96 h, pHrodo-labeled *E. coli* was applied to the respective cell culture. The incidence of pHrodo-positive red fluorescence, indicating phagocytized *E. coli* in unstimulated TLR4-KO BMM cells ($75.7 \pm 4.5\%$), was significantly lower than that in WT ($89.6 \pm 2.9\%$) ($p < 0.05$) or TLR2-KO mice ($90.0 \pm 5.6\%$) ($p < 0.05$) (Fig. 4B and D). The incidence of pHrodo-positive RANKL-stimulated TLR4-KO BMM cells (mononuclear OCp, 30.0%

$\pm 3.0\%$; multinuclear OCm, $1.3 \pm 2.4\%$) was lower than that in WT (mononuclear OCp, $62.8 \pm 4.7\%$, $p < 0.01$; multinuclear OCm, $9.4 \pm 2.2\%$, $p < 0.05$) or TLR2-KO mice (mononuclear OCp, $63.1 \pm 2.8\%$, $p < 0.01$; multinuclear OCm, $9.4 \pm 1.2\%$, $p < 0.05$) (Fig. 4B and E). On the other hand, no significant difference was observed in the incidence of pHrodo-positive cells between WT and TLR2-KO BMM cells (Fig. 4B, C and D). According to flow cytometry analysis (Fig. 4E), the mononuclear cell population (OCp) of RANKL-prestimulated TLR4-KO BMM showed a significantly lower level of phagocytization of pHrodo-positive *E. coli* than either WT or TLR2-KO mice, corresponding to the results from fluorescent microscopy.

In order to examine the possible engagement of TLR2 and/or TLR4 in the production of ROS and NO from primary culture of OCp in response to the exposure with *E. coli*, the production of *E. coli*-induced ROS (Fig. 5A and C) and NO (Fig. 5B and D) from M-CSF + RANKL-prestimulated OCp was examined using BMM cells derived from WT, TLR2-KO and TLR4-KO mice. Similar to the results from RAW264.7 cells (Fig. 3), the OCp induced from BMM by prestimulation with M-CSF + RANKL for 48 h showed significantly elevated production of ROS (Fig. 5A) and NO (Fig. 5B) in response to *E. coli* exposure. In terms of OCp cells derived from BMM cells




Fig. 3. Production of ROS and NO from *E. coli*-exposed osteoclasts differentiated from RAW264.7 cells. RAW264.7 cells that were prestimulated with or without RANKL for 48 h were exposed to live *E. coli* for one hour. After elimination of the extracellular bacteria with antibiotics, RAW264.7 cells were cultured in fresh medium in the presence or absence of NAC (1 mM), L-NAME (1 mM) or D-NAME (1 mM). After the additional 3 h incubation, the levels of total oxidant status (TOS) and nitric acid (NO) in the culture supernatants were measured. (A): Levels of TOS produced from RAW cells in response to *E. coli* that were prestimulated with/without RANKL for 2 and 4 days (B and C): Effects of NAC and L-NAME on *E. coli*-induced TOS production from RAW264.7 cells that were prestimulated without RANKL [B] or with RANKL [C]. (D): Levels of NO produced from RAW cells in response to *E. coli* that were prestimulated with/without RANKL for 2 and 4 days (E and F): Effects of NAC and L-NAME on *E. coli*-induced NO production from RAW264.7 cells that were prestimulated without RANKL [E] or with RANKL [F]. ** $p < 0.01$.

prestimulated with M-CSF+RANKL for 48 h (Fig. 5C), ROS production from TLR4-KO OCp cells (13.2 ± 2.1 μmol H₂O₂ equivalent/L), but not TLR2-KO OCp cells (15.1 ± 0.9 μmol H₂O₂ equivalent/L), was significantly lower level than that of WT OCp cells (16.8 ± 1.16 μmol H₂O₂ equivalent/L) ($p < 0.05$) in the group prestimulated with M-CSF and RANKL for 48 h (Fig. 5C). OCp cells derived from BMM cells prestimulated with M-CSF+RANKL for 96 h showed the same trend of significantly diminished *E. coli*-induced ROS production by TLR4-KO BMM cells, compared to that (data not shown). The amount of *E. coli*-induced NO production from TLR2-KO OCp cells (1.7 ± 0.1 [NO₂⁻] μmol/l) and TLR4-KO OCp cells (1.4 ± 0.1 [NO₂⁻] μmol/l) was significantly lower than that detected from WT OCp cells (2.4 ± 0.3 [NO₂⁻] μmol/l)

($p < 0.01$) in the group of OCp derived from BMM prestimulated with M-CSF and RANKL for 48 h (Fig. 5D). The group prestimulated with M-CSF+RANKL for 96 h demonstrated results of *E. coli*-induced NO production similar to those found in the 48 h M-CSF+RANKL prestimulation group (data not shown). The production of ROS and NO from *E. coli*-exposed OCp derived from WT, TLR2-KO and TLR4-KO mice was suppressed in all cases by the addition of NAC, as well as L-NAME to the base line level (Fig. 5C and D). We did not find any difference in the susceptibility to NAC and L-NAME-mediated ROS and NO suppression among the three strains of mice for their production.

The bactericidal effects on phagocytized bacteria were also observed in the primary culture of BMM-derived OCp (Fig. 6).




Fig. 4. The uptake of pHrodo™ Red *E. coli* BioParticles® by osteoclasts differentiated from BMM cells of WT, TLR2-KO and TLR4-KO mice. (A) TRAP staining of WT, TLR2-KO and TLR4-KO bone marrow monocytes (BMM) incubated with M-CSF alone or M-CSF/RANKL for 96 h is shown. (B) BMM cells derived from WT, TLR2-KO and TLR4-KO mice were preincubated without RANKL [A] or with RANKL [B] for 96-h and, then, exposed to pHrodo™ Red *E. coli* for one hour. Subsequently, the cells were counter-stained with AlexaFluor 488 conjugated phalloidin (green) and DAPI (blue). Images of magnification $\times 200$ are shown. (C) The incidences (%) of pHrodo positive cells were counted from the captured images of BMM cells that were not prestimulated with RANKL. (D) The incidences (%) of pHrodo positive cells were counted from the captured images of BMM cells that were prestimulated with M-CSF/RANKL are shown: (a) pHrodo positive (%) in OCp; (b) pHrodo positive (%) in OCm. (E) The incidences (%) of pHrodo positive cells in mononuclear OCp population were analyzed using a flow cytometer. * $p < 0.05$ ** $p < 0.01$.




Fig. 5. Production of ROS and NO from *E. coli*-exposed osteoclasts differentiated from BMM cells of WT, TLR2-KO and TLR4-KO mice. (A and B) Levels of TOS (A) and NO (B) produced from *E. coli*-stimulated BMMs that were preincubated with/without RANKL for 2 and 4 days. (C and D) Effects of NAC and L-NAME on *E. coli*-induced TOS (C) and NO (D) production from BMM that were prestimulated with RANKL in the presence of M-CSF for 48 h. Bone marrow monocytes (BMM) derived from WT, TLR2-KO and TLR4-KO mice were preincubated with or without RANKL in the presence of M-CSF for 48-h and, then, exposed to live *E. coli* for one hour. After killing the extracellular bacteria with antibiotics, respective BMM cells were cultured in fresh medium in the presence or absence of NAC (1 mM), L-NAME (1 mM) or D-NAME (1 mM). After the additional 3 h incubation, the levels of total oxidant status (TOS) and nitric acid (NO) in the culture supernatants were measured. Each column and bar represents mean \pm SD from three wells. ** $p < 0.01$, n.s. –not significant.

Bacteria killing rate of control BMM preincubated with M-CSF alone showed no difference among WT, TLR2-KO and TLR4-KO mice (Fig. 6A). However, the bacteria killing rate by TLR4-KO OCp

was significantly lower that of WT or TLR2-KO mice (Fig. 6B). Furthermore, NAC and L-NAME downregulated bactericidal effects on phagocytized bacteria in OCp (Fig. 6B) as well as control M-CSF-




Fig. 6. Engagement of ROS or NO in the killing of phagocytized bacteria by osteoclast precursors. The killing rates of phagocytized bacteria by control macrophages (A) and OCp (B) that were differentiated from BMM of WT, TLR2-KO and TLR4-KO mice via preincubation with M-CSF alone (A) and M-CSF/RANKL (B) for 48 h are shown. As described in Fig. 5 legend, after elimination of the extracellular bacteria with antibiotics, respective BMM cells were cultured in fresh medium in the presence or absence of NAC (1 mM), L-NAME (1 mM) or D-NAME (1 mM) for 3 h, and CFU of survived *E. coli* in the BMM were counted. The killing rates (%) of phagocytized bacteria after 3 h incubation was calculated in comparison to the phagocytized live bacteria counted at base line (0 h). Each column and bar represents mean \pm SD of killing rates (%) from three wells. ## $p < 0.01$, significantly different between columns indicated by a bracket for the respective strain of mice.

prestimulated BMM (macrophages, Fig. 6A), suggesting that OCp killed the phagocytized bacteria in an ROS/NO-dependent manner, similar to macrophages. Interestingly, the bacteria killing effects by OCp derived from TLR4-KO mice were suppressed significantly more by NAC and L-NAME than that detected in WT or TLR2-KO. These results indicated that ROS and NO produced by OCp via TLR4-activation are in part associated with the bacteria killing effects on the phagocytized bacteria.

3.5. Bacterial phagocytosis inhibits OCp from RANKL-dependent osteoclast maturation in a manner independent of ROS and NO productions

Finally, since ROS and NO are known to promote the RANKL-mediated osteoclastogenesis [19,20,32], we asked if phagocytizing OCp which increase production of ROS and NO could affect their RANKL-mediated differentiation into OCm. To examine this, osteoclast precursors developed in vitro from BMM of WT, TLR2-KO and TLR4-KO mice were exposed to *E. coli* and, subsequently, maintained in culture under continuous RANKL stimulation. More specifically, after prestimulation of BMM cells with M-CSF and

RANKL for 48 h, these OCp cells were incubated with or without fixed *E. coli* for 3 h, followed by continuous incubation with RANKL for an additional 5 days. In the absence of bacterial exposure, all OCp isolated from WT, TLR2-KO and TLR4-KO mice differentiated into TRAP+ multinucleated OCm at comparable levels (Fig. 7H). On the other hand, contrary to our expectation, exposure to *E. coli* nearly completely inhibited the differentiation of OCp derived from WT and TLR2-KO mice (Fig. 7B and H). However, the OCp derived from TLR4-KO mice did differentiate into TRAP+ multinucleated OCm, albeit osteoclast size was smaller (Fig. 7E and H) than that found in osteoclasts differentiated in the absence of *E. coli* (Fig. 7C, D and H). Incubation of OCp cells with LPS (TLR4 ligand) and lipoteichoic acid (LTA; TLR2 ligand) suppressed RANKL-induced osteoclastogenesis (Fig. 7I). Importantly, at least 100-fold higher amount of LTA than LPS was required to completely suppress osteoclast differentiation (Fig. 7F, G and I), indicating that TLR4 activation has a greater magnitude of impact on osteoclastogenesis than that by TLR2 activation. These results implicated that bacterial exposure to OCp inhibits their differentiation into OCm, in part from the signal elicited by TLR4 activation. Finally, we examined the core question as to whether ROS or NO is engaged in the altered osteoclastogenesis in bacteria phagocytizing OCp (Fig. 7J). When either NAC or L-NAME was added to the culture of *E. coli*-phagocytizing WT OCp which were continuously incubated with RANKL, neither one abrogated the suppression of osteoclastogenesis resulting from exposure to *E. coli* (Fig. 7J), suggesting that neither ROS nor NO is engaged in *E. coli*-mediated suppression of osteoclastogenesis.

4. Discussion

The present study demonstrated that mononuclear osteoclast precursor cells (OCp), but not mature multinucleated osteoclasts (OCm), can phagocytize *E. coli* and kill phagocytized *E. coli* in association with the production of reactive oxygen species (ROS) and nitric oxide (NO). Both phagocytosis of *E. coli* and ROS and NO production were significantly lower in OCp derived from TLR4-KO mice than compared to wild-type (WT) and TLR2-KO mice. Furthermore, the OCp of wild-type and TLR2-KO mice that did phagocytize *E. coli* did not differentiate into OCm, even with continuous exposure to RANKL, whereas *E. coli*-phagocytized OCp of TLR4-KO mice could differentiate into OCm. The addition of NAC (ROS inhibitor) and L-NAME (NO inhibitor) did not abrogate the diminished differentiation of *E. coli*-phagocytized OCp to OCm by continuous exposure to RANKL, suggesting that neither ROS nor NO is associated with such bacterial phagocytosis-related interruption of OCp differentiation. These results clearly demonstrated that TLR4 signaling not only induces ROS and NO production to kill phagocytized bacteria, but also downregulates differentiation toward OCm. Osteoclasts have always been considered bone-resorbing cells; therefore, the findings in this study may gain new insight into the etiology of osteoimmunological disorders where monocytes differentiate into osteoclast precursors.

Along with neutrophils, macrophages play a key role in innate immune responses toward bacterial infection based on their function to phagocytize and kill bacteria. Antibacterial activities by macrophages are, in part, mediated by their production of free radicals, including nitric oxide (NO) and reactive oxygen species (ROS) [18]. Bacteria phagocytized by macrophages are destroyed in the phagolysosome via oxygen-dependent and oxygen-independent mechanisms. As an oxygen-dependent bacteria-killing mechanism, free radicals, such as ROS and NO, are generated, while digestive enzymes, for example, lysozyme and cathepsin, are also utilized in the bacteria-killing mechanism [33,34]. Furthermore, lines of evince support ROS and NO as cell signaling




Fig. 7. Effects of bacteria phagocytosis by OCp on their RANKL-mediated osteoclastogenesis. Osteoclast precursors (OCp) developed in vitro from BMM of WT, TLR2-KO and TLR4-KO mice via prestimulation with M-CSF/RANKL for 2 days were exposed to fixed *E. coli* for 3 h. Subsequently, bacteria-phagocytized OCp were maintained in the culture under continuous M-CSF/RANKL stimulation for additional 5 days. (A–G) Images of TRAP-stained BMM cells after a total of 7 day incubation with M-CSF/RANKL (preincubation for 2 days + post-phagocytosis incubation for 5 days) are shown. (H–J) The number of TRAP+ multinuclear cells differentiated in the culture was measured. [H] RANKL-induced osteoclastogenesis was compared among three strains of mice, i.e., TLR2-KO, TLR4-KO and WT. [I] Effects of LPS and LTA on the RANKL-induced osteoclastogenesis from BMM of WT mice were shown. After prestimulation with M-CSF/RANKL for 2 days, WT BMM cells were incubated with or without LPS or LTA for additional 5 days in the presence of M-CSF/RANKL. [J] Effects of NAC, L-NAME (LN) and D-NAME (DN) on the RANKL-induced osteoclastogenesis from BMM of WT mice were shown. Each column and bar represents mean \pm SD of TRAP+ multinuclear cells from three wells. § $p < 0.05$, §§ $p < 0.01$, significantly different compared to the control M-CSF-stimulation alone (column with #). * $p < 0.05$, ** $p < 0.01$, significantly different between the columns indicated by brackets.

molecules in innate immune cells [35,36]. Interestingly, ROS produced by osteoclasts play a role in promoting the differentiation of these cells through Akt, NF- κ B and ERK activation [37,38], suggesting that ROS function as a cell signaling mediator in osteoclastogenesis. Recent study reported that ROS-induced promotion of osteoclastogenesis is downmodulated by an antioxidant response system involving Nrf2 in the homeostatic context [39,40], indicating the presence of a sophisticated regulatory mechanism for ROS-associated osteoclastogenesis. On the other hand, low level of NO increases the generation of mature osteoclasts by up-regulating actin remodeling in mononuclear preosteoclasts, thereby promoting cell fusion processes required for multinucleation of Ocm [32]. In contrast, high concentration of NO inhibits osteoclastogenesis and activity of Ocm, while same high level of NO induces apoptosis in OCp [41]. In sum, the effects NO on osteoclastogenesis are dependent on the concentration of NO released in the microenvironment surrounding OCp or Ocm. Surprisingly, antibacterial roles of NO and ROS produced by osteoclasts have not been addressed until the present study. Our results demonstrated that ROS and NO produced by OCp are engaged in

bactericidal effects. The chemical inhibitors that interrupt the generation of ROS and NO, i.e., NAC and L-NAME, respectively, suppressed their bactericidal effects on phagocytized bacteria in OCp (Fig. 6). The concentration of NAC and L-NAME that showed nearly complete suppression of ROS and NO production by osteoclasts (Fig. 3 and 5) could downregulate the bactericidal effects at the rate of 35–45% (Fig. 6), suggesting that other factors, such as lysozyme or cathepsin, are involved in bactericidal effects by OCp in addition to ROS and NO.

The activation of macrophages through Toll-like receptor (TLR) signaling pathways consists of a major component in the innate immune responses to bacterial infection [42]. Activation of Toll-like receptors, such as TLR1, TLR2 and TLR4, results in the generation of ROS and NO from macrophages by means of nicotinamide adenine dinucleotide phosphate (NADPH) oxidases, myeloperoxidase, mitochondria, or NO-producing enzymes [43]. However, as noted above, while ROS and NO produced by osteoclasts are known to be engaged in osteoclast differentiation [32,37,38], the role of TLRs in the induction of ROS and NO has been elusive. On the other hand, the impact of TLR activation on

osteoclastogenesis appears to be more complex than that on macrophages. That is, TLR signals inhibit RANKL-induced osteoclast differentiation in the early precursor stage, whereas differentiation of mid-late stage of RANKL-primed osteoclasts is promoted by TLR signaling [44,45]. Our results demonstrated that RANKL-stimulated mononuclear TRAP+ OCp can kill bacteria in a manner similar to that of macrophages, in which activation of TLR4, but little or no activation of TLR2, is involved in the induction of phagocytosis (Fig. 4) and ROS/NO production (Fig. 6). However, this same TLR4 signaling appeared to be engaged in the suppression of osteoclastogenesis in TRAP+ OCp cells challenged with *E. coli* in a manner independent of NO/ROS produced by stimulation with *E. coli* (Fig. 7). More specifically, *E. coli*-mediated suppression of osteoclastogenesis was partially abrogated in the TRAP+ OCp derived from TLR4-KO mice in response to continuous exposure to RANKL, whereas *E. coli* completely suppressed osteoclastogenesis in TRAP+ OCp derived from TLR2-KO or WT mice that were also continuously exposed to RANKL. That TLR4 is partially involved in *E. coli*-induced suppression of osteoclastogenesis seems incongruent with its promotion of RANKL-induced osteoclastogenesis by induction of ROS and NO production. This discrepancy indicates the presence of some unknown signaling pathway elicited by TLR4 which does not involve ROS/NO downregulation of osteoclastogenesis.

One of the key findings in this study is that phagocytosis of *E. coli* was observed in OCps, but not in OCms. According to our unpublished result, similar to *E. coli*, phagocytosis of Gram (+) bacterium, *Enterococcus faecalis* (*Ef*), was also found in OCps, but not in OCms (data not shown). However, as fluorescent micro-particle (fluorescent latex beads $\phi=0.75\ \mu\text{m}$) was applied to the osteoclast phagocytosis assay, both OCps and OCms phagocytized fluorescent micro-particle (data not shown). In correspond to the latter finding, latex and PMMA particles were reported to be phagocytized by OCms [46]. Meng et al. also reported that OCms can phagocytize titanium particle ($\phi < 1\ \mu\text{m}$) and that the expressions of osteoclast-maturation genes, including, TRAP and cathepsin K, were inhibited by a higher concentration of titanium particles, but enhanced at a lower concentration of titanium particles [47]. Since the sizes of fluorescent micro-particle and titanium are similar to that of *E. faecalis* ($\phi=0.6\text{--}2.0\ \mu\text{m}$, cocci) and *E. coli* ($0.5 \times 2\ \mu\text{m}$, rod), it appeared that molecules distinctly expressed on bacteria regulate selective phagocytosis by OCps. As noted above, at least, TLRs are engaged in the induction of phagocytosis of bacteria by OCps.

The addition of LTA, a TLR2 ligand, to TRAP+ OCp suppressed osteoclastogenesis to a lesser degree compared to LPS, a TLR4 ligand (Fig. 7), suggesting that TLR ligand(s) other than those of TLR4 are present on bacteria phagocytized by OCp and may also contribute to the suppression of osteoclastogenesis. Interestingly, it was demonstrated that *Ef* suppress the differentiation of macrophages into osteoclasts [48], and that activation of TLR2 by LTA produced by *Ef* is in part engaged in the attenuation of osteoclastogenesis [49]. Indeed, various TLR ligands inhibit RANKL-mediated osteoclastogenesis from OCp [50–52]. One of the theories underlying TLR-elicited inhibition of osteoclastogenesis is the autocrine production of IFN- β [53,54] which, in turn, suppresses the expression of c-Fos protein, a pivotal transcription factor for the formation of osteoclasts [55]. The latter studies were carried out using mouse OCp. However, in terms of human OCp, it is reported that TLR ligands suppress osteoclastogenesis by inhibiting expression of receptor activator of NF- κB (RANK), thereby making precursor cells refractory to the effects of RANKL [56]. The detailed molecular mechanism underlying the suppression of RANKL-induced osteoclastogenesis caused by phagocytized bacteria in OCp will be addressed in future studies.

Our current working hypothesis supports that phagocytic property by OCps may play a role in prevention of the bacterial

dissemination into the bone in the context of infectious bone lytic disease, such as periodontitis. Especially it has become evident that OCps in the circulation migrate to intact bone as well as inflammatory bone resorption lesion and participate in bone resorption [57,58], suggesting that OCps function similar to innate immune macrophages [59]. For this reason, it is plausible that any pharmaceutical intervention for bone lytic diseases that targets osteoclast differentiation, such as bisphosphonate and Denosumab [60,61], would attenuate the OCps-mediated anti-bacterial barrier mechanism for alveolar bone. As a consequence of attenuated anti-bacterial function by OCps which permits the invasion of bacteria into bone, osteonecrosis of jaw may be induced [62].

In conclusion, the present study showed that RANKL-stimulated, TRAP-positive OCp can phagocytize and then kill bacteria through TLR4 signaling and induction of ROS/NO, respectively.

Acknowledgements

This work was partially supported by NIH grants RO1 DE-018499, RO1 DE-019917, R56 DE023807 and RO1DE025255 and T32 DE 7327-12 from National Institute of Dental and Craniofacial Research (NIDCR) and a research fund from King Abdulaziz University (KAU).

Appendix A. Supplementary material

Supplementary data associated with this article can be found in the online version at <http://dx.doi.org/10.1016/j.freeradbiomed.2016.06.021>.

References

- [1] S.L. Teitelbaum, F.P. Ross, Genetic regulation of osteoclast development and function, *Nat. Rev. Genet.* 4 (8) (2003) 638–649.
- [2] L. Xing, E.M. Schwarz, B.F. Boyce, Osteoclast precursors, RANKL/RANK, and immunology, *Immunol. Rev.* 208 (2005) 19–29.
- [3] M. Asagiri, H. Takayanagi, The molecular understanding of osteoclast differentiation, *Bone* 40 (2) (2007) 251–264.
- [4] T. Suda, N. Takahashi, N. Udagawa, E. Jimi, M.T. Gillespie, T.J. Martin, Modulation of osteoclast differentiation and function by the new members of the tumor necrosis factor receptor and ligand families, *Endocr. Rev.* 20 (3) (1999) 345–357.
- [5] G. Hattersley, J. Owens, A.M. Flanagan, T.J. Chambers, Macrophage colony stimulating factor (M-CSF) is essential for osteoclast formation in vitro, *Biochem. Biophys. Res. Commun.* 177 (1) (1991) 526–531.
- [6] S. Tanaka, N. Takahashi, N. Udagawa, T. Tamura, T. Akatsu, E.R. Stanley, T. Kurokawa, T. Suda, Macrophage colony-stimulating factor is indispensable for both proliferation and differentiation of osteoclast progenitors, *J. Clin. Invest.* 91 (1) (1993) 257–263.
- [7] H. Yoshida, S. Hayashi, T. Kunisada, M. Ogawa, S. Nishikawa, H. Okamura, T. Sudo, L.D. Shultz, S. Nishikawa, The murine mutation osteopetrosis is in the coding region of the macrophage colony stimulating factor gene, *Nature* 345 (6274) (1990) 442–444.
- [8] W. Wiktor-Jedrzejczak, A. Bartocci, A.W. Ferrante Jr., A. Ahmed-Ansari, K. W. Sell, J.W. Pollard, E.R. Stanley, Total absence of colony-stimulating factor 1 in the macrophage-deficient osteopetrotic (op/op) mouse, *Proc. Natl. Acad. Sci. USA* 87 (12) (1990) 4828–4832.
- [9] E.R. Stanley, K.L. Berg, D.B. Einstein, P.S. Lee, Y.G. Yeung, The biology and action of colony stimulating factor-1, *Stem Cells* 12 (Suppl. 1) (1994) S15–S24 (discussion 25).
- [10] R.E. Carson, F.S. Sayegh, P.F. Fedji Jr., Osteoclastic resorption of alveolar bone affected by periodontitis—correlation of light microscopic and scanning electron microscopic observations, *J. Periodontol.* 49 (8) (1978) 406–414.
- [11] H. Okada, T. Kida, H. Yamagami, Identification and distribution of immunocompetent cells in inflamed gingiva of human chronic periodontitis, *Infect. Immun.* 41 (1) (1983) 365–374.
- [12] S. Thorbert-Mros, L. Larsson, T. Berglundh, Cellular composition of long-standing gingivitis and periodontitis lesions, *J. Periodontol. Res.* 50 (4) (2015) 535–543.
- [13] T. Gehrke, C. Sers, L. Morawietz, G. Fernald, J. Neidel, L. Frommelt, V. Krenn, Receptor activator of nuclear factor kappaB ligand is expressed in resident and inflammatory cells in aseptic and septic prosthesis loosening, *Scand. J.*

- Rheumatol. 32 (5) (2003) 287–294.
- [14] R. Medzhitov, Toll-like receptors and innate immunity, *Nat. Rev. Immunol.* 1 (2) (2001) 135–145.
- [15] C.A. Janeway Jr., R. Medzhitov, Innate immune recognition, *Annu. Rev. Immunol.* 20 (2002) 197–216.
- [16] A.P. West, I.E. Brodsky, C. Rahner, D.K. Woo, H. Erdjument-Bromage, P. Tempst, M.C. Walsh, Y. Choi, G.S. Shadel, S. Ghosh, TLR signalling augments macrophage bactericidal activity through mitochondrial ROS, *Nature* 472 (7344) (2011) 476–480.
- [17] C. Prolo, M.N. Alvarez, R. Radi, Peroxynitrite, a potent macrophage-derived oxidizing cytotoxin to combat invading pathogens, *BioFactors* 40 (2) (2014) 215–225.
- [18] G. Weiss, U.E. Schaible, Macrophage defense mechanisms against intracellular bacteria, *Immunol. Rev.* 264 (1) (2015) 182–203.
- [19] I.R. Garrett, B.F. Boyce, R.O. Oreffo, L. Bonewald, J. Poser, G.R. Mundy, Oxygen-derived free radicals stimulate osteoclastic bone resorption in rodent bone in vitro and in vivo, *J. Clin. Invest.* 85 (3) (1990) 632–639.
- [20] L.L. Key Jr., W.L. Ries, R.G. Taylor, B.D. Hays, B.L. Pitzer, Oxygen derived free radicals in osteoclasts: the specificity and location of the nitroblue tetrazolium reaction, *Bone* 11 (2) (1990) 115–119.
- [21] H. Kim, L.Y. Kim, S.Y. Lee, D. Jeong, Bimodal actions of reactive oxygen species in the differentiation and bone-resorbing functions of osteoclasts, *FEBS Lett.* 580 (24) (2006) 5661–5665.
- [22] S.J. Wimalawansa, Rationale for using nitric oxide donor therapy for prevention of bone loss and treatment of osteoporosis in humans, *Ann. N. Y. Acad. Sci.* 1117 (2007) 283–297.
- [23] N.K. Lee, Y.G. Choi, J.Y. Baik, S.Y. Han, D.W. Jeong, Y.S. Bae, N. Kim, S.Y. Lee, A crucial role for reactive oxygen species in RANKL-induced osteoclast differentiation, *Blood* 106 (3) (2005) 852–859.
- [24] G.C. Franco, M. Kajiya, T. Nakanishi, K. Ohta, P.L. Rosalen, F.C. Groppo, C. W. Ernst, J.L. Boyesen, J.D. Bartlett, P. Stashenko, M.A. Taubman, T. Kawai, Inhibition of matrix metalloproteinase-9 activity by doxycycline ameliorates RANK ligand-induced osteoclast differentiation in vitro and in vivo, *Exp. Cell Res.* 317 (10) (2011) 1454–1464.
- [25] M. Kajiya, H. Komatsuzawa, A. Papanonakis, M. Seki, S. Makihira, K. Ouhara, Y. Kusumoto, S. Murakami, M.A. Taubman, T. Kawai, Aggregatibacter actinomycetemcomitans Omp29 is associated with bacterial entry to gingival epithelial cells by F-actin rearrangement, *PLoS One* 6 (4) (2011) e18287.
- [26] D.W. Lawrence, J. Kornbluth, E3 ubiquitin ligase NKLAM is a macrophage phagosome protein and plays a role in bacterial killing, *Cell. Immunol.* 279 (1) (2012) 46–52.
- [27] E.K. McEachern, J.H. Hwang, K.M. Sladewski, S. Nicatia, C. Dewitz, D.P. Mathew, V. Nizet, L.E. Crotty Alexander, Analysis of the effects of cigarette smoke on staphylococcal virulence phenotypes, *Infect. Immun.* 83 (6) (2015) 2443–2452.
- [28] A. Matsui, J.O. Jin, C.D. Johnston, H. Yamazaki, Y. Hourii-Haddad, S.R. Rittling, Pathogenic bacterial species associated with endodontic infection evade innate immune control by disabling neutrophils, *Infect. Immun.* 82 (10) (2014) 4068–4079.
- [29] L.C. Green, D.A. Wagner, J. Glogowski, P.L. Skipper, J.S. Wishnok, S. R. Tannenbaum, Analysis of nitrate, nitrite, and [15N]nitrate in biological fluids, *Anal. Biochem.* 126 (1) (1982) 131–138.
- [30] H. Tsutsuki, K. Yahiro, K. Suzuki, A. Suto, K. Ogura, S. Nagasawa, H. Ihara, T. Shimizu, H. Nakajima, J. Moss, M. Noda, Subtilase cytotoxin enhances *Escherichia coli* survival in macrophages by suppression of nitric oxide production through the inhibition of NF- κ B activation, *Infect. Immun.* 80 (11) (2012) 3939–3951.
- [31] O. Erel, A new automated colorimetric method for measuring total oxidant status, *Clin. Biochem.* 38 (12) (2005) 1103–1111.
- [32] D. Nilforoushan, A. Gramoun, M. Glogauer, M.F. Manolson, Nitric oxide enhances osteoclastogenesis possibly by mediating cell fusion, *Nitric Oxide: Biol. Chem.* 21 (1) (2009) 27–36.
- [33] P. Elsbach, J. Weiss, Oxygen-dependent and oxygen-independent mechanisms of microbicidal activity of neutrophils, *Immunol. Lett.* 11 (3–4) (1985) 159–163.
- [34] P. Elsbach, J. Weiss, A reevaluation of the roles of the O₂-dependent and O₂-independent microbicidal systems of phagocytes, *Rev. Infect. Dis.* 5 (5) (1983) 843–853.
- [35] A. Matsuzawa, K. Saegusa, T. Noguchi, C. Sadamitsu, H. Nishitoh, S. Nagai, S. Koyasu, K. Matsumoto, K. Takeda, H. Ichijo, ROS-dependent activation of the TRAF6-ASK1-p38 pathway is selectively required for TLR4-mediated innate immunity, *Nat. Immunol.* 6 (6) (2005) 587–592.
- [36] A.P. Gobert, M. Asim, M.B. Piazzuelo, T. Verriere, B.P. Scull, T. de Sablet, A. Glumac, N.D. Lewis, P. Correa, R.M. Peek Jr., R. Chaturvedi, K.T. Wilson, Disruption of nitric oxide signaling by *Helicobacter pylori* results in enhanced inflammation by inhibition of heme oxygenase-1, *J. Immunol.* 187 (10) (2011) 5370–5379.
- [37] H. Ha, H.B. Kwak, S.W. Lee, H.M. Jin, H.M. Kim, H.H. Kim, Z.H. Lee, Reactive oxygen species mediate RANK signaling in osteoclasts, *Exp. Cell Res.* 301 (2) (2004) 119–127.
- [38] D.Z. Li, Q.X. Zhang, X.X. Dong, H.D. Li, X. Ma, Treatment with hydrogen molecules prevents RANKL-induced osteoclast differentiation associated with inhibition of ROS formation and inactivation of MAPK, AKT and NF- κ B pathways in murine RAW264.7 cells, *J. Bone Miner. Metab.* 32 (5) (2014) 494–504.
- [39] S. Hyeon, H. Lee, Y. Yang, W. Jeong, Nrf2 deficiency induces oxidative stress and promotes RANKL-induced osteoclast differentiation, *Free Radic. Biol. Med.* 65 (2013) 789–799.
- [40] H. Kanzaki, F. Shinohara, I. Kanako, Y. Yamaguchi, S. Fukaya, Y. Miyamoto, S. Wada, Y. Nakamura, Molecular regulatory mechanisms of osteoclastogenesis through cytoprotective enzymes, *Redox Biol.* 8 (2016) 186–191.
- [41] R.J. van't Hof, S.H. Ralston, Cytokine-induced nitric oxide inhibits bone resorption by inducing apoptosis of osteoclast progenitors and suppressing osteoclast activity, *J. Bone Miner. Res.* 12 (11) (1997) 1797–1804.
- [42] T. Kaisho, S. Akira, Toll-like receptors and their signaling mechanism in innate immunity, *Acta Odontol. Scand.* 59 (3) (2001) 124–130.
- [43] B. Brune, N. Dehne, N. Grossmann, M. Jung, D. Namgaladze, T. Schmid, A. von Knethen, A. Weigert, Redox control of inflammation in macrophages, *Antioxid. Redox Signal.* 19 (6) (2013) 595–637.
- [44] T. Krisher, Z. Bar-Shavit, Regulation of osteoclastogenesis by integrated signals from toll-like receptors, *J. Cell. Biochem.* 115 (12) (2014) 2146–2154.
- [45] M. Kajiya, G. Giro, M.A. Taubman, X. Han, M.P. Mayer, T. Kawai, Role of periodontal pathogenic bacteria in RANKL-mediated bone destruction in periodontal disease, *J. Oral Microbiol.* 2 (2010).
- [46] W. Wang, D.J. Ferguson, J.M. Quinn, A.H. Simpson, N.A. Athanasou, Osteoclasts are capable of particle phagocytosis and bone resorption, *J. Pathol.* 182 (1) (1997) 92–98.
- [47] B. Meng, X. Yang, Y. Chen, J. Zhai, X. Liang, Effect of titanium particles on osteoclast activity in vitro, *Mol. Med. Rep.* 3 (6) (2010) 1065–1069.
- [48] O.J. Park, J. Yang, J. Kim, C.H. Yun, S.H. Han, *Enterococcus faecalis* attenuates the differentiation of macrophages into osteoclasts, *J. Endod.* 41 (5) (2015) 658–662.
- [49] J. Yang, O.J. Park, J. Kim, J.E. Baik, C.H. Yun, S.H. Han, Lipoteichoic acid of *Enterococcus faecalis* inhibits the differentiation of macrophages into osteoclasts, *J. Endod.* 42 (4) (2016) 570–574.
- [50] M. Takami, N. Kim, J. Rho, Y. Choi, Stimulation by toll-like receptors inhibits osteoclast differentiation, *J. Immunol.* 169 (3) (2002) 1516–1523.
- [51] W. Zou, Z. Bar-Shavit, Dual modulation of osteoclast differentiation by lipopolysaccharide, *J. Bone Miner. Res.* 17 (7) (2002) 1211–1218.
- [52] W. Zou, H. Schwartz, S. Endres, G. Hartmann, Z. Bar-Shavit, CpG oligonucleotides: novel regulators of osteoclast differentiation, *FASEB J.* 16 (3) (2002) 274–282.
- [53] H. Ha, J.H. Lee, H.N. Kim, H.B. Kwak, H.M. Kim, S.E. Lee, J.H. Rhee, H.H. Kim, Z. H. Lee, Stimulation by TLR5 modulates osteoclast differentiation through STAT1/IFN- β , *J. Immunol.* 180 (3) (2008) 1382–1389.
- [54] H. Suzuki, A. Mochizuki, K. Yoshimura, Y. Miyamoto, K. Kaneko, T. Inoue, D. Chikazu, M. Takami, R. Kamijo, Bropiramine inhibits osteoclast differentiation through production of interferon- β , *Biochem. Biophys. Res. Commun.* 467 (1) (2015) 146–151.
- [55] H. Takayanagi, S. Kim, K. Matsuo, H. Suzuki, T. Suzuki, K. Sato, T. Yokochi, H. Oda, K. Nakamura, N. Ida, E.F. Wagner, T. Taniguchi, RANKL maintains bone homeostasis through c-Fos-dependent induction of interferon- β , *Nature* 416 (6882) (2002) 744–749.
- [56] J.D. Ji, K.H. Park-Min, Z. Shen, R.J. Fajardo, S.R. Goldring, K.P. McHugh, L. B. Ivashkiv, Inhibition of RANK expression and osteoclastogenesis by TLRs and IFN- γ in human osteoclast precursors, *J. Immunol.* 183 (11) (2009) 7223–7233.
- [57] T. Ishii, Y. Shimazu, I. Nishiyama, J. Kikuta, M. Ishii, The role of sphingosine 1-phosphate in migration of osteoclast precursors: an application of intravital two-photon microscopy, *Mol. Cells* 31 (5) (2011) 399–403.
- [58] A. Movila, T. Ishii, A. Albassam, W. Wisitrasameewong, M. Howait, T. Yamaguchi, M. Ruiz-Torruella, L. Bahammam, K. Nishimura, T. Van Dyke, T. Kawai, Macrophage Migration Inhibitory Factor (MIF) supports homing of osteoclast precursors to peripheral osteolytic lesions, *J. Bone Miner. Res.* (2016).
- [59] S. Nourshargh, R. Alon, Leukocyte migration into inflamed tissues, *Immunity* 41 (5) (2014) 694–707.
- [60] D.W. Dempster, C.L. Lambing, P.J. Kostenuik, A. Grauer, Role of RANK ligand and denosumab, a targeted RANK ligand inhibitor, in bone health and osteoporosis: a review of preclinical and clinical data, *Clin. Ther.* 34 (3) (2012) 521–536.
- [61] C. Reyes, M. Hitz, D. Prieto-Alhambra, B. Abrahamsen, Risks and benefits of bisphosphonate therapies, *J. Cell. Biochem.* 117 (1) (2016) 20–28.
- [62] M.S. Epstein, H.D. Ephros, J.B. Epstein, Review of current literature and implications of RANKL inhibitors for oral health care providers, *Oral Surg. Oral Med. Oral Pathol. Oral Radiol.* 116 (6) (2013) e437–e442.

TRAP-positive osteoclast precursors mediate ROS/NO-dependent bactericidal activity via TLR4

ORIGINALITY REPORT

20%
SIMILARITY INDEX

12%
INTERNET SOURCES

19%
PUBLICATIONS

0%
STUDENT PAPERS

PRIMARY SOURCES

- 1** Xinqiang Lai, Yanxia Ye, Chenghong Sun, Xiuyan Huang, Xiangao Tang, Xiangfeng Zeng, Pinghe Yin, Yaoying Zeng. "Icaritin exhibits anti-inflammatory effects in the mouse peritoneal macrophages and peritonitis model", *International Immunopharmacology*, 2013
Publication **1%**
- 2** Marcos Roberto de Oliveira, Seyed Fazel Nabavi, Seyed Mohammad Nabavi, Fernanda Rafaela Jardim. "Omega-3 polyunsaturated fatty acids and mitochondria, back to the future", *Trends in Food Science & Technology*, 2017
Publication **1%**
- 3** Murat Konak, Nuriye Tarakci, Hüseyin Altunhan, Ali Annagür, Aysun Toker, Rahmi Örs. "Total antioxidant, total oxidant and serum paraoxonase levels according to lipid administration method in parenterally fed **1%**

premature infants", The Journal of Maternal-Fetal & Neonatal Medicine, 2016

Publication

4	link.springer.com Internet Source	1 %
5	Maria Belén Herrero, Suvro Chatterjee, Linda Lefièvre, Eve de Lamirande, Claude Gagnon. "Nitric oxide interacts with the cAMP pathway to modulate capacitation of human spermatozoa", Free Radical Biology and Medicine, 2000 Publication	1 %
6	www.spandidos-publications.com Internet Source	1 %
7	www.frontiersin.org Internet Source	1 %
8	"Program & Abstracts", Journal of Bone and Mineral Research, 09/2002 Publication	<1 %
9	Seicho Makihira, Hiroki Nikawa, Mikihiro Kajiya, Toshihisa Kawai et al. "Blocking of sodium and potassium ion-dependent adenosine triphosphatase- α 1 with ouabain and vanadate suppresses cell-cell fusion during RANKL-mediated osteoclastogenesis", European Journal of Pharmacology, 2011 Publication	<1 %

10

www.biorxiv.org

Internet Source

<1 %

11

www.dent.meikai.ac.jp

Internet Source

<1 %

12

Qi Zhang, Decheng Wang, Guozhong Jiang, Wei Liu, Qing Deng, Xiujun Li, Wei Qian, Hugues Ouellet, Jianjun Sun. "EsxA membrane-permeabilizing activity plays a key role in mycobacterial cytosolic translocation and virulence: effects of single-residue mutations at glutamine 5", Scientific Reports, 2016

Publication

<1 %

13

Staab, Elizabeth, Geoffrey M. Thiele, Dillon Clarey, Todd A. Wyatt, Debra J. Romberger, Adam D. Wells, Anand Dusad, Dong Wang, Lynell W. Klassen, Ted R. Mikuls, Michael J. Duryee, and Jill A. Poole. "Toll-Like Receptor 4 Signaling Pathway Mediates Inhalant Organic Dust-Induced Bone Loss", PLoS ONE, 2016.

Publication

<1 %

14

X. Lin. "Antibody to Receptor Activator of NF-B Ligand Ameliorates T Cell-Mediated Periodontal Bone Resorption", Infection and Immunity, 02/01/2011

Publication

<1 %

- | | | |
|----|---|------|
| 15 | Gilson C.N. Franco, Mikihito Kajiya, Tadashi Nakanishi, Kouji Ohta et al. "Inhibition of matrix metalloproteinase-9 activity by doxycycline ameliorates RANK ligand-induced osteoclast differentiation in vitro and in vivo", <i>Experimental Cell Research</i> , 2011
Publication | <1 % |
| 16 | apps.dtic.mil
Internet Source | <1 % |
| 17 | synapse.koreamed.org
Internet Source | <1 % |
| 18 | García González Víctor Guadalupe. "Propiedades amiloidogénicas de un fragmento derivado del dominio C-terminal de la proteína transferidora de ésteres de colesterol (CETP)", TESIUNAM, 2013
Publication | <1 % |
| 19 | www.ncbi.nlm.nih.gov
Internet Source | <1 % |
| 20 | Amit R. Ashtekar, Ping Zhang, Jannet Katz, Champion C. S. Deivanayagam et al. " TLR4-mediated activation of dendritic cells by the heat shock protein DnaK from ", <i>Journal of Leukocyte Biology</i> , 2008
Publication | <1 % |
| 21 | Jian Wu, Feng-Ling Luo, Yan Xie, Huan Xiong, Yadong Gao, Guanghui Liu, Xiao-Lian Zhang. | <1 % |

"EST12 regulates Myc expression and enhances anti-mycobacterial inflammatory response via RACK1-JNK-AP1-Myc immune pathway", *Frontiers in Immunology*, 2022

Publication

22

S. M. Morris. "Arginine: Master and Commander in Innate Immune Responses", *Science Signaling*, 2010

Publication

23

www.freepatentsonline.com

Internet Source

24

Lee, N.K.. "RANKL-induced schlafen2 is a positive regulator of osteoclastogenesis", *Cellular Signalling*, 200812

Publication

25

Li, Changhong, Jinxia Zhao, Lin Sun, Zhongqiang Yao, Rui Liu, Jiansheng Huang, and Xiangyuan Liu. "RANKL downregulates cell surface CXCR6 expression through JAK2/STAT3 signaling pathway during osteoclastogenesis", *Biochemical and Biophysical Research Communications*, 2012.

Publication

26

Wittrant, Y.. "High d(+)glucose concentration inhibits RANKL-induced osteoclastogenesis", *Bone*, 200806

Publication

<1 %

<1 %

<1 %

<1 %

<1 %

27	www.researchgate.net Internet Source	<1 %
28	Kerri L. George. "Ajulemic acid, a nonpsychoactive cannabinoid acid, suppresses osteoclastogenesis in mononuclear precursor cells and induces apoptosis in mature osteoclast-like cells", <i>Journal of Cellular Physiology</i> , 03/2008 Publication	<1 %
29	T.C. Peixoto, E.G. Moura, E. Oliveira, V. Younes-Rapozo et al. "Neonatal tobacco smoke reduces thermogenesis capacity in brown adipose tissue in adult rats", <i>Brazilian Journal of Medical and Biological Research</i> , 2018 Publication	<1 %
30	Young-Chang Cho, Kwang Youl Lee, Bok Yun Kang, Haeng Soon Park. "Enhanced IL-12p40 production by phenylarsine oxide is mediated by cAMP response element in macrophages", <i>Archives of Pharmacal Research</i> , 2010 Publication	<1 %
31	"ASBMR 23rd annual meeting", <i>Journal of Bone and Mineral Research</i> , 2001 Publication	<1 %
32	Yuichi Mine, Takahiro Shuto, Hiroki Nikawa, Toshihisa Kawai et al. "Inhibition of RANKL-	<1 %

dependent cellular fusion in pre-osteoclasts by amiloride and a NHE10-specific monoclonal antibody", Cell Biology International, 2015

Publication

33

Lisa Mullen, Manuela Mengozzi, Eva-Maria Hanschmann, Ben Alberts, Pietro Ghezzi. "How the redox state regulates immunity", Free Radical Biology and Medicine, 2019

Publication

<1 %

34

Naoki Koide, Hiroyasu Ito, Mya Mya Mu, Tsuyoshi Sugiyama et al. "Inhibition of extracellular signal-regulated kinase 1/2 augments nitric oxide production in lipopolysaccharide-stimulated RAW264.7 macrophage cells", FEMS Immunology & Medical Microbiology, 2005

Publication

<1 %

35

Okito, Asuka, Ken-ichi Nakahama, Masako Akiyama, Takashi Ono, and Ikuo Morita. "Involvement of the G-protein-coupled receptor 4 in RANKL expression by osteoblasts in an acidic environment", Biochemical and Biophysical Research Communications, 2015.

Publication

<1 %

36

jnanobiotechnology.biomedcentral.com

Internet Source

<1 %

37 Nakamura, Kimiya, Yoshiaki Deyama, Yoshitaka Yoshimura, Kuniaki Suzuki, and Manabu Morita. "Toll-like receptor 3 ligand-induced antiviral response in mouse osteoblastic cells", International Journal of Molecular Medicine, 2007. <1 %

Publication

38 Wen-Fang Xia, Fu-Lei Tang, Lei Xiong, Shan Xiong, Ji-Ung Jung, Dae-Hoon Lee, Xing-Sheng Li, Xu Feng, Lin Mei, Wen-Cheng Xiong. "Vps35 loss promotes hyperresorptive osteoclastogenesis and osteoporosis via sustained RANKL signaling", The Journal of Cell Biology, 2013 <1 %

Publication

39 journals.sagepub.com <1 %

Internet Source

40 tvst.arvojournals.org <1 %

Internet Source

41 www.mdpi.com <1 %

Internet Source

42 www.plosone.org <1 %

Internet Source

43 www.prf.upol.cz <1 %

Internet Source

44 "Abstracts", Inflammation Research, 2014

<1 %

45

Lijuan Hu. "Retinoic Acid Increases Proliferation of Human Osteoclast Progenitors and Inhibits RANKL-Stimulated Osteoclast Differentiation by Suppressing RANK", PLoS ONE, 10/11/2010

Publication

<1 %

46

M. Emoto, T. Yoshida, T. Fukuda, I. Kawamura, M. Mitsuyama, E. Kita, R. Hurwitz, S. H. E. Kaufmann, Y. Emoto. "-Galactosylceramide Promotes Killing of Listeria monocytogenes within the Macrophage Phagosome through Invariant NKT-Cell Activation", Infection and Immunity, 2010

Publication

<1 %

47

Patrizia Bottoni, Roberto Scatena. "The Tangled Mitochondrial Metabolism in Cancer: An Innovative Pharmacological Approach", Current Medicinal Chemistry, 2020

Publication

<1 %

48

Seung-Hyun Ryu, Jong-Hyung Park, Soo-Young Choi, Hee-Yeon Jeon, Jin-Il Park, Jun-Young Kim, Seung-Hoon Ham, Yang-Kyu Choi. "The Probiotic Lactobacillus Prevents Citrobacter rodentium-Induced Murine Colitis in a TLR2-Dependent Manner", Journal of Microbiology and Biotechnology, 2016

<1 %

49

docksci.com

Internet Source

<1 %

50

Donald W. Lawrence, Jacki Kornbluth. "E3 ubiquitin ligase NKLAM is a macrophage phagosome protein and plays a role in bacterial killing", Cellular Immunology, 2012

Publication

<1 %

51

Eun-Kyeong Jo. "Role of apoptosis-regulating signal kinase 1 in innate immune responses by Mycobacterium bovis bacillus Calmette-Guérin", Immunology and Cell Biology, 10/14/2008

Publication

<1 %

52

Jae Lim. "Streptococcus pneumoniae synergizes with nontypeable Haemophilus influenzae to induce inflammation via upregulating TLR2", BMC Immunology, 2008

Publication

<1 %

53

Jisu Lee, Jeongmin Lee, Eunha Hwang, Hye-Jung Kim et al. "An Efficient and Rapid Two-step Purification Method for Active Human Macrophage Colony-stimulating Factor from Escherichia coli", Biotechnology and Bioprocess Engineering, 2019

Publication

<1 %

54

Jun Xu, Xiaomei Zhang, Marc Monestier, Naomi L. Esmon, Charles T. Esmon. "Extracellular Histones Are Mediators of Death through TLR2 and TLR4 in Mouse Fatal Liver Injury", *The Journal of Immunology*, 2011
Publication

<1 %

55

Liang, Xing. "Effect of titanium particles on osteoclast activity in vitro", *Molecular Medicine Reports*, 2010.
Publication

<1 %

56

Orli Even-Or, Shani Avniel-Polak, Yechezkel Barenholz, Gabriel Nussbaum. "The cationic liposome CCS/C adjuvant induces immunity to influenza independently of the adaptor protein MyD88", *Human Vaccines & Immunotherapeutics*, 2020
Publication

<1 %

57

Prachi Vilekar, Catherine King, Pallavi Lagisetty, Vibhudutta Awasthi, Shanjana Awasthi. "Antibacterial Activity of Synthetic Curcumin Derivatives: 3,5-Bis(benzylidene)-4-Piperidone (EF24) and EF24-Dimer Linked via Diethylenetriaminepentacetic Acid (EF2DTPA)", *Applied Biochemistry and Biotechnology*, 2014
Publication

<1 %

58

Woo Sik Kim, Jong-Seok Kim, Seung Bin Cha, Hongmin Kim et al. " Rv3628 drives Th1-type T

<1 %

cell immunity via TLR2-mediated activation of dendritic cells and displays vaccine potential against the hyper-virulent Beijing K strain ",
Oncotarget, 2016

Publication

59	bsdwebstorage.blob.core.windows.net Internet Source	<1 %
60	downloads.hindawi.com Internet Source	<1 %
61	iai.asm.org Internet Source	<1 %
62	journals.aai.org Internet Source	<1 %
63	nagasaki-u.repo.nii.ac.jp Internet Source	<1 %
64	oak.jejunu.ac.kr Internet Source	<1 %
65	people.uea.ac.uk Internet Source	<1 %
66	s-space.snu.ac.kr Internet Source	<1 %
67	www.aimspress.com Internet Source	<1 %
68	Christel Claes, Johanna Van Den Daele, Ruben Boon, Sarah Schoutedden et al. "Human stem	<1 %

cell-derived monocytes and microglia-like cells reveal impaired amyloid plaque clearance upon heterozygous or homozygous loss of TREM2", *Alzheimer's & Dementia*, 2019

Publication

69

George Pan. "Modulation of Osteoclastogenesis Induced by Nucleoside Reverse Transcriptase Inhibitors", *AIDS Research and Human Retroviruses*, 11/2006

Publication

70

Hiroto Nakajima, Naoki Shimbara, Yoko Shimonishi, Tatsuyuki Mimori, Shin-ichiro Niwa, Hideyuki Saya. "Expression of random peptide fused to invasins on bacterial cell surface for selection of cell-targeting peptides", *Gene*, 2000

Publication

71

Hongbin Deng, Na Zhang, Yan Wang, Jinjing Chen, Jiajia Shen, Zhen Wang, Rong Xu, Jingpu Zhang, Danqing Song, Diandong Li. "S632A3, a new glutarimide antibiotic, suppresses lipopolysaccharide-induced pro-inflammatory responses via inhibiting the activation of glycogen synthase kinase 3 β ", *Experimental Cell Research*, 2012

Publication

72

Jennifer M. Burke, Lisa M. Ganley-Leal, Asma Khatri, Lee M. Wetzler. "PorB, a TLR2 Ligand,

<1 %

<1 %

<1 %

<1 %

Induces an Antigen-Specific Eosinophil Recall Response: Potential Adjuvant for Helminth Vaccines? ", The Journal of Immunology, 2007

Publication

73

K.H. Szymczyk, B.A.E. Kerr, T.A. Freeman, C.S. Adams, M.J. Steinbeck. "Involvement of hydrogen peroxide in the differentiation and apoptosis of preosteoclastic cells exposed to arsenite", Biochemical Pharmacology, 2006

Publication

74

Kittaka, M., H. Shiba, M. Kajiya, K. Ouhara, K. Takeda, K. Kanbara, T. Fujita, H. Kawaguchi, H. Komatsuzawa, and H. Kurihara. "Antimicrobial peptide LL37 promotes vascular endothelial growth factor-A expression in human periodontal ligament cells", Journal of Periodontal Research, 2013.

Publication

75

Kondapalli Mrudula Spurthi, Mohsen Sarikhani, Sneha Mishra, Perumal Arumugam Desingu et al. "Toll-like receptor 2 deficiency hyperactivates the FoxO1 transcription factor and induces aging-associated cardiac dysfunction in mice", Journal of Biological Chemistry, 2018

Publication

76

Lari, R.. "Macrophage lineage phenotypes and osteoclastogenesis-Complexity in the control

<1 %

<1 %

<1 %

<1 %

77

Lee, T.H.. "Inhibitory effects of a spinasterol glycoside on lipopolysaccharide-induced production of nitric oxide and proinflammatory cytokines via down-regulating MAP kinase pathways and NF- κ B activation in RAW264.7 macrophage cells", *International Immunopharmacology*, 201207

Publication

<1 %

78

W Komatsu. "Docosahexaenoic acid suppresses nitric oxide production and inducible nitric oxide synthase expression in interferon- γ plus lipopolysaccharide-stimulated murine macrophages by inhibiting the oxidative stress", *Free Radical Biology and Medicine*, 2003

Publication

<1 %

79

Ya-Lin Huang. "Comparison of anti-inflammatory activities of ruscogenin, a major steroidal sapogenin from *Radix Ophiopogon japonicus*, and Its succinylated derivative, RUS-2HS", *Drug Development Research*, 06/2008

Publication

<1 %

80

api.research-repository.uwa.edu.au

Internet Source

<1 %

81

coek.info

Internet Source

<1 %

82	epdf.tips Internet Source	<1 %
83	ir.library.louisville.edu Internet Source	<1 %
84	libraetd.lib.virginia.edu Internet Source	<1 %
85	ndl.ethernet.edu.et Internet Source	<1 %
86	patents.google.com Internet Source	<1 %
87	pure.rug.nl Internet Source	<1 %
88	www.academia.dk Internet Source	<1 %
89	Hae-Ri Lee, Jun Ho Jeon, Ok-Kyu Park, Jeong-Hoon Chun, Jungchan Park, Gi-eun Rhie. "The poly-γ- d -glutamic acid capsule surrogate of the Bacillus anthracis capsule induces nitric oxide production via the platelet activating factor receptor signaling pathway", Molecular Immunology, 2015 Publication	<1 %
90	M. R. Murawski, G. N. Bowen, A. M. Cerny, L. J. Anderson, L. M. Haynes, R. A. Tripp, E. A. Kurt-Jones, R. W. Finberg. "Respiratory Syncytial Virus Activates Innate Immunity through Toll-	<1 %

Like Receptor 2", Journal of Virology, 2008

Publication

91

Mengqin Liu, Sha Liu, Qi Zhang, Yingqi Fang, Yanwei Yu, Lulin Zhu, Yuling Liu, Wan Gong, Luying Zhao, Luping Qin, Qiaoyan Zhang.
"Curculigoside attenuates oxidative stress and osteoclastogenesis via modulating Nrf2/NF- κ B signaling pathway in RAW264.7 cells", Journal of Ethnopharmacology, 2021

Publication

<1 %

92

S.W. Fox, K.E. Evans, A.C. Lovibond.
"Transforming growth factor- β enables NFATc1 expression during osteoclastogenesis", Biochemical and Biophysical Research Communications, 2008

Publication

<1 %

93

Sundaram, K.. "RANK ligand signaling modulates the matrix metalloproteinase-9 gene expression during osteoclast differentiation", Experimental Cell Research, 20070101

Publication

<1 %

94

Wang, Yu-Hsiung, Reza Nemati, Emily Anstadt, Yaling Liu, Young Son, Qiang Zhu, Xudong Yao, Robert B. Clark, David W. Rowe, and Frank C. Nichols. "Serine dipeptide lipids of Porphyromonas gingivalis inhibit osteoblast

<1 %

differentiation: Relationship to Toll-like receptor 2", Bone, 2015.

Publication

95

Yan Xu. "PAMM: A REDOX REGULATORY PROTEIN THAT MODULATES OSTEOCLAST DIFFERENTIATION", Antioxidants and Redox Signaling, 12/2009

Publication

<1 %

96

Yoshitaka Hosokawa. "Tea polyphenols inhibit IL-6 production in tumor necrosis factor superfamily 14-stimulated human gingival fibroblasts", Molecular Nutrition & Food Research, 05/11/2010

Publication

<1 %

97

Akira Takahashi, Masayuki Nakano, Keinosuke Okamoto, Yoshio Fujii, Kazuaki Mawatari, Nagakatsu Harada, Yutaka Nakaya. "Aeromonas sobria hemolysin causes diarrhea by increasing secretion of HCO₃⁻", FEMS Microbiology Letters, 2006

Publication

<1 %

98

Collazo, C.M.. "Myeloid differentiation factor-88 (MyD88) is essential for control of primary in vivo Francisella tularensis LVS infection, but not for control of intramacrophage bacterial replication", Microbes and Infection, 200603

Publication

<1 %

99

Jianxin Qiu, Ting Jiang, Guangyong Yang, Yuhang Gong, Weikang Zhang, Xiaohang Zheng, Zhenghua Hong, Haixiao Chen.

"Neratinib exerts dual effects on cartilage degradation and osteoclast production in Osteoarthritis by inhibiting the activation of the MAPK/NF- κ B signaling pathways", *Biochemical Pharmacology*, 2022

Publication

<1 %

100

Soo-Hyun Yoon. "Adenylate cyclase and calmodulin-dependent kinase have opposite effects on osteoclastogenesis by regulating the PKA-NFATc1 pathway", *Journal of Bone and Mineral Research*, 06/2011

Publication

<1 %

101

Yurtcu, Engin, Cihan Togrul, Sebnem Ozyer, Ozlem Uzunlar, Yeksin H. Karatas, Kerem D. Seckin, Muzaffer Caydere, Sema Hucumenoglu, and Nedim Cicek. "Dose dependent protective effects of vardenafil on ischemia-reperfusion injury with biochemical and histopathologic evaluation in rat ovary", *Journal of Pediatric Surgery*, 2015.

Publication

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

TRAP-positive osteoclast precursors mediate ROS/NO-dependent bactericidal activity via TLR4

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10

PAGE 11

PAGE 12
