

Neglected Orbitozygomaticomaxillary Fractures With Complications: A Case Report

by Indri Lakhsmi Putri

Submission date: 07-Apr-2023 04:36PM (UTC+0800)

Submission ID: 2058271570

File name: gomaticomaxillary_Fractures_With_Complications_A_Case_Report.pdf (1.64M)

Word count: 2799

Character count: 16109



ELSEVIER

Contents lists available at ScienceDirect

International Journal of Surgery Case Reports

journal homepage: www.casereports.com

Neglected orbitozygomaxillary fractures with complications: A case report

Siti Isya Wahdini^{a,*}, Ishandono Dachlan^a, Rosadi Seswandhana^a,
Magda Rosalina Hutagalung^b, Indri Lakshmi Putri^b, Dwiki Afandy^c

^a Division of Plastic, Aesthetic & Reconstructive Surgery, Department of Surgery, Faculty of Medicine, Public Health and Nursing, Universitas Gadjah Mada/Dr. Sardjito Hospital, Yogyakarta, Indonesia

^b Department of Plastic, Aesthetic & Reconstructive Surgery, Faculty of Medicine, Universitas Airlangga/Dr. Soetomo Hospital, Surabaya, Indonesia

^c Faculty of Medicine, Public Health and Nursing, Universitas Gadjah Mada, Yogyakarta, Indonesia

ARTICLE INFO

Article history:

Received 14 May 2019

Received in revised form 18 July 2019

Accepted 19 July 2019

Available online 26 July 2019

Keywords:

Orbitozygomaxillary fracture
Maxillofacial trauma
Surgical repair
Case report

ABSTRACT

INTRODUCTION: Zygomatic complex fracture is one of the most frequently occurring injuries to facial structures and often involves damage to the surrounding structures, including the orbital floor and/or medial orbital wall. These complicated injuries can cause both serious ophthalmic and aesthetic complications that make reconstructions more difficult.

CASE PRESENTATION: A 22-year old male presented with facial asymmetry. On examination, there were neglected fractures of left orbitozygomaxillary complex with various complications. We performed lateral orbitotomy, bone graft, medial and lateral cantopexy with 3D reconstruction models as a guidance. There were some appearance improvements including improved orbital dystopia and corrected facial asymmetry. However, his vision could not be corrected because the reconstruction was too late.

DISCUSSION: Several examinations before reconstruction of the injury has to be done, especially radiological examinations to decide proper diagnosis and reconstruction plans. Some complications may make difficulties in establishing precise time for surgery. Three-dimensional (3-D) reconstruction model can help increase accuracy of the reconstruction but consume more time.

CONCLUSION: Severe maxillofacial fracture should be treated immediately to avoid further complications. Multidisciplinary examinations can provide more accurate preliminary recommendations particularly when combine with properly calibrated CT scan imaging.

© 2019 The Authors. Published by Elsevier Ltd on behalf of IJS Publishing Group Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Facial reconstruction surgery is common because the face is a prominent point of injury due to sports injuries and vehicular accidents. The cheek bones and eye sockets often need repair requiring extensive rebuilding [1]. Several cosmetic procedures have been developed and the most frequently occurring injury is the zygomatic complex (ZMC) fractures characterized by fracture (s) of the zygoma or adjacent bones, such as the maxilla, orbit, or temporal bone [2,3]. Those ZMC fractures often involve damage to the orbital floor and/or the medial orbital wall [4].

The orbital floor has a fragile structure, making it easily damaged in a craniomaxillofacial trauma. The area located in medial to the infraorbital groove and canal are main locations of sustained fractures. Sometimes, the fractures involve damage to medial orbital wall because of the reduced bone thickness in that area [5]. These fractures can cause serious ophthalmic and aesthetic complications. Serious ophthalmic complications include visual disturbance, diplopia, and enophthalmos, while facial asymmetry is an aesthetic complication results from the injuries [6]. These complications can give significant challenges to the plastic surgeon to repair.

This case report follows SCARE criteria [7].

2. Case presentation

A 22 year old male presented to the Division of Plastic, Reconstruction, and Aesthetic Surgery, in the Department of Surgery of our institute with facial asymmetry. He was injured in a traffic accident 2 years ago and had underwent interdental and intermaxillary wiring by a surgeon at a district hospital. The patient did not feel any improvements and then came to our institute. On the physical

* Corresponding author at: Division of Plastic, Aesthetic & Reconstructive Surgery, Department of Surgery, Faculty of Medicine, Public Health and Nursing, Universitas Gadjah Mada/Dr. Sardjito Hospital, Jl. Kesehatan No. 1, Yogyakarta 55281, Indonesia.
E-mail addresses: tisya.wahdini@gmail.com (S. Isya Wahdini), ishandono@yahoo.co.id (I. Dachlan), rosadi_seswandhana@ugm.ac.id (R. Seswandhana), magdahutagalung@rocketmail.com (M.R. Hutagalung), indrilakshmiptutri@gmail.com (I.L. Putri), dwiki.afandy@mail.ugm.ac.id (D. Afandy).

¹ <https://doi.org/10.1016/j.ijscr.2019.07.055>

2210-2612/© 2019 The Authors. Published by Elsevier Ltd on behalf of IJS Publishing Group Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).




Fig. 1. The patient's physical appearance before reconstruction.



Fig. 2. Three-dimensional CT-scan before reconstruction.



Fig. 3. Three-dimensional (3-D) reconstructions model used as guidance during surgery.

examination, there was maxilla deformity on the left side (Fig. 1). Ophthalmology examination found light perception only in the left eye. This came along with orbital dystopia and obstruction of nasolacrimal duct. Then we performed 3-D CT scan and turned out that there was a fracture of left orbitozygomaticomaxillary complex. It showed shattered orbital plate and processus frontalis. Zygoma bone and maxilla bone also shifted to below of its proper position (Fig. 2).

We did the reconstruction using bicoronal approach and 3D reconstruction models as a guidance for the surgical procedures (Fig. 3). We performed zygomaticomaxilla osteotomy and Tessier's inferior orbital marginotomy. Bone graft was harvested from 8th rib bone with 6,5 cm total length. 3,5 cm was applied to frontozygoma and 3 cm to left maxilla (Fig. 4). The bone graft then was fixed with plate. Medial and lateral cantopexy were installed with screw. The schematic picture is showed in (Fig. 5).

In the 4th month after operation the cantopexy was still broken causing sagging of the cheek skin to be prominent. The 2nd operation was performed to fix the sagging cheek by mini face lift and orbital dystopia by adding bone wax and orbital mesh, and also

reconstruction of nasolacrimal duct. There were some appearance improvements including corrected orbital dystopia and improved facial asymmetry (Fig. 6). Also, he had no more epiphora. However, his vision could not be corrected because it was too late.

3. Discussion

Fractures of the orbitozygomaticomaxillary complex are involved in nearly 30% of all facial fractures [8], while orbital fractures represent over 40% of maxillofacial injuries, making them the most common trauma in the midface. Diagnosis of ZMC involves fracture of the lateral orbital wall and articulation of the zygoma with the greater wing of the sphenoid [9]. Manson et al. describe the injury around the zygomatic bone as low energy, middle energy, and high energy injuries [10]. In our report, the patient belongs to




Fig. 4. Bone graft installment in frontozygoma (left) and maxilla (right).




Fig. 5. Schematic picture of our reconstruction approach (left) and the result of reconstruction (right). (Courtesy of Siti Isya Wahdini).

the high energy injury due to the comminution in the lateral orbit and telescoping of the zygomatic arch.

Healthcare providers who take care of these patients have to understand the related physical examinations, including ophthalmology findings. Any misdiagnosis or incorrect reconstruction of the preinjury anatomy can cause post traumatic deformities of the orbit that can lead to serious complications, including enophthalmos, diplopia, and visual acuity disturbance [11]. Early diagnosis and management of any facial injury is essential to prevent late treatment and possible permanent disfigurement. Preoperative work on these patients has to be done properly including computed tomography (CT) scans [9]. Correct phantom calibration is essential for CT image readings and proper diagnosis.

A variety of defects in the orbital wall can be captured with CT imaging ranging from hairline fractures without displacement to full separation of orbital bones which can cause tissue prolapse in the eye socket and surrounding orbitomaxillary structures including the cheek bone and detached facial muscles. During repair of the injury to the zygoma, early exploration of the internal orbit can provide a foundation for later cosmetic surgery by rebuilding the orbital floor [12]. In this patient, we report a successful repair using a mesh floor to support the subsequent face lift and cosmetic tissue repair.

According to Nigel et al., orbital floor fractures are considered urgent with following complications: diplopia with restriction of gaze, entrapment of extraocular muscles, enophthalmos >2 mm, fracture that involved over half the orbital floor and the medial wall, and "trapdoor" orbital floor fractures in children [8]. A systematic review by Dubois et al. identified some patients who developed challenging enophthalmos or persistent diplopia 2 weeks post injury. These complications create difficulties in early treatment recommendations particularly when patients have sustained only small damage to the eye and may not need surgery [13].

Ophthalmologic findings and CT scan imaging can contribute significantly to recommendations for surgery and may need further evaluations from other disciplines such as neurosurgery for possible traumatic brain injury [5,13]. The goal of fracture reconstructions at the orbital area is different with fractures at long bones. They aim to restore the appearance and skeletal anatomy before injury instead of bone healing [14]. The time of surgery is still controversial. Burstine criteria is used often to decide the best time of surgery but must be revisited. Dubois et al. discussed recent studies of the effects of surgical timing on post-traumatic orbital reconstruction outcomes, which provide criteria for immediate repair and augmented guidelines to determine the ideal time for non-immediate surgery [13].



Fig. 6. The patient's physical appearance and three-dimensional CT-scan after reconstruction.

High energy injuries also often require extensive osteotomy, repositioning, and cranial bone graft replacement of displaced, partially resorbed, and comminuted bone. Extended open reduction and rigid fixation techniques in treating high-energy fractures are highly required, since the soft tissue deforming forces overwhelm the limited fixation of comminuted fractures, particularly fixation provided by interfragment wiring. A common combination of procedures is osteotomy and repositioning of the zygomatic arch to decrease facial width and custom onlay grafting of the malar prominence [14]. In the first treatment of our patient, he only underwent interdental wiring-intermaxillary wiring. It was an inadequate treatment that made him present to our hospital with complications. Then we performed lateral orbitotomy, bone graft, medial and lateral cantopexy with 3-D reconstruction model as a guidance for the surgeries.

Orbital reconstruction in orbital fracture treatment using implant materials is effective, but it is difficult to assess its accuracy during surgery. Therefore, a three-dimensional (3-D) reconstruction model was created as guidance during surgery. The use of 3-D models has significantly aided the surgical planning and treatment outcome of both pathologic and traumatic maxillofacial conditions. The main goal is restoration of the original orbital volume and prevention of long term complications [6,15]. Although this method easily reproduces accurate orbital positioning and shape, the prolonged time required to create and prepare these custom made pre-bending plate systems make it inapplicable during emergencies [13].

4. Conclusion

We successfully treated a neglected case of orbitozygomati-comaxillary fracture with 2 years onset that never reported in any literature before. Bone healing process normally involves callus resorption by osteoclast and formation of lamellar bone by

osteoblast. There is no any healing of injured bone in the result of imaging in this patient. Our approachment to this case give a satisfying outcome despite of inevitable vision loss. When complex and severe maxillofacial fractures occurred especially with high energy injury causing ophthalmology and aesthetic complications, these cases should be referred to a plastic surgeon immediately. Early reconstruction is recommended although its accuracy has to be performed carefully because the 3-D reconstruction model takes longer time. Multidisciplinary examinations can provide more accurate preliminary recommendations particularly when combine with properly calibrated CT scan imaging.

Sources of funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Ethical approval

Ethical clearance is not needed in the ethics commission at our institution for a case report.

Consent

Written informed consent was obtained from the patient for publication of this case report and accompanying images. A copy of the written consent is available for review by the Editor-in-Chief of this journal on request.

Author's contribution

SIW is the author conceived the study, and ID, RS, MRH, and ILP did the supervision. DA contributed in data collection.

Registration of research studies

We use registry body from Clinicaltrials.gov with registration number NCT00583856.

Guarantor

Siti Isya Wahdini.

Provenance and peer review

Not commissioned, externally peer-reviewed.

Declaration of Competing Interest

The authors declare that they have no competing interests.

Acknowledgments

We thank the surgical staff and nursing team who were involved in the patient's care. We are also thankful to a native speaker at English Services Center, Faculty of Medicine, Public Health and Nursing, Universitas Gadjah Mada, for proofreading of our manuscript.

References

- [1] S. Ramisetty, R. Gaddipati, R. Yamsani, B. Yamsani, N. Vura, Zygomaticomaxillary complex fractures: a review of 101 cases, *J. Maxillofac. Oral Surg.* 15 (2015) 417–424, <http://dx.doi.org/10.1007/s12663-015-0851-9>.
- [2] L. Ye, D. Dikhat, Y. Wang, W. Liu, L. Liu, H. Li, X. Zhang, Y. Chen, Surgical navigation improves reductions accuracy of unilateral complicated zygomaticomaxillary complex fractures: a randomized controlled trial, *Sci. Rep.* 8 (2018) 1–9, <http://dx.doi.org/10.1038/s41598-018-25053-z>.
- [3] X.G. Dds, Y.H. Dds, J.A. Dds, Y.Y. Dds, X.H. Dds, M.L. Dds, Y.Z. Dds, Y.Z. Dds, Application of a Computer-Assisted Navigation System (CANS) in the delayed treatment of zygomatic fractures: a randomized controlled, *J. Oral Maxillofac. Surg.* 75 (2016) 1450–1463, <http://dx.doi.org/10.1016/j.joms.2016.10.001>.
- [4] G. Bittemann, M.C. Metzger, S. Schlager, W.A. Lagrèze, N. Gross, C.P. Cornelius, R. Schmelzeisen, Orbital reconstruction: prefabricated implants, data transfer, and revision surgery, *Facial Plast. Surg.* 30 (2014) 554–560, <http://dx.doi.org/10.1055/s-0034-1395211>.
- [5] M. Maglione, Orbital floor reconstruction in facial asymmetry: a clinical case, *Trauma Cases Rev.* 1 (2017) 3–6, <http://dx.doi.org/10.23937/2469-5777/1510016>.
- [6] C. Jadhav, A.J. Hayes, S. Cunneen, D.A. Morrison, P.J. Meier, A novel individual reconstruction of a medial orbital wall blow-out fracture using a bone graft molded intraoperatively using a 3-D model: a case report, *Oral Maxillofac. Surg. Cases* 2 (2016) 19–21, <http://dx.doi.org/10.1016/j.omsc.2016.05.001>.
- [7] R.A. Agha, M.R. Borrelli, R. Farwana, K. Koshy, A. Fowler, D.P. Orgill, For the SCARE Group, The SCARE 2018 statement: updating consensus Surgical CARE RReport (SCARE) guidelines, *Int. J. Surg.* 60 (2018) 132–136.
- [8] N.R. Johnson, N.R. Singh, M. Oztef, V.N. Vangaveti, B.B. Rahmel, L. Ramalingam, Ophthalmological injuries associated with fractures of the orbitozygomaticomaxillary complex, *Br. J. Oral Maxillofac. Surg.* 56 (2018) 221–226, <http://dx.doi.org/10.1016/j.bjoms.2018.02.009>.
- [9] L.H. Hollier, J. Thornton, P. Pazmino, S. Stal, The management of orbitozygomatic fractures, *Plast. Reconstr. Surg.* 111 (2003) 2386–2392, <http://dx.doi.org/10.1097/01.PRS.0000061010.42215.23>.
- [10] P.N. Manson, B. Markowitz, S. Mirvis, M. Dunham, M. Yaremchuk, Toward CT-based facial fracture treatment, *Plast. Reconstr. Surg.* 85 (2) (1990) 202–212.
- [11] O. Lieger, R. Richards, M. Liu, T. Lloyd, Computer-assisted design and manufacture of implants in the late reconstruction of extensive orbital fractures, *Arch. Facial Plast. Surg.* 12 (2010) 186–191, <http://dx.doi.org/10.1001/archfacial.2010.26>.
- [12] R.J. McGalliard, J. Kimpton, N.M.H. McLeod, Ophthalmic outcomes of fractured zygomas, *Br. J. Oral Maxillofac. Surg.* 55 (2017) 363–366, <http://dx.doi.org/10.1016/j.bjoms.2016.10.026>.
- [13] L. Dubois, S.A. Steenen, P.J.J. Gooris, M.P. Mourits, A.G. Becking, Controversies in orbital reconstruction - II. Timing of post-traumatic orbital reconstruction: a systematic review, *Int. J. Oral Maxillofac. Surg.* 44 (2015) 433–440, <http://dx.doi.org/10.1016/j.ijom.2014.12.003>.
- [14] M. Yaremchuk, Orbital deformity after craniofacial fracture repair: avoidance and treatment, *J. Cranio-Maxillofac. Trauma* 5 (1999) 7–16.
- [15] S. Sukegawa, T. Kanno, Y. Koyama, A. Shibata, Intraoperative navigation-assisted surgical orbital floor reconstruction in orbital fracture treatment: a case report, *Shimane J. Med. Sci.* 33 (2017) 87–92.

Open Access

This article is published Open Access at sciedirect.com. It is distributed under the [IJSCR Supplemental terms and conditions](#), which permits unrestricted non commercial use, distribution, and reproduction in any medium, provided the original authors and source are credited.

Neglected Orbitozygomaticomaxillary Fractures With Complications: A Case Report

ORIGINALITY REPORT

18%

SIMILARITY INDEX

15%

INTERNET SOURCES

14%

PUBLICATIONS

8%

STUDENT PAPERS

PRIMARY SOURCES

1	explore.openaire.eu Internet Source	2%
2	apps.dtic.mil Internet Source	2%
3	gavinpublishers.com Internet Source	2%
4	www.ijhcr.com Internet Source	2%
5	Nathan Montoya Albrecht, Samuel Ostrower. "Case report of a laryngotracheal reconstruction with anterior and posterior costal cartilage graft and stent placement – Surgical technique", International Journal of Surgery Case Reports, 2019 Publication	1%
6	cemse.kaust.edu.sa Internet Source	1%
7	Xi Gong, Yang He, Jingang An, Yao Yang, Xiuling Huang, Meng Liu, Yangyang Zhao, Yi	1%

Zhang. "Application of a Computer-Assisted Navigation System (CANS) in the Delayed Treatment of Zygomatic Fractures: A Randomized Controlled Trial", Journal of Oral and Maxillofacial Surgery, 2017

Publication

8

"Corrigendum to "Hernia with spermatocele, cord and testis inside sac: Case report" [Int. J. Surg. Case Rep. 53 (2018) 397–399]", International Journal of Surgery Case Reports, 2019

Publication

1 %

9

"Smith and Nesi's Ophthalmic Plastic and Reconstructive Surgery", Springer Science and Business Media LLC, 2021

Publication

1 %

10

focusderma.com

Internet Source

1 %

11

Larry H. Hollier. "The Management of Orbitozygomatic Fractures", Plastic and Reconstructive Surgery, 06/2003

Publication

1 %

12

mafiadoc.com

Internet Source

1 %

13

Lieger, Olivier, Robin Richards, Mingjun Liu, and Tim Lloyd. "Computer-Assisted Design and Manufacture of Implants in the Late

1 %

Reconstruction of Extensive Orbital Fractures", Archives of Facial Plastic Surgery, 2010.

Publication

14 ejmcm.com 1 %
Internet Source

15 www.orl.fi 1 %
Internet Source

16 rsdjournal.org <1 %
Internet Source

17 M.C. Metzger, R. Schön, R. Tetzlaff, N. Weyer, A. Rafii, N.-C. Gellrich, R. Schmelzeisen. <1 %
"Topographical CT-data analysis of the human orbital floor", International Journal of Oral and Maxillofacial Surgery, 2007
Publication

18 Emanuel Mordechaev, Gabriel Shakarov, Deep Parikh. "Unilateral acute posterior multifocal placoid pigment epitheliopathy (APMPPE) with delayed contralateral eye involvement", Research Square Platform LLC, 2023 <1 %
Publication

19 Rodrigo Otávio Moreira Marinho, Belini Freire-Maia. "Management of Fractures of the Zygomaticomaxillary Complex", Oral and <1 %

Maxillofacial Surgery Clinics of North America, 2013

Publication

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

Neglected Orbitozygomaticomaxillary Fractures With Complications: A Case Report

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5
