

The Importance of Spirituality, physical Activity, and Sleep Duration to Prevent Hypertension among Elderly in Aceh-Indonesia

Sufyan Anwar, Loh Su Peng, Trias Mahmudiono

DOI: 10.31838/srp.2020.11.194



- > Baidu Scholar
- > Dimensions
- > Index Copernicus
- > JournalGuide
- > Microsoft Academic
- · ReadCuhe



Editorial Board

Editorial Board

FILIPIC Bratko, PhD

Professor, Department of Experimental and Translation Oncology

Croatian Institute for experimental and translation oncology, Zagreb, Croatia, Croatia

Dr Lucius, MBBS

General Practice

Ludwig Maximilians University Munich, Germany

Dr. Aygul Z. Ibatova

Department of Natural Sciences

Tyumen Industrial University, Scopus Author ID: 57191110632 http://orcid.org/0000-0003-0565-8533, Russia

Dr Ahmad Faisal Ismail

Kulliyyah of Dentistry

International Islamic University Malaysia, Kuantan Campus, 25200 Kuantan, Pahang, Scopus Author ID: 35388596700, Malaysia

Dr. Huiliang ZHAOÂ , Ph.D

Guizhou Minzu University, Huaxi District, Guiyang, China

Dr. Mohd Armi Abu Samah

International Islamic University Malaysia, (IIUM) 25200 Kuantan Pahang

Dr. Baded ramjiÂ

Sri Lanka

Dr. Chris randea

South Africa

Dr. Yingwen ZHAO

Researcher of Guizhou Rural Economic and Social Development Research Institute, China

Dr. Li Zihan, Ph.D

University of Glasgow, UK

Gabriela Cioca

Faculty of Medicine, Pharmacology Department

Lucian Blaga University of Sibiu, Romania, Lucian Blaga street, no 2A, Sibiu, Romania

Dariusz Nowak

Municipal Hospital, Mickiewicza street no 12, 42-200 Czestochowa, Poland

Aleksandra Zyska

Faculty of Medicine, Department of Physiology

Opole University, Oleska street no 48, 45-052 Opole, Poland

Katarzyna Sznajder

Faculty of Medicine, Clinical Department of Diagnostic Imaging

Opole University, Oleska street no 48, 45-052 Opole, Poland

Jacek Jó?wiak

Faculty of Medicine, Department of Family Medicine and Public Health

Opole University, Oleska street no 48, 45-052 Opole, Poland

Luciano Benedini

Universidad Nacional del Sur (National University of South-UNS), Bahía Blanca 8000, Argentina

Paula Messina

Departamento de Biolog

Universidad Nacional del Sur (National University of South-UNS), Bioquímica y Farmacia, Bahía Blanca 8000, Argentina

Michael Walsh

Washington State University, College of Pharmacy and Pharmaceutical Sciences (CPPS), USA

Prof. Dr. Kittisak Jermsittiparsert Â

Henan University, China

Amel Dawod Kamel Gudia, PhD

Faculty of nursing

Cairo University, Egypt

Arif Nur Muhammad Ansori

Airlangga University, Scopus Author ID: 57195995342, https://orcid.org/0000-0002-1279-3904, Indonesia

Mohammed Nader ShalabyÂ

Suez Canal University, Associate Professor of Biological Sciences and Sports Health, Egypt

Dr. Faten Abo-Aziza Mohamed, PhD

Associate Professor, Clinical Pathology and Stem Cell Research

National Research Centre, Manager of Veterinary Division Central Lab (605), 33 El-Behoos St, Dokki, Cairo, Egypt

Professor Asim Ahmed Elnour Ahmed

College of Pharmacy

Al-Ain University of Science and Technology, UAE

S. Parasuraman, M.Pharm., Ph.D

AIMST University, Malaysia

Ebenezer Wiafe, PhD

Pharmacy

University of Kwazulu-Natal, South Africa

Editor-in-Chief

Dr.Ayad F. Alkaim

University of Babylon, College of Science for Women, Babylon, Scopus Author ID: 55255310600, Iraq

☑ Submit Article

Login

Register

Most Viewed Articles


Dental Development between Assisted Reproductive Therapy (Art) and Natural Conceived Children: A Comparative Pilot Study Norzaiti Mohd Kenali, Naimah Hasanah Mohd Fathil, Norbasyirah Bohari, Ahmad Faisal Ismail, Roszaman Ramli SRP. 2020; 11(1): 01-06 » doi: 10.5530/srp.2020.1.01

Psychometric properties of the World Health Organization Quality of life instrument, short form: Validity in the Vietnamese healthcare context Trung Quang Vo*, Bao Tran Thuy Tran, Ngan Thuy Nguyen, Tram ThiHuyen Nguyen, Thuy Phan Chung Tran SRP. 2020; 11(1): 14-22 » doi: 10.5530/srp.2019.1.3

A Review of Pharmacoeconomics: the key to "Healthcare for All" Hasamnis AA, Patil SS, Shaik Imam, Narendiran K SRP. 2019; 10(1): s40-s42 » doi: 10.5530/srp.2019.1s.21

Deuterium Depleted Water as an Adjuvant in Treatment of Cancer Anton Syroeshkin, Olga Levitskaya, Elena Uspenskaya, Tatiana Pleteneva, Daria Romaykina, Daria Ermakova SRP. 2019; 10(1): 112-117 » doi: 10.5530/srp.2019.1.19

- > Baidu Scholar
- > Dimensions
- > Index Copernicus
- > JournalGuide
- > Microsoft Academic
- · ReadCuhe



2020: Volume 11, Issue 11

Research Article

<u>Gadget Use, Pocket Money, and Snacking Habits of Children with and without Overweight/Obesity</u>
<u>Problem in Surabaya, Indonesia</u>

Lirias Mahmudiono, Qonita Rachmah, Diah Indriani, Triska Susila Nindya, Calista Segalita, Susi Hidayah, Azizah Ajeng Pratiwi, Loh Su Peng

SRP. 2020; 11(11): 1087 - 1090

» Abstract » PDF DOI: 10.31838/srp.2020.11.155

Review Article

Ameliorative Effect Of Thiamine Pyrophosphate Against Cisplatin-Induced Reproductive System

<u>Damage Within Male Rats</u>

▲ Maitham Abd Ali Mnati, Bahir Abdul Razzaq Mshimesh, Mustafa Mohammed Ibraheem, Suzan Yousif Jasim SRP. 2020; 11(11): 396 - 403

» Abstract » PDF DOI: 10.31838/srp.2020.11.59

Research Article

Neutrophil Extracellular Traps in Coronavirus Infection: Interaction Network Analysis

📤 Amal Bouzid, Asmaa T Uthman, Noor N Al-Rawi, Natheer H Al-Rawi

SRP. 2020; 11(11): 1091 - 1101

» Abstract » PDF DOI: 10.31838/srp.2020.11.156

Review Article

<u>An Observational Study Of Coronavirus (Covid-19) In Iraqi Patients At Al-Shifa Medical Center In Baghdad's Capital, Al-Rusafa</u>

Saad Abdul Kareem Mohammed, Suha Hussein Ahmed, Ali I. Omran Al- Saadawi, Mohammed Mahmood Mohammed

SRP. 2020; 11(11): 404 - 411

» Abstract » PDF DOI: 10.31838/srp.2020.11.60

Review Article

<u>Pathogenic Microorganisms, Toxigenic Fungi, Heavy Metal accumulation, and Toxic Materials</u>
<u>Contamination in Natural Products</u>

A Huda S. Husni

SRP. 2020; 11(11): 1772 - 1776

» Abstract » PDF DOI: 10.31838/srp.2020.11.247

Review Article

<u>Immunophenotypic characterization of malignant lymphoma in Iraqi patients using immunohistochemical CD-marker study</u>

🖺 Hameda Abd Al-Mahdi Ghazi, Rajaa Ali Moheiseen Al-Taee, Hayder Abdul-Amir Makki Al-Hindy

SRP. 2020; 11(11): 412 - 417

» Abstract » PDF DOI: 10.31838/srp.2020.11.61

Review Article

<u>The Therapeutic Effect Of Eucalyptus Microtheca Against The Giardiasis Induced Renal Damage In Male Rats</u>

📤 Abdulrazzaq Mohammed Hammood, Muhannad Shweash, and Hazim Ghazzay

SRP. 2020; 11(11): 418 - 422

» Abstract » PDF DOI: 10.31838/srp.2020.11.62

Research Article

The Determinant of Exclusive Breastfeeding among Female Worker in Indonesia

L Ina Kusrini, Mara Ipa, Agung Dwi Laksono, Noviati Fuada, Sri Supadmi

SRP. 2020; 11(11): 1102 - 1106

» Abstract » PDF DOI: 10.31838/srp.2020.11.157

Review Article

Effect of COVID-19 on Bacterial Resistance

A Hussam H. Tizkam, Osama Q. Fadhil, Esraa Ghazy

SRP. 2020; 11(11): 423 - 427

» Abstract » PDF DOI: 10.31838/srp.2020.11.63

Research Article

<u>Next-Generation Sequencing Technologies for Environmental DNA as an Efficient Bio Indicator for Bacterial Biodiversity in Tigris River, Iraq</u>

🆺 Fikrat M Hassan, Warqaa Y. Salih and Halah H Al-Haideri

SRP. 2020; 11(11): 1107 - 1114

» Abstract » PDF DOI: 10.31838/srp.2020.11.158

Review Article

Analysis of the Prophet Saleh in Al Quran through the food Science Approach to the Miracle of She-Camel

💄 Endin Mujahidin, Nur Richana, Adian Husaini, Didin Hafidhuddin

SRP. 2020; 11(11): 771 - 775

» Abstract » PDF DOI: 10.31838/srp.2020.11.111

Review Article

<u>The Relationship of Family Support to the Success of the Treatment of Tuberculosis (TB) Patients in Kediri</u>

🆀 Fauzan Adima, Chatarina U. Wahjuni, Hari Basuki Notobroto, Shrimarti Rukmini Devy

SRP. 2020; 11(11): 776 - 778

» Abstract » PDF DOI: 10.31838/srp.2020.11.112

Research Article

<u>Levels of Myeloperoxidase, Malondialdehyde and Lipid Profile in Type 2 Diabetic Patients on Metformin Versus Glibenclamide Therapy</u>

📤 Zainab H. Fathi, Jehan A. Mohammad, Marwah H. Mohammed

SRP. 2020; 11(11): 1777 - 1782

» Abstract » PDF DOI: 10.31838/srp.2020.11.248

Research Article

Recent Development In Oxine Complexes And Their Medical Application: A Review

📤 Shatha M . H. Obaid, Waleed K . Mahdi, Falah H. Hussein, Yahya F. Al-Khafaji

SRP. 2020; 11(11): 428 - 437

» Abstract » PDF DOI: 10.31838/srp.2020.11.64

Review Article

<u>Cytotoxic Activity of Ethanol Extract Legundi Leaf (Vitex trifolia L.) and N-Hexan, Ethyl Acetate and Ethanol-Water Fraction against MCF-7 Breast Cancer Cells</u>

📤 Peni Indrayudha, Indah Gita Cahyani

SRP. 2020; 11(11): 779 - 785

» Abstract » PDF DOI: 10.31838/srp.2020.11.113

Review Article

<u>Detecting Mycobacterium Tuberculosis through Bacterioferritin Gene Amplification by Real-Time</u>
<u>PCR</u>

🆺 Raed Obaid Saleh, Nagham Maad Hamdi, Rana Kadhim Mohammed.

SRP. 2020; 11(11): 786 - 789

» Abstract » PDF DOI: 10.31838/srp.2020.11.114

Review Article

MIC of Some Plant Extracts against E. Coli Isolated from UTI

A Ibrahim Mohammed Saeed Hussein, Jalank Hameed Mahmoud, Halala Mohammed Abdulla

SRP. 2020; 11(11): 790 - 792

» Abstract » PDF DOI: 10.31838/srp.2020.11.115

Review Article

Acute Coronary Syndrome in Patients with Oncological Diseases

📤 N.A. Shanazarov, N.K. Seidalin, A.F. Khalirahmanov, G.D. Gatiyatullina, S.V. Zinchenko, K. F. Idrisova

» Abstract » PDF DOI: 10.31838/srp.2020.11.159

Review Article

<u>Effects of a Cellular Camel Urinary Bladder Sub Mucosa on Chronic Full Thickness Skin Wound</u> <u>Healing in Equine Species</u>

Ahmed Kassem, Ayad N.D. Alhaki, Abd-Alhadi Jaithom Marzok.

SRP. 2020; 11(11): 793 - 797

SRP. 2020; 11(11): 1115 - 1120

» Abstract » PDF DOI: 10.31838/srp.2020.11.116

Review Article

<u>Chromium (III) Ions Uptake from Aqueous Solutions by Chelating Resin Containing Pendant Multidentate Ligand</u>

Ahmed Dhary Saleh

SRP. 2020; 11(11): 798 - 804

» Abstract » PDF DOI: 10.31838/srp.2020.11.117

Review Article

<u>Stunting Prevention Through Participative Counselling on The Implementation of Balanced</u>
<u>Nutrition Toward Children by Involving Local Puppeteers in Central Lombok Regency, West Nusa Tenggara</u>

A Made Darawati, Andi Eka Yunianto, Ni Ketut Sri Sulendri, Omdah

SRP. 2020; 11(11): 805 - 810

» Abstract » PDF DOI: 10.31838/srp.2020.11.118

Research Article

<u>Phytochemical Screening and In Vivo Hepatoprotective Activity of the Cordia Myxa L Fruit Extracts</u>
<u>Against Paracetamol-Induced Hepatotoxicity in Rats</u>

📤 Ahmed Mohammed Hassan Murad and Majed H. Karbon

SRP. 2020; 11(11): 1782 - 1794

» Abstract » PDF DOI: 10.31838/srp.2020.11.249

Research Article

Role of Pseudomonas Fluorescens and Organic Matter in Controlling of Potato Black Scurf Disease Caused by Rhizoctonia Solani

📤 Ali A. Kadhum, Jasim M. Abed and Theyab A. Farhan

SRP. 2020; 11(11): 1121 - 1125

» Abstract » PDF DOI: 10.31838/srp.2020.11.160

Review Article

THE ROLE OF THE SUPREME COURT OF THE REPUBLIC OF INDONESIA IN REGULATING E-COURT DURING THE NEW

Muhammad Syarifuddin.

SRP. 2020; 11(11): 811 - 817

» Abstract » PDF DOI: 10.31838/srp.2020.11.119

Research Article

Breast Cancer in Patients with Gene Polymorphisms Associated with Metabolic Syndrome

🚨 N. K. Seidalin, V. V. Benberin, N. A. Shanazarov, Voshchenkova T. A., Aripzhanova G. O., Zhapparov E. I.

SRP. 2020; 11(11): 1126 - 1129

» Abstract » PDF DOI: 10.31838/srp.2020.11.161

Review Article

CONTRIBUTION OF FORENSIC LABORATORY TO SCIENTIFIC CRIME INVESTIGATION IN INVESTIGATING LEGAL CASE OF DRUG ABUSE: A SYSTEMATIC REVIEW

Rudy Heriyanto Adi Nugroho

SRP. 2020; 11(11): 818 - 823

» Abstract » PDF DOI: 10.31838/srp.2020.11.120

Review Article

SETTLEMENT OF TAX PAYABLE IN BANKRUPTCY CASE: A SYSTEMATIC REVIEW

4 Yeheskiel Minggus Tiranda

SRP. 2020; 11(11): 824 - 829

» Abstract » PDF DOI: 10.31838/srp.2020.11.121

Research Article

In Vitro Stiumlation of Ergosterol from Coelastrella Terrestris by Using Squalene and Studying Antioxidant Effect

Altaf AL-Rawi, Fikrat M. Hassan and Bushra M.J.Alwash

SRP. 2020; 11(11): 1795 - 1803

» Abstract » PDF DOI: 10.31838/srp.2020.11.250

Review Article

A SYSTEMATIC REVIEW OF THE ROLE OF APPLIED AND PURE SCIENCES IN THE DEVELOPMENT OF POLICE SCIENCE IN INDONESIA

A Chairuddin Ismail

SRP. 2020; 11(11): 830 - 836

» Abstract » PDF DOI: 10.31838/srp.2020.11.122

Review Article

<u>In Vitro and in Silico Analysis on the Bone Formation Activity of N-Hexane Fraction of Semanggi</u> (<u>Marsilea crenata Presl.</u>)

📤 Agnis Pondineka Ria Aditama, Burhan Ma'arif, Denis Mery Mirza, Hening Laswati, Mangestuti Agil.

SRP. 2020; 11(11): 837 - 849

» Abstract » PDF DOI: 10.31838/srp.2020.11.123

Review Article

Biochemistry and Molecular Cell Biology of Diabetic Complications

Athmar Jawad Ahmed, Suhad Rasheed Majeed, Hussein Mohammed Obaid.

SRP. 2020; 11(11): 850 - 860

» Abstract » PDF DOI: 10.31838/srp.2020.11.124

Review Article

<u>Pedagogical Conditions for Ensuring the Formation of Information and Communication</u> <u>Competences of Higher Education Applicants</u>

A Petro Boichuk, Vitalii Honcharuk, Olena Redko, Nataliya Dziubyshyna, Olena Vlasiuk, Valentuna Tiuska

SRP. 2020; 11(11): 1130 - 1134

» Abstract » PDF DOI: 10.31838/srp.2020.11.162

Review Article

The Use of Information Technology in the Invariant System of Presentation of Learning Material in the Learning Process

🏝 Nataliia Tytova, Volodymyr Bogoliubov, Vasyl Yefymenko, Lesia Makarenko, Liudmyla Mova, Natalia Kalientsova.

SRP. 2020; 11(11): 861 - 867

» Abstract » PDF DOI: 10.31838/srp.2020.11.125

Research Article

Relationship between Lumbar Herniated Disc with Flat Feet

A Haider Mekhlif Ali

SRP. 2020; 11(11): 1804 - 1806

» Abstract » PDF DOI: 10.31838/srp.2020.11.251

Review Article

<u>Analysis of the Organization and Features of the Implementation of Information Technologies in the Educational Process of Institutions of Higher Education</u>

📤 Ihor Bondar, Nadiia Bachynska, Tetiana Novalska, Vladyslav Kasian, Valerii Kuchnarov, Volodymyr Pylypiv.

SRP. 2020; 11(11): 868 - 872

» Abstract » PDF DOI: 10.31838/srp.2020.11.126

Review Article

<u>Acute Coronary Syndrome During Chemotherapy</u>

🚨 N. A. Shanazarov, V. V. Benberin, N.K. Seidalin, A.F. Khalirakhmanov, E.A. Gaziyev, S.V. Zinchenko

SRP. 2020; 11(11): 1135 - 1139

» Abstract » PDF DOI: 10.31838/srp.2020.11.163

Review Article

<u>The Ethnopsychological Peculiarities of Europeans and Russians as a Factor of Different Cultures</u>
<u>Representatives Interaction and Mutual Understanding Effectiveness</u>

🚨 Tatiana Malkova, Iryna Synhaivska, Hanna Abanina, Olga Vasilchenko, Olga Petrunko, Serhii Vitkalov.

SRP. 2020; 11(11): 873 - 879

» Abstract » PDF DOI: 10.31838/srp.2020.11.127

The Importance Of Spirituality, Physical Activity and Sleep Duration to Prevent Hypertension among Elderly in Aceh-Indonesia

Sufyan Anwar¹, Loh Su Peng^{2,3}, Trias Mahmudiono^{3*}

ABSTRACT

Introduction: Hypertension is a condition that is common in the elderly population. Unlike physical activity and sleep duration, only a small proportion of studies have assessed the association of spirituality with hypertension.

Aims: This study was aimed at looking at the relationship between spirituality, physical activity, and sleep duration with hypertension.

Methods: This research is a cross-sectional study and data collection was conducted through a survey of 483 elderly people aged 60 years or over in 2019 in Aceh, Indonesia. To answer the research hypothesis, we used a logistic regression test.

Results: Although we found an association between spirituality (p=0.007), physical activity (p<0.001) and sleep duration (p=0.006) and hypertension, we did not find an association between coffee drinking habits (p=0.204) and hypertension in the elderly. in Aceh. The study results showed that physical activity had a stronger relationship with hypertension

Conclusions: This study concludes that spirituality, physical activity, and sleep duration can prevent or reduce hypertension in the elderly in Aceh.

Keywords: Elderly, Hypertension, Physical Activity, Spirituality, Sleep Duration

Correspondence

Trias Mahmudiono

Department of Nutrition, Faculty of Public Health Universitas Airlangga,

Surabaya, East Java, Indonesia **Email**: trias-m@fkm.unair.ac.id

INTRODUCTION

It was recognized that hypertension is a risk factor for heart and blood vessel disease. It is a trigger for heart failure, stroke and kidney disease, and even death (1). It is estimated that 9.4 people die every year in the world as a result of complications rather than the number of hypertension (2). Globally in 2015, it is estimated that the adult population suffering from hypertension will be 1.13 billion and this number has increased compared to 1975 which only amounted to 594 million people (3). Most of them or two-thirds live in low and middle-income countries (2).

It is common knowledge that hypertension often occurs in the elderly. Age is considered as a risk factor for hypertension. The longer a person lives, the higher the chances of developing hypertension. Until now, hypertension is still a big challenge in the world of health This study aims to assess the relationship between spirituality, physical activity, and sleep duration with hypertension in the elderly in Aceh Province, Indonesia. We suspect that spirituality, physical activity, and sleep duration have a specific role in preventing hypertension and lowering blood pressure.

MATERIALS AND METHODS

Study Design

This study was conducted using a cross-sectional approach and a population of 60 years or more. The sample size required was based on the rule of thumb formula, where the sample was accounted for 5-10 times the indicator (observed variables), namely 43 indicators. Hence the minimum sample size was 430 respondents. The actual sample size taken was 483 respondents. The inclusion criteria for partisipants were: able to communicate, signing a statement of consent to be a respondent, and were not severely ill or admitted to a health service at the time of the study.

in Indonesia (4). Hypertension cases among elderly people in Indonesia are also increasing. Based on the results of the Basic Health Research in 2013 the incidence of hypertension in the elderly population was 57.6% and the results of Riskesdas 2018 hypertension in the elderly population was 63.5% (5). There is an increase in the incidence of hypertension in the elderly by 5.6% over five years. The Institute for Health Metrics and Evaluation (IHME) in 2017 stated that hypertension was a risk factor for the cause of death (23.7%) of a total of 1.7 million deaths in Indonesia in 2016 (5). To detect the incidence of hypertension, the Indonesian government through the ministry of health determines the detection/examination of blood pressure in the elderly population as an indicator of health services for the elderly. Besides, although drugs that are effective for hypertension are widely available, control of hypertension has not been adequate (6).

Measure And Instrument

Four independent variables were measured, namely spiritual well-being, physical activity, coffee drinking habits, and sleep duration. Assessment of spiritual wellbeing used was the spirituality orientation inventory previously validated (7). Likert scale was used for assessment, namely strongly disagree, disagree, agree, and strongly agree. The higher the score obtained, the better the respondent's spiritual welfare. Measurement of physical activity using questions about the activity carried out at least 30 minutes. Physical activity was divided into four groups, including no activity, light, moderate, and heavy physical activity. The form of each physical activity refered to the guidelines from the Ministry of Health of the Republic of Indonesia. Strenuous physical activity was activities that expend energy> 7 Kcal / minute (for example lifting / carrying wood, rice, sandstone; hoeing; cutting trees; fast cycling, single tennis/badminton, sprinting, marathon, pedaling pedicabs, climbing mountains. and cycling with loads. Moderate physical

¹Doctoral of Public Health Study Program, Faculty of Public Health Universitas Airlangga, Surabaya, East Java, Indonesia.

²Department of Nutrition, Faculty of Medicine and Health Science, Universiti Putra Malaysia, Selangor, Malaysia

³Department of Nutrition, Faculty of Public Health Universitas Airlangga, Surabaya, East Java, Indonesia

^{*}Author for Correspondence: Trias Mahmudiono Email: trias-m@fkm.unair.ac.id

activity was an activity that emits energy from 3.5 to 7 Kcal/minute (for example: sweeping the yard, mopping the floor, washing clothes, drawing water, farming, cleaning the bathroom, jogging, double badminton, swimming. Mild physical activity was an activity that releases energy < 3.5 Kcal/minute (for example: walking at home or shopping center, reading, writing, painting, cooking, fishing, washing dishes and ironing. Coffee drinking habit was measured using "yes" or "no" question. The sleep duration assessment also used an open question such as: "how many hours do you sleep on average at night?". Meanwhile, the measurement of hypertension, which was the dependent variable, was carried out by measuring systolic blood pressure. The tool used was the digital tension meter OMRON HEM-8712. The implementation was following the Posbindu technical guidelines, where measurements were taken twice with an

The presentation of demographic characteristics as seen in Table 1 showed that the average age of the respondents was 69.76 years and more in the 60-69 age group (56.5%). More respondents were women (68.9%) than men and did not work (58.2%) than those who worked. The results of step 1 logistic regression analysis (Table 2) showed that there were three variables associated with hypertension which were: spiritual well-being (p=0.007), physical activity (p=<0.001), and sleep duration (p=0.006). The

interval of 1-2 minutes. If the results of the first and second measurements was differed of >10 mmHg, then the third measurement was performed after resting for 10 minutes by removing the cuff on the arm (6).

Data Analysis

A descriptive analysis was conducted to see the demographic characteristics of the respondents. Logistic regression analysis was performed using SPSS version 21 to determine the correlation between spiritual well-being, physical activity, coffee drinking habits, and sleep duration with hypertension.

Ethical Approval

Ethical approval for this study was obtained from the health research ethics committee of the Faculty of Nursing, Airlangga University on December 18, 2018, No. 1229-KEPK.

habit of drinking coffee did not show a correlation with hypertension (p=0.204). Furthermore, the second stage logistic regression test was carried out by removing the habit of drinking coffee (Table 3). After the coffee drinking habit was excluded from the analysis, it was found that all variables had a significant value related to hypertension (spiritual well-being (p=0.007), physical activity (p=<0.001) and sleep duration (p=0.006). The three variables showed physical activity had the strongest relationship with hypertension (Figure 1 and Figure 2).

Table 1: Demography Characteristics of Respondents

Respondents Characteristics (n=483)	n	Mean/%	SD
Age		69.76	8.41
Age 60-69	273	56.5	
Age ≥ 70	210	43.5	
Gender			
Male	150	31.1	
Women	333	68.9	
Employment			
Not Working	281	58.2	
Farmer/ Laborer/Fisher	133	27.5	
Pensioners	24	5.0	
Private Employee	45	9.3	
Income		520559.01	856194.59
No Income	198	41.0	
IDR < 1,000,000, -	184	38.1	
IDR 1,000,000, - s.d <2,000,000,-	50	12.4	
IDR ≥ 2,000,000, -	41	8.5	

Table 2: Logistic Regression Analysis Stage 1

Variables	В	SE	Wald	p-value	Adj.	95%CI	
	Б				OR	Lower	Upper
Spiritual well-being	060	.022	7.244	0.007*	.941	.901	.984
Physical Activity	594	.145	16.881	<0.001*	.552	.416	.733
Coffee drinker	246	.194	1.614	0.204	.782	.535	1.143
Sleep duration	519	.190	7.457	0.006*	.595	.410	.864
Constant	3.513	.983	12.764	< 0.001	33.557		

Abbreviations: B = Beta coefficient, SE = Standard Error, CI = Confidence Interval, Adj OR = Adjusted odds ratio, * = significant p-value

Table 3: Logistic Regression Analysis Stage 2

Variables	D	SE	Wald	<i>p</i> -value	Adj. OR	95%	%CI
variables	В	SE	waiu			Lower	Upper
Spiritual well-being	061	.022	7.293	0.007*	.941	.901	.984
Physical Activity	584	.144	16.430	<0.001*	.558	.420	.740

Variables	D	SE	Wald	<i>p</i> -value	Adj. OR	959	%CI
variables	ь	3E	waiu	p-value		Lower	Upper
Sleep duration	523	.190	7.591	0.006*	.593	.409	.860
Constant	3.413	.977	12.201	< 0.001	30.351		

Abbreviations: B = Beta coefficient, SE = Standard Error, CI = Confidence Interval, Adj OR = Adjusted odds ratio, * = significant p-value




Figure 1: Scatter Plot of Systolic Blood Pressure and Spiritual Well-Being




Figure 2: Scatter Plot of Systolic Blood Pressure and Sleep Duration

DISCUSSION

Our study revealed significant association between spirituality and good management of hypertension. This finding alligned with many studies that showed a significant relationship between spirituality and clinical/laboratory results. Another study showed minor blood pressure reactivity in subjects with greater religiousness, suggesting that religious beliefs can be an important variable for the study of patients with arterial hypertension, particularly the elderly. In addition to spirituality, the act of praying can add optimism to the process of addressing chronic diseases. The practice of

praying was beneficial in several ways because, in addition to requests to God, prayers were often made to thank life, health and family, which results in greater feelings of gratitude.

Religious beliefs and practices can reduce loss of control, stress, and a sensation of helplessness, allowing the cognitive structure reduce suffering and making treatment more effective (8). The results showed that spirituality, religiousness, and faith have a positive influence whilst patients cope with life obstacles and hardships, in addition to increasing patients' resilience, thus, improving their condition. Spirituality can be used as a strategy to tackle

critical life circumstances, because it can increase one's sense of purpose and life meaning, associated with a greater resistance to stress-related diseases (9).

The result showed that physical activity had significant relationship with hypertansion. Hypertension affects more than 40% of adults worldwide and is strongly associated with coronary artery disease, stroke, and heart failure. One of the ways to reduce hypertension is by doing physical activity or exercise. Exercise is a one of the important things of life-style therapy for the number one prevention of high blood pressure. A range of research constantly revealed useful outcomes of exercising on high blood pressure with discounts in each systolic and diastolic blood stress with as plenty as 5-7 mmHg reductions in people with high blood pressure (10). Exercise has been related to on-the-spot huge reductions in systolic blood pressure. This on-the-spot reduction in blood stress after exercising can persist for nearly 24 hours and is known as post-exercising hypotension with the maximum reported outcomes visible in people with better baseline blood pressure (11).

The reduction in blood pressure was associated with the attenuation in peripheral vascular resistance, which can be because of neuro hormonal and structural responses with reductions in sympathetic nerve interest and a boom in arterial lumen diameters, respectively (12). The mechanisms underlying blood stress discount with exercising and its related results are nevertheless under research with much research confined with the aid of using length and marked heterogeneity (13). In fact, in comparison to men and women 25-forty-four years of age, mortality chance amongst older adults is increased with the aid of using 100-fold for stroke and continual lung disease, more or less 90-fold for coronary heart disease, pneumonia and influenza, and over 40-fold for cancer. Hypertension in the elderly patients represents a management dilemma to cardiovascular (CV) specialists and another practitioner. Furthermore, with the wide adoption of multiple drug strategies targeting subgroups of hypertensive patients with specific risk conditions to lower blood pressure (BP) beyond traditional goals, difficult questions arise about how aggressive elderly patients should be treated.

The latest data from the Global Burden of Disease Study in 2015 showed that high systolic blood pressure continues to be the largest contributor to global disability-adjusted life-years (DALYs), causing 211 million global DALYs each year (14). According to a previous study, hypertension was a major risk factor for cardiovascular disease (CVD), heart attack, heart failure, stroke, and kidney disease. Studies have shown that sleep duration was associated with hypertension (15). Short sleep duration, usually defined as less than 7 hours, 6 hours or 5 hours per night, was associated with an increased risk of prevalent hypertension (15).

The relationship between self-reported sleep duration and hypertension was first reported as a U-shaped association in the Sleep Heart Health Study (SHHS). In the study, Gottlieb et al found that sleep duration of less than 7 hours per night or more than 8 hours per night was associated with an increased prevalence of hypertension (16). The results from the NHANES also demonstrated an association between sleep duration of less than 5 hours per night and an increased risk of hypertension in the same year (17).

One of the cross-sectional studies concluded that they observed an association between short sleep durations

(<7 hours/day) and an increased risk of hypertension in young adults (18-44 years) (18). This association was attenuated by the inclusion in the multivariate models of socio-demographic covariates—BMI and lifestyle factors. Compared with the young adults, an association between short sleep duration and hypertension was not found for middle-aged participants (45–59 years), old participants (60–79 years) or the total sample (18). Furthermore, no association between sleep duration and hypertension was found when male or female participants were analyzed separately (18). A significant result shows that sleep duration and hypertension is having relationship with age especially in elderly. Compared with sleep durations of 7 to < 8 hours per night, self-reported usual sleep durations of less than 7 hours per night or 8 or more hours per night were associated with higher adjusted odds ratio (OR) for hypertension (19).

They suggest that levels of habitual sleep restriction that are common in the adult population may contribute to the high population prevalence of hypertension. Lusardi and colleagues found that a single night of experimental sleep restriction to 4 hours of sleep in the home setting resulted in a 4- to 7-mm Hg increase in mean morning SBP in normotensive and hypertensive subjects, respectively, with smaller increases in mean morning DBP (20).

CONCLUSION

In conclusion, spirituality, physical activity, and sleep duration can prevent or reduce hypertension among elderly in Aceh. This study showed that spirituality have a positive influence to cope with life obstacles and hardships, in addition to increasing patients' resilience, thus improving their condition.

ACKNOWLEDGEMENT

The authors would like to thank all participants of elderly in Aceh, and also lecturers in Doctoral Study Program of Public Health, Faculty of Public Health Universitas Airlangga for their guidance, suggestions and input so that the writing of this paper can be completed.

REFERENCES

- Chobanian, A. V., Bakris, G. L., Black, H. R., Cushman, W. C., Green, L. A., Izzo, J. L., Jr., et al. Seventh report of the joint national committee on prevention, detection, evaluation, and treatment of high blood pressure. Hypertension. 2003; 42: 1206–125.
- 2. WHO. A global brief on hypertension. Geneva; 2013. Available from: https://apps.who.int/iris/bitstream/handle/10665/7 9059/WHO_DCO_WHD_2013.2_eng.pdf;jsessionid=E4 90DD6C72928BBC5E5FE804A16D2F2B?sequence=1
- 3. NCD Risk Factor Collaboration (NCD-RisC). Worldwide trends in blood pressure from 1975 to 2015: a pooled analysis of 1479 population-based measurement studies with 19•1 million participants. Lancet (London, England). 2017; 389(10064): 37–55. https://doi.org/10.1016/S0140-6736(16)31919-5.
- 4. Kementerian Kesehatan RI. Profil Kesehatan Indonesia Tahun 2018. Jakarta; Kementerian Kesehatan RI: 2019. Available from: https://www.kemkes.go.id/resources/download/pusdatin/profil-kesehatan-indonesia/PROFIL_KESEHATAN_2018_1.pdf

- Kementerian Kesehatan RI. Hari hipertensi dunia 2019: "Know your number, kendalikan tekanan darahmu dengan CERDIK". 2019. Accessed in November 21st 2019 at http://p2ptm.kemkes.go.id/kegiatan-p2ptm/pusat-/hari-hipertensi-dunia-2019.
- Kementerian Kesehatan RI. Profil Kesehatan Indonesia 2016. Jakarta; Kementerian Kesehatan RI: 2017. Available from: https://www.kemkes.go.id/resources/download/pusdatin/profil-kesehatan-indonesia/Profil-Kesehatan-Indonesia-tahun-2017.pdf
- 7. Wahyuningsih H. Validitas konstruk alat ukur spirituality orientation inventory (SOI). Jurnal Psikologi. 2009; 36 (2): 116-129.
- 8. Peres, J. F. P., Newberg, A. B., Mercante, J. P., Sima o, M., Albuquerque, V. E., Peres, M. J. P., &Nasello, A. G. Cerebral blood flow changes during retrieval of traumatic memoriesbefore and after psychotherapy: A SPECT study. Psychological Medicine, in press. 2007.
- Elkins, D. N., Hedstrom, L. J., Hughes, L. L., Leaf, J. A., & Saunders, C. Toward a humanistic-phenomenological spirituality: Definition, description, and measurement. Journal of Humanistic Psychology. 1988; 28(4): 5–18
- 10.Díaz, A. Intercultural understanding and professional learning through critical engagement. Babel. 2013; 48(1): 12-19.
- 11.Pestacello LS, MacDonald HV, Lamberti L, Johnson BT. Exercise for hypertension: A prescription update integrating existing recommendations with emerging research. Curr Hypertens Rep. 2015; 17: 7
- 12. Hamer, Mark & Taylor, Adrian & Steptoe, Andrew. The effect of acute aerobic exercise on stress related blood pressure responses: A systematic review and meta-analysis. Biological psychology. 2006; 71: 183-90.
- 13.Semlitsch, T., Jeitler, K., Hemkens, L G., Horvath, K., Nagele, E., Schuermann, C., Siebenhofer, A. Increasing Physical Activity for The Treatment of Hypertension: A Systematic Review and meta-analysis. Sports Medicine. 2013; 43(10): 1009-1023
- 14.WHO. GLOBAL HEALTH RISKS: Mortality and burden of disease attributable to selected major risks. Geneva; 2009. Available from: https://www.who.int/healthinfo/global burden disease/GlobalHealthRisks report full.pdf
- 15.Grandner M, Mullington JM, Hashmi SD, Redeker NS, Watson NF, Morgenthaler TI. Sleep Duration and Hypertension: Analysis of > 700,000 Adults by Age and Sex. Journal of Clinical Sleep Medicine. 2018 June 15; 14(6): 1031-1039.
- 16.Gottlieb DJ, Redline S, Nieto FJ, Baldwin CM, Newman AB, Resnick HE, et al. Association of usual sleep duration with hypertension: The Sleep Heart Health Study. Sleep 2006; 29(8):1009–1014.
- 17.Gangwisch JE, Heymsfield SB, Boden-Albala B, Buijs RM, et al. Short Sleep Duration as a Risk Factor for Hypertension: Analyses of the First National Health and Nutrition Examination Survey. Hypertension 2006; 47(5): 833-839.
- 18.Li M, Yan S, Jiang S, Ma X, Gao T, Li B. Relationship between sleep duration and hypertension in northeast China: a cross-sectional study. BMJ Open 2019;9.
- 19. Hall MH, Muldoon MF, Jennings JR, Buysse DJ, Flory JD, Manuck SB, et al. Self-Reported Sleep Duration is Associated with the Metabolic Syndrome in Midlife Adults. Sleep 2008; 31(5): 635-643

20.Lusardi P, Zoppi A, Preti P, Pesce RM, Piazza E, Fogari R. Effects of insufficient sleep on blood pressure in hypertensive patients: a 24-hour study. Am J Hypertens 1999; 12:63-8.