

The clusters of health-risk behaviours and mental wellbeing and their sociodemographic correlates: a study of 15,366 ASEAN university students

by Trias Mahmudiono

Submission date: 09-Mar-2023 11:32AM (UTC+0800)

Submission ID: 2032663947

File name: A3.pdf (938.77K)

Word count: 9904

Character count: 51727

RESEARCH

Open Access

6

The clusters of health-risk behaviours and mental wellbeing and their sociodemographic correlates: a study of 15,366 ASEAN university students



Apichai Wattanapisit^{1,2}, Hanif Abdul Rahman^{3,4}, Josip Car⁵, Khadzah Haji Abdul-Mumin^{3,6}, Ma. Henrietta Teresa O. de la Cruz⁷, Michael Chia⁸, Michael Rosenberg^{9,10}, Moon-ho Ringo Ho¹¹, Surasak Chaiyasong¹², Trias Mahmudiono¹³, Yuvadee Rodjarkpai¹⁴, Ivo D. Dinov⁴, Mohammad Ottom^{4,15} and Areekul Amornsriwatanakul^{9,10*}

10

Abstract

Background: This study investigated, through cluster analysis, the associations between behavioural characteristics, mental wellbeing, demographic characteristics, and health among university students in the Association of Southeast Asian Nations (ASEAN) University Network – Health Promotion Network (AUN-HPN) member universities.

Methods: Data were retrieved from a cross-sectional self-administered online survey among undergraduate students in seven ASEAN countries. A two-step cluster analysis was employed, with cluster labels based on the predominant characteristics identified within the clusters. The 'healthy' cluster was assigned as the reference group for comparisons using multinomial logistic regression analysis.

Results: The analytic sample size comprised 15,366 university students. Five clusters of student-types were identified: (i) 'Healthy' ($n = 1957$; 12.7%); (ii) 'High sugary beverage consumption' ($n = 8482$; 55.2%); (iii) 'Poor mental wellbeing' ($n = 2009$; 13.1%); (iv) 'Smoker' ($n = 1364$; 8.9%); and (v) 'Alcohol drinker' ($n = 1554$; 10.1%). Being female (OR 1.28, 95%CI 1.14, 1.45) and being physically inactive (OR 1.20, 95%CI 1.04, 1.39) increased the odds of belonging to the 'High sugary beverage consumption' cluster. Being female (OR 1.21, 95%CI 1.04, 1.41), non-membership in a sports club (OR 1.83, 95%CI 1.43, 2.31) were associated with 'Poor mental wellbeing'. Obesity (OR 2.03, 95%CI 1.47, 2.80), inactively commuting to campus (OR 1.34, 95%CI 1.09, 1.66), and living in high-rise accommodation (OR 2.94, 95%CI 1.07, 8.07) were associated with membership in the 'Smoker' cluster. Students living in The Philippines, Singapore, Thailand, and Vietnam had a higher likelihood of being alcohol drinkers, compared with those who lived in Brunei.

Conclusions: ASEAN university students exhibited health-risk behaviours that typically clustered around a specific health behaviour and mental wellbeing. The results provided support for focusing interventions on one

*Correspondence: areekulk@gmail.com

¹⁰ College of Sports Science and Technology, 999 Mahidol University, Phutthamonthon Sai 4 Rd, Salaya, Phutthamonthon District, Nakhon Si Thammaraj 73170, Thailand

Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

dominant health-risk behaviour, with associated health-risk behaviours within clusters being potential mediators for consideration.

Keywords: Diet, Drinking, Mental wellbeing, Physical activity, Smoking, Students, University

Background

Health-risk behaviours are defined as behaviours with potentially negative effects on health such as risks of diseases and injuries [1, 2]. Health-risk behaviours vary in different age groups, environments and cultures [3–5]. Specifically, undergraduate students are in early adulthood and are in transition from high school to university. Health behaviours in university life may have a long-term impact on health conditions and incidence of chronic diseases in later adulthood [6]. Several health-risk behaviours, such as, tobacco smoking, alcohol drinking, physical inactivity, and unhealthy diet, and also mental wellbeing are highlighted as important factors of health in early adulthood [7–9]. The 60-d lifestyle behaviours are known risk factors for non-communicable diseases (NCDs) such as diabetes mellitus, stroke, coronary heart disease, and some forms of cancer [10]. These health-risk behaviours are potentially modifiable and preventable [11, 12]. Modification of these health-risk behaviours can improve health and reduce risks of health problems in later adulthood [13].

Apart from health care agencies, Higher Education Institutions like universities have an essential role in promoting health [14, 15]. In Southeast Asia, the Association of Southeast Asian Nations (ASEAN), consists of 10-member countries: Brunei Darussalam, Cambodia, Indonesia, Lao, Malaysia, Myanmar, The Philippines, Singapore, Thailand, and Vietnam, advocates policies that emphasise the important role that universities play in health promotion [16]. In 2014, a new thematic network of the ASEAN, the ASEAN University Network - Health Promotion Network (AUN-HPN), was established for the purpose of health promotion in the ASEAN region [17]. The AUN-HPN focuses on health promotion among university students and staff under the healthy university framework [16]. The framework includes building systems and infrastructures to support health promoting environments and covers the thematic areas as (i) zero tolerance areas (i.e., smoking, alcohol consumption, illicit drug use, gambling, violence, bullying and sexual harassment, and road safety violations) and (ii) health promotion areas (i.e., health literacy, mental wellbeing, social interaction (e.g., student clubs), physical activity and active mobility, healthy diet and balanced nutrition, safe sexual behaviour, and work-life balance and healthy ageing [15]. To achieve these targets, several strategies

have been implemented, and these included health education and health promotion research [16].

Individuals may have a single dominant health-risk behaviour or multiple health-risk behaviours. Previous research classified people with health-risk behaviours into single dominant or combined health-risk behavioural clusters [7–9, 18–20]. Health-risk behaviours often co-occur or cluster, and having many health-risk behaviours concurrently could increase the probability of mortality (e.g., from cancers), and therefore these lifestyle behaviours have significant public health implications. However, information about either single or combined health behaviours, such as smoking, alcohol drinking, fruit and vegetable consumption, and sugary beverage consumption is scarce and their associations with mental wellbeing among ASEAN university students are not clear. This first surveillance of the health behaviours among ASEAN university students is thus important for understanding the situation and better-informs the health promotion strategies of the AUN-HPN. The identification of the behavioural health-risk and mental wellbeing clusters including their sociodemographic correlates, provides helpful information for designing targeted health-enabling interventions that can tackle multiple health-risk behaviours at the same time for university students. The present study foregrounded the behavioural health-risk and mental wellbeing clusters among ASEAN university students and investigated the associations between the identified clusters and student sociodemographic information.

Methods

Study design and data source

Data analysed were retrieved from a cross-sectional online survey, called the AUN-HPN health behavioural survey. The survey was conducted between 2020 and 2021 and investigated the health-related behaviours and mental wellbeing of ASEAN students from 17 AUN-HPN member universities across seven ASEAN countries (Table 1). The online survey comprised seven sections: 1) Physical activity, 2) Social support for physical activity, 3) University's environment, 4) Health-related behaviours, 5) Mental wellbeing, 6) Opinion regarding university support, and 7) Sociodemographic information. The survey was developed based on previously tested instruments [21–24]. The survey, originally in English, was translated into

Table 1 Participating universities in seven ASEAN countries

Countries (n)	Total number of students n (%)	University names
Brunei Darussalam (1)	1020 (6.6)	Universiti Brunei Darussalam
Indonesia (3)	338 (2.2)	Universitas Airlangga
	3113 (20.3)	Universitas Indonesia
	979 (6.4)	Universitas Gadjah Mada
Malaysia (2)	76 (0.5)	University of Malaya
	213 (1.4)	Universiti Putra Malaysia
The Philippines (1)	322 (2.1)	Ateneo de Manila University
Singapore (1)	259 (1.7)	Nanyang Technological University
Thailand (8)	634 (4.1)	Chulalongkorn University
	253 (1.7)	Chiang Mai University
	265 (1.7)	King Mongkut's University of Technology North Bangkok
	312 (2.0)	Naresuan University
	267 (1.7)	Maharakham University
	619 (4.0)	Mahidol University
	1247 (8.1)	Thammasat University
	397 (2.6)	Walailak University
Vietnam (1)	5052 (32.9)	Vietnam National University

four languages: Bahasa Indonesia, Malaysian, Thai, and Vietnamese. A pilot test of the online survey that included garnering student feedback on the survey using Qualtrics platform (Qualtrics International Inc., WA, USA) was conducted a sub-sample of university students to ensure comprehension and functionality of the online survey.

Measures

Student demographic characteristics

Demographic characteristics including year of study (year 1, 2, 3, and 4 or above); age (18 years, 19 to 21 years, and ≥ 22 years); gender (male and female); country (Brunei Darussalam, Indonesia, Malaysia, The Philippines, Singapore, Thailand, and Vietnam); body mass index (BMI) ('underweight' ($< 18.5 \text{ kg/m}^2$), 'normal' (18.5 to 22.9 kg/m^2), 'overweight' (23.0 to 24.9 kg/m^2), and 'obese' ($\geq 25 \text{ kg/m}^2$) according to World Health Organization (WHO) Asian cut-offs) [25]; the different types of grading from each institution was standardised into a 5-point grade point average (GPA) scale, which was interpreted into three levels consisting of high GPA (> 3.9), moderate GPA (3.3 to 3.9), and low GPA (≤ 3.2); place of living (on-campus and off-campus); commute time to campus (< 15 min, 15 to 30 min, 30 to 45 min, 45 to 60 min, 60 to 90 min, and > 90 min); commute type (active and inactive transportation); housing type (single house, townhouse,

apartment, high-rise condo), and member of sports club (yes and no) were collected.

Exercise/sport participation, physical activity, sedentary time, and sleep

Exercise/sport participation was classified into four categories: none, 1 to 3 days/week, 4 to 6 days, and > 6 days/week.

The Global Physical Activity Questionnaire (GPAQ) version 2.0, which had an acceptable concurrent validity ($r = 0.54$) and high level of repeatability (0.67 – 0.81) was used to collect data on physical activity levels [26, 27]. Physical activity levels were classified into 'inactive' (< 600 Metabolic Equivalent (MET)-min/week) and 'active' (≥ 600 MET-min/week) [28].

Sedentary time was collected from the 40th item of GPAQ 2.0 and divided into three groups: < 4 hours/day, 4 to 8 hours/day, and > 8 hours/day). Sleep time were dichotomised into < 7 hours/day and ≥ 7 hours/day, based on recommendations on sleep hours per night for healthy adults (18–60 years) (i.e., 7 or more sleep hours per night) [29].

Smoking and alcohol drinking

Students were identified as smokers or drinkers when they reported that they are current smokers or drinkers (drink/smoke daily).

Fruit and vegetable consumption, snacking, sugary beverage consumption, and salt intake

Students were classified as healthy (≥ 5 servings/day) or unhealthy (< 5 servings/day) fruit and vegetable consumer. Students, who ate snacks or fast food every day, were categorised into at-risk snacking category; and otherwise, were categorised as at lower risk of snacking. Sugary beverage consumption was classified into at-risk consumption (drink sugary beverage every day) and lower risk of sugary beverage consumption (did not drink every day). At-risk salt intake was defined as when a student added salt to their food before eating ≥ 1 teaspoon/day, and at a lower risk of salt intake meant adding < 1 teaspoon/day of salt.

Mental wellbeing

The shortened Warwick-Edinburgh Mental Wellbeing Scale (WEMWBS) that contained seven items, is a reliable and valid tool for assessing the mental wellness of university students, was used to assess mental wellbeing [22]. The WEMWBS score was dichotomized into negative (poor) and positive (good) mental wellbeing.

Statistical analysis

The R v4.1.1 and RStudio v1.4.1717 for Mac (RStudio, MA, USA) were used for all analyses. Incomplete survey records (i.e., missing demographic characteristics

or relevant health-risk behaviours) were removed from the analysis. A two-step cluster analysis using k-means and hierarchical clustering were employed. In step-one, the number of clusters was determined using k-means algorithm, which indicated that a five-cluster model was optimal [30]. In step-two, hierarchical clustering using Euclidean distance was used to subset the data based on the five-clusters [31]. Descriptive statistics were used to present sociodemographic characteristics of the samples and characteristics of the health-risk behaviours clusters. The cluster labels were based on the predominant characteristics within the clusters. The 'healthy' cluster was identified based on the least number of risk factors and was used as the reference group for comparison using multinomial logistic regression analysis, which was performed to assess the associations between health-risk behaviour clusters and demographic characteristics. McFadden's R-square was used to check for overall model fit. Statistical significance was set at $p < 0.05$.

Results

Sociodemographic characteristics of participants

A final sample of 15,366 ASEAN university students was used for analyses. The sample consisted about equal distribution of male (47.4%) and female (52.6%) students. A majority of the university students were from Vietnam (33.2%), followed by Indonesia (28.8%), and Thailand (25.6%). The participants were mostly in the first year (64.7%) of university life, had normal BMI (61.5%), achieved moderate GPA (69.2%), lived off-campus (65.2%), and commuted to the university using a physically inactive means of transportation (82.9%) (Table 2).


Cluster outputs and characteristics

After data cleaning, the analytical sample comprised 15,366 students. The two-step cluster analysis grouped respondents with similar health-risk behaviours and mental wellbeing resulted in 4 to 15 solutions, and it emerged that the five-cluster model was the most optimal (Fig. 1). Table 3 presents the characteristics of each cluster where the greater proportion of health-risk behaviours or mental wellbeing exhibited by the cluster determined the cluster label.

The largest cluster ($n = 8482$; 55.2%) of the sample that exhibited the highest proportion of high sugary beverage consumption was labelled as 'High sugary beverage consumption'. The other four clusters according to size were labelled as 'Poor mental wellbeing' ($n = 2009$; 13.1%), 'Healthy' ($n = 1957$; 12.7%), 'Alcohol drinker' ($n = 1554$; 10.1%), and 'Smoker' ($n = 1364$; 8.9%). The 'Healthy' cluster exhibited mostly safe health behaviours and was used as the reference group

Table 2 Sociodemographic characteristics of the university students

Characteristics	n	%
Gender		
Male	7289	47.4
Female	8077	52.6
Age in years (n = 13,597)		
18	2496	18.4
19 to 21	9016	66.3
≥ 22	2085	15.3
Academic year		
1st	9940	64.7
2nd	2895	18.8
3rd	1800	11.7
4th or more	731	4.8
BMI (n = 13,097)		
Normal	8441	61.5
Underweight	2917	21.3
Overweight	1739	12.7
Obese	624	4.5
Country		
Brunei Darussalam	1020	6.6
Indonesia	4430	28.8
Malaysia	289	1.9
The Philippines	322	2.1
Singapore	259	1.7
Thailand	3940	25.7
Vietnam	5106	33.2
GPA (n = 12,151)		
≤ 3.2	2443	20.1
3.3 to 3.9	8406	69.2
> 3.9	1302	10.7
Place of living		
On-campus	5345	34.8
Off-campus	10,021	65.2
Commute time to university		
14 min	5917	38.5
15 to 30 min	4127	26.9
30 to 45 min	1973	12.8
45 to 60 min	1692	11.0
60 to 90 min	1059	6.9
> 90 min	598	3.9
Commute type to university		
Active	2639	17.1
Inactive	12,727	82.9
Housing type		
Single house	11,319	73.7
Townhouse	2773	18.0
Apartment	1201	7.8
High rise condo	73	0.5
Member of sports club		
Yes	9054	58.9
No	6312	41.1



for multinomial logistic regression. The healthy, high sugary beverage consumption, poor mental wellbeing and alcohol drinker clusters were each comprised entirely of non-smokers, with those who smoked clustering into their own smoking cluster. The healthy, high sugary beverage consumption and poor mental wellbeing clusters comprised all non-alcoholic drinkers, with the alcohol drinkers comprising all drinkers and 63% of the smoking group cluster also drinking alcohol. Members of the healthy cluster did not drink sugary beverages, and all had positive mental wellbeing. Members of the high sugary beverage consumption cluster were also non-smokers, non-drinkers and had positive mental wellbeing. Members of the smoking cluster also had high sugary beverage consumption with almost 80% having positive mental wellbeing. Members of the alcohol cluster were all non-smokers with 87% with high sugary beverage consumption, 83% positive mental wellbeing and 72% in the high physically active group. The level of exercise/sport participation, physical activity, sedentary time, sleeping hours and high salt intake did not appear to differentiate clusters.

Associations between clusters and demographic characteristics

To further understand the similarities and differences of the 5 clusters, multinomial logistic regression analyses were run with the demographic characteristics as the predictors and the cluster membership as the outcome variable. The 'Healthy' cluster was chosen as the reference group in the analysis. Table 4 shows the likelihood

of university students falling into the unhealthy or health-risk clusters. 51

Students in Year 2 (OR 1.24, 95%CI 1.04, 1.46) and Year 3 (OR 1.28, 95%CI 1.01, 1.63), being female (OR 1.28, 95%CI 1.14, 1.45) living in Thailand (OR 1.71, 95%CI 1.16, 2.52), and being inactive (OR 1.20, 95%CI 1.04, 1.39) had a higher likelihood of falling into 'High sugary beverage consumption' cluster (cluster 2), compared to the 'Healthy' cluster. Meanwhile, students living in The Philippines (OR 0.38, 95%CI 0.24, 0.69), being underweight (OR 0.85, 95%CI 0.74, 0.99) or overweight (OR 0.82, 95%CI 0.68, 0.97), and staying in a townhouse (OR 0.82, 95%CI 0.68, 1.00) had less likelihood of being in the 'High sugary beverage consumption' cluster.

36 comparison to the healthy group, being in 2nd year (OR 1.24, 95%CI 1.01, 1.52), females (OR 1.21, 95%CI 1.04, 1.41), non-membership to a sports club (OR 1.83, 95%CI 1.43, 2.34), and travelling between 30 to 45 minutes to campus (OR 1.33, 95%CI 1.95, 1.70) were significantly more likely to fall into the 'Poor mental wellbeing' cluster (cluster 3). Students who were underweight (OR 0.74, 95%CI 0.61, 0.90) and resided in Indonesia (OR 0.34, 95%CI 0.24, 0.49), Malaysia (OR 0.32, 95%CI 0.16, 0.67), The Philippines (OR 0.4, 95%CI 0.24, 0.67), or Vietnam (OR 0.62, 95%CI 0.44, 0.86) compared to students from Brunei were less likely to be in the 'Poor mental wellbeing' cluster.

Belonging to the 'Smoker' cluster (cluster 4), was associated with being in 2nd year (OR 1.37, 95%CI 1.09, 1.72), overweight (OR 1.58, 95%CI 1.23, 2.04), obese (OR 2.03, 95%CI 1.47, 2.80), travelling 30–45 minutes to campus (OR 1.40, 95%CI 1.08, 1.82), inactively commuting to campus (OR 1.34, 95%CI 1.09, 1.66), and living in

Table 3 Characteristics of the naturally occurring clusters extracted from the dataset (n = 15,366)

	Total (n = 15,366) n (%)	Cluster 1 Healthy (n = 1957) n (%)	Cluster 2 High sugary beverage consumption (n = 8482) n (%)	Cluster 3 Poor mental wellbeing (n = 2009) n (%)	Cluster 4 Smoker (n = 1364) n (%)	Cluster 5 Alcohol drinker (n = 1554) n (%)
Health behaviours and mental wellbeing						
Exercise/sport participation						
None	2131 (13.87)	309 (15.79)	1123 (13.24)	409 (20.36)	174 (12.76)	116 (7.46)
1–3 days/week	8081 (52.59)	1111 (56.77)	4596 (54.19)	1068 (53.16)	678 (49.71)	628 (40.41)
4–6 days/week	2937 (19.11)	319 (16.30)	1548 (18.25)	260 (12.94)	330 (24.19)	480 (30.89)
> 6 days/week	2217 (14.43)	218 (11.14)	1215 (14.32)	272 (13.54)	182 (13.34)	330 (21.24)
Physical activity						
Inactive (< 600 MET-min/week)	6097 (39.68)	807 (41.24)	3390 (39.97)	962 (47.88)	504 (36.95)	434 (27.93)
Active (≥ 600 MET-min/week)	9269 (60.32)	1150 (58.76)	5092 (60.03)	1047 (52.12)	860 (63.05)	1120 (72.07)
Sedentary time						
< 4 hours/day	2376 (15.46)	360 (18.40)	1272 (15.00)	278 (13.84)	226 (16.57)	240 (15.44)
4–8 hours/day	6213 (40.43)	742 (37.92)	3605 (42.50)	685 (34.10)	569 (41.72)	612 (39.38)
> 8 hours/day	6777 (44.10)	855 (43.69)	3605 (42.50)	1046 (52.07)	569 (41.72)	702 (45.17)
Sleeping hours						
< 7 hours/day	10,566 (68.76)	1232 (62.95)	5965 (70.33)	1377 (68.54)	875 (64.15)	1117 (71.88)
≥ 7 hours/day	4800 (31.24)	725 (37.05)	2517 (29.67)	632 (31.46)	489 (35.85)	437 (28.12)
Smoking						
Smoker	1365 (8.88)	0	0	0	1364 (100)	1 (0.06)
Non-smoker	14,001 (91.12)	1957 (100)	8482 (100)	2009 (100)	0	1553 (99.94)
Alcohol drinking						
Drinker	2420 (15.75)	0	0	0	866 (63.49)	1554 (100)
Non-drinker	12,946 (84.25)	1957 (100)	8482 (100)	2009 (100)	498 (36.51)	0
Fruit and vegetable consumption						
Unhealthy (< 5 servings/day)	7339 (47.76)	908 (46.40)	4052 (47.77)	797 (39.67)	683 (50.07)	899 (57.85)
Healthy (≥ 5 servings/day)	8027 (52.24)	1049 (53.60)	4430 (52.23)	1212 (60.33)	681 (49.93)	655 (42.15)
Snacking						
At-risk (every day)	10,019 (65.20)	1098 (56.11)	5539 (65.30)	1283 (63.86)	1027 (75.29)	1072 (68.98)
Lower risk (not every day)	5347 (34.80)	859 (43.89)	2943 (34.70)	726 (36.14)	337 (24.71)	482 (31.02)
High sugary beverage consumption						
Yes	12,598 (81.99)	0	8482 (100)	1622 (80.74)	1148 (84.16)	1346 (86.62)
No	2768 (18.01)	1957 (100)	0	387 (19.26)	216 (15.84)	208 (13.38)
High Salt intake						
Yes	8305 (54.05)	804 (41.08)	4682 (55.20)	1063 (52.91)	815 (59.75)	941 (60.55)
No	7061 (45.95)	1153 (58.92)	3800 (44.80)	946 (47.09)	549 (40.25)	613 (39.45)
Mental wellbeing						
Negative mental wellbeing	2559 (16.65)	0	0	2009 (100)	290 (21.26)	260 (16.73)
Positive mental wellbeing	12,807 (83.35)	1957 (100)	8482 (100)	0	1074 (78.74)	1294 (83.27)

Bold numbers represent dominant health behaviours and mental wellbeing in each cluster

high-rise accommodation (OR 2.94, 95%CI 1.07, 8.07) compared to members of the healthy cluster.

The alcohol drinker cluster (cluster 5) was associated with being 53 2nd year (OR 1.28, 95%CI 1.03, 1.60) or 3rd year (OR 1.46, 95%CI 1.08, 1.98) compared to the

healthy cluster and 1st year students. Students living in The Philippines (OR 3.11, 95%CI 1.47, 6.56), Singapore (OR 14.70, 95%CI 6.28, 34.42), Thailand (OR 7.06, 95%CI 3.64, 13.70), or Vietnam (OR 6.34, 95%CI 3.51, 11.43) had higher odds of being a drinker compared with students

Table 4 Factors associated with the healthy cluster (Cluster 1) using multinomial logistic regression analysis (n = 15,366)

Demographic characteristics	Cluster 2 High sugary beverage consumption OR (95%CI)	Cluster 3 Poor mental wellbeing OR (95%CI)	Cluster 4 Smoker OR (95%CI)	Cluster 5 Alcohol drinker OR (95%CI)
Gender (Ref. = Male)				
Female	1.28 (1.14, 1.45)**	1.21 (1.04, 1.41)*	0.93 (0.78, 1.10)	1.14 (0.96, 1.36)
Age (years) (Ref. = 18)				
19 to 21	0.94 (0.81, 1.10)	0.99 (0.81, 1.22)	1.06 (0.84, 1.34)	1.29 (0.98, 1.70)
≥ 22	0.90 (0.70, 1.15)	0.76 (0.55, 1.05)	1.09 (0.77, 1.54)	1.24 (0.85, 1.81)
Year of study (Ref. = Year 1)				
Year 2	1.24 (1.04, 1.46)*	1.24 (1.01, 1.52)*	1.37 (1.09, 1.72)**	1.28 (1.03, 1.60)*
Year 3	1.28 (1.01, 1.63)*	1.15 (0.85, 1.55)	1.19 (0.86, 1.64)	1.46 (1.08, 1.98)*
Year 4 or above	1.36 (0.92, 2.03)	1.37 (0.84, 2.25)	1.22 (0.72, 2.06)	1.57 (0.96, 2.59)
BMI (Ref. = Normal)				
Underweight	0.85 (0.74, 0.99)*	0.74 (0.61, 0.90)**	1.09 (0.86, 1.37)	1.01 (0.81, 1.25)
Overweight	0.82 (0.68, 0.97)*	0.87 (0.70, 1.09)	1.58 (1.23, 2.04)**	1.22 (0.95, 1.56)
Obese	0.91 (0.72, 1.17)	1.01 (0.75, 1.36)	2.03 (1.47, 2.80)**	0.99 (0.70, 1.40)
Country (Ref. = Brunei Darussalam)				
Indonesia	0.86 (0.63, 1.16)	0.34 (0.24, 0.49)**	1.17 (0.78, 1.75)	0.75 (0.40, 1.42)
Malaysia	0.45 (0.24, 0.77)**	0.32 (0.16, 0.67)**	0.23 (0.09, 0.57)**	0.20 (0.04, 0.98)*
Philippines	0.38 (0.24, 0.69)**	0.40 (0.24, 0.67)**	1.05 (0.58, 1.89)	3.11 (1.47, 6.56)**
Singapore	1.11 (0.58, 2.12)	0.91 (0.43, 1.92)	1.23 (0.53, 2.88)	14.70 (6.28, 34.42)**
Thailand	1.71 (1.16, 2.52)**	0.95 (0.60, 1.52)	1.18 (0.70, 1.99)	7.06 (3.64, 13.70)**
Vietnam	1.32 (0.98, 1.77)	0.62 (0.44, 0.86)**	0.86 (0.58, 1.30)	6.34 (3.51, 11.43)**
GPA (Ref. = ≤ 3.2)				
3.3 to 3.9	1.05 (0.90, 1.23)	0.97 (0.80, 1.18)	0.85 (0.69, 1.05)	0.76 (0.62, 0.92)**
> 3.9	0.91 (0.72, 1.14)	0.91 (0.68, 1.22)	0.72 (0.52, 1.01)	0.64 (0.48, 0.86)**
Place of living (Ref. = On-campus)				
Off-campus	1.18 (0.94, 1.49)	1.05 (0.79, 1.40)	1.10 (0.78, 1.56)	0.91 (0.67, 1.24)
Commute time (Ref. = less than 15 min)				
15 to 30 min	1.06 (0.92, 1.24)	1.04 (0.86, 1.26)	1.06 (0.86, 1.31)	1.04 (0.85, 1.27)
30 to 45 min	1.07 (0.88, 1.30)	1.33 (1.05, 1.70)*	1.40 (1.08, 1.82)*	1.04 (0.78, 1.38)
45 to 60 min	0.84 (0.69, 1.01)	0.83 (0.65, 1.07)	0.81 (0.61, 1.08)	0.76 (0.57, 1.02)
60 to 90 min	0.96 (0.76, 1.20)	0.95 (0.71, 1.27)	0.76 (0.53, 1.07)	0.57 (0.39, 0.83)**
> 90 min	0.88 (0.67, 1.16)	1.27 (0.90, 1.79)	0.71 (0.46, 1.10)	0.55 (0.32, 0.94)*
Commute type (Ref. = Active)				
Inactive	1.20 (1.04, 1.39)*	1.13 (0.93, 1.36)	1.34 (1.09, 1.66)**	1.05 (0.85, 1.30)
Housing type (Ref. = Single house)				
Townhouse	0.82 (0.68, 1.00)*	0.80 (0.63, 1.02)	1.03 (0.80, 1.33)	1.12 (0.89, 1.40)
Apartment	1.01 (0.81, 1.26)	0.92 (0.70, 1.22)	1.28 (0.95, 1.73)	1.25 (0.93, 1.68)
High rise condo	1.18 (0.48, 2.88)	1.20 (0.42, 3.44)	2.94 (1.07, 8.07)*	2.34 (0.78, 6.97)
Member of sports club (Ref. = Yes)				
No	0.99 (0.83, 1.18)	1.83 (1.43, 2.34)**	0.45 (0.35, 0.56)**	0.70 (0.54, 0.90)**

95%CI 95% Confidence interval, BMI Body Mass Index, GPA Grade Point Average, OR Adjusted odds ratio, Ref. Reference group

*p < 0.05, **p < 0.01

living in Brunei. Students with a higher GPA were less likely to belong to the alcohol cluster (OR 0.76, 95%CI 0.62, 0.92 for GPA 3.3 to 3.9; and OR 0.64, 95%CI 0.48, 0.86 for GPA > 3.9) compared to students in the healthy

cluster. Students who spent more time travelling to campus (> 60 min) were less likely to be in the alcohol drinker cluster (OR 0.57, 95%CI 0.39, 0.83 for 60 to 90 min; and OR 0.55, 95%CI 0.32, 0.94 for > 90 min) compared to

students in the healthy cluster. Students who were not members of a sports club had 30% lower odds of belonging to the alcohol drinker cluster than the 'healthy' cluster and students who belonged to a sports club.

Discussion

The research investigated the health-risk behaviours and mental wellbeing among university students in seven ASEAN countries and it emerged that five behavioural clusters (i.e., healthy, high sugary beverage consumption, smoker, and alcohol drinker) and mental wellbeing, were identified. The results showed that among ASEAN university students, consuming sugary beverages was the dominant health behaviour across all clusters. This finding is consistent with the findings reported by Pengpid and Pletzer (2019) who observed that a high proportion of ASEAN university students consumed sugary soft drinks and this was associated with other unhealthy behaviours, such as smoking and drinking alcohol [32]. The present finding added to a small body of evidence linking the consumption of sugary beverages with a range of unhealthy behaviours among ASEAN university students. Therefore, it may be important to consider sugary beverage consumption behaviour when targeting other health-risk behaviours for improvement.

This research showed that several characteristics of university students were related to health-risk behaviours and mental wellbeing. Students in the second year of university study had a higher likelihood of having poor mental wellbeing and falling into the 'high sugary beverage consumption', 'smoker', and 'alcohol drinker' clusters, compared to the 'healthy' cluster. These results were in line with the findings of a Chinese study that showed that second year students suffered relatively higher levels of depression and stress, compared to students in the first, third, and fourth years of university study [33]. Another study conducted in Korea also supported the present results, where Korean university students in the second year experienced more mental health problems than the students in other years of study [34]. The academic demands of year 2 of university life might present greater stress and mental health challenges for year 2 students. Many universities usually set general courses for freshmen and introduce more specialised courses from the second year onwards [33]. Higher levels of stress, coupled with mental health challenges in year 2 students may relatively predispose them to getting involved in other health-risk behaviours compared to first year students.

Our results showed that more than half of ASEAN university students consumed sugary beverages.

Paradoxically, both underweight and overweight students were less likely to consume sugary beverages compared to university peers of normal body weight. Our findings contrasted with those of a previous study on Saudi Arabian adolescents showing that intake of sugar-sweetened carbonated beverage was positively associated with BMI [35]. It appears that both socio-economic and environmental factors, are associated with high sugar diets and sugary beverage consumption in university students [36–38]. Of interest, university students' perceptions regarding sugar intake were explored in a qualitative study and although students perceived that excessive sugar intake affected their body image because of weight-gain, the students, nonetheless thought that they were not at risk of negative health outcomes as they were young [39]. Our findings alluded that the strategies to reduce sugary beverage consumption so as to reduce obesity among ASEAN university students should be applied to all students regardless of their body weight status.

The present study revealed that poor mental wellbeing was the second most prevalent health-risk characteristic among university students. Mental health issues among university students increased the burden on campus counselling resources, and have received great attention from educators within the AUN-HPN as poor mental wellbeing could lead to significant psychological problems and tragedies (i.e. suicide) [40]. We found that students who were not a member of any sports clubs had 83% increased risk of poor mental wellbeing. Non-membership to a sports club presented the highest odds of having poor mental health, albeit it this is only an association and may not causative. Nonetheless, mental health promotion and helping students become physically active on a regular basis are recommendations for improvement in mental wellbeing among students [40–42]. A US-based study showed that sports club participation in college students is associated with positive health-related outcomes [41]. Also aligned, Australian students with higher sports club involvement have a positive and significant association with social-emotional wellbeing indicators, such as happiness, resilience, and body image, whereas low involvement in sports club is associated with a greater incidence of mental health diagnosis [43]. There is compelling evidence for the benefits of sports involvement and participation in university students. Involvement in team sports is also associated with reduced depressive mood because sport participation protects students against social isolation [44, 45]. Other research shows that lower depression scores are reported in the moderate sports involvement group, compared to the low sports involvement group [46]. Students who are active in sports clubs have a better self-concept because sports participation helps students build confidence,

acquire competent behaviours such as social skills, and release energy and aggression in socially accepted ways [47]. Joining and playing on team sports is also associated with greater life satisfaction, higher self-image, and less distress than students who are not involved in sport [48–50]. By being a member of sports club, students are more psychologically resilient, confident, assertive, have better social skills, self-esteem, self-efficacy, self-control, self-concept, and competent [49]. Participating in sports clubs can be a means of improving mental wellbeing. Therefore, encouraging students to join sports clubs might be considered as part of strategies to promote mental wellbeing among the ASEAN university students.

Participation in sports clubs also seemed to influence the likelihood of being smokers or drinkers among ASEAN university students. The present study showed that not being sports club members was associated with less likelihood to smoke and drink alcohol. These results are consistent with evidence where university students who joined sports groups or organisational sports are more likely to smoke and drink alcohol [51–54]. A systematic review by Lisha et al. showed that university students in sports clubs reported higher levels of drinking and smoking than those who were not in any sports clubs [55]. Plausible explanations for these observations are that university students were in a transition period of their lives and were exposed to substantial changes in terms of environmental (living on their own away from families, gaining independence) and social aspects (making new friends with need to belong). This kind of adolescent-adulthood transition is often associated with an increase in heavy and risky alcohol use [56, 57]. Theory and empirical findings suggest that peer pressure is a combination of three distinct influences: overt offers of alcohol, modelling, and social norms [58]. Overt offers of alcohol can range from polite gestures to intense goading or commands to drink [58]. Modelling occurs when the student's behaviour corresponds to another student's concurrent drinking behaviour [58]. Perceived social norms can serve to make excessive alcohol use appear common and acceptable to the students [58]. In the present case, making a recommendation for university students to join a sports club may seem contradictory since on the one hand, it could safeguard mental wellbeing, yet on the other hand, this could expose students to pressures and risks of high drinking and smoking. Therefore, comprehensive strategies that address these paradoxical observations and promote health among ASEAN university students should be considered. For instance, having trained student ambassadors in sports clubs and incentivising healthier activities among sports teams that are in keeping with the culture and context of each country are ideas that are worthy of exploration.

In the present research, the country of residence was associated with specific health-risk behaviours such as alcohol drinking. University students in all countries except Indonesia and Malaysia had higher odds of being current drinkers, whereas university students in Indonesia and Malaysia had lower odds, compared to those in Brunei. A majority of the population in Brunei, Malaysia, and Indonesia is Islamic and alcohol drinking is considered haram (prohibited or sinful) for consumption and therefore alcohol consumption is not likely to be high [59]. Research shows that alcohol consumption is associated with religion [60]. Moreover, some research suggested that religious commitment among the pious and those who are faithful to the teachings of religion (e.g. advocating abstinence from alcohol consumption), is associated with reduced likelihood of alcohol drinking [61]. We are of the view that differences in culture and context within each country in ASEAN pose challenges in the implementation of a common intervention programme to address alcohol consumption in university students. Further research is required to tease out factors that might play a role in alcohol consumption among university students in each country and how these country and culture-specific contexts might be built into future interventions to reduce alcohol abuse for positive results to be accomplished and sustained.

Overall, our results showed that, during the COVID-19 pandemic, only 12.7% of university students exhibited multiple healthy behaviours, with over 85% exhibiting at least one dominant unhealthy behaviour (e.g., high sugary beverage consumption, alcohol consumption and smoking). These results showed that a majority of ASEAN university students was at risk of developing multiple health problems, if their present health behaviours are not improved over time. The AUN-HPN Healthy University Framework highlighted smoking, alcohol consumption, and mental wellbeing as important areas to address [16]. A majority of smokers also drank alcohol. Health promotion policies and interventions targeting smoking behaviours should combine strategies to prevent or reduce alcohol drinking concomitantly. In need of urgent attention and action is to reduce the consumption of sugary beverages, as this habit is most prevalent among other identified health-risk behaviours and should be considered as a priority health promotion issue among ASEAN university students. Although cause-and-effect cannot be determined in the associations between the cluster of high sugary consumption and other health-risk behaviour clusters identified in our cross-sectional analyses, the results suggested that clusters of health-risk behaviours tend to 'reinforce' each other, and future research should examine if moderating high sugary consumption

among university students might have a significant impact on other health-risk behaviours.

This cross-sectional descriptive research provided some insights into the potential areas of concern and actions for health promotion among university students in the seven ASEAN countries. First, university policy makers should focus on the transition from the 1st to 2nd year of university study. This period provides an opportunity for health advocates to intervene and prevent adolescents transiting into early adulthood from adopting high risk and unhealthy behaviours and habits. Data from the present research provided useful finer-grain information and specificity in terms of areas for intervention. Specifically, health promotion interventions should be targeted at reducing sugary beverage consumption and relatively poor mental health, especially among female students. To prevent and reduce the prevalence of smoking, interventions should focus on students who are overweight and obese as well as those who travelled to the campus by motorised transport. Some consideration and attention are needed to attend to alcohol drinking in students living in the non-Islamic countries to reduce the likelihood of alcohol abuse. While mental health may be safeguarded by joining sports clubs, the risks of picking up unhealthy behaviours of alcohol consumption and smoking in sports clubs also need to be moderated and monitored. The key results of our study established some demographic characteristics with specific health-risk behaviour and showed that health-risk behaviours were clustered and co-existed with each other. Policy makers should formulate health-enabling programmes that can best address multiple health risk behaviours at the same time such as adopting a socio-ecological model interventionist approach [62].

The present research had a number of strengths. Firstly, the research was conducted in seven countries in the Southeast Asian region and involved a sizeable sample. Secondly, the use of a self-reported online questionnaire allowed more university students to be polled in the COVID-19 pandemic when online interactions and travel restrictions were the norm. The research also had two notable limitations. The research design was cross-sectional and descriptive in nature and therefore only associations among demographic characteristics and health-risk behaviours could be established and cause-and-effect among the associations cannot be determined. Causal inference may require future research based on the longitudinal monitoring of university students. Another potential limitation was that the research was conducted during the COVID-19 pandemic, which could have a disproportionate impact on many health behaviours.

Conclusion

The majority of ASEAN university students exhibited one common health-risk behaviour. A large proportion of the university students had a habit of sugary beverage consumption. This was followed by poor mental coping strategies, alcohol drinking, and tobacco smoking. The transition from the 1st to 2nd year of university life was particularly challenging and 2nd year university students had higher odds of exhibiting several health-risk behaviours. Health promotion strategies for healthy universities should focus on comprehensive interventions in addressing the dominant health-risk behaviour as well as the other associated health-risk behaviours within the identified clusters.

Abbreviations

ASEAN: Association of Southeast Asian Nations; AUN-HPN: ASEAN University Network - Health Promotion Network; BMI: Body mass index; COVID-19: Coronavirus disease 2019; GPAQ: Global Physical Activity Questionnaire; GPA: Grade point average; NCDs: Non-communicable diseases; WEMWBS: Warwick Edinburgh Mental Wellbeing Scale; WHO: World Health Organization.

Acknowledgements

Authors thank administrative staffs of all participating universities for their assistance during sampling recruitment and data collection. We feel gratitude to all students who participated in the survey. Authors appreciate institutional support and advice from the AUN-HPN International Advisory Committee with special thanks to Dr. Wiwat Rojanapithayakorn and Dr. Viji Kasemsab during development of this project.

Authors' contributions

AW, HAR, MR, and AA initiated the concepts of the study. AW, HAR, MR, and AA analysed, interpreted the study results. AW wrote and revised the manuscript. All authors provided critical comments on the drafts. AW, MR, MC and AA edited the manuscript. All authors read and approved the final version of the manuscript.

Funding

This study was funded in part by the Thai Health Promotion Foundation through Children and Youth Physical Activity Studies (Ref: 61-00-1814), the Centre of Advanced Research, Universiti Brunei Darussalam (UBD/RSC/1.10/FICBF(b)/2019/005), and NIH training grant (T32 GM141746).

Availability of data and materials

The datasets generated and/or analysed during the current study are not publicly available due to restrictions on intellectual property regulations of the funding organization. Data are however available provided that an application is submitted at info@thaihealth.or.th or areekulk@gmail.com approved by the dataset custodians. No administrative permissions were required to access raw data.

Declarations

Ethics approval and consent to participate

This study protocols were undertaken in accordance with relevant local and international ethical guidelines and regulations; e.g., Declarations of Helsinki. The primary ethical approval for the whole study protocols was obtained from Mahidol University Central Institutional Review Board (MU-CIRB 2020/089.0704). Participating universities also obtained ethical approval from their institutional review board: University Research and Ethics Committee of Universiti Brunei Darussalam (UBD/OAVCR/UREC/OCTOBER2020-76) Universitas Gadjah Mada (KE/FK/1066/EC/2020), Universitas Indonesia (KET-126/UN2.F1/ETIK/PPM00.02/2021), Universiti Putra Malaysia (JKEUPM-2020-156), Ateneo de Manila University (AdMUREC-20-010), National Institute of

Education, Nanyang Technological University (IRB-2021-03-027), Naresuan University (P29936/63), Thammasat University (075/2563), Burapha University (HS035/2563), Chiang Mai University (AMSEC-63EX-019), Vajiravudh University (WUEC-20-172-01), Mahasarakham University (266/2020), Vietnam National University, Hanoi (01/2020/IRB-VN01.017-MEDVNU), Universitas Airlangga and King Mongkut's University of Technology North Bangkok used MU-CIRB. Data analysed in this study were anonymized before use. Participants in this study gave their informed online consent by clicking 'I agree to participate' before completing the survey. The informed online consent was approved by Mahidol University Central Institutional Review Board (MU-CIRB 2020/089.0704).

Consent for publication
Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹School of Medicine, Walailak University, Nakhon Si Thammarat, Thailand. ²Walailak University Hospital, Nakhon Si Thammarat, Thailand. ³Institute of Health Sciences, Universiti Brunei Darussalam, Brunei Darussalam. ⁴University of Michigan, Ann Arbor, MI, USA. ⁵Centre for Population Health Sciences, Lee Kong Chian School of Medicine, Nanyang Technological University, Singapore, Singapore. ⁶School of Nursing, Midwifery, La Trobe University, Melbourne, VIC, Australia. ⁷Faculty Ateneo School of Medicine and Public Health, Ateneo de Manila University, Quezon City, The Philippines. ⁸Physical Education & Sports Science, National Institute of Education, Nanyang Technological University, Singapore, Singapore. ⁹School of Human Sciences (Sport Science, Exercise and Health), University of Western Australia, Perth, WA, Australia. ¹⁰College of Sports Science and Technology, Mahidol University, Phuthamonthon Sai 4 Rd, Salaya, Phuthamonthon District, Nakhon Pathom 73170, Thailand. ¹¹School of Social Sciences, Nanyang Technological University, Singapore, Singapore. ¹²Alcohol and Health Promotion Policy Research Unit, Faculty of Pharmacy, Mahasarakham University, Kantharavichai, Mahasarakham, Thailand. ¹³Department of Nutrition, Faculty of Public Health, Universitas Airlangga, Surabaya, Indonesia. ¹⁴Faculty of Public Health, Burapha University, Saen Suk, Chon Buri, Thailand. ¹⁵Department of Information Systems, Yamouk University, Irbid, Jordan.

Received: 1 April 2022 Accepted: 22 September 2022
Published online: 01 October 2022

References

- Peltzer K, Pengpid S, Yung TKC, Aounallah-Skhiri H, Rehman R. Comparison of health risk behavior, awareness, and health benefit beliefs of health science and non-health science students: an international study. *Nurs Health Sci*. 2016;18(2):180–7.
- Suris JC, Michaud PA, Akre C, Sawyer SM. Health risk behaviors in adolescents with chronic conditions. *Pediatrics*. 2008;122(5):e1113–8.
- Duell N, Steinberg L, Icenogle G, Choin J, Chaudhary N, Di Giunta L, et al. Age patterns in risk taking across the world. *J Youth Adolesc*. 2018;47(5):1052–72.
- Rattay P, von der Lippe E, Mauz E, Richter F, Hölling H, Lange C, et al. Health and health risk behaviour of adolescents—differences according to family structure. Results of the German KiGGS cohort study. *Plos One*. 2018;13(3):e0192968.
- Wattanapit A, Jiraporncharoen W, Pinyopornpanish K, Jiraniramai S, Thakla K, Angkurawaranon C. Health-risk behaviours and injuries among youth and young adults in Chiang Mai, Thailand: a population-based survey. *Int J Environ Res Public Health*. 2020;17(10):3696.
- Corder K, Winpenny E, Love R, Brown HE, White M, Suijs EV. Change in physical activity from adolescence to early adulthood: a systematic review and meta-analysis of longitudinal cohort studies. *Br J Sports Med*. 2019;53(8):496–503.
- Bennasar-Veny M, Yañez AM, Pericas J, Ballester L, Fernandez-Dominguez JC, Tauler P, et al. Cluster analysis of health-related lifestyles in university students. *Int J Environ Res Public Health*. 2020;17(5):1776.
- Jao NC, Robinson LD, Kelly PJ, Ciecierski CC, Hitsman B. Unhealthy behavior clustering and mental health status in United States college students. *J Am Coll Heal*. 2019;67(8):790–800.
- Murphy JJ, MacDonncha C, Murphy MH, Murphy N, Timperio A, Leech RM, et al. Identification of health-related behavioural clusters and their association with demographic characteristics in Irish university students. *BMC Public Health*. 2019;19(1):121.
- Peters R, Ee N, Peters J, Beckett N, Booth A, Rockwood K, et al. Common risk factors for major noncommunicable disease, a systematic overview of reviews and commentary: the implied potential for targeted risk reduction. *Ther Adv Chronic Dis*. 2019;10:2040622319880392.
- Bolnick HJ, Bui AL, Bulchis A, Chen C, Chapin A, Lomsadze L, et al. Health-care spending attributable to modifiable risk factors in the USA: an economic attribution analysis. *Lancet Public Health*. 2020;5(10):e525–35.
- Budreviciute A, Damati S, Sabir DK, Onder K, Schuller-Goetzburg P, Plakys G, et al. Management and prevention strategies for non-communicable diseases (NCDs) and their risk factors. *Front Public Health*. 2020;8:574111.
- Dawson KA, Schneider MA, Fletcher PC, Bryden PJ. Examining gender differences in the health behaviors of Canadian university students. *JR Soc Promot Heal*. 2007;127(1):38–44.
- Bachert P, Wäsche H, Albrecht F, Hildebrand C, Kunz AM, Woll A. Promoting students' health at university: key stakeholders, cooperation, and network development. *Front Public Health*. 2021;9:680714.
- Faria MGA, Fernandes RC, Gallasch CH, Alves LW. Contributions of the health-promoting universities' movement: an integrative literature review. *J Educ Health Promot*. 2021;10:114.
- AUN-Health Promotion Network. AUN Healthy University framework. Bangkok: Mahidol University; 2017.
- ASEAN University Network. AUN Health Promotion Network (AUN-HPN) <https://www.aunsec.org/discover-aun/thematic-networks/aun-hpn>. Accessed 28 Dec 2021.
- French S, Rosenberg M, Knuiam M. The clustering of health risk behaviours in a Western Australian adult population. *Health Promot J Austr*. 2008;19(3):203–9.
- Noble N, Paul C, Turon H, Oldmeadow C. Which modifiable health risk behaviours are related? A systematic review of the clustering of smoking, nutrition, alcohol and physical activity ('SNAP') health risk factors. *Prev Med*. 2015;81:16–41.
- Meador N, King K, Moe-Byrne T, Wright K, Graham H, Petticrew M, et al. A systematic review on the clustering and co-occurrence of multiple risk behaviours. *BMC Public Health*. 2016;16:657.
- Centers for Disease Control and Prevention. Behavioral Risk Factor Surveillance System Survey Questionnaire, in Environmental module. Georgia: Centers for Disease Control and Prevention, U.S. Department of Health and Human Services; 2002.
- Fung SF. Psychometric evaluation of the Warwick-Edinburgh mental well-being scale (WEMWBS) with Chinese University students. *Health Qual Life Outcomes*. 2019;17(1):46.
- Noncommunicable Diseases and Mental Health, World Health Organization. WHO STEPS surveillance manual: the WHO STEPwise approach to chronic disease risk factor surveillance. Geneva: World Health Organization; 2005.
- Treiber FA, Baranowski T, Braden DS, Strong WB, Levy M, Knox W. Social support for exercise: relationship to physical activity in young adults. *Prev Med*. 1991;20(6):737–50.
- World Health Organization, Regional Office for the Western Pacific. The Asia-Pacific perspective: redefining obesity and its treatment. Sydney: Health Communications Australia; 2000.
- Bull FC, Maslin TS, Armstrong T. Global physical activity questionnaire (GPAQ): nine country reliability and validity study. *J Phys Act Health*. 2009;6(6):790–804.
- Keating XD, Zhou K, Liu X, Hodges M, Liu J, Guan J, et al. Reliability and concurrent validity of global physical activity questionnaire (GPAQ): a systematic review. *Int J Environ Res Public Health*. 2019;16(21):4128.
- Chu AH, Ng SH, Koh D, Müller-Riemenschneider F. Reliability and validity of the self- and interviewer-administered versions of the global physical activity questionnaire (GPAQ). *Plos One*. 2015;10(9):e0136944.
- Watson NF, Badr MS, Belenky G, Blhwise DL, Buxton OM, Buysse D, et al. Recommended amount of sleep for a healthy adult: a joint consensus statement of the American Academy of sleep medicine and Sleep Research Society. *Sleep*. 2015;38(6):843–4.
- Kodinariya TM, Makwana PR. Review on determining number of cluster in K-means clustering. *Int J*. 2013;1:90–5.

31. Nielsen F. Hierarchical clustering. In: Nielsen F, editor. Introduction to HPC with MPI for data science. Cham: Springer; 2016. p. 195–211.
32. Pengpid S, Peltzer K. High carbonated soft drink consumption is associated with externalizing but not internalizing behaviours among university students in five ASEAN states. *Psychol Res Behav Manag.* 2019;12:585–92.
33. Liu X, Ping S, Gao W. Changes in undergraduate students' psychological well-being as they experience university life. *Int J Environ Res Public Health.* 2019;16(16):2864.
34. Choi J. Impact of stress levels on eating behaviors among college students. *Nutrients.* 2020;12(5):1241.
35. Collison KS, Zaidi MZ, Subhani SN, Al-Rubeaan K, Shoukri M, Al-Mohanna FA. Sugar-sweetened carbonated beverage consumption correlates with BMI, waist circumference, and poor dietary choices in school children. *BMC Public Health.* 2010;10:234.
36. Bruce MA, Beech BM, Thorpe RJ Jr, Mincey K, Griffith DM. Racial and gender disparities in sugar consumption change efficacy among first-year college students. *Appetite.* 2017;109:33–9.
37. Kim Y, Chau TY, Rutledge JM, Erickson D, Lim Y. Factors that affect sugar sweetened beverage intake in rural, southern college students in the US. *Int J Vitam Nutr Res.* 2015;85(1–2):5–13.
38. Kim Y, Yang HY, Kim AJ, Lim Y. Academic stress levels were positively associated with sweet food consumption among Korean high-school students. *Nutrition.* 2013;29(1):213–8.
39. Prada M, Godinho CA, Garrido MV, Rodrigues DL, Coelho I, Lopes D. A qualitative study about college students' attitudes, knowledge and perceptions regarding sugar intake. *Appetite.* 2021;159:105059.
40. Association for University and College Counseling Center Directors. The Association for University and College Counseling Center Directors Annual Survey 2020. Available from: <https://www.ucccd.org/assets/documents/Survey/2019-2020%20Annual%20Report%20FINAL%204-2021.pdf>. Accessed 25 Jan 2022.
41. Chu TL, Zhang T. Sport club participation and health-related outcomes in college students: comparisons by sex and academic classification. *Recreat Sports J.* 2018;42(1):33–47.
42. Lipson SK, Lattie EG, Eisenberg D. Increased rates of mental health service utilization by U.S. college students: 10-year population-level trends (2007–2017). *Psychiatr Serv.* 2019;70(1):60–3.
43. Usher W, Curran C. Predicting Australia's university students' mental health status. *Health Promot Int.* 2019;34(2):312–22.
44. Barber BL, Eccles JS, Stone MR. Whatever happened to the jock, the brain, and the princess?: young adult pathways linked to adolescent activity involvement and social identity. *J Adolesc Res.* 2001;16(5):429–55.
45. Gore S, Farrell F, Gordon J. Sports involvement as protection against depressed mood. *J Res Adolesc.* 2001;11(1):119–30.
46. Sanders CE, Field TM, Diego M, Kaplan M. Moderate involvement in sports is related to lower depression levels among adolescents. *Adolescence.* 2000;35(140):793–7.
47. Donaldson SJ, Ronan KR. The effects of sports participation on young adolescents' emotional well-being. *Adolescence.* 2006;41(162):369–89.
48. Harrison PA, Narayan G. Differences in behavior, psychological factors, and environmental factors associated with participation in school sports and other activities in adolescence. *J Sch Health.* 2003;73(3):113–20.
49. Hiremath C. Impact of sports on mental health. *Int J Physiol Nutr Phys Educ.* 2019;4(S1):14–8.
50. Valois RF, Zullig KJ, Huebner ES, Drane JW. Physical activity behaviors and perceived life satisfaction among public high school adolescents. *J Sch Health.* 2004;74(2):59–65.
51. Inaç Y, Larivière Y, Hoque M, Van Hal G. Risk factors for hazardous drinking in university students from South Africa and Belgium: a cross-cultural comparison study. *Afr Health Sci.* 2021;21(1):123–31.
52. O'Brien KS, Kypri K. Alcohol industry sponsorship and hazardous drinking among sportspeople. *Addiction.* 2008;103(12):1961–6.
53. Partington S, Partington E, Heather N, Longstaff F, Allsop S, Jankowski M, et al. The relationship between membership of a university sports group and drinking behaviour among students at English universities. *Addict Res Theory.* 2013;21(4):339–47.
54. Quinn PD, Fromme K. Alcohol use and related problems among college students and their noncollege peers: the competing roles of personality and peer influence. *J Stud Alcohol Drugs.* 2011;72(4):622–32.
55. Lisha NE, Sussman S. Relationship of high school and college sports participation with alcohol, tobacco, and illicit drug use: a review. *Addict Behav.* 2010;35(5):399–407.
56. National Institute on Alcohol Abuse and Alcoholism. College drinking 2021. <https://www.niaaa.nih.gov/publications/brochures-and-fact-sheets/college-drinking>. Accessed 25 Jan 2022.
57. Lorant V, Nicaise P, Soto VE, d'Hoore W. Alcohol drinking among college students: college responsibility for personal troubles. *BMC Public Health.* 2013;13:615.
58. Borsari B, Carey KB. Peer influences on college drinking: a review of the research. *J Subst Abus.* 2001;13(4):391–424.
59. Michalak L, Trocki K. Alcohol and Islam: an overview. *Contemp Drug Probl.* 2006;33(4):523–62.
60. Ajayi AI, Owolabi EO, Olajire OO. Alcohol use among Nigerian university students: prevalence, correlates and frequency of use. *BMC Public Health.* 2019;19(1):752.
61. Luczak SE, Prescott CA, Dalais C, Raine A, Venables PH, Mednick SA. Religious factors associated with alcohol involvement: results from the Mauritian joint child health project. *Drug Alcohol Depend.* 2014;135:37–44.
62. Golden SD, McLeroy KR, Green LW, Earp JAL, Lieberman LD. Upending the social ecological model to guide health promotion efforts toward policy and environmental change. *Health Educ Behav.* 2015;42(1 Suppl):85–145.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions



The clusters of health-risk behaviours and mental wellbeing and their sociodemographic correlates: a study of 15,366 ASEAN university students

ORIGINALITY REPORT

20%

SIMILARITY INDEX

18%

INTERNET SOURCES

12%

PUBLICATIONS

8%

STUDENT PAPERS

PRIMARY SOURCES

1	www.burnet.edu.au Internet Source	2%
2	Submitted to Barry University Student Paper	1%
3	journals.plos.org Internet Source	1%
4	etd.uwc.ac.za Internet Source	1%
5	www.science.gov Internet Source	1%
6	www.semanticscholar.org Internet Source	1%
7	Hanif Abdul Rahman, Nurul Nazurah Julaini, Siti Nurzaimah Nazhirah Zaim, Nurfatin Amalina Masri, Khadizah H. Abdul-Mumin. "Mental wellbeing and Health-Risk behaviors of University students in Brunei: A cross-	1%

sectional study during COVID-19", Research Square Platform LLC, 2022

Publication

8	www.aunsec.org Internet Source	1 %
9	Submitted to CSU Northridge Student Paper	<1 %
10	essentials.ebsco.com Internet Source	<1 %
11	mafiadoc.com Internet Source	<1 %
12	www.frontiersin.org Internet Source	<1 %
13	Submitted to University of Greenwich Student Paper	<1 %
14	www.aftra.com Internet Source	<1 %
15	synapse.koreamed.org Internet Source	<1 %
16	www.journalofsports.com Internet Source	<1 %
17	www.ssph-journal.org Internet Source	<1 %
18	escholarship.mcgill.ca Internet Source	<1 %

19

[Submitted to University of Chichester](#)

Student Paper

<1 %

20

[aun-hpn.mahidol.ac.th](#)

Internet Source

<1 %

21

["Abstracts from the 30th Annual Meeting of the Society of General Internal Medicine", Journal of General Internal Medicine, 2007](#)

Publication

<1 %

22

[www.ukessays.com](#)

Internet Source

<1 %

23

[Linda Simoni-Wastila, Grant Ritter, Gail Strickler. "Gender and Other Factors Associated with the Nonmedical Use of Abusable Prescription Drugs", Substance Use & Misuse, 2004](#)

Publication

<1 %

24

[core.ac.uk](#)

Internet Source

<1 %

25

[sustainable.kmutt.ac.th](#)

Internet Source

<1 %

26

[www.jmir.org](#)

Internet Source

<1 %

27

[zaguan.unizar.es](#)

Internet Source

<1 %

28

[www.adscientificindex.com](#)

Internet Source

<1 %

29

aihd.mahidol.ac.th

Internet Source

<1 %

30

www.safetylit.org

Internet Source

<1 %

31

academic-accelerator.com

Internet Source

<1 %

32

dl.icdst.org

Internet Source

<1 %

33

theses.gla.ac.uk

Internet Source

<1 %

34

Elisabeth Rohwer, Natascha Mojtahedzadeh, Felix Alexander Neumann, Albert Nienhaus et al. "The Role of Health Literacy among Outpatient Caregivers during the COVID-19 Pandemic", International Journal of Environmental Research and Public Health, 2021

Publication

<1 %

35

e-journal.unair.ac.id

Internet Source

<1 %

36

pure.eur.nl

Internet Source

<1 %

37

Submitted to Saint Leo University

Student Paper

<1 %

38

bmcendocrdisord.biomedcentral.com

Internet Source

<1 %

39

en.wikipedia.org

Internet Source

<1 %

40

repositorioacademico.upc.edu.pe

Internet Source

<1 %

41

Submitted to 51347

Student Paper

<1 %

42

I Neethling, E V Lambert, A Cois, R A Roomaney, O F Awotiwon, R Pacella, D Bradshaw, V Pillay-van Wyk. "Estimating the changing burden of disease attributable to low levels of physical activity in South Africa for 2000, 2006 and 2012", South African Medical Journal, 2022

Publication

<1 %

43

lib.bioinfo.pl

Internet Source

<1 %

44

www.openagrar.de

Internet Source

<1 %

45

Submitted to Chulalongkorn University

Student Paper

<1 %

46

Dunyaporn Trachootham, Shizuko Satoh-Kuriwada, Aroonwan Lam-ubol, Chadamas Promkam et al. "Differences in Taste Perception and Spicy Preference: A Thai-Japanese Cross-cultural Study", *Chemical Senses*, 2018

Publication

<1 %

47

eprints.gla.ac.uk

Internet Source

<1 %

48

researchrepository.wvu.edu

Internet Source

<1 %

49

web.thepeninsulaqatar.com

Internet Source

<1 %

50

Eime, Rochelle M, Janet A Young, Jack T Harvey, Melanie J Charity, and Warren R Payne. "A systematic review of the psychological and social benefits of participation in sport for children and adolescents: informing development of a conceptual model of health through sport", *International Journal of Behavioral Nutrition and Physical Activity*, 2013.

Publication

<1 %

51

Li, Guowei, Alexandra Papaioannou, Lehana Thabane, Ji Cheng, and Jonathan D. Adachi. "Frailty change and major osteoporotic fracture in the elderly: data from the Global

<1 %

Longitudinal Study of Osteoporosis in Women
3-year Hamilton cohort", Journal of Bone and
Mineral Research, 2015.

Publication

52

tci-thaijo.org

Internet Source

<1 %

53

www.fao.org

Internet Source

<1 %

54

Chien Huey Teh, Ming Woey Teh, Kuang Hock Lim, Chee Cheong Kee et al. "Clustering of lifestyle risk behaviours and its determinants among school-going adolescents in a middle-income country: a cross-sectional study", BMC Public Health, 2019

Publication

<1 %

55

Tarek Tawfik Amin, Abudllah Mohammed Al-Hammam, Nasser Abdullah AlMulhim, Mohammed Ibrahim Al-Hayan et al. "Physical Activity and Cancer Prevention: Awareness and Meeting the Recommendations among Adult Saudis", Asian Pacific Journal of Cancer Prevention, 2014

Publication

<1 %

56

environhealthprevmed.biomedcentral.com

Internet Source

<1 %

57

sajcn.co.za

Internet Source

<1 %

58

tggs.kmutnb.ac.th

Internet Source

<1 %

59

tic.car.chula.ac.th

Internet Source

<1 %

60

www.globalhealthaction.net

Internet Source

<1 %

61

Choi Wan Chan, Sau Fong Leung. "Lifestyle Health Behaviors of Hong Kong Chinese", Asia Pacific Journal of Public Health, 2014

Publication

<1 %

62

Jae H. Kang, Walter C. Willett, Bernard A. Rosner, Susan E. Hankinson, Louis R. Pasquale. "Prospective Study of Alcohol Consumption and the Risk of Primary Open-Angle Glaucoma", Ophthalmic Epidemiology, 2009

Publication

<1 %

63

Li-Juan Liu, Xun Sun, Chun-Li Zhang, Yue Wang, Qiang Guo. "A survey in rural China of parent-absence through migrant working: the impact on their children's self-concept and loneliness", BMC Public Health, 2010

Publication

<1 %

64

Mohamed Hassan Elnaem, Naeem Mubarak, Mohammed Salim K. T., Muna Barakat et al. "Assessment of mental wellbeing of undergraduate pharmacy students from 14

<1 %

countries: The role of gender, lifestyle, health-related, and academic-related factors",
Frontiers in Public Health, 2022

Publication

65

Riaz Uddin, Nicola W. Burton, Asaduzzaman Khan. "Combined Effects of Physical Inactivity and Sedentary Behaviour on Psychological Distress Among University-Based Young Adults: a One-Year Prospective Study",
Psychiatric Quarterly, 2019

Publication

<1 %

66

Shuo Cheng, Cunxian Jia, Yongjie Wang. "Only Children Were Associated with Anxiety and Depressive Symptoms among College Students in China", International Journal of Environmental Research and Public Health, 2020

Publication

<1 %

67

aunsec.org

Internet Source

<1 %

68

bmjopen.bmj.com

Internet Source

<1 %

69

globalizationandhealth.biomedcentral.com

Internet Source

<1 %

70

pediatrics.aappublications.org

Internet Source

<1 %

repositorio.pgsskroton.com

71	Internet Source	<1 %
72	scholarworks.umt.edu Internet Source	<1 %
73	substanceabusepolicy.biomedcentral.com Internet Source	<1 %
74	wpc.org.pk Internet Source	<1 %
75	www.academicjournals.org Internet Source	<1 %
76	www.hindawi.com Internet Source	<1 %
77	www2.mdpi.com Internet Source	<1 %
78	Matthew YW Kwan, Guy EJ Faulkner, Kelly P Arbour-Nicitopoulos, John Cairney. "Prevalence of health-risk behaviours among Canadian post-secondary students: descriptive results from the National College Health Assessment", BMC Public Health, 2013 Publication	<1 %
79	Jay Thakkar, Julie Redfern, Aravinda Thiagalingam, Clara K Chow. "Patterns, predictors and effects of texting intervention on physical activity in CHD – insights from the	<1 %

TEXT ME randomized clinical trial", European Journal of Preventive Cardiology, 2016

Publication

80

Marit Müller De Bortoli, Anne Kristin Møller Fell, Martin Veel Svendsen, Paul K. Henneberger et al. "Lifestyle, sick leave and work ability among Norwegian employees with asthma—A population-based cross-sectional survey conducted in Telemark County, Norway", PLOS ONE, 2020

Publication

<1 %

81

bmcpalliatcare.biomedcentral.com

Internet Source

<1 %

82

Submitted to South Bank University

Student Paper

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On