

**MALAYSIAN JOURNAL OF
PUBLIC HEALTH
MEDICINE**



e-ISSN: 2590-3829

ISSN:1675-0306

Volume 22 No 1 2022

Official Publication of the

MALAYSIAN PUBLIC HEALTH PHYSICIANS' ASSOCIATION

Articles

HEALTH RELATED BEHAVIORAL CHANGES DURING COVID-19 MOVEMENT CONTROL ORDER (MCO) AMONG UNIVERSITY STUDENTS: A QUALITATIVE STUDY

Loganathan Salvaraji, Mohammad Saffree Jeffree, Ahmad Asyraf Abdul Rahim, Hemanath Sinnathamby, Mohd Fazeli Sazali, Nafsah Dulajis, Puteri Nureylia Amir, Haryati Abdul Karim, Freddie Robinson 1-15



PHARMQUIT: AN INTERACTIVE COMMUNITY PHARMACIST-ASSISTED SMARTPHONE APPLICATION FOR SMOKING CESSATION

Narong Asayut, Phayom Sookaneknun Olson, Juntip Kanjanasilp, Preut Thanarat, Bhattaraporn Senkraigul 16-27



FACTORS ASSOCIATED WITH ASYMPTOMATIC COVID-19 INFECTION IN A DISTRICT SETTING IN MALAYSIA

Asraf Qamruddin, Sophia Rasheeqa Ismail, Awanis Sharif, Reza Murad Qamruddin 28-37



KNOWLEDGE, ATTITUDE AND PRACTICES (KAP) THEORY TOWARDS PREVENTIVE MEASURES AMONG MALYSIAN IN EARLY OUTBREAK OF COVID-19

Nurhasniza Idham Abu Hasan, Shahrman Zainal Abidin, Mogana Darshini Ganggayah, Nur Faezah Jamal, Wan Noor Hayatie Wan Abdul Aziz 38-47



EXPLORING AND DEVELOPING ITEMS MEASURING QUALITY OF LIFE CONSTRUCT AMONG HEART FAILURE PATIENTS: A CASE OF MALAYSIA

Wan Marina Ismail, Zainuddin Awang, Idayubadilla Idris 48-55



PSYCHOLOGICAL DISTRESS AMONG MOTHERS OF CHILDREN WITH EMOTIONAL AND BEHAVIORAL PROBLEMS DURING COVID-19 PANDEMIC IN MALAYSIA

Ruziana Masiran, Normala Ibrahim, Hamidin Awang, Poh Ying Lim 56-67



RISK PERCEPTION IN CHILDHOOD VACCINATION QUESTIONNAIRE: A VALIDATION STUDY ON PREGNANT MOTHERS IN MALAYSIA

Mohd Nazir Mohd Nazori, Rohani Ismail, Nur Syahmina Rasudin, Rosminah Mohamed, Norhayati Mohd Noor, Zailiza Sulli, Masitah Mohamed, Noor Azliah Ahmad Zainuri, Ashvini Jayapalan 68-78



UTILIZATION AND PERCEPTION OF COMPLEMENTARY AND ALTERNATIVE THERAPIES (CATS) AMONG SECONDARY SCHOOL STUDENTS IN KUALA TERENGGANU

Pei Lin Lua, Nurul Afiedia Roslim, Aryati Ahmad, Mardiana Mansor, Myat Moe Thwe Aung, Farrahdilla Hamzah 79-85



CORRELATION BETWEEN APPETITIVE TRAITS AND DIET QUALITY OF YOUNG CHILDREN IN NORTH JAKARTA

Alfi Rahma Putri, Dian Novita Chandra, Luh Ade Ari Wiradnyani

86-94

 PDF

THE CORRELATION OF ENVIRONMENTAL FACTORS WITH THE PREVALENCE OF ATOPIC DISEASE IN ELEMENTARY SCHOOL CHILDREN

Asti Widuri

95-99

 PDF

CHARACTERISTICS OF PEOPLE WITH INFERTILITY IN INDONESIA REFLECTED IN FERTIQOL SCORES

Fitri Damayanti, Mohammad Hakim, Mochamad Anwar, Diah Ayu Puspandari

100-108

 PDF

NOVEL TECHNIQUE OF MASS TRIAGING OF COVID 19 POSITIVE PATIENT: SABAH EXPERIENCE

Khoo Elaine, Aethoosham Bin Suleman, Jolina Wilfred Nini, Mohd Ariff Hasreen Bin Ahlun, Shaik Farid Abdull Wahab

109-115

 PDF

IMPACT OF OCCUPATIONAL FACTORS ON SAFETY AND HEALTH THROUGH ERGONOMIC APPROACH AMONG OLDER TAXI DRIVERS IN MALAYSIA

Irwan syah Md Yusoff, Ismail, M.H, Rohaizahtulamni, R

116-123

 PDF

COPING STRATEGIES, SELF-EFFICACY, AND PERCEPTION AMONG FAMILY CAREGIVERS FOR POST-STROKE SURVIVORS IN INDONESIA

Vione Deisi Oktavina Sumakul, Hari Basuki Notobroto, Shrimarti Rukmini Devy

124-130

 PDF

USING THEORY OF PLANNED BEHAVIOR TO EXPLORE FACTORS RELATED TO CONTRACEPTIVE USE AMONG WOMEN IN REPRODUCTIVE AGE

Retno Heru Setyorini, Ismi Dwi Astuti Nurhaeni, Ari Probandari, Eny Lestari

139-145

 PDF

Malaysian Journal of Public Health Medicine

[Current](#) [Archives](#) [About](#) [Authors Guidelines](#) [Ethics Statement](#) [Contact](#)

[Home](#) / [Editorial Team](#)

Chief Editor:

Professor Dato' Dr. Syed Mohamed Aljunid
National University of Malaysia

Deputy Chief Editor:

Professor Dr. Mohamed Rusli Abdullah
Perdana University - Royal College of Surgeons in Ireland

Editorial Board Members:

Professor Datuk Dr. Lokman Hakim Sulaiman
International Medical University

Professor Dr. Retneswari Masilamani
Universiti Tunku Abdul Rahman Malaysia

Professor Dr. Saperi Sulong
National University of Malaysia

Assistant Professor Dr. Amrizal Muhd Nur
Kuwait University

Professor Dr. Maznah Dahlui
University of Malaya

Dr. Roslan Johari
Ministry of Health Malaysia

Dr. Othman Warijo
Ministry of Health Malaysia

Dr. Amin Sah bin Ahmad
Ministry of Health Malaysia

Dr. Ghazali bin Chik
Ministry of Health Malaysia

Dr. Sabrina binti Che Abd Rahman
Ministry of Health Malaysia

Dr. Mariam binti Mohamad
Universiti Teknologi MARA (UITM)

Dr. Tahir bin Aris
Ministry of Health Malaysia

Dr. Mahanim Md. Yusof
Ministry of Health Malaysia

Dr. Mohd Azahadi bin Omar
Ministry of Health Malaysia

Professor Dr. Jamalludin Ab Rahman
International Islamic University Malaysia

Indexed By

Scopus®

REVIEW ARTICLE

COPING STRATEGIES, SELF-EFFICACY, AND PERCEPTION AMONG FAMILY CAREGIVERS FOR POST-STROKE SURVIVORS IN INDONESIA

Vione Deisi Oktavina Sumakul¹, Hari Basuki Notobroto*² and Shrimarti Rukmini Devy²

¹Stikes Gunung Maria, Tomohon, Indonesia.

²Department of Epidemiology, Biostatistics, Population Studies, and Health Promotion, Faculty of Public Health, Universitas Airlangga, Surabaya, Indonesia.

*Correspondence author: Hari Basuki Notobroto

E-mail: haribasuki.n@fkm.unair.ac.id.

ABSTRACT

Family caregivers experience stress and burden while caring for post-stroke survivors. Coping strategies are needed to deal with the stress, especially problem emotional-focused coping. The stress adaptation model states that coping was influenced by self-efficacy. Meanwhile, ABC Hill model explained that perception also had effects on coping. The purpose of this study was to analyze factors that affected coping among family caregivers for post-stroke survivors, especially self-efficacy and caregiver perception. This study used cross-sectional design and also used self-report questionnaire. It was conducted from October 2020 - April 2021. Sample of this study was family members who take care the family member with post-stroke, and the post-stroke survivors already post hospitalized between 6 months to 3 years. Total sample was 200 family caregivers in 7 health care centers, Tomohon City, Indonesia. We used PLS to estimate the hypothesized fit with the data. T statistic was used to include or exclude the hypotheses (t -statistic >1.96). All path coefficients were significant. Self-efficacy had a positive effect on problem emotional-focused coping ($\beta = 0.177$; t -value = 0.030, $R^2 = 0.078$). In addition, the caregiver perception had a positive effect on problem emotional-focused coping ($\beta = 0.188$, t -value = 2.248, $R^2 = 0.175$). Managing self-efficacy and perception among caregivers are needed to improve problem emotional-focused coping.

Keywords: coping strategies, self-efficacy, perception, family caregivers, stroke.

INTRODUCTION

The global prevalence of stroke remains high. There were people who died from stroke every 3 minutes 33 seconds. Also, a total of 3.3 million people died from ischemic stroke, 2.9 million died from intracerebral hemorrhage, and 0.4 million died from subarachnoid hemorrhage¹. Each year, 60% of all strokes occur in people under the age of 70 years, 8% occur in people under the age of 44 years. In addition, 52% of all strokes occur in men, 48% of all strokes occur in women².

Basic Health Research (RISKESDAS) in Indonesia showed that the highest prevalence of stroke was recorded in Yogyakarta (14.6%), followed by North Sulawesi (14.0%), the highest prevalence by age was 75 years old (50.2%); based on gender was 11.0% male and 10.9% female, In addition, the residence in the city 12.6% was higher than in the village³. In North Sulawesi, the highest prevalence of age group suffering from stroke was at the age of 65-74 years⁴.

Stroke is the fourth leading cause of death and disability. Nearly 50% of post-stroke patients

experienced moderate to severe disorders, paralysis and were unable to care for themselves⁵. Stroke survivors need a caregiver after being discharged from the hospital, such as a spouse or family member. Family caregiver assists their daily living, health care, finance, as well as social interaction⁶⁻⁸. Furthermore, family caregivers are the key to caring for post-stroke survivor⁹. The important aspect of caring for post-stroke survivors are to promote early recovery and to prevent complications. If the patient receives the appropriate rehabilitation within the golden period, the stroke recovery will improve significantly¹⁰⁻¹³. However, caring for post-stroke survivors was complicated and different from other chronic diseases. Family caregivers must have knowledge and skills regarding prevention of complications, activities of daily living (ADL), mobility, rehabilitation, and management of stress¹⁴⁻¹⁶. Family caregivers experience stress and burden while caring for post-stroke survivors¹⁷⁻¹⁹. The prevalence of mental health problems among caregiver was 3% to 62% for post traumatic stress

disorder, 4% to 94% for depression, and 2% to 80% for anxiety²⁰. Coping is needed to face these situations. Coping refers to problem-emotion focused coping. Problem focused coping is an active caregiver's effort to deal with stressful conditions and eliminate sources of stress through individual behavior. While emotion focused coping was passive effort to reduce stressful condition through emotional²¹.

Based on this background, the purpose of this study was to analyze factors that affected coping among family caregivers for post-stroke survivors. This research is needed as a theory to improve the family caregiver's quality of life as well as to reduce stress.

The stress adaptation model states that coping was influenced by self-efficacy. Self-efficacy is our ability to cope with the demands of life²². Meanwhile, ABC Hill model developed by McCubbin and Patterson (1983) explained that perception also had effects on coping²³. Perception is a positive or negative interpretation regarding stress when caring for family members with post-stroke²⁴. In addition, the ability to cope with the demands of life had effect on negative or positive perception. More self efficacy influences the effort to achieve the goals and had effect on positive perception²⁵. We combined these two theories for developing new theory.




Figure 1. Study framework

METHODS

All of procedure in this study was granted by ethical clearance commission from Faculty of Dental Medicine, Universitas Airlangga, number: 463/HRECC.FODM/X/2020. This study used cross-sectional design with explanatory research and also used self-report questionnaire. It was conducted from October 2020 - April 2021. Sample of this study was family members who take care for family member post-stroke, and the family members were already post hospitalized between 6 months to 3 years. Inclusion criteria were family caregiver who is living together in the same house with post-stroke survivor and only caring for 1 family member with post-stroke. Total sample was 200 family caregiver respondents in 7 health care centers in Tomohon City, Indonesia: Puskesmas Pangolombian, Puskesmas Lansot, Puskesmas Matani, Puskesmas Rurukan, Puskesmas Tara Tara, Puskesmas Tinoor, Puskesmas Kakaskasen. We distributed the offline questionnaires door to door based on data from these primary health cares. The family members who agreed to join this study must signed the informed consent and the allowed to withdraw after reading the questionnaire. The family members who finished and completed to fill the questionnaire got

reward. Only researcher who can access the result of these questionnaires.

Measures

The questionnaires consisted of demographic data, age, level of education, economic status, level of knowledge, status caregiver in the family, number of family members. We used a self-made questionnaire. We had 23 parameters, so we recruited 115 caregivers. Five cases per free parameter in the model²⁶⁻²⁸. We also used caregiver perception questionnaire. This questionnaire was used to measure the caregiver's interpretation of the situations while taking care the family members post-stroke. Perception had 6 items. The Cronbach's alpha of this questionnaire was 0.91. We also used self-efficacy which was used to measure the caregiver's ability to achieve the goals. Self-efficacy had 3 domains: belief, consistency and capability. Belief had 6 items, consistency had 1 item, and capability had 3 items. The Cronbach's alpha of this questionnaire was 0.92. We used 7 items for assessing problem emotional-focused coping. This questionnaire was used to ask how the care giver solves problems, adaptation, and responds to threatening situations. The Cronbach's alpha of this questionnaire was 0.95. The cronbach

alpha of this questionnaire was 0.95. Previous studies suggested that cronbach's alpha > 0.5 is considered accepted reliability²⁹⁻³¹.

All the data were analyzed using SPSS computer software. Descriptive statistics were used for the variables. We used PLS to estimate the hypothesized fit with the data. T statistic was used to include or exclude the hypotheses (t-statistic >1.96).

RESULTS

Table 1 presents the characteristic of all participants. Regarding participant's age was predominant by participant's age more than 40 age years old (43.5%). Most of the participants had senior high school degree (47.5%). Regarding economic status, most of the participants had region minimum wage (55.5%). In addition, 26% of respondents had three members in their family.

Table 1. Distribution of participant characteristics (N= 200)

Characteristic	n	%
Age (years)		
< 19	30	15.0 %
20-30	37	18.5 %
31-40	46	23.0 %
>40	87	43.5 %
Total	200	100.0 %
Education degree		
Elementary school	41	20.5 %
Junior high school	43	21.5 %
Senior high school	95	47.5 %
High school degree	21	10.5 %
Total	200	100.0 %
Economic status		
< region minimum wage	111	55.5 %
> region minimum wage	89	44.5 %
Total	200	100.0 %
Level of knowledge		
Low	31	15.5 %
Moderate	78	39.0 %
Good	91	45.5 %
Total	200	100.0 %
Caregiver		
Husband	59	29.5 %
Wife	69	34.5 %
Children	48	24.0 %
Sibling	13	6.5 %
Others	11	5.5 %
Total	200	100.0 %
Number of family members		
2	30	15.0 %
3	52	26.0 %
4	51	25.5 %
5	29	14.5 %
6	26	13.0 %
7	12	6.0 %
Total	200	100.0 %

Table 2 provides information on the self-efficacy, caregiver perception and coping strategies. Most of the respondents experienced positive self-efficacy

(81.5%), positive experience (positive perception) (86%), and used problem-based coping strategies (95%).

Table 2. Description of the observed variables among family caregivers who take care their family post-stroke

Variables	n	%
Self-efficacy		
Negative	37	18.5 %
Positive	163	81.5 %
Total	200	100.0 %
Perception		
Negative	28	14.0 %
Positive	172	86.0 %
Total	200	100.0 %
Problem Emotional -Focused Coping		
Emotional-based coping	10	5.0 %
Problem-based coping	190	95.0 %
Total	200	100.0 %

Table 3 lists path coefficient, and t-statistic value coefficient. All path coefficients were significant, except self-efficacy →caregiver perception. Figure 1 shows the final model, with the statistical goodness of fit was 0.380.

Table 3. Summary of model graph result

Hypotheses test	Coefficient	t-value	p-value	Remark
Self-efficacy →caregiver perception	0.039	1.86	> 0.05	Not significant
<i>Self-Efficacy</i> → coping strategy	0.177	2.183	0.030	Significant
Caregiver perception→ coping strategy	0.188	2.248	0.025	Significant




Figure 1. Final model of investigated self-efficacy and caregiver perception that affect coping strategamong caregiver

DISCUSSION

The aim of this study was to analyze factors that affecting coping strategies among family caregiver.

This study involved 200 family caregivers who take care the family members with post-stroke in Tomohon City, Indonesia. The results of this study indicated that most of family caregivers were 40

age years old. This data was similar with the data in the United States, the average caregiver was over 40 years old³². In addition, family caregivers who take care of family members with post-stroke were generally spouses and children. Data in the United States shows that most family caregivers are spouses, parents, or children³². The majority of family caregivers had income less than the minimum wage. This situation had effect on financial burden among caregivers and also affected their ability to obtain adequate access to health care facilities³³. This study showed that the level of knowledge of family caregiver to take care the family member with post stroke was good. Knowledge, abilities and skills are needed to take care the family members with post-stroke³⁴. So that it can reduce caregiver stress³⁵.

The Effect of Self Efficacy on Problem-Emotion Coping Strategies

Self-efficacy had a positive effect on problem-emotion coping strategies ($\beta = 0.177$; $t\text{-value} = 2.183$, $R^2 = 0.078$). The results also showed that most caregivers who have positive self-efficacy had problem-based coping strategies (77%). Problem-based coping strategies were strongly influenced by self-efficacy. This study is in line with previous research which describes a significant positive relationship between self-efficacy and problem-based coping³⁶. The higher the self-efficacy, the higher the problem-based coping. Individuals who have positive self-efficacy will tend to perceive the problem as a challenge, being tolerant, and able to overcome the obstacles. Individuals with positive self-efficacy and problem-based coping tend to improve the situation, while individuals who have negative self-efficacy tend to choose emotional-based coping strategies³⁷.

The Effect of Caregiver Perception on Problem-Emotion Coping Strategies

The caregiver perception had a positive effect on problem-emotion coping strategy ($\beta = 0.188$, $t\text{-value} = 2.248$, $R^2 = 0.175$). The results showed that the caregiver's perception was positive when caring for family members who stroke survivors for 11-20 months and had a mild level of dependence. Perception is the way we understand or interpret the stressor. Negative perceptions can influence behavior³⁸. When the caregiver had a positive perception, it increases the motivation for caring the family members with post-stroke. The caregiver will make good efforts so the family can recover. Limitation of this study was our study used a cross-sectional design, therefore, the causal direction implied by the path analysis must be interpreted carefully. Longitudinal studies are needed to examine these factors and interventions are needed to manage self-efficacy and family caregiver

perception. In addition, these caregivers only take care 1 family member with post stroke, so it cannot be generalized.

CONCLUSION

Coping strategies among family caregivers were influenced by perception and self-efficacy. Increasing perception and self-efficacy among caregivers are needed to improve problem-emotion coping strategies. Through these coping strategies, caregivers tend to tolerate with life stressors.

ACKNOWLEDGEMENT

We would like to thank for the funding from Stikes Gunung Maria, Tomohon, Indonesia.

REFERENCES

1. Association AH. Heart Disease and Stroke Statistics Update Fact Sheet At-a-Glance 2021 [cited 2021 13 July]. Available from: https://www.heart.org/-/media/phd-files-2/science-news/2/2021-heart-and-stroke-statupdate/2021_heart_disease_and_stroke_statistics_update_fact_sheet_at_a_glance.pdf.
2. Lindsay MP, Norrving B, Sacco RL, et al. World Stroke Organization (WSO): global stroke fact sheet 2019. SAGE Publications Sage UK: London, England; 2019.
3. RI K. Laporan hasil riset kesehatan dasar (Riskesmas) Indonesia tahun 2018. Jakarta: Badan Penelitian dan Pengembangan Kesehatan Kemenkes RI. 2018:5-10.
4. Riskesdas T. Riskesdas 2018: Laporan Provinsi Sulawesi Utara. Lembaga Penerbit Badan Penelitian dan Pengembangan Kesehatan Kementerian Kesehatan RI. 2018.
5. Kuptniratsaikul V, Kovindha A, Suethanapornkul S, et al. Long-term morbidities in stroke survivors: a prospective multicenter study of Thai stroke rehabilitation registry. *BMC geriatrics*. 2013;13(1):1-9.
6. Blake H, Lincoln NB, Clarke DD. Caregiver strain in spouses of stroke patients. *Clinical rehabilitation*. 2003;17(3):312-7.
7. Kaminishi KS, Safavi R, Hirsch CH. Caregiver burnout. *Geriatric Psychiatry: Springer*; 2018. p. 691-708.

8. Association AH. What Is Caregiver Burnout? 2012 [cited 2021 13 July]. Available from: <https://www.stroke.org/-/media/files/health-topics/answers-by-heart/what-is-caregiver-burnout.pdf?la=en>.
9. Han B, Haley WE. Family caregiving for patients with stroke: review and analysis. *Stroke*. 1999;30(7):1478-85.
10. Bernhardt J, Dewey H, Thrift A, et al. Inactive and alone: physical activity within the first 14 days of acute stroke unit care. *Stroke*. 2004;35(4):1005-9.
11. Belagaje SR. Stroke rehabilitation. *CONTINUUM: Lifelong Learning in Neurology*. 2017;23(1):238-53.
12. Stinear CM, Lang CE, Zeiler S, et al. Advances and challenges in stroke rehabilitation. *The Lancet Neurology*. 2020;19(4):348-60.
13. Good DC, Bettermann K, Reichwein RK. Stroke rehabilitation. *CONTINUUM: Lifelong Learning in Neurology*. 2011;17(3):545-67.
14. Rhoda A. Knowledge of nurses regarding dysphagia in patients post stroke in Namibia. *curationis*. 2015;38(2):1-7.
15. Mackenzie A, Perry L, Lockhart E, et al. Family carers of stroke survivors: needs, knowledge, satisfaction and competence in caring. *Disability and rehabilitation*. 2007;29(2):111-21.
16. Zeiler SR. Should we care about early post-stroke rehabilitation? Not yet, but soon. *Current neurology and neuroscience reports*. 2019;19(3):1-9.
17. Menon B, Salini P, Habeeba K, et al. Female caregivers and stroke severity determines caregiver stress in stroke patients. *Annals of Indian Academy of Neurology*. 2017;20(4):418.
18. Yigitalp G, Surucu HA, Gumus F, et al. Predictors of caregiver burden in primary caregivers of chronic patients. *International Journal of Caring Sciences*. 2017;10(3):1168.
19. Caro CC, Costa JD, Da Cruz DMC. Burden and quality of life of family caregivers of stroke patients. *Occupational therapy in health care*. 2018;32(2):154-71.
20. Johnson CC, Suchyta MR, Darowski ES, et al. Psychological sequelae in family caregivers of critically ill intensive care unit patients. A systematic review. *Annals of the American Thoracic Society*. 2019;16(7):894-909.
21. Lazarus RS, Folkman S. *Stress, appraisal, and coping*: Springer publishing company; 1984.
22. Stuart G. *Psychiatric Of Nursing Edisi 9*. Mosby: USA Elsevier. 2013.
23. McCubbin HI, Patterson JM. The family stress process: The double ABCX model of adjustment and adaptation. *Marriage & family review*. 1983;6(1-2):7-37.
24. Yeh PM, Chang Y. Use of Zarit Burden Interview in analysis of family caregivers' perception among Taiwanese caring with hospitalized relatives. *International journal of nursing practice*. 2015;21(5):622-34.
25. Eydi H, Aghaie N. The Study of Effect of Hardiness and Self-Efficacy on Perception of Success of Women's karate in Kermanshah Province. *Sport Management Studies*. 2021;13(65).
26. Tabachnick BG, Fidell LS, Ullman JB. *Using multivariate statistics*: Pearson Boston, MA; 2007.
27. Kline RB. *Principles and practice of structural equation modeling*: Guilford publications; 2015.
28. Chou C-P, Bentler PM. *Estimates and tests in structural equation modeling*. 1995.
29. Morera OF, Stokes SM. Coefficient α as a measure of test score reliability: Review of 3 popular misconceptions. *American journal of public health*. 2016;106(3):458-61.
30. Streiner DL. Starting at the beginning: an introduction to coefficient alpha and internal consistency. *Journal of personality assessment*. 2003;80(1):99-103.
31. Setyowati A, Chung M-H, Yusuf A, et al. Psychometric of the curiosity and exploration inventory-ii in Indonesia. *Journal of Public Health Research*. 2020;9(3).

32. Alliance FC. Caregiver Statistics: Demographics 2021 [Available from: <https://www.caregiver.org/resource/care-giver-statistics-demographics/>].
33. Andrén S, Elmståhl S. Relationships between income, subjective health and caregiver burden in caregivers of people with dementia in group living care: a cross-sectional community-based study. *International Journal of Nursing Studies*. 2007;44(3):435-46.
34. Lutz BJ, Young ME. Rethinking intervention strategies in stroke family caregiving. *Rehabilitation Nursing*. 2010;35(4):152-60.
35. Gallagher-Thompson D, Wang P-C, Liu W, et al. Effectiveness of a psychoeducational skill training DVD program to reduce stress in Chinese American dementia caregivers: results of a preliminary study. *Aging & mental health*. 2010;14(3):263-73.
36. Salas BL, Rodríguez VY, Urbieta CT, et al. The role of coping strategies and self-efficacy as predictors of life satisfaction in a sample of parents of children with autism spectrum disorder. *Psicothema*. 2017;29(1):55-60.
37. Devonport TJ, Lane AM. Relationships between self-efficacy, coping and student retention. *Social Behavior and Personality: an international journal*. 2006;34(2):127-38.
38. Liu Y, Wei M, Guo L, et al. Association between illness perception and health behaviour among stroke patients: The mediation effect of coping style. *Journal of Advanced Nursing*. 2021;77(5):2307-18.