

Clinical characteristics and outcomes of hospitalized COVID-19 patients with diabetes mellitus in East Java, Indonesia A cross-sectional study

by Erwin Astha Triyono

Submission date: 28-Dec-2022 11:23AM (UTC+0800)

Submission ID: 1987012549

File name: tes_mellitus_in_East_Java,_Indonesia_A_cross-sectional_study.pdf (488.14K)

Word count: 5633

Character count: 31444



RESEARCH ARTICLE

Clinical characteristics and outcomes of hospitalized COVID-19 patients with diabetes mellitus in East Java, Indonesia: A cross-sectional study [version 1; peer review: 2 approved]

Erwin Astha Triyono¹, Joni Wahyuhadi², Jongky Hendro Prajitno¹,
 Hermina Novida¹, Nenci Siagian¹, Cupuwatie Cahyani¹, Arinditia Triasti Putri¹,
 Michael Austin Pradipta Lusida¹, Amal Arifi Hidayat¹,
 Karisma Septari Idamusaga¹, Nastiti Imana Intansari³, Jose Asmara⁴,
 Agrasenfani Hadi⁴, I Ketut Mega Purnayasa Bandem⁴

¹Department of Internal Medicine, Faculty of Medicine, Dr. Soetomo Teaching Hospital, Airlangga University, Surabaya, 60115, Indonesia

²Department of Neurosurgery, Faculty of Medicine, Dr. Soetomo Teaching Hospital, Airlangga University, Surabaya, 60115, Indonesia

³Department of Cardiology, Faculty of Medicine, Dr. Soetomo Teaching Hospital, Airlangga University, Surabaya, 60115, Indonesia

⁴Indrapura Forefront Hospital, Surabaya, 60175, Indonesia

V1 First published: 21 Jun 2022, 11:684
<https://doi.org/10.12688/f1000research.111047.1>
 Latest published: 21 Jun 2022, 11:684
<https://doi.org/10.12688/f1000research.111047.1>

Abstract

Introduction: Diabetes mellitus has been perceived as the worsening factor for coronavirus disease 2019 (COVID-19), where diabetes mellitus patients with pre-existing inflammatory condition could develop acute respiratory disease syndrome as well as multi-organ dysfunction. Managing diabetes mellitus amidst severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection is also a matter of concern as several antidiabetic therapies could affect the progression of COVID-19. This study aimed to provide the clinical characteristics and outcomes of patients with both COVID-19 and diabetes mellitus receiving blood glucose lowering therapies and COVID-19 symptomatic treatments.

Methods: This retrospective study was performed on 260 medical records of patients hospitalized between May 2020 to February 2021 in East Java, Indonesia. Patients were confirmed COVID-19 positive based on the results from real time polymerase chain reaction (RT-PCR) using nasal swab samples collected on hospital admission. Data included were demographic characteristics, COVID-19 symptoms, severity of COVID-19, comorbidities (other than diabetes mellitus), fasting blood glucose (FBG), and 2-hours post-prandial blood glucose (2hPBG), and outcomes.

Results: Most of the patients had age range of 41–60 years old

Open Peer Review

Approval Status

	1	2
version 1	✓	✓
21 Jun 2022	view	view

1. **Kurnia Fitri Jamil**, Universitas Syiah Kuala, Banda Aceh, Indonesia
2. **Kuldeep Dhama**, ICAR-Indian Veterinary Research Institute, Bareilly, India

Any reports and responses or comments on the article can be found at the end of the article.

(76.1%) with more than a half of the subjects (60%) were obese. Patients with uncontrolled diabetes were distributed evenly among the COVID-19 severities (74.3% in asymptomatic group, 73.6% in mild group, and 74.1% in moderate group). There were reductions in FBG and 2hPBG levels measured before (210.75 ± 81.38 and 271.19 ± 100.7 mg/dL, respectively) and after the treatment (181.03 ± 68.9 and 222.01 ± 86.96 mg/dL, respectively). All patients received multivitamin and symptomatic treatment for COVID-19. Oral antidiabetic drug (57.6%) and insulin (28.8%) were administered to lower the blood glucose level of the patients. As many as 96.9% patients survived, while 3.1% died.

Conclusion: COVID-19 could affect the blood glucose level, suggesting the importance of antihyperglycemic therapies among patients with both COVID-19 and diabetes mellitus.

Keywords

ACE2, blood glucose, diabetes mellitus, hyperglycemia, SARS-CoV-2



23

This article is included in the **Emerging Diseases and Outbreaks** gateway.

48

Corresponding author: Erwin Astha Triyono (erwintriyono@yahoo.com)

Author roles: **Triyono EA:** Conceptualization, Data Curation, Investigation, Methodology, Project Administration, Resources, Supervision, Visualization, Writing – Original Draft Preparation; **Wahyuhadi J:** Data Curation, Investigation, Resources, Validation, Writing – Review & Editing; **Prajitno JH:** Data Curation, Methodology, Resources, Writing – Original Draft Preparation; **Novida H:** Data Curation, Methodology, Resources, Writing – Review & Editing; **Siagian N:** Data Curation, Methodology, Resources, Writing – Review & Editing; **Cahyani C:** Data Curation, Methodology, Resources, Writing – Review & Editing; **Putri AT:** Data Curation, Methodology, Resources, Writing – Review & Editing; **Lusida MAP:** Data Curation, Methodology, Resources, Writing – Review & Editing; **Hidayat AA:** Data Curation, Methodology, Resources, Writing – Review & Editing; **Idamusaga KS:** Data Curation, Methodology, Resources, Writing – Review & Editing; **Intansari NI:** Data Curation, Methodology, Resources, Writing – Review & Editing; **Asmara J:** Data Curation, Methodology, Resources, Writing – Review & Editing; **Hadi A:** Data Curation, Methodology, Resources, Writing – Review & Editing; **Bandem IKMP:** Data Curation, Methodology, Resources, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2022 Triyono EA *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Triyono EA, Wahyuhadi J, Prajitno JH *et al.* **Clinical characteristics and outcomes of hospitalized COVID-19 patients with diabetes mellitus in East Java, Indonesia: A cross-sectional study [version 1; peer review: 2 approved]** F1000Research 2022, 11:684 <https://doi.org/10.12688/f1000research.1111047.1>

First published: 21 Jun 2022, 11:684 <https://doi.org/10.12688/f1000research.1111047.1>

Introduction

A novel coronavirus severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) that causes coronavirus disease 2019 (COVID-19) has been responsible to almost 153,000 mortalities in Indonesia as of March 2022 based on Indonesian Government database and has caused many disruptions in the communities.^{1,2} SARS-CoV-2 uses angiotensin-converting enzyme 2 (ACE2) as the receptor, in which the enzyme is available in various organs (such as lungs, heart, kidneys, intestines, and so on).³ Diabetes mellitus is a condition responsible for high number of global morbidities, especially due to vascular diseases it induces through chronic inflammation. As any other underlying diseases, diabetes mellitus could contribute to the poor prognosis of COVID-19.⁴⁻⁶ This is ascribed to the role of inflammation in the pathogenesis of severe COVID-19, where chronic inflammation is a common condition in individuals with diabetes mellitus.⁴ Common comorbidities found in diabetes mellitus, such as obesity and hypertension, have been evidenced to be responsible for acute respiratory disease syndrome (ARDS) as well as multi-organ dysfunction.⁵

Understanding clinical characteristics of diabetes mellitus of patients who are infected with SARS-CoV-2 is important to provide proper management. For example, those with poor glycemic control could have worse viral infections, as proven by SARS and influenza H1N1 cases.⁴ Respiratory distress induced by the viral infection could lead to the apoptosis of pancreatic beta cells that consequently causes insulin insufficiency.⁷ Therefore, glucose-lowering therapies should be continued or performed during the COVID-19 management. Nonetheless, clinicians should also consider the pro-inflammatory effect of some antidiabetic drugs which could contribute to the progression of severe COVID-19. Thiazolidinediones (TZDs) is one of the antidiabetic therapies that has been found to induce inflammation by elevating ACE2 and angiotensin 1-7 expressions.⁸ Moreover, in a meta-analysis of 13 trials, increased risk of developing pneumonia was found in TZD group.⁹ These explanations suggest that there are strong associations between COVID-19 and both diabetes mellitus and its management.

In the case of COVID-19, based on a meta-analysis, the number of patients with diabetes mellitus could reach 8% of the total patients.¹⁰ However, the prevalence was dramatically higher (36%) in Italian population, where 34% of which died during the treatment.¹¹ In Indonesia itself, the data from Jakarta province (n=20,481) revealed that the prevalence was only 3.4%, but the mortality rate was higher in diabetes mellitus group (21.28%) than that in non-diabetes mellitus group (2.77%).¹² By using a larger data set from the Indonesian COVID-19 Task Force, a study revealed that diabetes mellitus as the second most common comorbidity (33.6%) after hypertension (52.1%).¹³ Taken altogether, it is still uncertain whether the prevalence of diabetes mellitus among COVID-19 patients, especially in Indonesia, is high. Herein, we reported the data from East Java Province, Indonesia, regarding the clinical characteristics of the COVID-19 and diabetes mellitus patients and the treatment they received during the hospitalization. Moreover, we also reported the outcomes from symptomatic management for COVID-19 in combination with diabetes mellitus therapies which could be recommended for hospitals in developing countries with limited medical and financial resources.

Methods

Study design and setting

This study was a retrospective study using the medical records of COVID-19 patients with diabetes mellitus who underwent a hospitalization from May 2020 to February 2021 in Indrapura Forefront Hospital Surabaya, East Java Province – Indonesia (n=260). Patients were confirmed COVID-19 positive by real time polymerase chain reaction (RT-PCR) through nasal swab admission. Diabetes mellitus was confirmed by the officially recorded medical history. The obtained data were collected from electronic medical records including demographic, treatments, laboratory results as well as clinical outcomes. All patients with COVID-19 enrolled in this study were diagnosed and managed according to the national guidelines. This study had received ethical approval from the Ethics Committee of the Faculty of Medicine, Airlangga University, Surabaya, Indonesia (registration number: 37/EC/KEPK/FKUA/2021). Since this study collected the data from the available medical records, the Ethics Committee waived of consent from the patients and this allows by the Indonesian law.

Study measures

Data used in this study were collected from electronic medical records. Demographic characteristics of the patients included age, sex, and occupation. Clinical characteristics extracted from the medical records were body mass index (BMI), comorbidities, and COVID-19 symptoms. Comorbidities included hypertension, cardiovascular diseases, chronic kidney diseases, and asthma. As for the COVID-19 symptoms, they were cough, fever, rhinorrhea, anosmia, dyspnea, nausea, and headache. Severity of the COVID-19 was classified following the national guideline; asymptomatic, mild, moderate, and severe. Patients showing no COVID-19 symptoms were assigned to asymptomatic groups. Mild COVID-19 was labeled to those who were presenting mild symptoms (such as fever, cough, and nausea) without dyspnea. Patients having manifestation of pneumonia and oxygen saturation $\geq 93\%$ fell under moderate category. The patients with severe COVID-19 presented with pneumonia accompanied by one of the followings: respiratory rate > 30 times/minute, severe respiratory distress, or oxygen saturation $< 93\%$. Fasting blood glucose (FBG) and 2-hours

post-prandial blood glucose level (2hPBG) of the patients were measured during the 24 hours of the admission (first measurement) and the last day prior to hospital discharge (second measurement). Changed values of FBG (Δ FBG) and 2hPBG (Δ 2hPBG) were determined by subtracting the value obtained from the first measurement with that from the second measurement. Herein, we also collected the data pertaining to the previous treatment and antihyperglycemic agents received by the patients including their changes during the hospitalization. Length of stay was assigned as the outcome of the treatment, in addition with the COVID-19 positivity. Patients were grouped based on the RT-PCR results on the hospital discharge; patients recovered with COVID-19 negative result and patients discharged with COVID-19 positive results and required additional self-quarantine or referred to another healthcare facility.

Data analysis

Results of this study were processed on SPSS software version 24 (SPSS Inc., Chicago, IL, USA) (SPSS, RRID: SCR_019096) and expressed as descriptive data. All continuous data were presented in mean \pm standard deviation (SD). Meanwhile, categorical data were presented in numbers and percentages.

Results

Demographic characteristics of the COVID-19 patients with diabetes mellitus (n=260) have been presented in Table 1. The average patients' age was 51.33 ± 8.85 years old, predominated by those who were 41–60 years old (76.1%) and followed by 61–80 years old group (13.1%) and 21–40 years old group (10.8%). The numbers of female and male patients were almost similar with percentages of 46.2% and 53.8%, respectively. Most of the patients were self-employed (45%), civil servants (19.2%), and housewives (13.8%).

As many as 156 (60%) of the total patients were categorized as obese, whereas 55 (21.2%), 36 (13.8%), and 11 (4.2%) others fell into overweight, normal, and underweight categories, respectively (Table 2). Hypertension was observed as the most common comorbidity (47.7%) recorded upon the hospital admission. There were close numbers of patients between those who were having 1 comorbidity (46.2%) and those who were having 2-3 comorbidities (53.8%). From the highest to the lowest frequency, the COVID-19 symptoms presented included cough (47.7%), fever (23.8%), rhinorrhea (15.4%), dyspnea (13.1%), anosmia (11.5%), nausea (10.0%), and headache (9.6%). As many as 65 (25.0%) patients had COVID-19 symptoms other than those aforementioned (Table 2).

According to the severity, most of the patients suffered mild COVID-19 (55.4%), followed by asymptomatic (26.9%), and moderate (10.4%). There were only 19 (7.3%) patients who were diagnosed with severe COVID-19. Individuals with asymptomatic, mild, and moderate levels of COVID-19 had similar proportion of patients with uncontrolled diabetes (74.3%, 73.6%, and 74.1%, respectively). Meanwhile, the proportion of patients with uncontrolled diabetes was fewer in severe COVID-19 group (57.9%). There was reduction of FBG and 2hPBG levels observed before (210.75 ± 81.38 and 271.19 ± 100.7 mg/dL, respectively) and after the treatment (181.03 ± 68.9 and 222.01 ± 86.96 mg/dL, respectively).

Table 1. Demographic characteristics of COVID-19 patients with diabetes mellitus.

Variable	n (%)
Age, mean \pm SD (years old)	51.33 \pm 8.85
21–40 years old	28 (10.8)
41–60 years old	198 (76.1)
61–80 years old	34 (13.1)
Sex	
Male	140 (53.8)
Female	120 (46.2)
Occupation	
Student	10 (3.84)
Civil servant	50 (19.2)
Teacher	16 (6.2)
Healthcare provider	10 (3.84)
Self-employed	117 (45.0)
Housewife	36 (13.8)
Others	21 (8.07)

Table 2. Clinical characteristic and blood glucose level of COVID-19 patients with diabetes mellitus (n=260).

Variable	n (%)
Body mass index (BMI)	
Obese	156 (60.0)
Overweight	55 (21.2)
Normal	36 (13.8)
Underweight	11 (4.2)
Comorbidity	
Hypertension	124 (47.7)
Cardiovascular disease	10 (3.8)
Chronic kidney disease	1 (0.4)
Asthma	5 (1.9)
Others	14 (5.3)
1 comorbidity	120 (46.2)
2–3 comorbidities	140 (53.8)
COVID-19 Symptoms	
Cough	124 (47.7)
Fever	62 (23.8)
Rhinorrhea	40 (15.4)
Anosmia	30 (11.5)
Dyspnea	34 (13.1)
Nausea	26 (10.0)
Headache	25 (9.6)
Others	65 (25.0)
Severity, n (%) [uncontrolled diabetes, n (%)]	
Asymptomatic	70 (26.9) [52 (74.3)]
Mild	144 (55.4) [106 (73.6)]
Moderate	27 (10.4) [20 (74.1)]
Severe	19 (7.3) [11 (57.9)]
Blood glucose level, mean \pm SD (mg/dL)	
First FBG/last FBG	210.75 \pm 81.38/181.03 \pm 68.9
First 2hPBG/last 2hPBG	271.19 \pm 100.7/222.01 \pm 86.96
Δ FBG	33.14 \pm 66.0
Δ 2hPBG	49.5 \pm 89.6

Higher mean reduction value was obtained in Δ 2hPBG (49.5 \pm 89.6 mg/dL) as compared with that in Δ FBG (33.14 \pm 66.0 mg/dL).

Therapies performed on the patients during the hospitalization to manage the COVID-19 symptoms as well as comorbidities have been presented (Figure 1). All patients herein were prescribed with multivitamin. Most of the patients were treated for their COVID-19 symptoms with antitussive/mucolytic agents (47.3%) and antipyretic agents (24.5%), where 31 of the patients (11.9%) received oxygen support. Frequencies of diabetes mellitus treatments received by the patients before and during the hospitalization have been presented (Figure 2). There were 150 (57.6%) and 75 (28.8%) patients who were given oral antidiabetic drug and insulin, respectively, where 22 (8.4%) others received both therapies. None of the patients included in this study were given antivirals.

Clinical outcomes of the COVID-19 patients with diabetes mellitus have been presented in Table 3. A total of 202 (77.7%) patients were treated in the hospital for more than 10 days. There were 22 (8.5%) patients who underwent




Figure 1. Therapies performed to treat COVID-19 symptoms and comorbidities of the patients during the hospital admission.




Figure 2. Diabetes mellitus therapies performed on the patients before and during hospital admission. DPP4 = dipeptidyl peptidase-4; OAD = oral antidiabetic drug; SGLT = sodium-glucose linked transporter; TZD = thiazolidinedione.

Table 3. Clinical outcomes of COVID-19 patients with diabetes mellitus (n=260).

Variable	n (%)
Length of stay	
<10 days	36 (13.8)
10 days	22 (8.5)
>10 days	202 (77.7)
Outcomes	
Recovered (negative RT-PCR)	229 (88.1)
Recovered (self-quarantine)	2 (0.8)
Referred & survive	21 (8.1)

the hospital treatment within 10 days. Meanwhile, 36 (13.8%) patients had the total hospital stay shorter than 10 days. As for the outcome, most of the patients were discharged from the hospital with COVID-19 negative results (88.1%). Only 2 (0.8%) patients who were discharged from the hospital and required for self-quarantine following the COVID-19 positive results. There were few individuals referred to the other healthcare facility who then survived (8.1%) and died (3.1%).

Discussion

Herein, patients presented with cough and fever as the most common symptoms during the onset of COVID-19.¹⁴ These results were similar to that of a previous report investigating 904 patients with COVID-19 and diabetes mellitus in China.¹⁵ According to multiple reports, combination of COVID-19 and diabetes is fatal because the diseases could complement one another.⁵ Most patients in this present study were in age group of 41–60 years old (76.1%) with an average of 51.33 ± 8.85 years old. In combination with pre-existing health conditions, such as obesity (60%) and hypertension (47.7%), COVID-19 patients with diabetes mellitus were at higher risk of poor prognosis.¹⁶ Obesity could cause poor outcomes in COVID-19 patients with diabetes mellitus because of its association with chronic inflammation.⁴ Hypertension could downregulate the expression of ACE2, which subsequently increases levels of angiotensin-2 and decreases angiotensin 1-7, leading to the worsening of ARDS.¹⁷ Upon the SARS-CoV-2 infection, ACE2 was also found to have decreased,¹⁸ becoming an interplay between COVID-19 and hypertension in causing multiple organ failures.¹⁹

Despite the fact that individuals with diabetes mellitus could have higher risk to develop severe COVID-19,⁵ our current findings suggest that only a small percentage of patients reported developing severe COVID-19 (7.3%). Most of the patients had mild (55.4%) and asymptomatic COVID-19 (26.9%). When glycemic control was observed, the majority of patients in asymptomatic, mild, and moderate COVID-19 groups were predominated by individuals with poor blood glucose control (>70%). It is worth mentioning that hyperglycemia could induce the glycation of ACE2, contributing to increased entry of SARS-CoV-2 into the host cells.²⁰ Our obtained data suggest that there is no association between diabetes mellitus or hyperglycemic condition with the severity of COVID-19. During the time frame of this study, people have been already aware of COVID-19 and massive testing was carried out.²¹ We argue that early detection of SARS-CoV-2 infection could prevent the development and the progression of the disease.

Prior to the hospital admission, more than 80% of the patients have received glucose-lowering agents. However, the FBG and 2hPBG were found to be in high levels during the admission. A study found that COVID-19 could cause a dysregulation of lipid metabolism which eventually contributes to insulin resistance.²² Another study reported that as a result of ACE2 downregulation following the SARS-CoV-2 infection, insulin resistance could be developed owing to the exaggeration of angiotensin II.²³ During the treatment in the hospital, there was a significant increase of patients receiving sodium-glucose linked transporter (SGLT) inhibitors and insulin therapies. Following the recovery from COVID-19 symptoms, patients showed a reduction of FBG and 2hPBG levels, though they were still far higher than the normal ranges (80–130 mg/dL and <180 mg/dL for FBG and 2hPBG, respectively).²⁴ Here, we could conclude that it is important to control the blood glucose level and maintain diabetes mellitus treatments during the course of COVID-19, as advised by the international panel of diabetes experts.⁴ Secondly, our data suggests that COVID-19 could influence the level of the blood glucose in diabetic patients.

In this study, there was no specific treatment for COVID-19, only vitamins and symptomatic treatment. None of the patients were given antivirals as there is currently no antiviral specified to treat SARS-CoV-2 infection. The majority of patients (97%) survived after the hospital treatment. As the limitation, with a small number of patients included in this study, it is impossible to draw conclusions that have clinical implications. Within the time frame of the observation, COVID-19 variants have emerged,²⁵ which could contribute to the biased results.

Conclusions

Most of the patients with both COVID-19 and diabetes mellitus were over 50 years old, and hypertension and obesity were commonly found preexisting conditions in them. Blood glucose level could be increased by the COVID-19, suggesting the importance of blood glucose lowering therapies in diabetic patients with COVID-19. Symptomatic management without antiviral and antibiotic therapy followed by blood glucose lowering therapies contribute to the survivability of the patients.

Data availability

Underlying data

Figshare: 'Clinical characteristics and outcomes of hospitalized COVID-19 patients with diabetes mellitus in East Java, Indonesia: A cross-sectional study.' DOI: <https://doi.org/10.6084/m9.figshare.19388771>.²⁶

Data are available under the terms of the [Creative Commons Attribution 4.0 International license \(CC-BY 4.0\)](#).

This project contains the following underlying data:

- Master Data.xls [Table containing the raw data of the study].
- Master Data.sav [Table containing the raw data of the study and the code book].

Reporting guidelines

Figshare: STROBE checklist for “Clinical characteristics and outcomes of hospitalized COVID-19 patients with diabetes mellitus in East Java, Indonesia: A cross-sectional study” - <https://doi.org/10.6084/m9.figshare.19388975>.

All data are available under the terms of the [Creative Commons Attribution 4.0 International license \(CC-BY 4.0\)](#).

Acknowledgement

The authors wish to thank the chairman of Indrapura Forefront Hospital, I.D.G. Nalendra Djaya Iswara and Djanuar Firtriadi for their supports throughout the making of this paper.

References

1. Fahrhani M, Anwar S, Yufika A, et al.: **Disruption of childhood vaccination during the COVID-19 pandemic in Indonesia.** *Narra. J.* 2021; **1**(1): e7. [PubMed Abstract](#) | [Publisher Full Text](#)
2. Bintari DC, Sudibyo DA, Karimah A: **Correlation between depression level and headache severity: A study among medical students during the COVID-19 pandemic.** *Narra. J.* 2021; **1**(3): e64. [PubMed Abstract](#) | [Publisher Full Text](#)
3. Hamming I, Timens W, Bulthuis ML, et al.: **Tissue distribution of ACE2 protein, the functional receptor for SARS coronavirus. A first step in understanding SARS pathogenesis.** *J. Pathol.* 2004 Jun; **203**(2): 631–637. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
4. Lim S, Bae JH, Kwon HS, et al.: **COVID-19 and diabetes mellitus: from pathophysiology to clinical management.** *Nat. Rev. Endocrinol.* 2021 Jan; **17**(1): 11–30. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
5. Landstra CP, de Koning EJP: **COVID-19 and Diabetes: Understanding the Interrelationship and Risks for a Severe Course.** *Front. Endocrinol. (Lausanne).* 2021; **12**: 649525. Epub 2021/07/06. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
6. Zahra Z, Ramadhani C, Mamfaluti T, et al.: **Association between depression and HbA1c levels in the elderly population with type 2 diabetes mellitus during COVID-19 pandemic.** *Narra. J.* 2022; **2**(3): e51. [PubMed Abstract](#) | [Publisher Full Text](#)
7. Wu CT, Lidsky PV, Xiao Y, et al.: **SARS-CoV-2 infects human pancreatic beta cells and elicits beta cell impairment.** *Cell Metab.* 2021 Aug 3; **33**(8): 1565–1576.e5. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
8. Consoli A, Devangelio E: **Thiazolidinediones and inflammation.** *Lupus.* 2005; **14**(9): 794–797. [PubMed Abstract](#) | [Publisher Full Text](#)
9. Singh S, Loke YK, Furberg CD: **Long-term use of thiazolidinediones and the associated risk of pneumonia or lower respiratory tract infection: systematic review and meta-analysis.** *Thorax.* 2011 May; **66**(5): 383–388. [PubMed Abstract](#) | [Publisher Full Text](#)
10. Yang J, Zheng Y, Gou X, et al.: **Prevalence of comorbidities and its effects in patients infected with SARS-CoV-2: a systematic review and meta-analysis.** *Int. J. Infect. Dis.* 2020 May; **94**: 91–95. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
11. Onder G, Rezza G, Brusaferro S: **Case-fatality rate and characteristics of patients dying in relation to COVID-19 in Italy.** *JAMA.* 2020 May 12; **323**(18): 1775–1776. [PubMed Abstract](#) | [Publisher Full Text](#)
12. Harbuwono DS, Handayani DOTL, Wahyuningsih ES, et al.: **Impact of diabetes mellitus on COVID-19 clinical symptoms and mortality: Jakarta's COVID-19 epidemiological registry.** *Prim. Care Diabetes.* 2022; **16**(1): 65–68. [PubMed Abstract](#) | [Publisher Full Text](#)
13. Karyono DR, Wicaksana AL: **Current prevalence, characteristics, and comorbidities of patients with COVID-19 in Indonesia.** *J. Community Empowerment Health.* 2020; **3**(2): 82–89.
14. Larsen JR, Martin MR, Martin JD, et al.: **Modeling the onset of symptoms of COVID-19.** *Front. Public Health.* 2020; **8**: 473. Epub 2020/09/10. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
15. Chen Y, Yang D, Cheng B, et al.: **Clinical characteristics and outcomes of patients with diabetes and COVID-19 in association with glucose-lowering medication.** *Diabetes Care.* 2020 Jul; **43**(7): 1399–1407. [PubMed Abstract](#) | [Publisher Full Text](#)
16. Zhang W, Du RH, Li B, et al.: **Molecular and serological investigation of 2019-nCoV infected patients: implication of multiple shedding routes.** *Emerg. Microbes. Infect.* 2020; **9**(1): 386–389. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
17. Revercomb L, Hanmandlu A, Wareing N, et al.: **Mechanisms of Pulmonary Hypertension in Acute Respiratory Distress Syndrome (ARDS).** *Front. Mol. Biosci.* 2020; **7**: 624093. Epub 2021/02/05. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
18. Ni W, Yang X, Yang D, et al.: **Role of angiotensin-converting enzyme 2 (ACE2) in COVID-19.** *Crit. Care.* 2020 Jul 13; **24**(1): 422. Epub 2020/07/15. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
19. Tsiouffis C, Dimitriadis K, Tousoulis D: **The interplay of hypertension, ACE-2 and SARS-CoV-2: Emerging data as the “Ariadne’s thread” for the “labyrinth” of COVID-19.** *Hell. J. Cardiol.* 2020 Jan - Feb; **61**(1): 31–33. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
20. D’Onofrio N, Scisciola L, Sardu C, et al.: **Glycated ACE2 receptor in diabetes: open door for SARS-CoV-2 entry in cardiomyocyte.** *Cardiovasc. Diabetol.* 2021 May 7; **20**(1): 99. Epub 2021/05/09. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
21. Harapan H, Anwar S, Nainu F, et al.: **Perceived Risk of Being Infected With SARS-CoV-2: A Perspective From Indonesia.** *Disaster Med. Public Health Prep.* 2020 Sep; **10**: 1–5. Epub 2020/09/11. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
22. He X, Liu C, Peng J, et al.: **COVID-19 induces new-onset insulin resistance and lipid metabolic dysregulation via regulation of secreted metabolic factors.** *Signal Transduct. Target. Ther.* 2021 Dec 16; **6**(1): 427. Epub 2021/12/18. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
23. Govender N, Khaliq OP, Moodley J, et al.: **Insulin resistance in COVID-19 and diabetes.** *Prim. Care Diabetes.* 2021 Aug; **15**(4):

629–634.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

24. Tanaka M: **Relationship between fasting and 2-hour postprandial plasma glucose levels and vascular complications in patients with type 2 diabetes mellitus.** *J. Int. Med. Res.* 2012; **40**(4): 1295–1303.
[PubMed Abstract](#) | [Publisher Full Text](#)
25. Korber B, Fischer WM, Gnanakaran S, *et al.*: **Tracking Changes in SARS-CoV-2 Spike: Evidence that D614G Increases Infectivity of the COVID-19 Virus.** *Cell.* 2020 Aug 20; **182**(4): 812–827.e19.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
26. Triyono E: **Clinical characteristics and outcomes of hospitalized COVID-19 patients with diabetes mellitus in East Java, Indonesia: A cross-sectional study.** *Figshare. Journal Contribution.* 2022.
[Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:  

Version 1

Reviewer Report 09 August 2022

<https://doi.org/10.5256/f1000research.122720.r144898>

© 2022 Dhama K. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Kuldeep Dhama 

Division of Pathology, ICAR-Indian Veterinary Research Institute, Bareilly, Uttar Pradesh, India

The study "Clinical characteristics and outcomes of hospitalized COVID-19 patients with diabetes mellitus in East Java, Indonesia: A cross-sectional study" described the clinical characteristics and determined the outcomes of 260 COVID-19 patients with comorbid diabetes mellitus who received the blood glucose lowering therapies. The study used the medical reports between May 2020 and February 2021 in Indonesia.

There are some suggestions / recommendations to improve the quality of the article:

1. The diabetes mellitus diagnosis should be provided in more detail. In the article written: "Diabetes mellitus was confirmed by the officially recorded medical history." This criteria used should be explained.
2. "All patients with COVID-19 enrolled in this study were diagnosed and managed according to the national guidelines." Please explain the guidelines. Provide the references here.
3. It is not clear the outcome groups. It is written: "Patients were grouped based on the RT-PCR results on the hospital discharge; patients recovered with COVID-19 negative result and patients discharged with COVID-19 positive results and required additional self-quarantine or referred to another healthcare facility". Please revise this sentence.
4. Please provide the reference of how the BMI was classified into obese, overweight, normal, and underweight.
5. It is not clear the criteria of "uncontrolled diabetes". Please explain in Method and provide the references.
6. Please explain how previous treatment of diabetes was assessed in this study.
7. Minor: Please define "SARS" in the second paragraph of Introduction.

Summary

Diabetes mellitus is an important comorbidity with COVID-19, and COVID-19 also affects blood glucose levels, hence antihyperglycemic therapies for lowering blood glucose levels and COVID-19 symptomatic treatments are of importance for treating such patients.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Infectious diseases, emerging pathogens.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 07 July 2022

<https://doi.org/10.5256/f1000research.122720.r141584>

© 2022 **Jamil K.** This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Kurnia Fitri Jamil 

Universitas Syiah Kuala, Banda Aceh, Indonesia

The suggestions that I can give to the manuscript with the title "Clinical characteristics and outcomes of hospitalized COVID-19 patients with diabetes mellitus in East Java, Indonesia: A cross-sectional study", are:

1. There are some grammatical errors in the text. Authors need to re-examine the manuscript.
2. In my opinion, some of the information provided in the Introduction section are more

suitable to be included in the Discussion instead. A lengthy introduction compared to the Discussion might give the impression that available studies in the literature on this matter is more than adequate in contributing the information without the need of the current study.

3. I suggest displaying the figures (figures 1 and 2), in a more attractive form.
4. In my opinion, data or comparison groups are needed for users of single oral anti-diabetic drugs only, and users of antidiabetic drugs with insulin, especially in obese patients.
5. It is better if the authors can provide more updates or additional facts that can be obtained from this study compared to what already been published in the literature, and also presents data from others Asian countries, especially Southeast Asia, as comparison data.
6. I would advise you in the future to use a prospective research design.
7. Reference: correct or cite in full reference numbers 14, 17, 26.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Partly

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Internist, Tropical Diseases and Infectious Consultant, Professor in Internal Medicine

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research

Clinical characteristics and outcomes of hospitalized COVID-19 patients with diabetes mellitus in East Java, Indonesia A cross-sectional study

ORIGINALITY REPORT

19%

SIMILARITY INDEX

10%

INTERNET SOURCES

18%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Tom Beneke, Ulrich Dobramysl, Carolina Moura Costa Catta-Preta, Jeremy Charles Mottram, Eva Gluenz, Richard Wheeler. "Genome sequence of *Leishmania mexicana* MNYC/BZ/62/M379 expressing Cas9 and T7 RNA polymerase", Wellcome Open Research, 2022
Publication 1%
- 2 Enkeleint A. Mechili, Krzysztof Przewoźniak, Pete Driezen, Christina N Kyriakos et al. "Smokers' support for the ban on sale of slim cigarettes in six European countries: findings from the EUREST-PLUS ITC Europe surveys", Open Research Europe, 2021
Publication 1%
- 3 Suzanne Karan, William A. Voter, Denham Ward. "Myofunctional training with negative airway pressure for obstructive sleep apnea: a prospective non-randomized cohort feasibility study", F1000Research, 2021 1%

4

Boon Zhan Sia, Wan Xin Boon, Yoke Yee Yap, Shalini Kumar, Chong Han Ng. "Prediction of the effects of the top 10 nonsynonymous variants from 30229 SARS-CoV-2 strains on their proteins", F1000Research, 2022

Publication

5

Mustanir Yahya, Binawati Ginting, Nurdin Saidi. "Anticancer and antiretroviral activities of methanolic extract from Theobroma cacao L pod husk: focusing on the ethyl acetate partition", F1000Research, 2022

Publication

6

Shahid S Memon, Dalia A Biswas. "COVID-19 and Diabetes Mellitus: From Pathophysiology to Clinical Management", Cureus, 2022

Publication

7

Lin Miao, Kyle R. Batty, Ayana N. Jackson, Heather A. Pieno, Maisy W. Rhoades, Shihoko Kojima. "Genetic and environmental perturbations alter the rhythmic expression pattern of a circadian long non-coding RNA, Per2AS, in mouse liver", F1000Research, 2022

Publication

8

Ravi Philip Rajkumar. "Post-traumatic stress in the wake of the COVID-19 pandemic: a scoping review", F1000Research, 2020

Publication

1 %

1 %

1 %

1 %

1 %

9

Dante S. Harbuwono, Dwi O.T.L. Handayani, Endang S. Wahyuningsih, Novita Supraptowati et al. "Impact of diabetes mellitus on COVID-19 clinical symptoms and mortality: Jakarta's COVID-19 epidemiological registry", Primary Care Diabetes, 2022

Publication

<1 %

10

Gusti Noor Ramadany Saputra, Resti Yudhawati, Munawaroh Fitriah. "Association of soluble receptor for advanced glycation end-products (sRAGE) serum on COVID-19 severity: A cross-sectional study", Annals of Medicine and Surgery, 2022

Publication

<1 %

11

jennymeehan.wordpress.com

Internet Source

<1 %

12

Banafsheh Bigdelou, Mohammad Reza Sepand, Sahar Najafikhoshnoo, Jorge Alfonso Tavares Negrete et al. "COVID-19 and Preexisting Comorbidities: Risks, Synergies, and Clinical Outcomes", Frontiers in Immunology, 2022

Publication

<1 %

13

Tanaka, M.. "Relationship between Fasting and 2-Hour Postprandial Plasma Glucose Levels and Vascular Complications in Patients with Type 2 Diabetes Mellitus", Journal of International Medical Research, 2012.

<1 %

14

Mochammad Thaha. "Intravenous N-Acetylcysteine during Haemodialysis Reduces the Plasma Concentration of Homocysteine in Patients with End-Stage Renal Disease", *Clinical Drug Investigation*, 2006

Publication

15

expeditiorepositorio.utadeo.edu.co

Internet Source

16

www.slideshare.net

Internet Source

17

"Abstracts of 51st EASD Annual Meeting", *Diabetologia*, 2015

Publication

18

Charlotte Steenblock, Peter E H Schwarz, Barbara Ludwig, Andreas Linkermann et al. "COVID-19 and metabolic disease: mechanisms and clinical management", *The Lancet Diabetes & Endocrinology*, 2021

Publication

19

Rakesh Kumar, Cut Meurah Yeni, Niken Asri Utami, Rupali Masand et al. "SARS-CoV-2 infection during pregnancy and pregnancy-related conditions: Concerns, challenges, management and mitigation strategies—a narrative review", *Journal of Infection and Public Health*, 2021

Publication

<1 %

<1 %

<1 %

<1 %

<1 %

<1 %

20	encyclopedia.pub Internet Source	<1 %
21	www.science.org Internet Source	<1 %
22	www.sid.ir Internet Source	<1 %
23	Taufik Faturohman, Giofella Adesta Navaky Kengsiswoyo, Harapan Harapan, Suhaiza Zailani, R. Aswin Rahadi, Neneng Nurlaela Arief. "Factors influencing COVID-19 vaccine acceptance in Indonesia: an adoption of Technology Acceptance Model", F1000Research, 2021 Publication	<1 %
24	bip.covid19.athenarc.gr Internet Source	<1 %
25	dokumen.pub Internet Source	<1 %
26	www.intechopen.com Internet Source	<1 %
27	www.scielo.br Internet Source	<1 %
28	"RHEUMATOID ARTHRITIS ? TREATMENT", APLAR Journal of Rheumatology, 8/2006 Publication	<1 %

29

Aman Rajpal, Leili Rahimi, Faramarz Ismail - Beigi. " Factors Leading to High Morbidity and Mortality of - 19 in Patients with Type 2 Diabetes ", Journal of Diabetes, 2020

Publication

<1 %

30

Bruce Larson, Amir Shroufi, Charles Muthoga, Rita Oladele et al. "Induction-phase treatment costs for cryptococcal meningitis in high HIV-burden African countries: New opportunities with lower costs", Wellcome Open Research, 2021

Publication

<1 %

31

Hyun Lee, Hayoung Choi, Bumhee Yang, Sun-Kyung Lee et al. "Interstitial lung disease increases susceptibility to and severity of COVID-19", European Respiratory Journal, 2021

Publication

<1 %

32

Iswan Nusi, Amie Vidyani, Putu Niken Ayu Amrita, Herry Purbayu, Ulfa Kholili. "A Patient With Colon Perforation Due to Amebiasis: A Rare and Fatal Complication", American Journal of Gastroenterology, 2014

Publication

<1 %

33

Nurshad Ali, Khaled Hossain. "Liver injury in severe COVID-19 infection: current insights and challenges", Expert Review of Gastroenterology & Hepatology, 2020

<1 %

34 Saminan Saminan, Linda Julisafrida, Muhammad Ridwan, Nurul Fajri. "COVID-19 Pandemic: What Considerations Should Be Taken during the Assessment and Management of COPD Exacerbation?", Open Access Macedonian Journal of Medical Sciences, 2022
Publication

35 medicinescience.org
Internet Source

36 nonpareilonline.com
Internet Source

37 ogscience.org
Internet Source

38 www.dovepress.com
Internet Source

39 "Clinical Vignettes/Case Reports - Esophagus", The American Journal of Gastroenterology, 2014.
Publication

40 Chun, Young-Soo, Seung-Hyuk Lee, Sang-Hoon Lee, Yoon-Je Cho, and Kee Hyung Rhyu. "Clinical Implication of Diabetes Mellitus in Primary Total Hip Arthroplasty", Hip & Pelvis, 2014.
Publication

41

Erik Olofsen, Albert Dahan. "Using Akaike's information theoretic criterion in population analysis: a simulation study", F1000Research, 2013

Publication

<1 %

42

I.Made Cahyadi Agastiya, Endar Kurianto, Hanifati Akalili, Anggi Lukman Wicaksana. "The impact of telehealth on self-management of patients with type 2 diabetes: A systematic review on interventional studies", Diabetes & Metabolic Syndrome: Clinical Research & Reviews, 2022

Publication

<1 %

43

Zakaria Zakariaei, Ali Sharifpour, Mahdi Fakhar, Mostafa Soleymani, Elham Sadat Banimostafavi, Amirmasoud Taheri. "Detection of in pericardial effusion sample in a COVID-19 patient with systemic sclerosis: An unusual case report ", SAGE Open Medical Case Reports, 2022

Publication

<1 %

44

co.whatcom.wa.us

Internet Source

<1 %

45

en.wikipedia.org

Internet Source

<1 %

46

journal.unhas.ac.id

Internet Source

<1 %

47

journals.plos.org

Internet Source

<1 %

48

pagepress.org

Internet Source

<1 %

49

pdfs.semanticscholar.org

Internet Source

<1 %

50

Hongyin Chen, Jiangyun Peng, Tengyao Wang, Jielu Wen, Sifan Chen, Yu Huang, Yang Zhang. "Counter-regulatory renin-angiotensin system in hypertension: Review and update in the era of COVID-19 pandemic", *Biochemical Pharmacology*, 2023

Publication

<1 %

51

Li-Gang Song, Su-Rong Bai, Deng-Hua Hui, Li-Ping Ding, Lu Sun. "Association of COVID-19 patient's condition with fasting blood glucose and body mass index: A retrospective study", *Technology and Health Care*, 2022

Publication

<1 %

52

Ningfei Ji, Mingshun Zhang, Liang Ren, Yunyun Wang et al. "SARS-CoV-2 in the pancreas and the impaired islet function in COVID-19 patients", *Emerging Microbes & Infections*, 2022

Publication

<1 %

53

Yuan-Hui Wang, Yue-Ying Yang, Hao-Qi Li, Qi-Dong Zhang, Fei Xu, Zhi-Jian Li.

<1 %

"Characterization of aroma-active compounds in steamed breads fermented with Chinese traditional sourdough", LWT, 2021

Publication

54

Berrin Ergun-Longmire, Ethel Clemente, Patricia Vining-Maravolo, Cheryl Roberts, Koby Buth, Donald E. Greydanus. "Diabetes education in pediatrics: How to survive diabetes", Disease-a-Month, 2021

Publication

<1 %

55

www.eidogen.com

Internet Source

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Clinical characteristics and outcomes of hospitalized COVID-19 patients with diabetes mellitus in East Java, Indonesia A cross-sectional study

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10

PAGE 11

PAGE 12

PAGE 13
