

Journal of Public Health Research

[Journal indexing and metrics](#)

Editorial board

[Hide all](#)

Editor in Chief

**Luigi
Barberini**

Department of Medical Sciences and Public Health, University of Cagliari,
Italy

Associate Editors

Laura Atzori

Department of Medical Sciences and Public Health, University of
Cagliari, Italy

**Antonella
Balestrieri**

Department of Medical Sciences and Public Health, University of
Cagliari, Italy

Doris Barcellona

Department of Medical Sciences and Public Health, University of
Cagliari, Italy

Cristina Cabras

Department of Pedagogy, Psychology, Philosophy, University of
Cagliari, Italy

**Francesco
Campanella**

Department of Occupational and Environmental Medicine,
Epidemiology and Hygiene, Worker Compensation Authority (INAIL)

 Privacy

**Emanuele
Cannizzaro**

Department of Health Promotion, Maternal and Infant Care, University
of Palermo, Italy

**Alessandro
Carriero**

Faculty of Medicine and Surgery, University of Eastern Piedmont, Italy

Mauro Carta

Department of Medical Sciences and Public Health, University of
Cagliari, Italy

Flaviana Cau

Department of Medical Sciences and Public Health, University of
Cagliari, Italy

Giulia Cossu

Department of Medical Sciences and Public Health, University of
Cagliari, Italy

Daniela Fanni

Department of Medical Sciences and Public Health, University of
Cagliari, Italy

Claudia Fattuoni

University of Cagliari Department of Chemical and Geological Sciences,
University of Cagliari, Italy

**Pierpaolo
Ferrante**

Department of Occupational and Environmental Medicine,
Epidemiology and Hygiene, Worker Compensation Authority (INAIL),
Italy

Clara Gerosa

Department of Medical Sciences and Public Health, University of
Cagliari, Italy

Dmitri Grapov

University of California, Davis, CA, USA

Danial Kahrizi

Razi University, Iran

Jutta Lindert

University of Emden, Germany

Federico Meloni

Department of Medical Sciences and Public Health, University of
Cagliari, Italy

Olga MulasDepartment of Medical Sciences and Public Health, University of
Cagliari, Italy**Marcello Nonnis**Department of Pedagogy, Psychology and Philosophy, University of
Cagliari, Italy**Germano Orrù**Department of Medical Sciences and Public Health, University of
Cagliari, Italy**Monica Piras**Department of Medical Sciences and Public Health, University of
Cagliari, Italy**Luca Saba**Department of Medical Sciences and Public Health, University of
Cagliari, Italy**Salvatore Sardo**Department of Medical Sciences and Public Health, University of
Cagliari, Italy**Alessandra
Scano**Department of Medical Sciences and Public Health, University of
Cagliari, Italy

Browse journal

[All articles](#)[Browse by year](#)

Journal information

[Journal description](#)[Aims and scope](#)[Editorial board](#)[Submission guidelines](#)[Journal indexing and metrics](#)



Open Access

Research article

| First published Apr 27, 2022

[Instrumental Music Therapy Reduced Depression Levels in Stroke Patients](#)

Vione Deisi Oktavina Sumakul, Hari Basuki Notobroto, Ni Luh Widani, M. Havidz Aima

Journal of Public Health Research

[Preview abstract](#)

PDF / EPUB



Open Access

Research article

| First published Jul 14, 2022

[The Effect of a Smartphone-Based Perioperative Nursing Intervention: Prayer, Education, Exercise Therapy, Hypnosis, and Music toward Pain, Anxiety, and Early Mobilization on Cardiac Surgery](#)

Sidik Awaludin, Elly Nurachmah, Tri Wisesa Soetisna, Jahja Umar

Journal of Public Health Research

[Preview abstract](#)

PDF / EPUB



Open Access

Review article

| First published May 16, 2022

[Informal Caregivers and Psychosocial Support: Analysis of European Dementia Policy documents](#)

Miroslava Tokovska, Magde Mohamed Nour, Anette Sørensen, Ursula Småland Goth

Journal of Public Health Research

[Preview abstract](#)

ARTICLE

Instrumental music therapy reduced depression levels in stroke patients

Vione Deisi Oktavina Sumakul,¹ Hari Basuki Notobroto,² Ni Luh Widani,³ M. Havidz Aima³

¹Doctoral Program of Public Health; ²Department of Biostatistics and Population, Faculty of Public Health, Universitas Airlangga, Mulyorejo, Surabaya; ³Master of Medical Surgical Nursing, Sint Carolus School of Health Sciences, Jakarta, Indonesia

Abstract

Background: Stroke is the fifth cause of death and disability, leading also to depression. However, depression in stroke patients is hardly handled optimally. The purpose of this study therefore is to determine the effectiveness of instrumental music therapy in reducing depressive symptoms in stroke patients.

Design and methods: It used a quasi-experiment pre-post design with a simple random sampling with 59 respondents. The respondents were divided into 3 groups as follows; group A (standard treatment), group B (instrumental music therapy), and group C (combined treatment).

Results: The results show that the combined treatment provided the most significant influence on reducing the level of depression ($P=0.001$) with a contribution of 68.6% compared to the group A which was given standard treatment ($P=0.001$) with a contribution of 61.7%. Instrumental music therapy had no effect ($P=0.986$), though it contributed most among the three interventions, specifically 82.6%.

Conclusions: The study recommended further improvement to include music as treatment options for reducing depression among stroke patients.

Introduction

Stroke is a neurological disorder caused by the interruption of blood flow to parts of the brain. There are two main types of stroke, ischemic and hemorrhagic stroke. In general, ischemic accounts for 83% of all stroke cases while the remaining 17% is hemorrhagic.¹ Stroke is the most significant cause of complex disability, with at least half of all stroke patients suffering disabilities. In Indonesia, the highest prevalence of stroke was recorded in Yogyakarta, Central Java (14.6%), followed by North Sulawesi area (14%). The highest prevalence of stroke based on age is ≥ 75 years (50.2%), while according to sex, men account for 11.0% and women 10.9%. Based on the residence, stroke patients who live in

the city accounts for 12.6% more than people live in the village.² In UK, more than one-third of sufferers' were dependent on others, with 1 in 5 being treated by family and friends. Also, one third (33%) of sufferers developed post-stroke depression.³

Post Stroke Depression (PSD) is a common complication that causes disability to stroke patients due to a lack of rehabilitation therapy and a decreased quality of life.³ As a secondary type of depression, PSD inhibits the recovery of neurological and cognitive function, and increase mortality and disability rates. It often leads to marked mental and physical pain, increasing family's and social burden. The effects of stroke are unpredictable, as it can also lead to emotional and behavioural changes in the patient, as well as family issues that are not resolved peacefully and can trigger depression.⁴⁻⁶

Several strategies can be used to overcome depression in stroke, both non-pharmacological and pharmacological treatment. Non-pharmacological treatment includes music therapy and supportive therapy from caregivers, or people assisting stroke patients. People involved in helping sufferers are often called as caregivers, it can be a partner, family members, or friends. Partners, children, or parents provides most of the care to help stroke patients regain their physical and psychological abilities.¹ Pharmacological treatment is associated with medication therapy for stroke patients in managing the impacts of disease.

As one of non-pharmacological therapy, neurologic music therapy (NMT) has been developed as a new model for music therapy. The NMT techniques in stroke rehabilitation are mainly used to improve motor functions, especially in gait functions and upper limb motor skills. It was investigated that there was a decrease in depressive symptoms in stroke patients who received music therapy.⁷ Therapeutic interventions include listening to music in groups, body movements, and painting with music and improvising singing.⁸ Existing literatures show that listening to the preferred music could promote physical and cognitive recovery of the patient, along with mood improvement, thus helping for recovery after cerebral artery stroke.^{9,10} In caring for stroke patients, nurses are expected to include innovations in nursing interventions, especially aiming at functional and cognitive recovery.

Significance for public health

Stroke is the most significant cause of complex disability, with at least half of all stroke patients suffering disabilities, including depression. Depression often occurs in these patients and is hardly handled optimally, therefore, it is vital to identify appropriate interventions to help them adapt to the long terms effects of the stroke. Instrumental music therapy has been shown to promote cognitive recovery and mood improvement. This study describes the effectiveness of instrumental music therapy in reducing depressive symptoms in stroke patients.

ery after a stroke. Depression often occurs in stroke patients, and therefore it is vital to identify appropriate interventions to help them adapt to the long terms effects of the stroke.¹¹

The aim of this study, therefore, is to determine the effectiveness of instrumental music therapy in reducing depressive symptoms in stroke patients.

Design and methods

This research was conducted on stroke patients with moderate and severe depression classification based on the Hamilton Depression Scale (HAMD) criteria. It was conducted on April-June 2015 at public hospital in Manado, Indonesia. The study used a quasi-experiment research with a pretest-posttest group design. A total of 59 respondents were divided into three groups as follows: 20 respondents in group A (standard treatment), 19 respondents in group B (instrumental music therapy), and 20 respondents in group C (combined treatment). In the standard treatment group, the family member who became a primary caregiver was given training, education, and counseling on the way to care for stroke patients and to meet the daily needs of the patient. In the instrumental music therapy group, the patients also listen to instrumental music for 15 minutes, then rest, and continue listening to music again for 15 minutes. The combined intervention group was a blend of standard treatment and instrumental music therapy.

The study used MP-3 and questionnaires as the main instrument of analysis. Before the intervention, the level of depression was measured and the questionnaire was repeated afterwards. Data analysis consisted of univariate analysis on descriptive statistical data for the independent variables (age, sex, duration of the attack, and functional status). The dependent variable (level of depression) was analyzed through bivariate analysis with a paired different test (Wilcoxon test) and multivariate analysis using the Ordinal Logistic Regression test.

The clinical efficacy was evaluated using the HAMD-17 scores in all groups before and after treatment. The HAMD-17 score reduction rate as follows: score reduction rate = (score before treatment - score after treatment) / score before treatment × 100%. The interpretation criteria of HAMD-17 score: cured >75 %, excellent 51-75%, effective 25-50%, and non-effective <25%.¹²

Results and discussion

Data was analyzed using univariate test, bivariate test, and multivariate test. Table 1 shows the data about the patients; it emerges that the respondents mostly aged 45-57 years (50.8%), male and female were almost the same, the duration of the attack was 1-4 days (67.8%), and functional status with a moderate level of independence was 64.4%. Older age was characterized by more complex physical and psychological conditions. In the study emerges that almost half of respondents were middle aged. However, another study stated that younger age group had worse functional outcomes compared with older patients, and 1 in 8 stroke patients still was dependent on others 10 years after the stroke.¹³ Depression has adverse effects on the functional outcome of stroke patients. Yet, individuals with lesions in the right hemisphere were less likely to experience depression. This is attributed to the incidence of sub-acute PSD, though the family also helped to reduce its frequency.¹⁴

The main components while dealing with depression include patients struggling with a new phase of life, having lost their

independence and utilizing available resources from family and friends to help manage the impacts of stroke. Family members of individuals who survived a stroke responded practically to receive support from health care staff, friends and relatives for their daily needs. Family caregivers need a supportive program to encourage their positive behaviour and to get overall assistance to function properly at home.¹⁵⁻¹⁷

Stroke patients often feel helpless and worried about their prevailing conditions. Support from family and closest people helps them to obtain information about their health and manage their daily needs. Family caregivers are the primary source of support for stroke patients and are the first people to respond to the changes in status in the disease phase. Nurses also help in several ways, including preventing complications, reducing risk factors for recurrent strokes, identifying the needs for discharge planning according to family needs, providing the information needed, and encouraging families to be more effective and helpful. However, the central role of nurses in stroke patients is to offer health education to the families, describing pharmacological and non-pharmacological treatments, such as music therapy, to reduce patients'

Table 1. Characteristics of respondents.

Characteristics	N=59	%
Age		
30 – 45 years old	6	10.2
45 – 59 years old	30	50.8
60 – 74 years old	15	25.4
>74 years old	8	13.6
Sex		
Female	29	49.2
Male	30	50.8
Duration of attack		
1 – 4 days	40	67.8
5 – 8 days	10	16.9
9 – 12 days	3	5.1
13 – 16 days	5	8.5
17 – 20 days	1	1.7
Functional status		
Severe (40-60)	6	10.2
Moderate (>20-<40)	38	64.4
Full (0-20)	15	25.4

Table 2. Paired different test.

No.	Group categories	P-value
1.	Group A (standard treatment)	0.000
2.	Group B (instrumental music therapy)	0.000
3.	Group C (combined treatment)	0.000

Table 3. Pseudo R-square test between intervention group.

Intervention group	Cox and Snell
Group A (standard treatment)	0.671
Group B (instrumental music therapy)	0.826
Group C (combined treatment)	0.686

depression.¹⁸

Based on Table 2, $P < 0.005$ were obtained in all three intervention groups. This showed that statistically, there was a significant decrease in the level of depression before and after the interventions in all three groups.

As can be seen in Table 3, the group that contributed the most to the dependent variable was the instrumental music therapy (82.6%). This means that simultaneous musical therapy interventions have the most significant contribution to depression rates. The results of $P < 0.10$ were obtained in both the standard treatment and combined treatment groups. Both interventions had a significant influence on the level of depression. However, instrumental music therapy interventions alone had no statistically significant effect on depression levels. Judging from the magnitude of the estimated value obtained, standard treatment had the highest, 4.997. Statistically, the standard treatment had the most significant influence on the level of depression.

Music therapy has a positive influence on the mood of patients after stroke. Music is produced from stimuli sent from nerves and ascended to the reticular activating system (RAS) neurons. This stimulus is transmitted by specific nuclei from the thalamus through the cerebral cortex, limbic system, and corpus coliseum through the autonomic and neuroendocrine nervous system areas. The limbic system is formed by rings connected by cingulate gyrus, hippocampus, fornix, mammary bodies, hypothalamus, anterior thalamus, and olfactory bulb. When music is played, all areas related to the limbic system is stimulated to produce feelings and expressions.¹⁹⁻²¹

Music has complex effects on humans, both physiologically and spiritually. Individual responses to music are affected by personality, environment, education, and cultural factors. Music causes changes in the status of brain waves and the stress hormones of patients. There is an increase in frequency in the alpha rhythm group and a more significant (coherent) equation between different regions within the cerebral cortex, which most often occurs in the frontal lobe. Activation of the right frontal lobe decreases, reducing the secretion of cortisol and stress hormones to the normal range.²²⁻²⁴

The burden, anxiety, and depression of caregivers reflect the quality of life of stroke survivors. Financial issues ($F=6.12$, $P=0.04$), depression in stroke patients ($F=20.8$, $P < 0.001$), and the burden of family caregivers ($F=7.36$, $P=0.009$) were significantly correlated with decreased patients' quality of life.²⁵ On the other hand, other studies revealed that poverty, low educational attainment, poor health status, being a caregiver for a spouse, and longer length of stay in hospital could reduce the caregivers' quality of life. Thus, family caregivers education programs should be more focused on enhancing health practices and behaviours for stroke survivors and caregivers too, such as developing and promoting adaptation techniques, accessing community services and support from stakeholders.^{26,27}

Conclusions

Statistical tests show that standard treatment and the combined treatment have a significant effect on reducing the level of depression in stroke patients. However, standard treatment had the most significant effect in this study. Therefore, there is a need for primary caregivers to receive training, education, and counseling on treating stroke patients and for hospitals to provide media for listening to instrumental music during hospitalization.

Correspondence: Hari Basuki Notobroto, Department of Biostatistics and Population, Faculty of Public Health, Universitas Airlangga, Jl. Mulyorejo, Surabaya, Jawa Timur 60115, Indonesia. Tel.: +62315920948 - Fax: +62315924618, E-mail: haribasuki.n@fkm.unair.ac.id.

Key words: instrumental music therapy; depression levels; stroke patients.

Contributions: All authors contributed equally.

Conflict of interests: The authors declare no potential conflict of interests.

Funding: This study was supported by Faculty of Public Health, Universitas Airlangga.

Acknowledgments: The authors thanks to Faculty of Public Health, Universitas Airlangga for the continuous support during the study.

Clinical trials: This research has been approved by research ethics committee of Sint Carolus School of Health Sciences, Jakarta, Indonesia.

Conference presentation: Part of this paper was presented at the 4th International Symposium of Public Health, 2019 October 29-31, Griffith University, Gold Coast, Australia.

Received for publication: 6 March 2020.
Accepted for publication: 13 June 2020.

©Copyright: the Author(s), 2020

Licensee PAGEPress, Italy

Journal of Public Health Research 2020;9:1847

doi:10.4081/jphr.2020.1847

This work is licensed under a Creative Commons Attribution NonCommercial 4.0 License (CC BY-NC 4.0).

References

1. Benjamin EJ, Muntner P, Alonso A, et al. Heart Disease and Stroke Statistics 2019 Update A Report From the American Heart Association. *Circulation* 2019;139.
2. Kementerian Kesehatan Republik Indonesia. Hasil Riset Kesehatan Dasar. Jakarta: Kemenkes RI; 2018.
3. Zhang W-N, Pan Y-H, Wang X-Y, et al. A Prospective Study of the Incidence and Correlated Factors of Post-Stroke Depression in China. *PLoS ONE* 2013;8:e78981.
4. Ibeneme SC, Nwosu A, Anyachukwu CC, et al. Burden and factors associated with post-stroke depression in East central Nigeria. *Afr Health Sci* 2017;17:859-67.
5. Duan PX, Wu SY. Advances of integrated Chinese and Western research on mechanism of post-stroke depression. *Zhong Xi Yi Jie He Xin Nao Xue Guan Bing Za Zhi* 2011;9:104-6.
6. Kim JS. Post-stroke Mood and Emotional Disturbances: Pharmacological Therapy Based on Mechanisms. *J Stroke* 2016;18:244-55.
7. Thaut MH, McIntosh GC. Neurologic Music Therapy in Stroke Rehabilitation. *Curr Phys Med Rehab Rep* 2014;2:106-13.
8. Kim DS, Park YG, Choi JH, et al. Effects of Music Therapy on Mood in Stroke Patients. *Yonsei Med J* 2011;52:977.
9. Sarkamo T, Tervaniemi M, Laitinen S, et al. Music listening enhances cognitive recovery and mood after middle cerebral artery stroke. *Brain* 2008;131:866-76.

10. Särkämö T. Music for the ageing brain: Cognitive, emotional, social, and neural benefits of musical leisure activities in stroke and dementia. *Dementia* 2017;17:670–85.
11. Fatoye FO, Mosaku SK, Komolafe MA, et al. Depressive symptoms and associated factors following cerebrovascular accident among Nigerians. *J Mental Health* 2009;18:224–32.
12. Facai L, Dehong H, Nana H, et al. Effect of music therapy derived from the five elements in Traditional Chinese Medicine on post-stroke depression. *J Tradit Chin Med* 2017;37:675–80.
13. Synhaeve NE, Arntz RM, Maaijwee NAM, et al. Poor Long-Term Functional Outcome After Stroke Among Adults Aged 18 to 50 Years: Follow-Up of Transient Ischemic Attack and Stroke Patients and Unelucidated Risk Factor Evaluation (FUTURE) Study. *Stroke* 2014;45:1157–60.
14. Ahn D-H, Lee Y-J, Jeong J-H, et al. The Effect of Post-Stroke Depression on Rehabilitation Outcome and the Impact of Caregiver Type as a Factor of Post-Stroke Depression. *Ann Rehabil Med* 2015;39:74.
15. Hesamzadeh A, Dalvandi A, Bagher Maddah S, et al. Family Adaptation to Stroke: A Metasynthesis of Qualitative Research based on Double ABCX Model. *Asian Nurs Res* 2015;9:177–84.
16. Creasy KR, Lutz BJ, Young ME, et al. Clinical Implications of Family-Centered Care in Stroke Rehabilitation. *Rehabil Nurs* 2015;40:349-59.
17. Krieger T, Feron F, Dorant E. Developing a complex intervention programme for informal caregivers of stroke survivors: The Caregivers' Guide. *Scand J Caring Sci* 2016;31:146–56.
18. Given BA, Sherwood P, Given CW. Support for Caregivers of Cancer Patients: Transition After Active Treatment. *Cancer Epidemiol Biomarkers Prev* 2011;20:2015–21.
19. Ahtisaari M, Karanam K. Music and Emotion. 2015. Available from: <http://syncproject.co/blog/2015/7/21/music-and-emotion>. Accessed on: 9 September 2019.
20. Raglio A. Effects of music and music therapy on mood in neurological patients. *World J Psychiatry* 2015;5:68.
21. Guétin S, Soua B, Voiriot G, et al. The effect of music therapy on mood and anxiety–depression: An observational study in institutionalised patients with traumatic brain injury. *Ann Physic Rehabil Med* 2009;52:30–40.
22. Kučikienė D, Praninskienė R. The impact of music on the bio-electrical oscillations of the brain. *Acta Med Litu* 2018;25:101-6.
23. Schaefer H-E. Music-Evoked Emotions-Current Studies. *Frontiers Neurosci* 2017;11:600.
24. Ramdinmawii E, Mittal VK. The effect of music on the human mind: A study using brainwaves and binaural beats. 2nd International Conference on Telecommunication and Networks (TEL-NET) 2017. Available from: <https://doi.org/10.1109/telnet.2017.8343514>. Accessed on: 6 September 2019.
25. Isaac V, Stewart R, Krishnamoorthy E. Caregiver Burden and Quality of Life of Older Persons With Stroke: A Community Hospital Study in South India. *J Appl Gerontol* 2011;30:643-654.
26. Tsai Y-H, Lou M-F, Feng T-H, et al. Mediating effects of burden on quality of life for caregivers of first-time stroke patients discharged from the hospital within one year. *BMC Neurology* 2018;18:50.
27. Jeong Y-G, Myong J-P, Koo J-W. The modifying role of caregiver burden on predictors of quality of life of caregivers of hospitalized chronic stroke patients. *Disabil Health J* 2015;8:619–25.