


**The concentration of potentially toxic elements (PTEs) in the coffee products: a systematic review and meta-analysis**

Amin Mousavi Khaneghah, Trias Mahmudiono, Fardin Javanmardi, Behrouz Tajdar-oranj , Amene Nematollahi, Mohadeseh Pirhadi, Yadolah Fakhri

DOI: <https://doi.org/10.1007/s11356-022-23110-9>





# Environmental Science and Pollution Research



[Environmental Science and Pollution Research](#) > [Editors](#)

## Editors

### Editor-in-Chief

#### Philippe Garrigues

Institut des Sciences Moléculaires

Université de Bordeaux 1, France

E-mail: [philippe.garrigues@u-bordeaux.fr](mailto:philippe.garrigues@u-bordeaux.fr)

### Editorial Assistants

#### Fanny Creusot, Florence Delavaud, Giulia Marinaccio

Institut des Sciences Moléculaires

Université de Bordeaux 1, France

E-mail: [edito-ism@u-bordeaux.fr](mailto:edito-ism@u-bordeaux.fr)

### Editors

**Mohamed M. Abdel-Daim**, Pharmacology Department, Faculty of Veterinary Medicine, Suez Canal University, Ismailia, Egypt (*Environmental Pharmacology and Toxicology, Aquatic Toxicology, Nanotoxicology, Nanopharmacology, Nanomedicine, Alternative Medicine*)

**Lotfi Aleya**, National Center for Scientific Research (CNRS), Franche-Comté University, Faculty of Biology, la Bouloie and Faculty of Medicine, Hauts du Chazal, France (*Epidemiology, Protistology, Toxicology, Parasitology, Medicine, Embryotoxicity, Environmental Neurology, Freshwater and Marine Ecosystems*)

**Nicholas Apergis**, Department of Banking and Financial Management, University of Piraeus, Greece (*Economics of Energy, Energy Finance, Renewable Eenergy*)

**Kitae Baek**, School of Civil/Environmental/Resources and Energy Engineering (Major in Environmental Engineering), Jeonbuk National University, Jeollabukdo, Republic of Korea

*(Remediation of Metals-contaminated Soil, Adsorption of Metals, Oxidation Process, Geochemistry of Arsenic)*

**Giovanni Benelli**, Department of Agriculture, Food and Environment, University of Pisa, Italy  
*(Invertebrate ecotoxicology and behavior; Pesticides and nanoparticles)*

**Luděk Bláha**, RECETOX, Faculty of Science, Masaryk University, Brno, Czech Republic *(Ecotoxicology, Environmental Toxicology, Water quality, Organic pollutants, Endocrine disruption, Adverse Outcome Pathways)*

**Angeles Blanco**, Department of Chemical Engineering, Complutense University of Madrid, Spain  
*(Environmental Engineering)*

**Thomas Bucheli**, Agroscope, Zurich, Switzerland *(Environmental organic chemistry, analytical chemistry)*

**Tito Cadaval Jr.**, School of Chemistry and Food, Federal University of Rio Grande - FURG, Brazil  
*(Physicochemical treatment of water and wastewater and unity)*

**Eyup Dogan**, Department of Economics, Abdullah Gül University, Kayseri, Turkey *(Energy Economics, Environmental Economics, Applied Econometrics)*

**Guilherme Luiz Dotto**, Chemical Engineering Department, Federal University of Santa Maria-UFSM, Brazil *(Physicochemical treatments for wastewaters)*

**Robert Duran**, Research Institute for Materials and Environment (IPREM), University of Pau, France  
*(Microbiology)*

**Christian Gagnon**, Environment and Climate Change Canada, Montreal, Canada *(Aquatic biogeochemistry, trace metals, nanomaterials, pharmaceuticals, wastewater)*

**Baojing Gu**, College of Environmental and Resource Sciences / School of Public Affairs, Zhejiang University, China *(biogeochemical cycle, environmental management, cost and benefit, policy)*

**Ester Heath**, Department of Environmental Sciences, Jožef Stefan Institute, Slovenia *(Environmental Organic Chemistry)*

**Thomas Hein**, Institute of Hydrobiology and Aquatic Ecosystem Management, University for Natural Resources and Life Sciences, Vienna, Austria *(Water Ecology and Nutrient Dynamics)*

**Roula Inglesi-Lotz**, Department of Economics, University of Pretoria, South Africa *(Energy Economics, Electricity demand, Energy efficiency and economic growth)*

**Roland Kallenborn**, Norwegian University of Life Sciences (NMBU), Ås, Norway *(Chemical Technology and Engineering)*

**Ioannis A. Katsoyiannis**, Department of Chemistry, Aristotle University of Thessaloniki, Greece  
*(Water and wastewater treatment, water reuse, adsorption, oxidation, advanced oxidation processes, ozonation, membrane water treatment, arsenic removal, chromate removal, antimony removal, heavy metals removal)*

**George Z. Kyzas**, Department of Chemistry, International Hellenic University, Kavala, Greece  
*(Adsorption, Photocatalysis, Nanomaterials, Chemical Technology, Wastewater treatment, Graphene)*

**Gerhard Lammel**, Research Centre for Toxic Compounds in the Environment, Masaryk University, Brno, Czech Republic/Max Planck Institute for Chemistry, Mainz, Germany *(Atmospheric Processes and Air Pollution)*

**Séverine Le Faucheur**, IPREM-University of Pau and Pays de l'Adour, France *(Trace metals, Mercury,*

*Aquatic ecosystems, Microorganisms, Bioavailability)*

**Shimin Liu**, John and Willie Leone Family, Department of Energy and Mineral Engineering, University Park, PA, USA (*Coal Sciences*)

**Wei Liu**, Department F.-A. Forel for Environmental and Aquatic Sciences, University of Geneva, Genève, Switzerland (*Metals, Nanoparticles, Microorganisms, Invertebrates, Ecotoxicology, Aquatic Chemistry*)

**Philippe Loubet**, Bordeaux INP/ University of Bordeaux, France (*Life cycle assessment, Eco-design*)

**Christopher Lowe**, UK (*Soil Ecology, Earthworm Ecotoxicology, Biological Land Remediation*)

**Elena Maestri**, Department of Chemistry, Life Sciences and Environmental Sustainability, University of Parma, Italy (*Phytoremediation and Ecosystem Restoration*)

**Luke Mosley**, Faculty of Sciences, University of Adelaide, Australia (*Biogeochemistry of soil and water (metals, nutrients, organic matter, pH), effects of drought and climate change on soil and water quality, acid sulfate soils*)

**Bruno Nunes**, Centre for Environmental and Marine Studies (CESAM), University of Aveiro, Portugal (*Ecotoxicology, Environmental Toxicology, Aquatic ecotoxicology, Toxicology, Pharmacology, Biomarkers, Pharmaceutical drugs*)

**Ilhan Ozturk**, Faculty of Economics and Business, Cag University, Mersin, Turkey (*Energy Economics, Environmental Economics, International Economics*)

**Cinta Porte**, Department of Environmental Chemistry, Institute of Environmental Assessment and Water Research, Barcelona, Spain (*Aquatic Toxicology ; Ecotoxicology ; Impact and interactions of Organic Compounds*)

**Diane Purchase**, Middlesex University, London, UK (*Environmental Microbiology, Bioremediation, Biotechnology, Contaminated Land, E-Waste*)

**Sami Rtimi**, Swiss Federal Institute of Technology, Lausanne, Switzerland (*Photocatalytic materials for environmental remediation (wastewater and indoor air) )*

**Michel Sablier**, Muséum National d'Histoire Naturelle, Centre de Recherche sur la Conservation, Sorbonne Universités, Paris, France (*Conservation science, degradation processes, separation techniques and mass spectrometry*)

**Constantini Samara**, Department of Chemistry, Aristotle University of Thessaloniki, Greece (*Atmospheric Processes and Air Pollution*)

**VVSS Sarma**, CSIR-National Institute of Oceanography, Visakhapatnam, India (*Marine Science, Marine Chemistry/Chemical Oceanography*)

**Marcus Schulz**, University of Osnabrueck, Germany (*Environmental modelling, Reactive transport modelling, Hydrology, Biogeochemistry, Limnology, Sedimentology, Applied Statistics*)

**Arshian Sharif**, UCSI Graduate Business School, UCSI University, Kuala Lumpur, Malaysia (*Energy Finance, Energy Economics, Environmental Economics, Tourism Economics*)

**Gangrong Shi**, School of Life Sciences, Huaibei Normal University, China (*Plant ecotoxicology; Plant physiology, phytoremediation of heavy metals*)

**José Dinis Silvestre**, Civil Engineering Research and Innovation for Sustainability (CERIS), Instituto Superior Técnico, Universidade de Lisboa, Portugal (*Environmental impacts and toxicity of construction materials; Life cycle assessment; Sustainable construction*)

**Alexandros I. Stefanakis**, School of Environmental Engineering, Technical University of Crete, Chania, Greece (*Nature-based solutions, Constructed Wetlands, Natural treatment systems, Sustainable water and wastewater treatment, Ecological Engineering, Municipal and industrial wastewater management, Water reuse, Sludge dewatering*)

**Georg Steinhauser**, Institut für Angewandte Synthesechemie & TRIGA Center Atominstitut, Vienna, Austria (*Environmental radioactivity, nuclear forensics, radioanalytical methods*)

**Hongwen Sun**, College of Environmental Science and Engineering, Nankai University, China (*Environmental Organic Chemistry*)

**Roberto Terzano**, Department of Soil, Plant and Food Sciences, University of Bari "Aldo Moro", Italy (*Trace Metals, Soil, Plant, Rhizosphere, X-Ray Analyses, Soil Remediation*)

**Gérald Thouand**, Laboratory for Environment and Food Process Engineering, University of Nantes, France (*Biodegradation; Microbial Biosensor; Microbiology; Biosystems*)

**Ricardo A. Torres Palma**, Instituto de Química, Universidad de Antioquia, Medellín, Colombia

**Rongrong Wan**, Nanjing Institute of Geography and Limnology, Chinese Academy of Sciences, China (*Remote Sensing; Land Use/Land Cover; Vegetation Cover; Wetland; Ecosystem Services; Carbon Sequestration; Watershed Geography*)

**Ta Yeong Wu**, Chemical Engineering Discipline, School of Engineering, Monash University, Malaysia (*Biofuel production; Biomass valorization; Biorefinery; Waste reuse; Waste treatment*)

**Zhihong Xu**, Environmental Futures Research Institute and School of Biomolecular and Physical Sciences, Griffith University, Australia (*Soil Organic Matter Dynamics and Nutrient Cycling; Biogeochemical Cycles in Terrestrial Ecosystems; climate Change Adaptation and Mitigation*)

**Xianliang Yi**, School of Ocean Science and Technology, Dalian University of Technology, China (*Water Quality, Ecological Risk assessment, Aquatic Toxicology*)

**Weiming Zhang**, School of Environment, State Key Laboratory of Pollution Control and Resource Reuse, State Environmental Protection Engineering Center for Organic Chemical Industrial Waste Water Disposal and Resources Reuse, Nanjing University, China (*Water Pollution Control and Resources Reuse; Adsorption and Membrane Filtration; Advanced Oxidation Technologies*)

## **Editorial Board**

**Marijan Ahel**, Zagreb, Croatia

**Imran Ali**, New Delhi, India

**José Luis Balcázar**, Girona, Spain

**Åke Bergman**, Stockholm, Sweden

**Elena Bernardi**, Bologna, Italy

**Thierry Burgeot**, Nantes, France

**Boguslav Buszewski**, Torun, Poland

**Xinde Cao**, Shanghai, China

**I-Cheng Chang**, Ylan City, Taiwan

**Yi-Ping Chen**, Xi'an, China

**Teresa Cutright**, Akron, OH, USA

**Mitchell D. Erickson**, Piscataway, NJ, USA

**Lee Ferguson**, Durham, NH, USA  
**Shashikanth Gajaraj**, Chesterfield, MO, USA  
**Stanislaw Gawronski**, Warsaw, Poland  
**Christian George**, Lyon, France  
**Jizheng He**, Beijing, China  
**Henner Hollert**, Aachen, Germany  
**Tomohiko Isobe**, Tsukuba-City, Japan  
**Rajeev Jain**, Gwalior, India  
**Allan Astrup Jensen**, Copenhagen, Denmark  
**Ulrike Kammann**, Hamburg, Germany  
**Hidetoshi Kuramochi**, Tsukuba, Japan  
**Kenneth M.Y. Leung**, Hong Kong, China  
**Filippo Maggi**, Camerino, Italy  
**Laura Maxim**, Paris, France  
**Laura McConnell**, Research Triangle Park, NC, USA  
**Costantinos M. Michael**, Nicosia, Cyprus  
**Maurice Millet**, Strasbourg, France  
**Jochen Mueller**, Brisbane, Australia  
**Margaret Murphy**, Hong Kong, China  
**Takeshi Nakano**, Suma-ku, Kobe, Japan  
**Spiros Pergantis**, Heraklion, Crete, Greece  
**José Paulo Pinheiro**, Vandoeuvre-les-Nancy, France  
**Susan Richardson**, Columbia, SC, USA  
**Andreas Schäffer**, Aachen, Germany  
**Peter Schroeder**, Munich, Germany  
**Staci Simonich**, Corvallis, OR, USA  
**Marie-Odile Simonnot**, Nancy, France  
**Panayotis Siskos**, Athens, Greece  
**Cafer Turgut**, Aydin, Turkey  
**Vitor Vilar**, Porto, Portugal  
**Emmanuelle Vulliet**, Villeurbanne, France  
**Roland Weber**, Göppingen, Germany  
**Steve Wiseman**, Lethbridge, Canada  
**Fengchang Wu**, Beijing, China  
**Harald G. Zechmeister**, Vienna, Austria  
**Hao Zhang**, Lancaster, United Kingdom  
**Jiabao Zhang**, Nanjing, China  
**Qixing Zhou**, Nankai, China  
**Yong-Guan Zhu**, Xiamen, China  
**Uri Zoller**, Haifa, Israel



# Environmental Science and Pollution Research



Environmental Science and Pollution Research > Volumes and issues > Volume 29, issue 52

Search within journal


## Volume 29, issue 52, November 2022

103 articles in this issue

### [Fuel cell-based electric vehicles technologies and challenges](#)

Tarek Selmi, Ahmed Khadhraoui & Adnen Cherif


Trend Editorial | Published: 29 September 2022 | Pages: 78121 - 78131



### [Potential implementation of trees to remediate contaminated soil in Egypt](#)

Heba Bedair, Soumya Ghosh ... Samar Sami AlKafaas


Review Article | Published: 30 September 2022 | Pages: 78132 - 78151



### [The concentration of potentially toxic elements \(PTEs\) in the coffee products: a systematic review and meta-analysis](#)

Amin Mousavi Khaneghah, Trias Mahmudiono ... Yadolah Fakhri


Review Article | Published: 30 September 2022 | Pages: 78152 - 78164



### [Association between solid fuel combustion and diabetes mellitus: a systematic review and meta-analysis](#)

Ning Li, Wanying Su ... Yehuan Sun


Review Article | Published: 01 October 2022 | Pages: 78165 - 78177



## [The toxic cosmopolitan cyanobacteria \*Moorena producens\*: insights into distribution, ecophysiology and toxicity](#)

Emily Curren, Chui Pin Leaw ... Sandric Chee Yew Leong


Review Article | Published: 03 October 2022 | Pages: 78178 - 78206



## [Sustainable conversion of saturated adsorbents \(SAs\) from wastewater into value-added products: future prospects and challenges with toxic per- and poly-fluoroalkyl substances \(PFAS\)](#)

Amrita Nighojkar, Vikas Kumar Sangal ... Balasubramanian Kandasubramanian


Review Article | Published: 03 October 2022 | Pages: 78207 - 78227



## [Sustainable green roofs: a comprehensive review of influential factors](#)

Mohsen Shahmohammad, Majid Hosseinzadeh ... Nasrin Aghamohammadi


Review Article | Published: 03 October 2022 | Pages: 78228 - 78254



## [Photocatalytic dye degradation using nickel ferrite spinel and its nanocomposite](#)

Parth Shah, Kartik Joshi ... Femina J. Patel


Research Article | Published: 11 June 2022 | Pages: 78255 - 78264



## [A novel PM2.5 concentrations probability density prediction model combines the least absolute shrinkage and selection operator with quantile regression](#)

Shaomei Yang & Haoyue Wu


Research Article | Published: 11 June 2022 | Pages: 78265 - 78291



## [Hexavalent chromium reduction and bioremediation potential of \*Fusarium proliferatum\* S4 isolated from chromium-contaminated soil](#)

Bing Shan, Ruixia Hao ... Yubo Ye


Research Article | Published: 11 June 2022 | Pages: 78292 - 78302



## [Assessment of microcystins in surface water and irrigated vegetables in Kwaru stream, Hayin Danmanji, Kaduna-Nigeria](#)

Halima Abdullahi, Yahuza Tanimu ... Mathias Ahii Chia


Research Article | Published: 11 June 2022 | Pages: 78303 - 78313



## [Utilization of nanosize spent oil shale for water treatment: application of top-down nanonization technology for solid residues](#)

Adnan Abu-Surrah & Yahya Al-Degs

Research Article | Published: 11 June 2022 | Pages: 78314 - 78329





## [An empirical assessment of electricity consumption and environmental degradation in the presence of economic complexities](#)


Elma Satrovic & Festus Fatai Adedoyin

Research Article | [Open Access](#) | Published: 11 June 2022 | Pages: 78330 - 78344

## [Carbon emission of China's power industry: driving factors and emission reduction path](#)

Xiaoyuan Wu, Changxin Xu ... Chenjun Zhang


Research Article | Published: 11 June 2022 | Pages: 78345 - 78360



## [Panel data analysis of energy conservation and emission reduction on high-quality development of logistics industry in Yangtze River Delta of China](#)

Linbang Fan, Hui Liu ... Cunfang Li


Research Article | Published: 11 June 2022 | Pages: 78361 - 78380



## [Assessing the relationship between air pollution, agricultural insurance, and agricultural green total factor productivity: evidence from China](#)

Houjian Li, Mengqian Tang ... Lili Guo


Research Article | Published: 11 June 2022 | Pages: 78381 - 78395



## [Impacts of engineered iron nanoparticles on oxidative stress, fatty acid composition, and histo-architecture of the smooth scallop \*Flexopecten glaber\*](#)

Imen Rabeh, Khaoula Telahigue ... Lazhar Mhadhbi


Research Article | Published: 11 June 2022 | Pages: 78396 - 78413



## [Future refrigerants with low global warming potential for residential air conditioning system: a thermodynamic analysis and MCDM tool optimization](#)

Rajendran Prabakaran, Vinothkumar Sivalingam ... Govindasamy Praveen Kumar


Research Article | Published: 11 June 2022 | Pages: 78414 - 78428



## [Moderate exercise relieves fluoride-induced liver and kidney inflammatory responses through the IKK \$\beta\$ /NF \$\kappa\$ B pathway](#)

Yanghuan Yu, Ruiyan Niu ... Zilong Sun


Research Article | Published: 11 June 2022 | Pages: 78429 - 78443



## [Integrated ozone-sono-Fenton for the enhanced degradation of acid orange 7: process optimization and kinetic evaluation](#)

Jalal Basiri Parsa, Mahya Alamdar & Farnaz Jafari


Research Article | Published: 11 June 2022 | Pages: 78444 - 78456



## [Analysis of the decoupling state and driving forces of China's construction industry under the carbon neutrality target](#)

Shenghua Yan & Weigong Chen


Research Article | Published: 12 June 2022 | Pages: 78457 - 78471



## [Mesoporous BiVO<sub>4</sub>/TiO<sub>2</sub> heterojunction: enhanced photoabsorption and photocatalytic ability through promoted charge transfer](#)

Nada Y. Tashkandi, Soha M. Albukhari & Adel A. Ismail

Research Article | Published: 13 June 2022 | Pages: 78472 - 78482



## [Relationship between meteorological factors and mortality from respiratory diseases in a subtropical humid region along the Yangtze River in China](#)

Lingxiang Yu, Junjun Zhu ... Faming Pan


Research Article | Published: 13 June 2022 | Pages: 78483 - 78498



## [Adsorption properties and mechanism of ginkgo biloba leaf-based materials for Cd \(II\) in aqueous solution](#)

Yan-rong Cai, Yang Song & Chun Chang


Research Article | Published: 13 June 2022 | Pages: 78499 - 78508



## [Prediction of black carbon in marine engines and correlation analysis of model characteristics based on multiple machine learning algorithms](#)

Ying Sun, Lin Lü ... Peng Lee


Research Article | Published: 13 June 2022 | Pages: 78509 - 78525



## [Urbanization and carbon emissions: looking at the role of mobile phone adoption in Sub-Saharan African countries](#)

Lasbrey I. Anochiwa, Tobechei F. Agbanike ... Nkama Nnachi Otta


Research Article | Published: 13 June 2022 | Pages: 78526 - 78541



## [Effects of light source and inter-species mixed culture on the growth of microalgae and bacteria for nutrient recycling and microalgae harvesting using black odorous water as the medium](#)

Xiaoyan Wang, Shaoxuan Ding ... Jie Lu


Research Article | Published: 13 June 2022 | Pages: 78542 - 78554



## [Effects of chemical mixtures on liver function biomarkers in the Korean adult population: thresholds and molecular mechanisms for non-alcoholic fatty liver disease involved](#)

Hai Duc Nguyen & Min-Sun Kim

Research Article | Published: 13 June 2022 | Pages: 78555 - 78587



## [The asymmetric effect of temperature, exchange rate, metals, and investor sentiments on solar stock price performance in China: evidence from QARDL approach](#)

Chien-Chiang Lee, Farzan Yahya & Asif Razzaq

Research Article | Published: 13 June 2022 | Pages: 78588 - 78602

## [Nitrate in groundwater and agricultural products: intake and risk assessment in northeastern Iran](#)

Mohammad Zendeabad, Majid Mostaghelchi ... Willibald Loiskandl


Research Article | [Open Access](#) | Published: 13 June 2022 | Pages: 78603 - 78619



## [Sol-gel synthesis of TiO<sub>2</sub> on Co<sub>3</sub>O<sub>4</sub>-coated sporopollenin exine microcapsules \(SECs\) and photocatalytic performance of new semiconductor heterojunction material](#)

Hatice Çağlar Yılmaz, Funda Ersoy Atalay ... Sema Erdemoğlu


Research Article | Published: 13 June 2022 | Pages: 78620 - 78636



## [Improvements in SO<sub>2</sub> pollution in India: role of technology and environmental regulations](#)

Jayanarayanan Kuttippurath, Vikas Kumar Patel ... Ajay Singh


Research Article | Published: 13 June 2022 | Pages: 78637 - 78649



## [Improved tunicate swarm search-based MPPT for photovoltaic on a “grid-connected” inverter system](#)

Godina Venkata Lakshmi & K. Harinadha Reddy


Research Article | Published: 13 June 2022 | Pages: 78650 - 78665



## [The nonlinear effect of land freight structure on carbon emission intensity: new evidence from road and rail freight in China](#)

Rujia Chen, Xiaoning Wang ... Qian Luo


Research Article | Published: 13 June 2022 | Pages: 78666 - 78682



## [Spillover connection between oil prices, energy risk exposure, and financial stability: implications for the COVID-19 pandemic](#)

Haoming Shi & Haiyang Zheng


Research Article | Published: 13 June 2022 | Pages: 78683 - 78697



## [Bioavailability of inhaled or ingested PFOA adsorbed to house dust](#)

Åsa Gustafsson, Bei Wang ... Leo W. Y. Yeung

Research Article | [Open Access](#) | Published: 14 June 2022 | Pages: 78698 - 78710



## [Handheld two-stroke engines as an important source of personal VOC exposure for olive farm workers](#)

Akif Ari, Pelin Ertürk Ari ... Eftade O. Gaga

Research Article | Published: 14 June 2022 | Pages: 78711 - 78725



## [Sustainability reporting and corporate reputation of Malaysian IPO companies](#)


Yasir Abdullah Abbas, Waqas Mehmood ... Attia Aman-Ullah

Research Article | Published: 14 June 2022 | Pages: 78726 - 78738

## [Augmenting the productivity of tubular solar still using low-cost energy storage materials](#)

Ritesh Krishna Sambare, Satish Kumar Dewangan ... Sandeep Joshi

Research Article | Published: 14 June 2022 | Pages: 78739 - 78756



## [Do Twitter sentiments really effective on energy stocks? Evidence from the intercompany dependency.](#)

Emrah Sitki Yilmaz, Asli Ozpolat & Mehmet Akif Destek


Research Article | Published: 14 June 2022 | Pages: 78757 - 78767

Attitude

## [Economic, social, and environmental assessment of farming Nile tilapia in net-cages in a reservoir in hot semi-arid region during an extended drought event](#)

Gustavo Gonzaga Henry-Silva, Júlio César da Silva Cacho ... Wagner C. Valenti


Research Article | Published: 14 June 2022 | Pages: 78768 - 78779



## [Research on the impact of circular economy on total factor carbon productivity in China](#)

Tiening Cui & Yang Zhang

Research Article | Published: 14 June 2022 | Pages: 78780 - 78794



## [A comparative analysis of renewable and non-renewable energy generation to relegate CO<sub>2</sub> emissions and general costs in household systems](#)

Mudassar Hussain, Manzoor Sultan ... Muhammad Imran


Research Article | Published: 14 June 2022 | Pages: 78795 - 78808



## [Flocculation of \*Chlorella vulgaris\*-induced algal blooms: critical conditions and mechanisms](#)

Ping Zhang, Sihan Zhu ... Han Wang


Research Article | Published: 14 June 2022 | Pages: 78809 - 78820



## [Short-term associations of ambient air pollution with hospital admissions for ischemic stroke in 97 Japanese cities](#)

Kohei Hasegawa, Teruomi Tsukahara & Tetsuo Nomiyama


Research Article | Published: 14 June 2022 | Pages: 78821 - 78831



## [Activation of peroxydisulfate by modified coagulation sludge for bisphenol A degradation](#)

Jie Bai, Yihui Li ... Qing Wang


Research Article | Published: 14 June 2022 | Pages: 78832 - 78847



## [Utilization of zinc-ferrite/water hybrid nanofluids on thermal performance of a flat plate solar collector—a thermal modeling approach](#)

Prakasam Michael Joseph Stalin, Thottipalayam Vellingiri Arjunan ... Pasupathi Manoj Kumar


Research Article | Published: 14 June 2022 | Pages: 78848 - 78861



## [Construction and carbon source optimization of a microbial-plant coupled reactor for treating acid mine drainage](#)

Hai Lin, Yalu Tang & Yingbo Dong


Research Article | Published: 14 June 2022 | Pages: 78862 - 78873



## [α-\(Fe, Cu\)OOH/RGO nanocomposites for heterogeneous photo-Fenton-like degradation of ciprofloxacin under visible light irradiation](#)

Junge Xu, Die Hu ... Ziwei Zhang


Research Article | Published: 14 June 2022 | Pages: 78874 - 78886



## [Influences of a novel cylindrical solar dryer on farmer's income and its impact on environment](#)

Kirtika Sharma, Surendra Kothari ... Maga Ram Patel

Research Article | Published: 14 June 2022 | Pages: 78887 - 78900



✓ You have access to our articles

For authors



# The concentration of potentially toxic elements (PTEs) in the coffee products: a systematic review and meta-analysis

Amin Mousavi Khaneghah<sup>1</sup> · Trias Mahmudiono<sup>2</sup> · Fardin Javanmardi<sup>3</sup> · Behrouz Tajdar-oranj<sup>4</sup> · Amene Nematollahi<sup>5</sup> · Mohadeseh Pirhadi<sup>6</sup> · Yadolah Fakhri<sup>7</sup>

Received: 12 May 2022 / Accepted: 14 September 2022 / Published online: 30 September 2022  
© The Author(s), under exclusive licence to Springer-Verlag GmbH Germany, part of Springer Nature 2022

## Abstract

Coffee is one of the most consumed products globally, and its contamination with potentially toxic elements (PTEs) occurs throughout the production chain and production. Therefore, the current meta-analysis study aimed to estimate the concentration of essential elements (Cu and Co) and the contamination of PTEs (Ni, Cr, Pb, As, and Cd) in coffee. The recommended databases, including PubMed, Scopus, and ScienceDirect, were investigated to collect data regarding the contamination of PTEs in coffee products from 2010 to 2021. Among 644 retrieved citations in the identification step, 34 articles were included in the meta-analysis. The pooled mean concentration of essential elements in coffee products is much higher than that of toxic elements (Co (447.106 µg/kg, 95% CI: 445.695–448.518 µg/kg) > Ni (324.175 µg/kg, 95% CI: 322.072–326.278 µg/kg) > Cu (136.171 µg/kg, 95% CI: 134.840–137.503 µg/kg) > Cr (106.865 µg/kg, 95% CI: 105.309–108.421 µg/kg) > Pb (21.027 µg/kg, 95% CI: 20.824–21.231 µg/kg) > As (3.158 µg/kg, 95% CI: 3.097–3.219 µg/kg) > Cd (0.308 µg/kg, 95% CI: 0.284–0.332 µg/kg)). Results showed high differences between pooled concentrations of all PTEs in coffee products of different countries.

**Keywords** Potential toxic elements · Coffee products · PTEs · Systematic review · Meta-analysis

---

Responsible Editor: Philippe Garrigues

✉ Amin Mousavi Khaneghah  
amin.mousavi@ibprs.pl; mousavi.amin@gmail.com

- 1 Department of Fruit and Vegetable Product Technology, Prof. Waclaw Dąbrowski Institute of Agricultural and Food Biotechnology, Warsaw, Poland
- 2 Department of Nutrition, Faculty of Public Health, Universitas Airlangga, Surabaya, Indonesia
- 3 Department of Food Science and Technology, Faculty of Nutrition Science and Food Technology, Shahid Beheshti University of Medical Sciences, Tehran, Iran
- 4 Food and Drug Administration, Iran Ministry of Health and Medical Education, Tehran, Iran
- 5 Department of Food Safety and Hygiene, School of Health, Fasa University of Medical Sciences, Fasa, Iran
- 6 Department of Environmental Health Engineering, Division of Food Safety & Hygiene, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran
- 7 Food Health Research Center, Hormozgan University of Medical Sciences, Bandar Abbas, Iran

## Introduction

Chemical pollution of the environment (Guan et al. 2021; Li et al. 2021; Lin et al. 2021; Liu et al. 2011, 2020; Quan et al. 2021; Shi et al. 2021) followed by food contamination (Liu et al. 2022; Sun et al. 2022; Wang et al. 2022) has become a global concern. Coffee is an attractive beverage worldwide because of its pleasant aroma and taste. Coffee consumption varies according to geographical location (Godos et al. 2014). Flanders in Northern Europe and Bosnia and Herzegovina in Southern Europe are the largest consumers of coffee. The World Coffee Organization states that 500 billion cups of coffee are consumed annually (Dieng et al. 2017). Coffee and tea have been consumed for hundreds of years and have become an important part of cultural traditions and social life. In addition, people use coffee drinks to increase alertness and work productivity (Samet et al. 2018). The highest caffeine consumption in a typical meal is coffee, energy drinks, and caffeinated pills. The intermediate level is tea, and the lowest amount of soda (Rattan et al. 2015; van Dam et al. 2020). Caffeine is an alkaloid belonging to methylxanthine. It is also a chemical stimulant found naturally (Wolde 2014). Studies showed that coffee consumption has

chronic and acute effects on human health (Nędzarek et al. 2013). Caffeine can act as an antioxidant to prevent diseases. Caffeine also can reduce the risk of several chronic diseases such as liver cancer, diabetes, coronary heart disease, osteoporosis, and gastritis (Nartea et al. 2022). Moreover, the absorption of many vital nutrients such as vitamin B<sub>6</sub> is reduced by caffeine intake and the absorption of essential minerals such as magnesium (Mg), iron (Fe), and calcium (Ca) (de Mejia and Ramirez-Mares 2014; Escott-Stump 2008; Wolde 2014). However, consuming large amounts of this drink is not recommended because of some contaminants, including mycotoxins and potentially toxic elements (PTEs), besides to adverse effects of caffeine (Barrea et al. 2021; Batrinou et al. 2020; Berman et al. 2022; Khaneghah et al. 2019; Nourbakhsh and Tajbakhsh 2021; Pavithra 2021; Yazdanpanah et al. 2022).

It needs about 20 essential elements for the proper functioning of the human body. The list of metallic elements that are essential for the human body include sodium (Na), potassium (K), Mg, Ca, manganese (Mn), Fe, cobalt (Co), copper (Cu), zinc (Zn), molybdenum (Mo), and elements like tin (Sn), nickel (Ni), and vanadium (V), and are assumed not to be necessary for humans, according to chromium (Cr); based on experimental animal results, these metals were considered essential for the human body in the past (Zoroddu et al. 2019). Cu and Co are essential elements in the human body, a small amount of which plays a vital role in the body. The amount of Co we need is minimal, and only about 2 mg of it is needed for humans. High levels of Cu can cause cardiovascular disease such as cardiac arrhythmias. In addition, high levels of Cu lead to dementia, lung cancer, and lymphoma (Bost et al. 2016). The presence of Cu in the body interferes with the nervous system and causes Wilson's disease, and also, the presence of Cu in the brain leads to brain damage (Kalita et al. 2019).

PTEs are a serious and worrying environmental problem (Boumar et al. 2020; Gao et al. 2022; Mirmahdi et al. 2022). They are widely distributed in the environment and thus intake by humans through water, air, soil, and even the food chain (Alkheraz et al. 2019; Heshmati et al. 2020). Accumulating PTEs in the body has negatively affected human health (Fu and Xi 2020; Mirmahdi et al. 2021). These effects include interfering with the immune and nervous systems, endocrine disorders, and malformations (Khaneghah et al. 2020). Organic compounds of lead (Pb) can be absorbed through the skin and enter the brain, causing toxins in the central nervous system. Pb also causes oxidative stress and causes cells to lose antioxidants (Fu and Xi 2020). Adverse effects of Pb include chromosome aberrations, mutations, DNA breakdown, and inhibition of DNA synthesis (Ibrahim et al. 2020). Cadmium (Cd) is a toxic and dangerous metal. The World Health Organization limited the tolerable weekly Cd exposure to 50 µg

and recommended it as 0.007 mg/kg body weight (World Health Organization 2000). Depending on the route of exposure and duration, Cd can cause damage to the lung, liver, kidney, bone, testicle, and placenta (Hocaoğlu-Özyiğit and Genç 2020). Arsenic (As) is another PTE that causes damage and changes in DNA synthesis and cancer, cardiovascular and lung diseases, reproductive outcomes, and cognitive impairment in adults and children (Hu et al. 2022). Studies show that chronic exposure to As causes lung cancer, breast cancer, and bladder, kidney, and larynx diseases. It causes respiratory problems, infertility in adults, and premature birth in infants (Delgado Quezada et al. 2020; Khan et al. 2020; Pullella and Kotsopoulos 2020). Cr has a different effect on human health depending on the target dose and organ. Effects include respiratory problems, asthma, lung cancer, skin diseases such as dry skin, allergies, dizziness, general weakness, eye irritation, kidney stones, liver problems, gastrointestinal disorders, heart problems, blood disorders, reproduction problems, growth problems, nostril problems, and corneal damage (Ameri et al. 2021). Dental malformations such as discoloration and erosion are accompanied by high concentrations of Cr in the tongue papilla (Achmad and Auerkari 2017; Martineli et al. 2022; Teklay 2016). Ni is an element that may be widely derived from natural resources and human activities in the environment, air, water, and soil. Ni exposure causes digestive diseases, lung problems, lung cancer, and cardiovascular disease (Genchi et al. 2020).


This study systematically reviewed the prevalence of essential elements (Cu and Co) and potentially toxic elements (As, Cr, Cd, Pb, and Ni) in coffee products. Also, another goal of this study was to meta-analyze the results and identified the elements and geographical areas with the highest prevalence.

## Material and method

### Searching strategy

According to PRISMA guidelines, our study conducted the search strategy (Fig. 1) (Moher et al. 2010). International databases, including Scopus and PubMed (January 2000 and 20 June 2021), were screened to get papers on PTE concentrations in coffee products. Keywords referred to in the title and abstract of citations were ["heavy metal" OR "potentially toxic elements" OR "toxic elements" OR "elements" OR "toxic metal"] AND ["coffee products" OR "coffee" OR "caffeine" OR "cappuccino" OR "espresso"]. The reference list of papers was assessed for possible papers missed in the title and abstract screening step.

**Fig. 1** Process selection of papers based on PRISMA



### Inclusion and exclusion criteria and extraction of data

Our criteria included the following: detection of PTE concentrations in coffee products, full text available, contamination or prevalence investigation, no language limitation, no restriction for the study location, and present of mean, standard deviation, and/or range concentration of PTEs. One study was excluded because it did not meet the above criteria. One of the authors extracted the data in the retrieved paper. Data extracted from each article included the country, sample size, the mean and standard deviation of concentration, and measurement method.

### Meta-analysis of data

The concentration of PTEs in coffee products was meta-analyzed based on the standard error (SE) of the equation:

$$SE = \frac{SD}{\sqrt{n}} \quad (1)$$

where SD is the standard deviation and  $n$  is the number of samples when the  $I^2$  index is higher than 50% and heterogeneity is considerable (Higgins et al. 2008; Higgins and Thompson 2002; Quan and Zhang 2003); hence, the random-effect model was used for the meta-analysis of the concentration of PTEs in coffee products. PTE concentration's mean and standard deviation were converted to  $\mu\text{g}/\text{kg}$ .

A data meta-analysis was performed using Stata, version 14 (Stata Corporation, College Station, TX).

### Result and discussion

Our study included thirty-four papers with 151 data reports (Fig. 1). The overall rank order of PTEs in the coffee products was as follows: Co (447.106  $\mu\text{g}/\text{kg}$ , 95% CI: 445.695–448.518  $\mu\text{g}/\text{kg}$ ) > Ni (324.175  $\mu\text{g}/\text{kg}$ , 95% CI: 322.072–326.278  $\mu\text{g}/\text{kg}$ ) > Cu (136.171  $\mu\text{g}/\text{kg}$ , 95% CI: 134.840–137.503  $\mu\text{g}/\text{kg}$ ) > Cr (106.865  $\mu\text{g}/\text{kg}$ , 95% CI: 105.309–108.421  $\mu\text{g}/\text{kg}$ ) > Pb (21.027  $\mu\text{g}/\text{kg}$ , 95% CI: 20.824–21.231  $\mu\text{g}/\text{kg}$ ) > As (3.158  $\mu\text{g}/\text{kg}$ , 95% CI: 3.097–3.219  $\mu\text{g}/\text{kg}$ ) > Cd (0.308  $\mu\text{g}/\text{kg}$ ; 95% CI: 0.284–0.332  $\mu\text{g}/\text{kg}$ ). As it is obvious from this part of the result, the pooled mean concentration of essential elements in coffee products is much higher than that of toxic elements. Thus, the distribution of PTEs in coffee products is uneven because their content depends on numerous factors (Bost et al. 2016).

It should bear in mind that coffee plants need essential elements (like Co, Ni, and Cu) for their development, so they absorb them from water and soil selectively during growth which is influenced by the soil characteristics (chemical composition) in addition to the overall environmental growth conditions (climate, temperature, and agricultural practices) (Barbosa et al. 2014; Habte et al. 2016; Pigozzi et al. 2018). Petrović et al. (2020) also reported that the accumulation of



PTEs in plants relates to the plant's affinity to take up the appropriate element. Furthermore, there is usually a higher concentration of essential elements in the soil than toxic elements since they exist naturally in the rocks and release into water and soil through rock weathering and human activities (like mining) (Atamaleki et al. 2020; Pigozzi et al. 2018). Higher Co, Ni, and Cu pooled concentrations in the coffee products could also be attributed to the greater availability of these PTEs in the soils used for coffee plantation. Therefore, due to the active transmission of essential elements through the soil-root interface and their high content in the soil used for coffee cultivation, they might be readily absorbed by the plant and accumulated in the coffee beans compared with other PTEs (da Silva et al. 2017). As also evident in a study of da Silva et al. (2017), the elements of Cu, Ni, and Zn are necessary to the growth of the coffee plant, being utilized in the soil or coffee leaf, and are subsequently present in the coffee beans. Furthermore, some PTEs like Cu exist in pesticide ingredients applied in cultivation which assists the uptake of these PTEs by plants, validating its presence in coffee beans (da Silva et al. 2017). Albals et al. (2021) also noted that Co and Cu mainly exist at high concentrations in most plants when cultivated in contaminated soil. Cd is found at lower concentrations in such products than in fish.

PTE contamination in coffee products also has diverse forms associated with size, charge, hydrophobicity, activity, bioaccessibility, coffee variety, and type of processing (Alkherraz et al. 2019; Pohl et al. 2014; Şemen et al. 2017). The concentration of PTEs in coffee is also related to its processing, especially the roasting of coffee beans and coffee beverages' preparation method (Tiep 2014). When it comes

to coffee infusions, several researchers reported that the releasing amount of different PTEs to coffee infusions from the solid sample is different based on coffee type, preparation method, and nature of the metal (leachability and rate of complex formation or interaction with other components) (Ashu and Chandravanshi 2011; da Silva et al. 2017; Jeszka-Skowron et al. 2016; Welna et al. 2013; Winiarska-Mieczan et al. 2021).

### Concentration of As

As, a widespread toxic environmental contaminant, enters into the environment (air, water, and soil) through natural and anthropogenic (mainly pesticides used in agricultural practices) sources and is found in the natural world in organic, inorganic, and gaseous forms (Fu and Xi 2020; Petrović et al. 2020). Inorganic and gaseous forms of As are highly toxic, while organic As is considered relatively non-toxic in very low concentrations (Pullella and Kotsopoulos 2020). Exposure to As is of great concern. It has been related to adverse health effects, including cancer (skin, bladder, and lungs), damaged neurological growth, cardiovascular disorder, and diabetes (Pullella and Kotsopoulos 2020).

The rank order of countries based on the concentration of As in the coffee products was observed in the following: Brazil (178.882 µg/kg) > Netherland (74.981 µg/kg) > Greece (40.000 µg/kg) > Austria (20.050 µg/kg) > France (16.451 µg/kg) > Turkey (10.000 µg/kg) > Ethiopia (6.000 µg/kg) > Italy (1.250 µg/kg) > Germany (0.259 µg/kg) > Indonesia (0.100 µg/kg) ~ Romania (0.100 µg/kg) ~ Rwanda (0.100 µg/kg) ~ Slovenia (0.100 µg/kg)

**Table 1** Meta-analysis concentration of As in the coffee products (µg/kg)

Country	ES	Lower	Upper	Weight (%)	Heterogeneity statistic	Degrees of freedom	<i>P</i> value	<i>I</i> <sup>2</sup> (%)
Austria	20.050	2.350	59.152	4.43	58,897.79	1	<0.001	100.00
Brazil	178.882	128.720	229.043	0.39	4232.54	2	<0.001	100.00
Ethiopia	6.000	5.689	6.311	1.64	0.00	0		
France	16.451	15.290	17.612	7.27	92,121.16	2	<0.001	100.00
Germany	0.259	0.181	0.336	11.55	6968.40	5	<0.001	99.90
Greece	40.000	39.678	40.322	1.59	0.00	0		
Indonesia	0.100	0.094	0.106	2.84	0.00	0		
Italy	1.250	1.035	1.465	6.07	9419.07	4	<0.001	100.00
Netherland	74.981	6.382	143.580	1.62	1795.05	1	<0.001	99.90
Romania	0.100	0.097	0.103	5.68	0.00	1	1	0.00
Rwanda	0.100	0.096	0.104	2.84	0.00	0		
Slovenia	0.100	0.098	0.102	5.68	0.00	1	1	0.00
Turkey	10.000	8.303	11.697	0.12	0.00	0		
Western Balkan	0.100	0.094	0.106	48.28	554.78	18	<0.001	96.80
Overall	3.158	3.097	3.219	100.00	300,000.00	48	<0.001	100.00

ES effect size (mean of concentration)

kg) ~ Western Balkan (0.100 µg/kg) (Table 1). Since the maximum permitted level (MPL) of As in coffee is 1000 µg/kg (Petrović et al. 2020), it could be concluded that the As concentration of coffee products in all investigated countries does not show a risk to human health concerning this metal.

### Concentration of Ni

Ni is naturally present in the soil and widespread distribution in the environment (Genchi et al. 2020). This metal has commercial and industrial applications, including metallurgy, electronics, batteries, jet turbines, and medical tools (Şaylan et al. 2020). As an essential element for all living things, Ni has several beneficial impacts on different body organs' functions and metabolism, up to the permitted level. In this regard, Ni influences the liver and pancreas performances, the Fe absorption/metabolism, and the hematopoietic process (Gogoasa et al. 2013). However, in elevated concentrations, Ni can be dangerous for human health by creating respiratory disorders, including lung embolism, respiratory malfunction, asthma, chronic bronchitis, and allergic reactions (Nędzarek et al. 2013). As high long-time exposure to Ni is related to the development of different kinds of cancer such as lung, larynx,

and prostate, it is recognized by the International Agency for Research on Cancer (IARC) as a carcinogen (Nędzarek et al. 2013; Voica et al. 2016).

The rank order of countries based on the concentration of Ni in the coffee products was as follows: Turkey (9695.861 µg/kg) > France (8615.085 µg/kg) > Netherlands (5881.104 µg/kg) > India (3665.463 µg/kg) > Italy (2846.081 µg/kg) > Greece (2540.000 µg/kg) > Austria (2274.536 µg/kg) > Romania (1782.844 µg/kg) > Kenya (1630.881 µg/kg) > Brazil (1128.809 µg/kg) > Poland (1046.963 µg/kg) > Lebanon (842.300 µg/kg) > Bosnia (780.000 µg/kg) > Ethiopia (777.455 µg/kg) > Honduras (580.784 µg/kg) > Pakistan (410.000 µg/kg) > Hawaii (356.879 µg/kg) > Colombia (34.796 µg/kg) > Germany (5.047 µg/kg) > Western Balkan (4.471 µg/kg) > Slovenia (0.100 µg/kg) (Table 2).

There is no MPL for Ni in coffee beans, although Brazilian legislation authorizes an MPL of 5000 µg/kg for Ni in typical foods (da Silva et al. 2017; Pigozzi et al. 2018). Our results showed that the Ni concentrations in the coffee samples of all countries do not surpass this maximum level except for the coffee samples of Turkey, France, and the Netherlands.

**Table 2** Meta-analysis concentration of Ni in the coffee products (µg/kg)

Country	ES	Lower	Upper	Weight (%)	Heterogeneity statistic	Degrees of freedom	P value	I <sup>2</sup> (%)
Austria	2274.536	344.647	4204.426	2.63	23,640.73	1	<0.001	100.00
Bosnia	780.000	731.668	828.332	0.18	0.00	0		
Brazil	1128.809	1033.256	1224.361	5.40	7,300,000.00	8	<0.001	100.00
Colombia	34.796	3.540	101.104	3.95	146.58	1	<0.001	99.30
Ethiopia	777.455	774.516	780.394	13.63	1,400,000.00	7	<0.001	100.00
France	8615.085	1490.616	16,000.000	0.03	3581.61	1	<0.001	100.00
Germany	5.047	4.391	5.704	7.05	87,862.69	5	<0.001	100.00
Greece	2540.000	2514.223	2565.777	0.53	0.00	0		
Hawaii	356.879	124.475	589.284	0.18	48.95	4	<0.001	91.80
Honduras	580.784	565.420	596.147	1.16	0.12	1	0.732	0.00
India	3665.463	3429.303	3901.623	0.01	0.04	1	0.836	0.00
Italy	2846.081	2002.296	3689.867	3.33	35,250.89	5	<0.001	100.00
Kenya	1630.881	1584.530	1677.232	0.19	0.05	1	0.83	0.00
Lebanon	842.300	833.284	851.316	1.71	0.00	0		
Netherlands	5881.104	5516.598	6245.610	0.07	4.38	1	0.036	77.20
Pakistan	410.000	300.242	519.758	0.04	0.00	0		
Poland	1046.963	698.597	1395.328	4.60	13,874.64	8	<0.001	99.90
Romania	1782.844	1304.206	2261.481	2.59	1295.84	5	<0.001	99.60
Slovenia	0.100	0.097	0.103	5.01	0.00	1	1	0.00
Turkey	9695.861	1993.787	17,000.000	0.17	43,968.77	3	<0.001	100.00
Western Balkan	4.471	4.339	4.604	47.56	360,000.00	18	<0.001	100.00
Overall	324.175	322.072	326.278	100.00	310,000,000.00	91	<0.001	100.00

ES effect size (mean of concentration)

### Concentration of Cr

Both kinds of Cr ( $\text{Cr}^{3+}$  and  $\text{Cr}^{6+}$ ) could enter the air, water, and soil through natural processes and industrial activities (like steel, leather, chemical, electronic, painting, and textile manufacturing) (Teklay 2016).  $\text{Cr}^{3+}$ , an essential element in the body, influences metabolism by adjusting blood sugar levels in relatively small amounts (Gogoasa et al. 2013; Neđzarek et al. 2013). It is worth noting that  $\text{Cr}^{6+}$  is more active and toxic than  $\text{Cr}^{3+}$ . This element occurs at many high levels in industrial activities, which could release  $\text{Cr}^{6+}$  into the environment (Achmad and Auerkari 2017). However,  $\text{Cr}^{3+}$  is ubiquitous in the environment (water and soil) and biological systems (Voica et al. 2016). It is reported that exposure to a high concentration of Cr could result in several physiological effects, including skin harm; respiratory injuries; gastrointestinal disorders; cardiovascular diseases; hepatic failure; carcinogenic, genotoxic, and neurotoxic mutagenic effects; and reproductive and developmental toxicity (Teklay 2016).

The rank order of countries based on the concentration of Cr in the coffee products was as follows: Netherland (34,000.000  $\mu\text{g}/\text{kg}$ ) > France (20,000.000  $\mu\text{g}/\text{kg}$ ) > Egypt (2500.000  $\mu\text{g}/\text{kg}$ ) > Kenya (714.922  $\mu\text{g}/\text{kg}$ ) > India (610.321  $\mu\text{g}/\text{kg}$ ) > Indonesia (600.000  $\mu\text{g}/\text{kg}$ ) > Honduras (290.000  $\mu\text{g}/\text{kg}$ ) > Ethiopia (246.854  $\mu\text{g}/\text{kg}$ ) > Hawaii (239.039  $\mu\text{g}/\text{kg}$ ) > Brazil (150.930  $\mu\text{g}/\text{kg}$ ) > Colombia

(121.610  $\mu\text{g}/\text{kg}$ ) > Poland (60.000  $\mu\text{g}/\text{kg}$ ) > Bosnia (40.000  $\mu\text{g}/\text{kg}$ ) > Lebanon (35.000  $\mu\text{g}/\text{kg}$ ) > Germany (0.643  $\mu\text{g}/\text{kg}$ ) > Italy (0.134  $\mu\text{g}/\text{kg}$ ) > Austria (0.100  $\mu\text{g}/\text{kg}$ ) ~ Greece (0.100  $\mu\text{g}/\text{kg}$ ) (Table 3).

Similar to Ni, there are no reported maximum levels for the existence of the Cr metal in coffee products. At the same time, Brazilian regulation has established an MPL of 100  $\mu\text{g}/\text{kg}$  for this element in the food (da Silva et al. 2017). Our results showed that the pooled concentrations of Cr in coffee products of all investigated countries exceed the maximum permitted level except Poland, Bosnia, Lebanon, Germany, Italy, Austria, and Greece.

### Concentration of Cu

Cu is a vital element in numerous human body functions like antioxidant defense, Fe metabolism, enzyme functions, energy production, bone formation, and skin health in trace amounts (Bost et al. 2016; Gogoasa et al. 2017). Furthermore, lacking Cu may cause anemia, hair whitening, and bone distortion (Petrović et al. 2020). However, exposure to high concentrations of Cu is toxic and could result in stomach pain, nausea, diarrhea, peroxidation of macromolecules, and tissue injury (Anosike and Oranusi 2018; Bost et al. 2016).

The rank order of countries based on the concentration of Cu in the coffee products was Germany (38,000.000  $\mu\text{g}/\text{kg}$ )

**Table 3** Meta-analysis concentration of Cr in the coffee products ( $\mu\text{g}/\text{kg}$ )

Country	ES	Lower	Upper	Weight (%)	Heterogeneity statistic	Degrees of freedom	P value	I <sup>2</sup> (%)
Austria	0.100	0.098	0.102	11.70	0.00	1.00	1	0.00
Bosnia	40.000	28.181	51.819	1.34	0.00	0.00		
Brazil	150.930	139.951	161.908	13.08	54,518.83	8.00	<0.001	100.00
Colombia	121.610	102.443	140.778	11.82	3178.71	5.00	<0.001	99.80
Egypt	2500.000	2428.432	2571.568	0.05	0.00	0.00		
Ethiopia	246.854	209.473	284.236	19.86	1759.86	7.00	<0.001	99.60
France	20,000.000	220.000	46,000.000	0.01	4284.30	1.00	<0.001	100.00
Germany	0.643	0.258	1.028	12.37	30,137.84	5.00	<0.001	100.00
Greece	0.100	0.097	0.103	5.85	0.00	0.00		
Hawaii	239.039	54.928	423.150	2.30	142.07	4.00	<0.001	97.20
Honduras	290.000	279.191	300.809	1.65	0.00	1.00	1	0.00
India	610.321	308.647	911.996	1.62	744.43	3.00	<0.001	99.60
Indonesia	600.000	511.357	688.643	0.03	0.00	0.00		
Italy	0.134	-0.170	0.438	11.71	18,762.12	4.00	<0.001	100.00
Kenya	714.922	691.221	738.624	0.41	0.52	1.00	0.472	0.00
Lebanon	35.000	34.409	35.591	5.80	0.00	0.00		
Netherlands	34,000.000	233.240	79,000.000	0.00	4397.72	1.00	<0.001	100.00
Poland	60.000	36.362	83.638	0.40	0.00	0.00		
Overall	106.865	105.309	108.421	100.00	28,000,000.00	58.00	<0.001	100.00

ES effect size (mean of concentration)

kg) > Austria (20,000.000 µg/kg) > Kenya (19,000.000 µg/kg) ~ Netherland (19,000.000 µg/kg) > Honduras (17,000.000 µg/kg) > India (14,000.000 µg/kg) > Colombia (13,000.000 µg/kg) ~ Costa Rica (13,000.000 µg/kg) ~ Guatemala (13,000.000 µg/kg) ~ Poland (13,000.000 µg/kg) > Nicaragua (12,000.000 µg/kg) > Peru (12,000.000 µg/kg) > Ethiopia (9097.608 µg/kg) > Italy (6598.028 µg/kg) > Greece (6520.000 µg/kg) > Nigeria (3970.000 µg/kg) > France (3538.513 µg/kg) > Brazil (2804.322 µg/kg) > Egypt (2200.000 µg/kg) > Hawaii (146.762 µg/kg) > Indonesia (90.000 µg/kg) (Table 4).

Our results showed that the concentrations of Cu in coffee products of all studied countries meet the MPL for Cu, determined by Brazilian legislation (30,000 µg/kg) (Alkherraz et al. 2019; Anosike and Oranusi 2018), except in Germany. Cu is usually used in the composition of some pesticides used in coffee fields as an active substance in different forms (Pigozzi et al. 2018). It is necessary for plant growth and is absorbed by plants during cultivation. Senkondo et al. (2014) observed that Cu concentrations in the coffee samples grown in the Cu-based pesticide soils were much higher compared with those in untreated soils, probably due to more availability of Cu for plant uptake, which could be the main reason for observed differences in Cu concentrations in different

countries in the present study. However, the soil properties, climate conditions, and differences in coffee bean varieties are also influential factors (Alkherraz et al. 2019; Senkondo et al. 2014).

### Concentration of Co

Co, another essential element, is a constituent of vitamin B<sub>12</sub>, which is vital in creating red blood cells (RBCs). It also participates in the appropriate functions of the nervous system, the self-defense of the body, activation of specific enzymes, and prevention of anemia (Gogoasa et al. 2013; Voica et al. 2016). However, a high Co intake could produce toxic effects on human and animal health, especially in the heart and liver (Nędzarek et al. 2013; Zoroddu et al. 2019).

The rank order of countries based on the concentration of Co in the coffee products was Ethiopia (2686.563 µg/kg) > France (865.526 µg/kg) > Austria (589.968 µg/kg) > Poland (428.061 µg/kg) > Italy (419.919 µg/kg) > Netherland (314.897 µg/kg) > Germany (308.238 µg/kg) > Hawaii (221.329 µg/kg) > Turkey (140.000 µg/kg) > Slovenia (124.982 µg/kg) > Greece (110.000 µg/kg) > Lebanon (66.000 µg/kg) > Bosnia (60.000 µg/kg) > Brazil (34.913 µg/kg) > Romania (0.102 µg/kg) (Table 5). There are no safety

**Table 4** Meta-analysis concentration of Cu in the coffee products (µg/kg)

Country	ES	Lower	Upper	Weight (%)	Heterogeneity statistic	Degrees of freedom	P value	I <sup>2</sup> (%)
Austria	20,000.000	6970.172	34,000.000	0.00	2702.34	1	<0.001	100.00
Brazil	2804.322	2712.544	2896.099	25.44	33,702.51	11	<0.001	100.00
Colombia	13,000.000	10,000.000	15,000.000	0.00	345.48	6	<0.001	98.30
Costa Rica	13,000.000	11,000.000	15,000.000	0.00	0.00	0		
Egypt	2200.000	2164.216	2235.784	0.14	0.00	0		
Ethiopia	9097.608	9004.660	9190.556	34.79	350,000.00	18	<0.001	100.00
France	3538.513	2483.860	4593.166	13.73	17,261.65	3	<0.001	100.00
Germany	38,000.000	27,000.000	48,000.000	0.35	66,115.44	5	<0.001	100.00
Greece	6520.000	6420.113	6619.887	0.02	0.00	0		
Guatemala	13,000.000	9995.342	16,000.000	0.00	5.05	1	0.025	80.20
Hawaii	146.762	53.569	239.955	0.02	0.05	4	1	0.00
Honduras	17,000.000	16,000.000	17,000.000	0.00	0.08	1	0.783	0.00
India	14,000.000	11,000.000	16,000.000	0.00	184.80	4	<0.001	97.80
Indonesia	90.000	76.704	103.296	0.93	0.00	0		
Italy	6598.028	6425.142	6770.915	23.38	64,400.57	6	<0.001	100.00
Kenya	19,000.000	19,000.000	20,000.000	0.00	0.14	1	0.704	0.00
Netherlands	19,000.000	16,000.000	21,000.000	0.00	115.12	1	<0.001	99.10
Nicaragua	12,000.000	10,000.000	13,000.000	0.00	1.39	1	0.238	28.20
Nigeria	3970.000	3958.240	3981.760	1.17	0.00	0		
Peru	12,000.000	11,000.000	14,000.000	0.00	0.00	0		
Poland	13,000.000	12,000.000	14,000.000	0.00	0.00	0		
Overall	136.171	134.840	137.503	100.00	1,300,000.00	83	<0.001	100.00

ES effect size (mean of concentration)

**Table 5** Meta-analysis concentration of Co in the coffee products ( $\mu\text{g}/\text{kg}$ )

Country	ES	Lower	Upper	Weight (%)	Heterogeneity statistic	Degrees of freedom	<i>P</i> value	<i>I</i> <sup>2</sup> (%)
Austria	589.968	2.354	1217.157	4.34	8914.82	1	<0.001	100.00
Bosnia	60.000	53.556	66.444	2.23	0.00	0		
Brazil	34.913	2.478	103.251	5.80	256.51	1	<0.001	99.60
Ethiopia	2686.563	2464.899	2908.227	20.99	4,300,000.00	10	<0.001	100.00
France	865.526	738.133	992.919	1.20	75.70	1	<0.001	98.70
Germany	308.238	257.686	358.790	17.02	5516.66	5	<0.001	99.90
Greece	110.000	106.778	113.222	3.41	0.00	0		
Hawaii	221.329	140.513	302.144	0.48	56.44	4	<0.001	92.90
Italy	419.919	291.202	548.637	13.43	16,640.50	4	<0.001	100.00
Lebanon	66.000	63.256	68.744	3.59	0.00	0		
Netherlands	314.897	89.501	540.293	4.24	1003.74	1	<0.001	99.90
Poland	428.061	1.654	863.858	2.41	333.91	2	<0.001	99.40
Romania	0.102	0.047	0.156	12.72	815.60	5	<0.001	99.40
Slovenia	124.982	56.383	193.581	7.50	1693.64	1	<0.001	99.90
Turkey	140.000	123.831	156.169	0.64	0.00	0		
Overall	447.106	445.695	448.518	100.00	4,800,000.00	49	<0.001	100.00

ES effect size (mean of concentration)

limits established for Co concentration in coffee products. However, there is an MPL of 10  $\mu\text{g}/\text{kg}$  for this element in tea (Yaqub et al. 2018). Our results revealed that Co concentration in coffee samples of all countries is above this limit (except Romania), representing possible health risks to the population through coffee consumption.

### Concentration of Cd

Cd is the most toxic PTE (Gogoasa et al. 2013; Voica et al. 2016). Research has not revealed any positive impacts of Cd on human and animal health (Alkheraz et al. 2019), while they could be exposed to Cd through food and water consumption or work conditions. This element could accumulate a high amount in the human organs during lifespan. It is detrimental to kidney function and could raise bone demineralization, even at low concentrations. Cd also could cause lung cancer with long-term exposure (Alkheraz et al. 2019). Given its water solubility trait, Cd could be released and spread quickly to the environment through natural rock weathering, agricultural practices, and industrial activities, including batteries, refining ores, metal alloys, and plastic, dye, ceramic, and glass production. This element is also present in phosphate manures and pesticides at small levels and subsequently could be taken up by plants from treated soils with such compounds (Alkheraz et al. 2019).

The rank order of countries based on the concentration of Cd in the coffee products was Indonesia (300.000  $\mu\text{g}/\text{kg}$ ) > Brazil (61.541  $\mu\text{g}/\text{kg}$ ) > Honduras (50.000  $\mu\text{g}/\text{kg}$ ) > India (40.000  $\mu\text{g}/\text{kg}$ ) > Colombia (25.010  $\mu\text{g}/\text{kg}$ )

> Poland (13.000  $\mu\text{g}/\text{kg}$ ) > Kenya (10.000  $\mu\text{g}/\text{kg}$ ) > Netherlands (5.041  $\mu\text{g}/\text{kg}$ ) > Italy (1.160  $\mu\text{g}/\text{kg}$ ) > Germany (0.676  $\mu\text{g}/\text{kg}$ ) > Ethiopia (0.183  $\mu\text{g}/\text{kg}$ ) > Austria (0.100  $\mu\text{g}/\text{kg}$ ) ~ France (0.100  $\mu\text{g}/\text{kg}$ ) ~ Greece (0.100  $\mu\text{g}/\text{kg}$ ) ~ Jamaica (0.100  $\mu\text{g}/\text{kg}$ ) ~ Lebanon (0.100  $\mu\text{g}/\text{kg}$ ) ~ Malawi (0.100  $\mu\text{g}/\text{kg}$ ) ~ Pakistan (0.100  $\mu\text{g}/\text{kg}$ ) (Table 6).

The MPL for Cd set by the European Union (EU) is 100  $\mu\text{g}/\text{kg}$  in coffee products (da Silva et al. 2017). The results showed that the pooled concentration of Cd in coffee products of all studied countries meets the regulation limit except Indonesia.

To this end, geographical location in waste zones and industrial units that release Cd to the environment are critical in Cd concentration in food products. In this regard, synthetic fertilizers in farming sections, contaminated water, and agricultural fields with Cd could increase human exposure to this toxic element through food consumption (Fu and Xi 2020; Hocaoglu-Özyiğit and Genç 2020).

### Concentration of Pb

Pb, another highly toxic element, could accumulate in the body (Gogoasa et al. 2013). It is a non-essential metal for living things (Wadhwa et al. 2014). Pb is present everywhere in the environment (Lee et al. 2020). Plants like coffee beans could easily absorb Pb from the soil, water, and air (Lee et al. 2020). The significant harmful effects of these PTEs on human and animal health are carcinogenicity, neurotoxicity, hepatotoxicity, endocrine toxicity, reproductive and developmental toxicity, cardiovascular diseases, and gastrointestinal

**Table 6** Meta-analysis concentration of Cd in the coffee products ( $\mu\text{g}/\text{kg}$ )

Country	ES	Lower	Upper	Weight (%)	Heterogeneity statistic	Degrees of freedom	<i>P</i> value	$I^2$ (%)
Austria	0.100	0.098	0.102	7.28	0.00	1	1	0.00
Brazil	61.541	45.578	77.503	0.26	3208.74	7	< 0.001	99.80
Colombia	25.010	3.658	73.714	3.64	622.50	1	< 0.001	99.80
Ethiopia	0.183	0.158	0.207	42.52	2660.06	13	< 0.001	99.50
France	0.100	0.098	0.102	7.28	0.00	1	1	0.00
Germany	0.676	0.593	0.759	9.01	1743.32	5	< 0.001	99.70
Greece	0.100	0.097	0.103	3.64	0.00	0		
Honduras	50.000	48.544	51.456	0.03	0.00	1	1	0.00
India	40.000	36.494	43.506	0.00	0.00	1	1	0.00
Indonesia	300.000	255.679	344.321	0.00	0.00	0		
Italy	1.160	0.982	1.339	7.61	7633.71	4	< 0.001	99.90
Jamaica	0.100	0.096	0.104	3.64	0.00	0		
Kenya	10.000	9.649	10.351	0.44	0.00	1	1	0.00
Lebanon	0.100	0.094	0.106	3.64	0.00	0		
Malawi	0.100	0.096	0.104	3.64	0.00	0		
Netherlands	5.041	2.647	14.739	3.73	580.20	1	< 0.001	99.80
Pakistan	0.100	0.091	0.109	3.63	0.00	0		
Poland	13.000	10.581	15.419	0.01	0.00	0		
Overall	0.308	0.284	0.332	100.00	25,346.57	53	< 0.001	99.80

ES effect size (mean of concentration)

and hepatic disorders (da Silva et al. 2017; Pigozzi et al. 2018).

The rank order of countries based on the concentration of Pd in the coffee products was Indonesia (3900.000  $\mu\text{g}/\text{kg}$ ) > Syria (2492.013  $\mu\text{g}/\text{kg}$ ) > Egypt (1900.000  $\mu\text{g}/\text{kg}$ ) > Bosnia (1050.000  $\mu\text{g}/\text{kg}$ ) > Lebanon (693.000  $\mu\text{g}/\text{kg}$ ) > Austria (350.000  $\mu\text{g}/\text{kg}$ ) > Netherland (274.986  $\mu\text{g}/\text{kg}$ ) > Italy (221.898  $\mu\text{g}/\text{kg}$ ) > Brazil (203.618  $\mu\text{g}/\text{kg}$ ) > Greece (200.000  $\mu\text{g}/\text{kg}$ ) > Germany (183.279  $\mu\text{g}/\text{kg}$ ) > Turkey (120.000  $\mu\text{g}/\text{kg}$ ) > India (92.630  $\mu\text{g}/\text{kg}$ ) > Slovenia (70.000  $\mu\text{g}/\text{kg}$ ) > Poland (60.934  $\mu\text{g}/\text{kg}$ ) > France (23.829  $\mu\text{g}/\text{kg}$ ) > Pakistan (20.000  $\mu\text{g}/\text{kg}$ ) > Honduras (15.039  $\mu\text{g}/\text{kg}$ ) > Colombia (13.48  $\mu\text{g}/\text{kg}$ ) > Kenya (9.949  $\mu\text{g}/\text{kg}$ ) > Hawaii (8.895  $\mu\text{g}/\text{kg}$ ) > Romania (0.459  $\mu\text{g}/\text{kg}$ ) > South Korea (0.240  $\mu\text{g}/\text{kg}$ ) > Western Balkan (0.137  $\mu\text{g}/\text{kg}$ ) > Ethiopia (0.104  $\mu\text{g}/\text{kg}$ ) (Table 7).

Polish food legislation, Brazilian legislation, and the EU have set the MPLs for Pb in coffee products as 1000  $\mu\text{g}/\text{kg}$ , 500  $\mu\text{g}/\text{kg}$ , and 200  $\mu\text{g}/\text{kg}$ , respectively (Albals et al. 2021; Azam et al. 2021; Jeszka-Skowron et al. 2016). It could be concluded that the investigated coffees of all countries are not dangerous to human health except for Indonesia, Syria, Egypt, and Bosnia. However, using Brazil and EU safe limits, one country (Lebanon) and 5 other countries (Lebanon, Austria, Netherland, Italy, Brazil, and Greece) do not meet these regulation limits, respectively.

As observed previously, there are high differences between pooled concentrations of all PTEs in coffee products of different countries. Different factors are influential in these differences, which are discussed in the following.

First of all, coffee characteristics include variety, species (Arabica or Robusta), ripeness, type (brand/raw or roasted/ground or instant/infusion or powder), size, and processing conditions such as grinding, drying, roasting, mixing, and brewing methods which are effective in the observed difference in the content of PTEs in coffee products in different countries (Albals et al. 2021; Gure et al. 2017). Furthermore, the origin of the coffee regarding environmental conditions, including soil properties (type of soil, organic matter concentration, pH, drainage grade), geographical location (climate condition), and agricultural practices (using fertilizers and pesticides, organic or conventional farming), has a crucial impact on the content of PTEs in coffee beans (da Silva et al. 2017; Habte et al. 2016).

It is worth noting that both PTEs' gains and losses may occur during coffee processing which can be another source of observed differences in their concentration in different countries. The type of water used for infusion is also effective (Ashu and Chandravanshi 2011). In the case of the type of coffee products, it is reported that mixtures of coffee substitutes or surrogates (cereal grains) with coffee (20–66%) intermediate exhibit contents. At the same time,

**Table 7** Meta-analysis concentration of Pb in the coffee products ( $\mu\text{g}/\text{kg}$ )

Country	ES	Lower	Upper	Weight (%)	Heterogeneity statistic	Degrees of freedom	P value	$I^2$ (%)
Austria	350.000	310.801	389.199	0.19	74.00	1	<0.001	98.60
Bosnia	1050.000	919.991	1180.009	0.00	0.00	0		
Brazil	203.618	199.095	208.142	4.41	2,700,000.00	11	<0.001	100.00
Colombia	13.488	2.091	24.886	3.46	853.99	5	<0.001	99.40
Egypt	1900.000	1896.422	1903.578	0.28	0.00	0		
Ethiopia	0.104	0.084	0.123	23.89	2152.79	12	<0.001	99.40
France	23.829	22.977	24.681	4.94	23,091.15	3	<0.001	100.00
Germany	183.279	117.326	249.233	3.94	41,156.25	5	<0.001	100.00
Greece	200.000	196.778	203.222	0.33	0.00	0		
Hawaii	8.895	3.437	14.352	0.46	7.36	4	0.118	45.70
Honduras	15.039	3.457	44.322	2.85	1396.88	1	<0.001	99.90
India	92.630	46.659	138.602	2.19	575.47	3	<0.001	99.50
Indonesia	3900.000	3323.822	4476.178	0.00	0.00	0		
Italy	221.898	125.044	318.752	2.39	24,069.10	4	<0.001	100.00
Kenya	9.949	3.546	29.371	2.19	99.00	1	<0.001	99.00
Lebanon	693.000	681.772	704.228	0.03	0.00	0		
Netherlands	274.986	4.560	617.980	0.80	10,395.51	1	<0.001	100.00
Pakistan	20.000	9.265	30.735	0.04	0.00	0		
Poland	60.934	38.272	83.597	3.04	1529.02	3	<0.001	99.80
Romania	0.459	0.338	0.579	10.41	11,222.07	6	<0.001	99.90
Slovenia	70.000	69.089	70.911	2.19	0.00	1	1	0.00
South Korea	0.240	0.198	0.282	1.95	0.00	0		
Syria	2492.013	2231.473	2752.552	0.00	0.56	1	0.453	0.00
Turkey	120.000	103.026	136.974	0.01	0.00	0		
Western Balkan	0.137	0.116	0.158	30.01	4431.90	18	<0.001	99.60
Overall	21.027	20.824	21.231	100.00	4,100,000.00	104	<0.001	100.00

ES effect size (mean of concentration)

lower amounts are determined in coffee surrogates without coffee addition (Oliveira et al. 2012).

In the case of the brewing method, it is reported that the extraction levels of most PTEs are highest in the Turkish coffee method, in which coffee is boiled together with water (Nędzarek et al. 2013; Şemen et al. 2017). Regarding the type of coffee in terms of raw or roasted, it is noteworthy that the concentrations of PTEs in green coffee are typically lower than those determined in roasted coffees, probably due to removing water and volatile constituents (Amorim Filho et al. 2007; da Silva et al. 2017). Moreover, Şemen et al. (2017) concluded that studied PTEs are leached to the infusion more difficult from roasted coffee than from green coffee. Ashu and Chandravanshi (2011) also observed that the infusions contain lower PTEs than the roasted coffee. Moreover, Welna et al. (2013) noted that the transfer degree of PTEs into brew might vary probably related to coffee type (instant and ground coffees). Amorim Filho et al. (2007) also observed that the concentration of Fe and Cr is highest in steel grinders because of releasing these metals into

the coffee samples via the mill overheating. Additionally, Nędzarek et al. (2013) reported that potential contamination with PTEs may originate from packaging materials containing metals during coffee storage. There is also a negative relationship between the size of the coffee powder and the extraction rate of PTEs into the infusion (Pigozzi et al. 2018).

Moreover, Ashu and Chandravanshi (2011) reported that the concentration of PTEs in coffee varies among coffee varieties cultivated in diverse parts of the world with different environments, soil, and fertilizers.

Further, differences in sampling, sample preparation, and analytical techniques used in different studies significantly impact the concentration of PTEs (Ashu and Chandravanshi 2011). The contents of PTEs in the ground coffee are different from those in instant (soluble) coffee, probably due to processing differences (Voica et al. 2016). There are different kinds of methods used for the determination of PTEs in coffee products, including atomic absorption spectrometry (AAS), inductively coupled plasma-optical emission

spectrometry (ICP-OES), inductively coupled plasma-mass spectrometry (ICP-MS), atomic fluorescence spectrometry (AFS), and electrodes. Among these methods, ICP-OES and ICP-MS are the most remarkably applied because of their high sensitivity, specificity, selectivity, low detection limits (LODs), ease of setup, less time, and low cost (Albals et al. 2021; Alkherraz et al. 2019). However, to overcome the disadvantages of AFS methods, some pre-concentration methods could be utilized during sample preparation, including solid-phase extraction (SPE), liquid–liquid extraction (LLE), as well as microextraction procedures like dispersive liquid–liquid microextraction (DLLME) (Şaylan et al. 2020).

## Conclusion

This study meta-analyzed the prevalence of potentially toxic elements in coffee products in different countries. The pooled mean concentration of essential elements in coffee products is much higher than that of toxic elements (Co > Ni > Cu > Cr > Pb > As > Cd). The result showed that the pooled mean concentration of essential elements in coffee products is much higher than that of toxic elements. Also, high differences between pooled concentrations of all PTEs were reported in coffee products of different countries. Various concentrations of potentially toxic elements in different studies may be due to several factors, including environmental growth conditions (climate, temperature, and agricultural practices), type of processing, coffee variety, preparation method, agricultural practices (using fertilizers and pesticides, organic or conventional farming), and analytical techniques. Therefore, more studies are needed to investigate the effect of mentioned factors on the prevalence of PTEs in coffee products in the future.

**Supplementary Information** The online version contains supplementary material available at <https://doi.org/10.1007/s11356-022-23110-9>.

**Author contribution** Trias Mahmudiono, Fardin Javanmardi, and Behrouz Tajdar-oranj: investigation, data curation, resources, conceptualization, methodology, and writing of original draft. Amene Nematollahi and Mohadeseh Pirhadi: literature searching, writing, and editing. Amin Mousavi Khaneghah and Yadolah Fakhri: supervision, conceptualization, data analysis, and review and editing.

**Data availability** Not applicable.

## Declarations

**Ethics approval and consent to participate** Ethics approval is not applicable. The authors declare their consent to participate in this article.

**Consent for publication** The authors declare their consent for publication of this article.

**Competing interests** The authors declare no competing interests.

## References

- Achmad RT, Auerkari EI (2017) Effects of chromium on human body. *Ann Res Rev Biol* 13:1–8
- Albals D, Al-Momani IF, Issa R, Yehya A (2021) Multi-element determination of essential and toxic metals in green and roasted coffee beans: a comparative study among different origins using ICP-MS. *Sci Prog* 104:00368504211026162
- Alkherraz A, Hashad O, Elsherif K (2019) Heavy metals contents in some commercially available coffee, tea, and cocoa samples in Misurata City -Libya. *Prog Chem Biochem Res* 2:99–107
- Ameri Z, Hoodaji M, Rajaie M, Ataabadi M (2021) Optimizing modified rice bran for treating aqueous solutions polluted by Cr (VI) ions: isotherm and kinetics analyses. *Qual Assur Saf Crops Foods* 13:1–11
- Amorim Filho VR, Polito WL, Gomes Neto JA (2007) Comparative studies of the sample decomposition of green and roasted coffee for determination of nutrients and data exploratory analysis. *J Braz Chem Soc* 18:47–53
- Anosike S, Oranusi S (2018) Assessment of microbiological and chemical qualities of selected cocoa, tea and coffee brands in Nigerian markets. *Afr J Clin Exp Microbiol* 19:186–194
- Ashu R, Chandravanshi BS (2011) Concentration levels of metals in commercially available Ethiopian roasted coffee powders and their infusions. *Bull Chem Soc Ethiopia*. <https://doi.org/10.4314/bcse.v25i1.63356>
- Atamaleki A, Sadani M, Raoofi A, Miri A, Bajestani SG, Fakhri Y, Heidarinejad Z, Khaneghah AM (2020) The concentration of potentially toxic elements (PTEs) in eggs: a global systematic review, meta-analysis and probabilistic health risk assessment. *Trends Food Sci Technol* 95:1–9
- Azam K, Akhtar S, Gong YY, Routledge MN, Ismail A, Oliveira CA, Iqbal SZ, Ali H (2021) Evaluation of the impact of activated carbon-based filtration system on the concentration of aflatoxins and selected heavy metals in roasted coffee. *Food Control* 121:107583
- Barbosa RM, Batista BL, Varriquer RM, Coelho VA, Campiglia AD, Barbosa F Jr (2014) The use of advanced chemometric techniques and trace element levels for controlling the authenticity of organic coffee. *Food Res Int* 61:246–251
- Barrea L, Pugliese G, Frias-Toral E, El Ghoch M, Castellucci B, Chapela SP, Carignano MdA, Laudisio D, Savastano S, Colao A (2021) Coffee consumption, health benefits and side effects: a narrative review and update for dietitians and nutritionists. *Crit Rev Food Sci Nutr* 1–24
- Batrinou A, Houhoula D, Papageorgiou E (2020) Rapid detection of mycotoxins on foods and beverages with enzyme-linked immunosorbent assay. *Qual Assur Saf Crops Foods* 12:40–49
- Berman N, Honig S, Cronstein B, Pillinger M (2022) The effects of caffeine on bone mineral density and fracture risk. *Osteoporos Int* 33:1235–1241
- Bost M, Houdart S, Oberli M, Kalonji E, Huneau J-F, Margaritis I (2016) Dietary copper and human health: current evidence and unresolved issues. *J Trace Elem Med Biol* 35:107–115
- Boumar A, Boukaka K, Leghouchi E (2020) Determination of heavy metals in tomatoes cultivated under green houses and human health risk assessment. *Qual Assur Saf Crops Foods* 12:76–86
- da Silva SA, Mendes F, Queiroz I, Reis MR, Passos FR, de Carvalho AMX, de Oliveira Rocha KR, Pinto FG (2017) Determination of heavy metals in the roasted and ground coffee beans and brew. *Afr J Agric Res* 12:221–228
- de Mejia EG, Ramirez-Mares MV (2014) Impact of caffeine and coffee on our health. *Trends Endocrinol Metab* 25:489–492



- Delgado Quezada V, Altamirano Espinoza M, Bundschuh J (2020) Arsenic in geoenvironments of Nicaragua: exposure, health effects, mitigation and future needs. *Sci Total Environ* 716:136527
- Dieng H, Ellias SB, Satho T, Ahmad AH, Abang F, Ghani IA, Noor S, Ahmad H, Zuharah WF, Morales Vargas RE (2017) Coffee, its roasted form, and their residues cause birth failure and shorten lifespan in dengue vectors. *Environ Sci Pollut Res* 24:14782–14794
- Escott-Stump S (2008) Nutrition and diagnosis-related care. *Lippincott Williams & Wilkins*
- Fu Z, Xi S (2020) The effects of heavy metals on human metabolism. *Toxicol Mech Methods* 30:167–176
- Gao L, Huang X, Wang P, Chen Z, Hao Q, Bai S, Tang S, Li C, Qin D (2022) Concentrations and health risk assessment of 24 residual heavy metals in Chinese mitten crab (*Eriocheir sinensis*). *Qual Assur Saf Crops Foods* 14:82–91
- Genchi G, Carocci A, Lauria G, Sinicropi MS, Catalano A (2020) Nickel: human health and environmental toxicology. *Int J Environ Res Public Health* 17:679
- Godos J, Pluchinotta FR, Marventano S, Buscemi S, Li Volti G, Galvano F, Grosso G (2014) Coffee components and cardiovascular risk: beneficial and detrimental effects. *Int J Food Sci Nutr* 65:925–936
- Gogoasa I, Pirvu A, Alda L, Velciov A, Rada M, Bordean D, Moigradean D, Simion A, Gergen I (2013) The mineral content of different coffee brands. *J Hortic For Biotechnol* 17:68–71
- Gogoasa I, Adina B, Adina N, Maria R, Antoanela C, Liana A (2017) Determination of trace elements in commercially available instant coffees. *J Hortic For Biotechnol* 21:1–5
- Guan Q, Zeng G, Song J, Liu C, Wang Z, Wu S (2021) Ultrasonic power combined with seed materials for recovery of phosphorus from swine wastewater via struvite crystallization process. *J Environ Manage* 293:112961
- Gure A, Chandravanshi BS, Godeto TW (2017) Metals in green coffee beans from major coffee-growing regions of Ethiopia. *Chem Int* 3:359–369
- Habte G, Hwang IM, Kim JS, Hong JH, Hong YS, Choi JY, Nho EY, Jamila N, Khan N, Kim KS (2016) Elemental profiling and geographical differentiation of Ethiopian coffee samples through inductively coupled plasma-optical emission spectroscopy (ICP-OES), ICP-mass spectrometry (ICP-MS) and direct mercury analyzer (DMA). *Food Chem* 212:512–520
- Heshmati A, Mehri F, Karami-Momtam J, Khaneghah AM (2020) The concentration and health risk of potentially toxic elements in black and green tea—both bagged and loose-leaf. *Qual Assur Saf Crops Foods* 12:140–150
- Higgins JPT, Thompson SG (2002) Quantifying heterogeneity in a meta-analysis. *Stat Med* 21:1539–1558
- Higgins J, White IR, Anzures-Cabrera J (2008) Meta-analysis of skewed data: combining results reported on log-transformed or raw scales. *Stat Med* 27:6072–6092
- Hocaoğlu-Özyiğit A, Genç BN (2020) Cadmium in plants, humans and the environment. *Front Life Sci Relat Technol* 1:12-21
- Hu L, Wang X, Zou Y, Wu D, Gao G, Zhong Z, Liu Y, Hu S, Fan H, Zhang B (2022) Effects of inorganic and organic selenium intervention on resistance of radish to arsenic stress. *Ital J Food Sci* 34:44–58
- Ibrahim S, Hassan M, Ibraheem Q, Arif K (2020) Genotoxic effect of lead and cadmium on workers at wastewater plant in Iraq. *J Environ Public Health* 20:1–9
- Jeszka-Skowron M, Staniszc E, De Peña MP (2016) Relationship between antioxidant capacity, chlorogenic acids and elemental composition of green coffee. *LWT* 73:243–250
- Kalita J, Kumar V, Misra UK, Bora HK (2019) Movement disorder in copper toxicity rat model: role of inflammation and apoptosis in the corpus striatum. *Neurotox Res* 37:904–912
- Khan KM, Chakraborty R, Bundschuh J, Bhattacharya P, Parvez F (2020) Health effects of arsenic exposure in Latin America: an overview of the past eight years of research. *Sci Total Environ* 710:136071
- Khaneghah AM, Fakhri Y, Abdi L, Coppa CFSC, Franco LT, de Oliveira CAF (2019) The concentration and prevalence of ochratoxin A in coffee and coffee-based products: a global systematic review, meta-analysis and meta-regression. *Fungal Biol* 123:611–617
- Khaneghah AM, Fakhri Y, Nematollahi A, Pirhadi M (2020) Potentially toxic elements (PTEs) in cereal-based foods: a systematic review and meta-analysis. *Trends Food Sci Technol* 96:30–44
- Lee J-G, Hwang J-Y, Lee H-E, Choi J-D, Kang G-J (2020) Comparative analysis of lead content during food processing. *Food Sci Biotechnol* 29:1063–1069
- Li W, Shi Y, Zhu D, Wang W, Liu H, Li J, Shi N, Ma L, Fu S (2021) Fine root biomass and morphology in a temperate forest are influenced more by the nitrogen treatment approach than the rate. *Ecol Ind* 130:108031
- Lin X, Lu K, Hardison AK, Liu Z, Xu X, Gao D, Gong J, Gardner WS (2021) Membrane inlet mass spectrometry method (REOX/MIMS) to measure  $^{15}\text{N}$ -nitrate in isotope-enrichment experiments. *Ecol Ind* 126:107639
- Liu W, Huang F, Wang Y, Zou T, Zheng J, Lin Z (2011) Recycling  $\text{Mg}(\text{OH})_2$  nano-adsorbent during treating the low concentration of  $\text{Cr}^{\text{VI}}$ . *Environ Sci Technol* 45:1955–1961
- Liu W, Li J, Zheng J, Song Y, Shi Z, Lin Z, Chai L (2020) Different pathways for Cr(III) oxidation: implications for Cr(VI) reoccurrence in reduced chromite ore processing residue. *Environ Sci Technol* 54:11971–11979
- Liu G, Nie R, Liu Y, Mehmood A (2022) Combined antimicrobial effect of bacteriocins with other hurdles of physicochemical and microbiome to prolong shelf life of food: a review. *Sci Total Environ* 825:154058
- Martinelli L, da Silva Berilli S, Amaro de Sales R, da Cunha M, Monaco PAVL, de Jesus Freitas S, Martinelli M, Gabriel Berilli APC, Pireda S, da Silva Oliveira D (2022) Influence of chromium and sodium on development, physiology, and anatomy of Conilon coffee seedlings. *Environ Sci Pollut Res* 29:44986–44997
- Mirmahdi RS, Zoghi A, Mohammadi F, Khosravi-Darani K, Jazaieri S, Mohammadi R, Rehman Y (2021) Biodecontamination of milk and dairy products by probiotics: boon for bane. *Ital J Food Sci* 33:78–91
- Mirmahdi RS, Mofid V, Zoghi A, Khosravi\_Darani K, Mortazavian AM (2022) Risk of low stability *Saccharomyces cerevisiae* ATCC 9763-heavy metals complex in gastrointestinal simulated conditions. *Heliyon* 8:e09452
- Moher D, Liberati A, Tetzlaff J, Altman DG, Group P (2010) Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Int J Surg* 8:336-341
- Nartea A, Lucci P, Loizzo MR, Tundis R, Leporini M, Gervasi L, Fanesi B, Núñez O, Frega NG, Fiorini D (2022) Is coffee powder extract a possible functional ingredient useful in food and nutraceutical industries? *Ital J Food Sci* 34:140–148
- Nędzarek A, Tórz A, Karakiewicz B, Clark JS, Laszczyńska M, Kaleta A, Adler G (2013) Concentrations of heavy metals (Mn, Co, Ni, Cr, Ag, Pb) in coffee. *Acta Biochim Pol* 60:623–627
- Nourbakhsh F, Tajbakhsh E (2021) Neurotoxicity mechanism of ochratoxin A. *Qual Assur Saf Crops Foods* 13:34–45
- Oliveira M, Casal S, Morais S, Alves C, Dias F, Ramos S, Mendes E, Delerue-Matos C, Oliveira MBP (2012) Intra- and interspecific mineral composition variability of commercial instant coffees and coffee substitutes: contribution to mineral intake. *Food Chem* 130:702–709

- Pavithra V (2021) Review article on caffeine activity. *J Nanotechnol Res* 3:1–5
- Petrović SM, Savić SR, Zvezdanović JB, Mladenović-Ranisavljević I, Cvetković DJ, Cvetanović AD (2020) Benefits and risks of commercially available coffee beverages from Western Balkan. *Chem Pap* 74:847–857
- Pigozzi MT, Passos FR, Mendes FQ (2018) Quality of commercial coffees: heavy metal and ash contents. *J Food Qual* 18:1–8
- Pohl P, Stelmach E, Szymczycha-Madeja A (2014) Simplified sample treatment for the determination of total concentrations and chemical fractionation forms of Ca, Fe, Mg and Mn in soluble coffees. *Food Chem* 163:31–36
- Pullella K, Kotsopoulos J (2020) Arsenic exposure and breast cancer risk: a re-evaluation of the literature. *Nutrients* 12:3305
- Quan H, Zhang J (2003) Estimate of standard deviation for a log-transformed variable using arithmetic means and standard deviations. *Stat Med* 22:2723–2736
- Quan Q, Gao S, Shang Y, Wang B (2021) Assessment of the sustainability of *Gymnocypis eckloni* habitat under river damming in the source region of the Yellow River. *Sci Total Environ* 778:146312
- Rattan S, Parande AK, Nagaraju VD, Ghiwari GK (2015) A comprehensive review on utilization of wastewater from coffee processing. *Environ Sci Pollut Res* 22:6461–6472
- Samet M, Charfeddine M, Kamoun L, Nouri-Ellouze O, Gargouri-Bouziid R (2018) Effect of compost tea containing phosphogypsum on potato plant growth and protection against *Fusarium solani* infection. *Environ Sci Pollut Res* 25:18921–18937
- Şaylan M, Zaman BT, Bakırdere EG, Bakırdere S (2020) Determination of trace nickel in chamomile tea and coffee samples by slotted quartz tube-flame atomic absorption spectrometry after preconcentration with dispersive liquid-liquid microextraction method using a Schiff base ligand. *J Food Compos Anal* 88:103454
- Şemen S, Mercan S, Yayla M, Açıkkol M (2017) Elemental composition of green coffee and its contribution to dietary intake. *Food Chem* 215:92–100
- Senkondo YH, Tack F, Semu E (2014) Copper accumulations in soils, coffee, banana, and bean plants following copper-based fungicides in coffee farms in Arusha and Kilimanjaro regions, Tanzania. *Commun Soil Sci Plant Anal* 45:2032–2045
- Shi C, Wu Z, Yang F, Tang Y (2021) Janus particles with pH switchable properties for high-efficiency adsorption of PPCPs in water. *Solid State Sci* 119:106702
- Sun Y, Li J, Zhu L, Jiang L (2022) Cooperation and competition between CRISPR-and omics-based technologies in foodborne pathogens detection: a state of the art review. *Curr Opin Food Sci* 44:100813
- Teklay A (2016) Physiological effect of chromium exposure: a review. *Int J Food Sci Nutr Diet S* 7:1–11
- Tiep TD (2014) Effect of roasting conditions on concentration in elements of Vietnam robusta coffee. *Acta Univ Cibinensis E Food Technol* 18:19–34
- van Dam RM, Hu FB, Willett WC (2020) Coffee, caffeine, and health. *N Engl J Med* 383:369–378
- Voica C, Feher I, Iordache A, Cristea G, Dehelean A, Magdas D, Mirel V (2016) Multielemental analysis of coffee by inductively coupled plasma-mass spectrometry. *Anal Lett* 49:2627–2643
- Wadhwa SK, Tuzen M, Kazi TG, Soylak M, Hazer B (2014) Polyhydroxybutyrate-b-polyethyleneglycol block copolymer for the solid phase extraction of lead and copper in water, baby foods, tea and coffee samples. *Food Chem* 152:75–80
- Wang Y, Wu X, Liu J, Zhai Z, Yang Z, Xia J, Deng S, Qu X, Zhang H, Wu D (2022) Mo-modified band structure and enhanced photocatalytic properties of tin oxide quantum dots for visible-light driven degradation of antibiotic contaminants. *J Environ Chem Eng* 10:107091
- Welna M, Szymczycha-Madeja A, Zyrnicki W (2013) Applicability of ICP-OES, UV-VIS, and FT-IR methods for the analysis of coffee products. *Anal Lett* 46:2927–2940
- Winiarska-Mieczan A, Kwiatkowska K, Kwiecień M, Zaricka E (2021) Assessment of the risk of exposure to cadmium and lead as a result of the consumption of coffee infusions. *Biol Trace Elem Res* 199:2420–2428
- Wolde T (2014) Effects of caffeine on health and nutrition: a review. *Food Sci Qual Manag* 30:59–65
- World Health Organization (2000) Evaluation of certain food additives and contaminants: fifty-third report of the Joint FAO/WHO Expert Committee on Food Additives. World Health Organization, Geneva
- Yaqub G, Ilyas F, Idrees M, Mariyam V (2018) Monitoring and risk assessment due to presence of heavy metals and pesticides in tea samples. *Food Sci Technol* 38:625–628
- Yazdanpanah S, Jocar S, Zandilak T, Haghighi M (2022) Evaluation and comparison of ochratoxin a contamination and metallic elements lead and arsenic in commercial instant coffee powders. *Iran Food Sci Technol Res J* 18:81–95
- Zoroddu MA, Aaseth J, Crisponi G, Medici S, Peana M, Nurchi VM (2019) The essential metals for humans: a brief overview. *J Inorg Biochem* 195:120–129

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.