



# Indian Journal of Forensic Medicine & Toxicology

Website: [www.ijfmt.com](http://www.ijfmt.com)



Official Organ of Indian Association of Medico-Legal Experts (Regd.)

# Indian Journal of Forensic Medicine & Toxicology

## EDITOR in Chief

**Prof. R K Sharma**

Formerly at All India Institute of Medical Sciences, New Delhi, E-mail: editor.ijfmt@gmail.com

## EDITOR

**Prof. Dr. Adarsh Kumar**

Forensic Medicine & Toxicology, AIIMS, New Delhi

### INTERNATIONAL EDITORIAL ADVISORY BOARD

1. **Prof Mete Gulmen** Cukurova University, TURKEY
2. **Prof. Leandro Duarte De Carvalho**, Minas Gerais, Belo Horizonte, Brazil
3. **Prof. Donata Favretto** (Full Professor) Forensic Toxicology at University of Padova, Italy
4. **Prof. Babak Mostafazadeh** Department of Forensic Medicine & Toxicology, Shahid Beheshti University of Medical Sciences, Tehran-Iran
5. **Prof Halis Dokgoz**, Mersin University, TURKEY
6. **Prof Jozef Sidlo**, Comenius University, Bratislava, SLOVAKIA
7. **Dr. Rahul Pathak** (Lecturer) Forensic Science, Dept of Life Sciences Anglia Ruskin University, Cambridge, United Kingdom
8. **Dr. Hareesh** (Professor & Head) Forensic Medicine, Ayder Referral Hospital, College of Health Sciences, Mekelle University, Mekelle Ethiopia East Africa
9. **Dr. Mokhtar Ahmed Alhrani** (Specialist) Forensic Medicine & Clinical Toxicology, Director of Forensic Medicine Unit, Attorney General's Office, Sana'a, Yemen
10. **Dr. Sarathchandra** Kodikara (Senior Lecturer) Forensic Medicine, Department of Forensic Medicine, Faculty of Medicine, University of Peradeniya, Sri Lanka
11. **Dr Noha A. Magdie El Rafie**, Forensic Toxicology, Ain Shams University, Cairo, EGYPT

### SCIENTIFIC COMMITTEE

1. **Prof Udai Pratap Singh**, Department of Anthropology Lucknow University Lucknow
2. **Dr Anil Rahule** (Associate Professor) Dept of Anatomy, Govt Medical College Nagpur
3. **Dr Shankar Bakkanwar** (Associate Professor) Forensic Medicine, Kasturba Medical College, Manipal, Karnatakad
4. **Dr K. Ravikumar** Raksha Shakti University, Ahmedabad, Gujrat.
5. **Dr. Pragmesh Parmar** (Associate Professor) Forensic Medicine, Valsad, Gujrat
6. **Dr Vandana Mudda** (Awati) (Associate Prof) Dept of FMT, M.R. Medical College, Gulbarga, Karnataka,
7. **Dr. Asha Srivastava** (Senior Scientific Officer) Forensic Psychology, Central Forensic Science Laboratory, CBI, Delhi
8. **Dr. Lav Kesharwani** (Asst.Prof.) School of Forensic Science, Sam Higginbottom Institute of Agriculture Technology & Sciences, Allahabad U.P,
9. **Dr. Anu Sharma** (Associate Prof) Dept of Anatomy, DMCH, Ludhiana (PB)
10. **Dr. Shalini Gupta** ( Prof ) Oral Pathology and Microbiology, Dental Sciences King George Medical University, Lucknow, UP
11. **Dr Rituja Sharma**, Associate Prof, Law Banasthali Vidyapeeth Jaipur

"Indian Journal of Forensic Medicine & Toxicology" is peer reviewed quarterly journal. It deals with Forensic Medicine, Forensic Science, Toxicology, DNA fingerprinting, sexual medicine and environment medicine. It has been assigned International standard serial No. p-0973-9122 and e- 0973-9130. The Journal has been assigned RNI No. DELENG/2008/21789. The journal is indexed with Index Copernicus (Poland) and is covered by EMBASE (Excerpta Medica Database). The journal is also abstracted in Chemical Abstracts (CAS) database (USA). The journal is also covered by EBSCO (USA) database. The Journal is now part of UGC, DST and CSIR Consortia. It is now official publication of Indian Association of Medico-Legal Experts (Regd.).

### NATIONAL EDITORIAL ADVISORY BOARD

#### Chairman

**Prof Sudhir K Gupta** - Head, Department of Forensic Medicine  
All India Institute of Medical Sciences, New Delhi

#### Members

1. **Prof. SK Dhatarwal**, Forensic Medicine, PGIMS, Rohtak, Haryana
2. **Prof. N K Aggrawal** Forensic Medicine, UCMS, Delhi
3. **Prof Ajay Ghangale** Forensic Medicine Dr DY Patil Medical College, Pune, Maharashtra
4. **Dr. Amar Jyoti** Patwory Professor, Forensic Medicine NEIGRIHMS, Shillong
5. **Dr S. Venkata Raghava** Professor, Forensic Medicine, Bangalore Medical College, Bengaluru
6. **Prof Praveen Arora**, Professor Department of Forensic Medicine & Toxicology, SAIMS, Indore
7. **Dr. Pankaj Datta** (Principal & Head) Department of Prosthodontics, Indraprastha Dental College & Hospital, Ghaziabad
8. **Dr. Mahindra Nagar** (Head) Department of Anatomy, UCMS & GTB Hospital, Delhi
9. **Dr. Virender Kumar Chhoker** Professor Forensic Medicine and Toxicology, Santosh Medical College, Ghaziabad, UP
10. **Dr. Dayanand G Gannur** (Professor) Department of Forensic Medicine & Toxicology, Shri BM Patil Medical College, Hospital & Research centre, Bijapur, Karnataka
11. **Dr. Alok Kumar** Professor Department of Forensic Medicine & Toxicology, UP Rural Institute of Medical Sciences and Research, Saifai, Etawah, U.P.

Print-ISSN:0973-9122 Electronic - ISSN: 0973-9130

Frequency: Quarterly, © All Rights reserved The views and opinions expressed are of the authors and not of the Indian Journal of Forensic Medicine & Toxicology. Indian Journal of Forensic Medicine & Toxicology does not guarantee directly or indirectly the quality or efficacy of any products or service featured in the advertisement in the journal, which are purely commercial.

Website: [www.ijfmt.com](http://www.ijfmt.com)

#### Editor

**Dr. R.K. Sharma**  
Institute of Medico-legal Publications  
Logix Office Tower, Unit No. 1704, Logix City Centre Mall, Sector- 32, Noida  
- 201 301 (Uttar Pradesh)

#### Printed, published and owned by

**Dr. R.K. Sharma**  
Institute of Medico-legal Publications  
Logix Office Tower, Unit No. 1704, Logix City Centre Mall, Sector- 32, Noida  
- 201 301 (Uttar Pradesh)

#### Published at

**Institute of Medico-legal Publications**  
Logix Office Tower, Unit No. 1704, Logix City Centre Mall, Sector- 32, Noida  
- 201 301 (Uttar Pradesh)

576. Identification of DNA Corpse in Ordo Diptera Familia Calliphoridae Larvae in CSF1PO, TH01, and TPOX Locus Using Polymerase Chain Reaction (PCR) Methods..... 3538  
*Tutik Purwanti, Ahmad Yudianto, Mochammad Soekry Erfan Kusuma*
577. Children Death with Child Abuse: A Case Report at Bhayangkara Hospital, Kediri Indonesia ..... 3544  
*Mochammad Soekry Erfan Kusuma, Tutik Purwanti*
578. Effectiveness of WhatsApp Group on Breastfeeding Practices ..... 3550  
*Wahidah Sukriani, Greiny Arisani*
579. The Effect of the Blood Cupping Therapy on High Density Lipoprotein (HDL) and Low Density Lipoprotein (LDL) in Hypercholesterol Patients ..... 3556  
*Wahyudi Widada, Teddy Ontoseno, Bambang Purwanto, Suhartati, Imam Subadi, Hari Basuki Notobroto, Ahsan*
580. Impact of High Fidelity Simulation Training on the Quality of Cardiopulmonary Resuscitation Performance among Emergency Medical Services (EMS) Professionals in Pune, India ..... 3563  
*Anand Hinduja, Parag Rishipathak, Shrimathy Vijayaraghavan*
581. Relationship of Smartphone Usage and Academic Performance among Undergraduate students – A Systematic Review ..... 3567  
*Ancy Albert; Nancy Netto; Neelkant Pawar; Fenilkumar Patel, Sharadha Ramesh; Manisha Mistry*
582. A Study to Assess the Effectiveness of Mindful Meditation on Health and Psychological Adjustment Faced by Nursing Students in Selected Colleges of Pune City ..... 3571  
*Anu Alexander, Ms. Feba Josemon, Shreeti Gahare, Neha Jadhav, Sonopant G Joshi, Ranjana Chavan*
583. Health and Lifestyle Practices of Under Graduate Students Studying in Selected Nursing College .. 3576  
*Anagha.M, Ritu Gaikwad, Ronald Masumba, Kraus, Asha Varghese, Deepak Sethi, Merlin Suji*
584. Osteoporosis Risk Analysis in Women Residing in Selected Urban Areas..... 3582  
*Milu Anna Laji , Phale Diksha Ramesh, Gumi Riram, Sahil Chavan, Akash Khade, Sharadha Ramesh, Merlin Suji*
585. A Study to Assess the use of Body Mechanics Practices among Nursing Students in Selected Colleges of Nursing in Pune City ..... 3586  
*Dipak Sethi, Jasneet Kaur, Jennifer William Anthony*
586. A Descriptive Study to Assess the Stress and Coping Mechanism Due to Lockdown among School Going Children..... 3590  
*Dipali Dumbre, Sharadha Ramesh, Ranjana Chavan, Mangesh Jabade*
587. Knowledge and Practices on Intra Venous Cannulation among Student Nurses..... 3598  
*Sharadha Ramesh, MerlinSuji*
588. Effect of lockdown on Students Studying in Various Colleges of Nursing..... 3603  
*Jennifer W. Anthony, Deepak Sethi, Mangesh Jabade, Poonam Yadav, Stephina Fernandes*
589. A Study to Assess the Effectiveness of Pranic Healing Therapy on Backache among Adults At Pranic Healing Centre in Pune City ..... 3609  
*Ms Kaguiphullu Sylvia Gangmei, Sheela Upendra*

# The Effect of the Blood Cupping Therapy on High Density Lipoprotein (HDL) and Low Density Lipoprotein (LDL) in Hypercholesterol Patients

Wahyudi Widada<sup>1</sup>, Teddy Ontoseno<sup>2</sup>, Bambang Purwanto<sup>3</sup>, Suhartati<sup>4</sup>, Imam Subadi<sup>5</sup>,  
Hari Basuki Notobroto<sup>6</sup>, Ahsan<sup>7</sup>

<sup>1</sup>Researcher, Universitas Muhammadiyah Jember / Doctoral Student, Universitas Airlangga, <sup>2</sup>Researcher, Universitas Airlangga, <sup>3</sup>Researcher, Universitas Airlangga, <sup>4</sup>Faculty of Medicine, Universitas Wijaya Kusuma, Surabaya, <sup>5</sup>Physician, Hospital of Universitas Airlangga, <sup>6</sup>Associate Professor, Faculty of Public Health, Universitas Airlangga, <sup>7</sup>Researcher, Department of Nursing, Universitas Brawijaya

## Abstract

**Introduction:** Hypercholesterol is a high cholesterol level in the blood. Blood cupping therapy is a technique of excreting metabolic waste in the blood through the skin. The study aimed to measure any effect of blood cupping therapy to HDL and LDL cholesterol. **Method:** It was a Quasi-Experimental study using humans as research subjects. Non random consecutive sampling. Patients with hypercholesterol after 12 hours of fasting were treated with blood cupping, 9 points. Research subjects were 51 men divided into three groups, the cupping group only consisted of 17 men, the drug group without cupping consisted of 17 men, and the cupping and drug consisted 17 men. Collecting data from third of the groups as pre and post data. The data were analyzed by Mann Whitney, Wilcoxon and Kruskal Wallis. The research took place at the Laboratory of Biochemistry, Universitas Jember. **Results:** Measurement of HDL cholesterol on the cupping group was pretest=33.03±3.73; posttest=37.58±6.54; p=0.000. The drug group was pretest=33.40±3.18; posttest=33.56±3.50; p=0.788. The cupping and drug group was pretest=34.32±3.38; posttest=37.61±2.01; p=0.002. Kruskal Wallis test on pretest group was 0.534 and on posttest group was 0.002. Measurement of LDL cholesterol on the cupping group was pretest=154.70±39.68; posttest=123.89±41.86; p=0.000. The drug group was pretest=151.24±44.17; posttest=151.24±44.17; p=0.019. The cupping and drug group was pretest=147.48±62.66; posttest=105.57±57.94; p=0.001. Kruskal Wallis test on pretest group was 0.439 and on posttest group was 0.082. This means that there was no difference in the average pretest and posttest LDL levels in the cupping, drug, cupping and drug groups. **Conclusion:** The intervention of blood cupping therapy can increase HDL cholesterol level, and reduce LDL cholesterol level. Further research needs to be done to measure the potential prevention of atherosclerosis.

**Keywords:** blood cupping; HDL; LDL; cholesterol

## Introduction

Blood cupping therapy or the wet cupping therapy has been used for medication since ancient times. Even Hipocrates used blood cupping in cases of internal drug.

<sup>(1)</sup> The long history of blood cupping therapy proves that the blood cupping therapy when performed correctly is a safe and effective method.

There is a misperception in interpreting blood cupping therapy. The depth of the needle into the skin is only 0.05mm. Injuries with such needles do not cause blood flow. Blood comes out after being pulled with a pump of negative strength -200 mmHg.<sup>(2,3)</sup> Actually, blood cupping therapy is not a treatment for removing blood circulation but for removing metabolic waste or

---

### Corresponding author:

Wahyudi Widada

E-mail: wahyudiwidada@unmuhjember.ac.id

Address: Campus "Universitas Airlangga" Indonesia

the so-called causative pathological substances.<sup>(4)</sup> In other words, blood cupping therapy does not reduce circulating blood volume. The bloodlike substance that comes out of the injury is cholesterol metabolism wastes, old erythrocytes, etc. The blood amount due to the conduct of a correctly performed blood cupping therapy does not reduce hemoglobin.<sup>(5)</sup>

We have understood that high levels of total blood cholesterol, especially in the form of LDL cholesterol, are a major risk factor for developing coronary heart diseases.<sup>(6)</sup> Some patients who receive LDL cholesterol reduction therapy still have complaints of coronary heart disease.<sup>(7)</sup> The aim of the study was to measure effect of blood cupping therapy to HDL and LDL cholesterol. The existence of HDL cholesterol in the reserve cholesterol transport is a mechanism to protect endothelium against the risk of atherosclerosis. HDL has anti-inflammatory, antioxidant, antithrombotic properties. It is also antiaterogenic.<sup>(8)</sup>

### Method

It was a Quasi-Experimental study using humans as

research subjects. Non random consecutive sampling. Patiens with hypercholesterol after 12 hours of fasting were treated with blood cupping, 9 points. Research subjects were 51 men divided into three groups, the cupping group only consisted of 17 men, the drug group without cupping consisted of 17 men, and the cupping and drug consisted 17 men. Collecting data from third of the groups as pretest and posttest. The research subjects were selected based on sample inclusion criteria, such as being aged between 45-55 years old, and having total cholesterol > 160mg/dl. After fasting for 12 hours and still consuming statin anti-cholesterol drugs, blood is taken through the cubiti vein as much as 5ml. To measure the HDL and LDL cholesterol, the enzymatic colourimetry method, diasys reagent, Biolyzer100 spectrophotometric device, in units of mg/dl were implemented. Data analysis was performed with the Wilcoxon, Mann Whitney, Kruskal wallis test with a significance level of 5%, in which the data were pretest-posttest and compared fom thid groups. The research was carried out at the Laboratory of Biochemistry, Universitas Jember.

### Findings and Discussion

**Table 1. Result of analysis HDL level on the cupping, drug, cupping and drug group**

| Groups           | n  | Levels HDL Mean±SD<br>Median ± IQR (Min-Max)  |   | p     |
|------------------|----|---|---|-------|
|                  |    | Pretest                                       | Posttest  |       |
| Cupping          | 17 | 33.03 ± 3.73<br>32.49 ± 5.54<br>(27.81-39.97) | 37.58 ± 6.54<br>35.64 ± 4.34ab<br>(31.59-55.35) | 0.000 |
| Drug             | 17 | 33.40 ± 3.18<br>32.67 ± 5.26<br>(29.27-39.96) | 33.56 ± 3.50<br>34.56 ± 5.13a<br>(25.92-38.61)  | 0.788 |
| Cupping and drug | 17 | 34.32 ± 3.38<br>34.84 ± 5.20<br>(26.19-39.16) | 37.61 ± 2.01<br>37.57 ± 3.25b<br>(35.11-42.17)  | 0.002 |
| p                |    | 0,534   | 0.002   |       |

Add : \* significan on a = 0,05

IQR = Inter Quartile Range

(13.8%).

Measurement of HDL cholesterol on the cupping group was pretest= 33.03±3.73; posttest=37.58±6.54; p=0.000. The drug group was pretest=33.40±3.18; posttest=33.56±3.50; p=0.788. The cupping and drug group was pretest=34.32±3.38; posttest 37.61±2.01; p=0.002. Kruskal Wallis test on pretest group was 0.534 and on posttest group was 0.002.

This data shows an average increase in HDL levels from 33.03 to 37.58. There was an increase of 4.55 point

Measurement of LDL on the cupping group was pretest=154.70±39.68; posttest=123.89±41.86; p=0.000. The drug group was pretest=151.24±44.17; posttest=151.24±44.17; p=0.019. The cupping and drug group was pretest=147.48±62.66; posttest=105.57±57.94 ; p=0.001. Kruskal Wallis test on pretest was 0.439 and on posttest was 0.082. This means that there was no difference in the average pretest and posttest LDL levels in the cupping, drug, cupping and drug groups.

**Table 2. Result of analysis LDL Level on the cupping, drug, cupping and drug group**

| Groups           | n  | LDL level<br>Mean±SD<br>Median ± IQR (Min-Maks)     |  | p     |
|------------------|----|---|--|-------|
|                  |    | Pretest   | Posttest   |       |
| Cupping          | 17 | 154.70 ± 39.68<br>151.03 ± 79.77<br>(94.34-214.92)  | 123.89 ± 41.86<br>116.45 ± 81.13<br>(67.45-193.18) | 0.000 |
| Drug             | 17 | 151.24 ± 44.17<br>138.34 ± 40.19<br>(112.16-284.03) | 136.40 ± 42.12<br>130.71 ± 42.40<br>(83.08-264.89) | 0.019 |
| Cupping and Drug | 17 | 147.48 ± 62.66<br>136.96 ± 28.65<br>(74.84-337.66)  | 105.57 ± 57.94<br>95.41 ± 74.37<br>(38.08-256.47)  | 0.001 |
| p                |    | 0,439   | 0.082  |       |

Add : \* significan on a = 0,05

IQR = Inter Quartile Range

This data shows an a decrease in LDL levels from an average of 154 to 123. There was a decrease of 31 point (20%).

The National Cholesterol Education Program Adult Treatment Panel III (NCEP ATP III) guidelines recommend checking non-HDL cholesterol levels containing apolipoprotein B when TG levels elevated.<sup>(9)</sup>

The main action of statin anti cholesterol drugs in reducing blood cholesterol is inhibiting HMG CoA reductase, which reduces the concentration of hepatocyte cholesterol and stimulates absorption of liver LDL cholesterol from the circulation.<sup>(10)</sup> However, there is an increasing evidence stating that statins can also have pleiotropic effects. Statins can alter the synthesis and metabolism of lipoproteins and accelerate the degradation of intracellular apolipoprotein B, resulting in a reduction in apopoprotein B secretion.<sup>(11)</sup> The Atorvastatin Comparative Cholesterol Efficacy and Safety Study

(ACCESS) in patients with hypercholesterolaemia reported reductions in plasma apoprotein B levels following statin therapy.<sup>(12)</sup>

This measurement follows the research conducted by Saryono<sup>(13)</sup>, that blood cupping therapy can reduce cholesterol levels. As a result of keratinocyte enrichment in the skin the body can experience hypoxia and induce Hypoxia Inducible Factor (HIF-1 $\alpha$ ) as a self-defense effort.<sup>(14)</sup> HIF-1 $\alpha$  will activate macrophages in the skin which subsequently induce proinflammatory genes such as IL-1, IL-4, IL-6 and TNF- $\alpha$ .<sup>(3)</sup> Interleukin-6 which is secreted by macrophages acts to stimulate the body's immune response, for example after trauma or tissue damage that leads to inflammation. The release of IL-6 stimulates young macrophage cells to mature and be able to do phagocytosis more efficiently. IL-6 also stimulates monocytes to produce inflammatory cytokines that play a role in local and systemic inflammation, resulting in accelerated proliferation and differentiation of macrophages.<sup>(1)</sup>

LDL is a cholesterol source for extrahepatic tissue. When LDL is very excessive, its capture system will be saturated so that excessive LDL can be taken by macrophages. Some LDL cholesterol is captured by macrophages before being oxidized. The more LDL

cholesterol levels exist in the plasma, the more cells will be captured by macrophages. Furthermore, macrophages will experience efflux and nascent HDL will approach the macrophage to take LDL cholesterol. Then, nascent HDL becomes adult HDL. After taking cholesterol free from macrophage cells, free cholesterol will be esterified to cholesterol ester by the enzyme Lecithin Cholesterol Acyl Transferase (LCAT). So HDL here serves as an absorber of LDL cholesterol from macrophages and as a carrier of LDL cholesterol back to the liver so that cholesterol levels in plasma decrease.<sup>(7)</sup>

According to El-Sayed et al.<sup>(4)</sup>, blood cupping therapy is an excretory minor surgical procedure that has medical and scientific basis for cleaning blood and interstitial space from cholesterol causative pathological substances (CPS) as metabolic waste production. Many studies have reported that blood cupping therapy can reduce LDL cholesterol. HDL cholesterol functions as an absorber of LDL cholesterol from macrophages and as an LDL cholesterol carrier back to the liver with the help of pre-HDL.<sup>(15)</sup> Pre-HDL has a role in the reverse cholesterol transport to increase excess cholesterol efflux from peripheral tissue back to the liver to be excreted through bile. The acceleration of migration of macrophages also increases due to stimulation of IL-6.<sup>(16,17)</sup>





Figure 1. The process of taking out metabolic waste from the blood

Source: Elsayed, 2013<sup>(4)</sup>

HDL is released as small cholesterol-poor particles containing apolipoprotein (apo) A, C, and E, and is called HDL nascent. Nascent HDL comes from the small intestine and liver, has a flat shape and contains apolipoprotein A1. Nascent HDL will approach macrophages to take cholesterol stored in macrophages. After taking cholesterol in macrophages, the nascent HDL turns into a rounded adult HDL. In order to be retrieved by nascent HDL, cholesterol (free cholesterol) on the inside of the macrophage must be brought to the surface of the macrophage cell membrane by a transporter called Adenosine Triphosphate-Binding Cassette Transporter-1

(ABC-1). After taking free cholesterol from macrophage cells, free cholesterol is esterified into cholesterol ester by the enzyme Lecithin Cholesterol Acyl Transferase (LCAT). Furthermore, some cholesterol esters carried by HDL take two pathways. The first path is to the heart and is captured by a class B type 1 receptor known as SR-B1. The second pathway is the cholesterol esters in HDL will be exchanged with TG from VLDL and IDL with the help of Cholesterol Esters Transfer Protein (CETP). Thus the function of HDL as a cholesterol absorber from macrophages has two pathways namely direct to the liver and indirect pathways through VLDL and IDL to carry cholesterol back to the liver.



Source: Kwiterovich PO, 2000<sup>(18)</sup>

### Figure 2. Reverse Cholesterol Transport Pathway

Metabolism of apolipoprotein is related to the development of atherosclerosis. Increased levels of apolipoprotein B or low levels of apolipoprotein A-I related to cardiovascular disease. Because apolipoprotein B and apolipoprotein A-I have opposite effects namely atherogenic and atheroprotective.<sup>(19)</sup>

Currently, the inflammatory mediators implicated in the pathogenesis of atherosclerosis include cytokines, chemokines, vasoactive molecules and growth factors. The anti-inflammatory effects of statins are attributed to multifaceted mechanisms including inhibition of cell cycle progression, induction of apoptosis, reduction of cyclooxygenase-2 activity and a biphasic, dose-dependent effect on angiogenesis.<sup>(1)</sup>

Cholesterol in nascent HDL is then esterified as fatty acids derived from lecithin by lecithin acyl cholesterol



transferase (LCAT) and its co-factor, apo A-I, producing spherical, mature HDL particles. Cholesterol esters in the HDL nucleus are then returned to the liver, either through the interaction of HDL with SR-B1 receptors, or transferred to lipoproteins containing B apo by cholesterol transfer proteins (CETP). (Reproduced with permission from H. Bryan Brewer Jr., personal communication, National Heart, Lung, and Blood Institute.<sup>(18)</sup>

### Conclusion

The intervention of blood cupping therapy has the potential to reduce levels of Apolipoprotein-B and total cholesterol in the blood. Bloodcupping therapy can be considered as an intervention that can reduce cholesterol, in addition to the use of anti-cholesterol drugs. Further research needs to be done to measure the potential prevention of atherosclerosis.

The main effect of statins decreasing the serum level of low-density lipoprotein (LDL) cholesterol, due to the inhibition of intracellular cholesterol biosynthesis. A minor effect is the decrease of serum triglycerides. Statins inhibit HMG-CoA reductase and decrease the production of mevalonate, geranyl pyrophosphate, and farnesyl pyrophosphate, and subsequent products on the way to construction of the cholesterol molecule. Thus, statins could inhibit inflammation, by inhibition of the cholesterol pathway and intracellularly interfering with Ras superfamily protein function. Mutations in transporter molecules or receptors can lead to an accumulation of cholesterol and a breakdown in the normal process of reverse cholesterol transport and cholesterol metabolism.

**Ethical Clearance:** The research ethical approval was obtained from the Ethics Committee of the University of Jember in Agustus 2019

**Source of Funding:** Authors

**Conflict of Interest:** No

### References

1. Mahdavi V, Reza M, Ghazanfari T. Evaluation of the effects of traditional cupping on the biochemical, hematological, and immunological factors of human venous blood. *A Compend Essays Altern Ther.* 2012:67-88.
2. Mahmoud HS, Abou-el-naga M, Ayoub N, Omar A, El-ghazzawy HA, Fathy YM. Alternative & integrative drug anatomical sites for practicing wet cupping therapy (al-blood cupping therapy): in light of modern drug and prophetic drug. 2013;2(8). doi:10.4172/2327-5162.1000138
3. Subadi I, Boya Nugraha, Hening Laswati, and Harjanto Josomuljono, 2017. Pain Relief with Wet Cupping Therapy in Rats is Mediated by Heat Shock Protein 70 and  $\beta$ -Endorphin. *Iran J Med Sci.* 2017;42(4):384–391.
4. El-Sayed SM, Mahmoud HS and Nabo MMH. Medical and Scientific Bases of Wet Cupping Therapy; in Light of Modern Drug and Prophetic Drug. *Alternative and Integrative Drug.* 2013;2:1-16.
5. Mourad SA, Soad K Al-Jaouni. The Effect of Wet Cupping on Blood Haemoglobin Level. *Alternative Integration Meditoin.* 2016;5:2.
6. Grundy SM. Low density lipoprotein, non-high density lipoprotein, and apolipoprotein B as targets of lipid-lowering therapy. *Circulation.* 2002;106:2526–9.
7. Mustafa L, Dawood M. Rukzan, Al-Sabaawy M. Osama. Effect of Wet Cupping on Serum Lipids Profile Levels of Hyperlipidemic Patients and Correlation with some Metal Ions. *Raf. J. Sci.* 2012;23(3):128-136.
8. Yan Xiaotong, Mingjie Yao, Xijie Wen, Yiwei Zhu, Erjiang Zhao, Xiangjun Qian, Xiangmei Chen, Weiquan Lu, Quanjun Lv, Ling Zhang, Fengmin Lu. Elevated apolipoprotein B predicts poor postsurgery prognosis in patients with hepatocellular carcinoma *OncoTargets and Therapy* 2019:12.
9. Steer P, Hulthe J, Millg<sup>o</sup>rd J et al. Endothelial vasodilatory function is predicted by circulatory apolipoprotein B and HDL in healthy humans. *Lipids.* 2002;37:1135–40.
10. Srivastava RAK, Srivastava N. High density lipoprotein, apolipoprotein A-I, and coronary artery disease. *Mol Cell Biochem.* 2000;209:131–44.
11. Wilcox LJ, Barrett PHR, Huff MW. Differential regulation of apolipoprotein B secretion from HepG2 cells by two HMGCoA reductase inhibitors, atorvastatin and simvastatin. *J Lipid Res.* 1999;40:1078–89.
12. Walldius G, Jungner I, Aastveit A, Holme I, Sniderman AD. Increased risk of fatal myocardial

- infarction related to high triglycerides, high apoB and low apoA-I in the AMORIS study. *Diabetes*. 2002;51(Suppl.):A20.
13. Saryono. Decrease in total cholesterol levels in hypertensive patients who received blood cupping therapy at the An-Nahl Purwokerto clinic. *Sudirman Nursing Journal (JKS)*. 2010;5(2):66–73.
  14. Barriga EH, Maxwell PH, Reyes AE, Mayor R. The hypoxia factor Hif-1 $\alpha$  controls neural crest chemotaxis and epithelial to mesenchymal transition. *J Cell Biol*. 2013;201(5):759-76. doi: 10.1083/jcb.201212100.
  15. Praningsih S. The effectiveness of spiritual cupping care compared to cupping care in reducing LDL cholesterol. Thesis. Unair Repository. 2017. <http://repository.unair.ac.id/70326/>
  16. Mehta P, Dhapte V. Cupping therapy: A prudent remedy for a plethora of medical ailments. *J Tradit Complement Med*. 2015;5(3):127-134. doi:10.1016/j.jtcme.2014.11.036
  17. Tagil SM, Celik HT, Ciftci S, Kazanci FH, Arslan M, Erdamar N, dkk. Wet-cupping removes oxidants and decreases oxidative stress. *Complement Ther Med*. 2014;22(6):1032-1036. doi:10.1016/j.ctim.2014.10.008
  18. Kwiterovich, PO Jr., MD, 2000, The metabolic pathways of high-density lipoprotein, low-density lipoprotein, and triglycerides: a current review. *Am J Cardiol* 2000,86:5L-10L.
  19. Bolibar I, von Eckardstein A, Assmann G, Thompson S. Short-term prognostic value of lipid measurements in patients with angina pectoris. The ECAT Angina Pectoris Study Group: European Concerted Action on Thrombosis and Disabilities. *Thromb Haemost* 2000; 84: 955–60.