

Preconditioning of Hypoxic Culture Increases The Therapeutic Potential of Adipose Derived Mesenchymal Stem Cells

by Tito Sumarwoto

Submission date: 18-Aug-2022 04:41PM (UTC+0800)

Submission ID: 1883877087

File name: artikel-Preconditioning_of_Hypoxic.pdf (2.84M)

Word count: 7180

Character count: 41653



Preconditioning of Hypoxic Culture Increases The Therapeutic Potential of Adipose Derived Mesenchymal Stem Cells

Tito Sumarwoto^{1,2}, Heri Suroto³, Ferdiansyah Mahyudin³, Dwikora Novembri Utomo³, Romaniyanto², Andhi Prijosedjati², Pamudji Utomo², Cita Rosita Sigit Prakoeswa⁴, Fedik Abdul Rantam^{5,6}, Damayanti Tinduh⁷, Hari Basuki Notobroto⁸, Sholahuddin Rhatomy^{9,10,*}

¹Doctoral Program, Faculty of Medicine, Airlangga University, Surabaya, Indonesia; ²Department of Orthopaedics and Traumatology, Faculty of Medicine, Prof Soeharso Orthopaedic Hospital, Sebelas Maret University, Surakarta, Indonesia; ³Department of Orthopaedic and Traumatology, Faculty of Medicine, Dr. Soetomo General Hospital, Airlangga University, Surabaya, Indonesia; ⁴Department of Dermatology and Venereology, Faculty of Medicine, Dr. Soetomo General Hospital, Airlangga University, Surabaya, Indonesia; ⁵Department of Microbiology, Virology and Immunology Laboratory, Faculty of Veterinary Medicine, Airlangga University, Surabaya, Indonesia; ⁶Stem Cell Research and Development Center, Airlangga University, Surabaya, Indonesia; ⁷Department of Physical Medicine and Rehabilitation, Faculty of Medicine, Dr. Soetomo General Hospital, Airlangga University, Surabaya, Indonesia; ⁸Faculty of Public Health, Airlangga University, Surabaya, Indonesia; ⁹Department of Orthopaedics and Traumatology, Dr. Soeradji Tirtonegoro General Hospital, Klaten, Indonesia; ¹⁰Faculty of Medicine, Public Health, and Nursing, Gadjah Mada University, Yogyakarta, Indonesia

Abstract

Stem cell-based therapy is one of the most promising strategies in promoting tissue regeneration. Mesenchymal stem cells (MSCs) derived from adipose are widely used as a source of stem cells. However, most donor stem cells die before their regenerative effects appear. Thereby, various strategies are carried out to enhance the therapeutic potential. It is hoped that the survival, engraftment, and function of implanted MSCs by preconditioning the cells will be improved. *In vitro* preconditioning strategies can improve the survival, engraftment, and paracrine properties of MSCs, thus optimize their reparative and regenerative capacities. One such attempt is to propagate these stem cells in a hypoxic environment, which mimics the natural tissue oxygen environment. Hypoxia preconditioning represents a comprehensive increase in regenerative gene expression in stem cell therapy. There are significant advantages in stem cell therapy with hypoxia preconditioning.

Edited by: Eli Djulejic

Citation: Sumarwoto T, Suroto H, Mahyudin F, Utomo DN, Romaniyanto, Prijosedjati A, Utomo P, Prakoeswa CRS, Rantam FA, Tinduh D, Notobroto HB, Rhatomy S. Preconditioning of Hypoxic Culture Increases The Therapeutic Potential of Adipose Derived Mesenchymal Stem Cells. Open Access Maced J Med Sci. 2021 Oct 31; 9(F):505-515. <https://doi.org/10.3889/oamjms.2021.5870>

Keywords: Hypoxic; Culture; Adipose; Derived Mesenchymal Stem Cells

*Correspondence: Sholahuddin Rhatomy, Department of Orthopaedics and Traumatology, Faculty of Medicine, Public Health, and Nursing, Universitas Gadjah Mada, Yogyakarta, Indonesia. KRT. Dr. Soeradji Tirtonegoro Street Number 1, Tegalyoso, South Klaten, Klaten, Indonesia 57424. Tel: +62272-321102, Fax: +62272-321104. E-mail: sholahuddin.rhatomy@mail.ugm.ac.id

Received: 09-Feb-2021

Revised: 15-Apr-2021

Accepted: 21-Oct-2021

Copyright: © 2021 Tito Sumarwoto, Heri Suroto, Ferdiansyah Mahyudin, Dwikora Novembri Utomo, Romaniyanto, Andhi Prijosedjati, Pamudji Utomo, Cita Rosita Sigit Prakoeswa, Fedik Abdul Rantam, Damayanti Tinduh, Hari Basuki Notobroto, Sholahuddin Rhatomy

Funding: This research did not receive any financial support

Competing Interest: The authors have declared that no competing interest exists

Open Access: This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0)

Introduction

Stem cell-based therapy is one of the most promising strategies in the medical field to promote tissue regeneration. Over the past few decades, stem cells have been widely used as regenerative therapy and mesenchymal stem cells (MSCs) emerged as a promising therapeutic option. Currently, stem cells are effective agents as therapeutic agents for several diseases due to their tissue protective and reparative mechanisms [1].

MSCs derived from adipose tissue are widely used as a source of stem cells. This is due to the fact they are accessible, abundant, and painless collection procedure in comparison to other sources. These cells can be maintained and cultured for long periods of time without losing their differentiation capacity [2].

Unfortunately, several recent studies have shown that donor stem cells die before their regenerative effects appear. Most of the stem cells did not survive the first 24 h *in vivo*, mainly due to the hypoxia-induced apoptosis of stem cells [3].

For this reason, various strategies are carried out to increase the therapeutic potential. Thus, it will improve the survival, engraftment, and function of implanted MSCs by conditioning the cells prior to implantation, known as preconditioning. To obtain optimal therapeutic potential, an adipose-derived MSCs (AdMSC) with maximum regenerative capacity is required and *in vitro* initial conditioning strategy can optimize its therapeutic potential [4].

One such attempt is to propagate these stem cells in a hypoxic environment prior to implantation (hypoxia preconditioning), which mimics the natural tissue oxygen environment (1–7%), unlike standard culture conditions (21%). This will increase the viability of these cells and will stimulate the secretion of growth factors required in the regenerative process [5], [6].

The following review aims to provide an overview of hypoxia preconditioning to enhance the therapeutic potential of AdMSCs.

Characteristics of MSCs

According to the MSC and Tissue Committee of the International Society for Cellular Therapy, MSCs have both self-renewing and differentiating capabilities. These cells showed positive expression for stromal markers CD73 and CD 105 and negative for the hematopoietic markers CD14 or CD 11b, CD34 and CD45; CD79a or CD19 and HLA class II. These cells have the ability to be plastic adherent and possess fibroblast-like morphology. These cells can be maintained in culture for a long period of time. Nevertheless, their potential for osteogenic, adipogenic, and chondrogenic differentiation must be demonstrated [2], [7].

The most useful characteristic reported for MSC is the non-immunogenic profile. MSCs are reported to exhibit Major Histocompatibility Complex (MHC) I, but deficiency of MHC II results in T-cell inactivation and immunosuppressive properties. MSCs secrete a variety of growth factors, chemokines, cytokines, and extracellular matrix (ECM) proteins that are involved in various biological processes including hematopoiesis, angiogenesis, leukocyte multiplication, immunity, and inflammatory responses. Multiple growth factors, including basic fibroblast growth factor (bFGF), vascular endothelial growth factor (VEGF), insulin-like growth factor 1, hepatocyte growth factors (HGF), and transforming growth factors (TGF)- β 1 are also secreted. This evidence holds a great promise in MSC-based therapies which allow allogeneic transplantation without the need for immunosuppression [8], [9], [10].

AdMSCs

Stem cells in adipose tissue, called AdMSC, have been shown to have broadly the same biological capabilities as bone marrow MSC (BM-MSCs) [11]. The advantages of AdMSC over BM-MSCs and other types of adult stem cells are that they are relatively easy to obtain. It can be obtained in large quantities through liposuction performed under local anesthesia. These cells are able to be maintain long-term phenotype and plasticity of *in vitro* culture. These cells also possess low immunogenicity characteristic. As a result, AdMSC was able to attract greater interest and is the most preferred cell type for tissue engineering and regenerative medicine [12], [13], [14].

Stem cells found in adipose tissue have a higher number than those in bone marrow. One gram of adipose tissue yields approximately 3.5×10^5 to 1×10^6 AdMSC, whereas isolation from one gram of bone marrow aspiration yields only 5×10^2 to 5×10^4 BM-MSC. In addition, AdMSC has advantages in terms of proliferation and differentiation, as well as the fact that age and location of origin do not have a significant difference in the therapeutic effect of these adipose-derived stem cells [9], [15].

Zuk *et al.*, were the first to investigate whether human adipose could be an alternative source of MSCs. They obtained human adipose from liposuction aspiration and used the enzyme collagenase to release stromal cells from the ECM by processing the stromal vascular fraction, which contains various cell types including AdMSC (Figure 1). Isolated adipose stromal cells are cultured with specific media to induce adipogenic, osteogenic, or chondrogenic differentiation. Adipose stromal cells are capable of developing intracellular lipid stores, markers representing adipose tissue, bone, and cartilage, respectively, as well as expressing alkaline




Figure 1: Isolation and utilization of adipose-derived mesenchymal stem cells [13]

phosphatase or proteoglycan [16]. To determine whether isolated adipose stromal cells were truly stem cells, Zuk *et al.*, investigated surface antigen expression and differentiation capacity against clonogenic cultures [17].

Using flowcytometry, it was shown that clonogenic cells express a surface antigen similar to BM-MSCs. In addition to the differentiation of mesenchymal lineages, clonal cells are also able to differentiate into neuron-like cells, which are proven by morphology and expression of phenotypic markers. AdMSCs also tend to stimulate angiogenesis as an important feature for regenerative [18], [19]. Its neurotropic and angiogenic properties have been shown to be due to neural growth factor (NGF), brain-derived neurotrophic factor (BDNF), glial cell-derived neurotrophic factor (GDNF), VEGF-A, and angiopoietin-1 [20]. The secretome of AdMSC is very complex. AdMSC has the ability to secrete proteins involved in angiogenesis, injury healing, tissue regeneration, and immunomodulation [21]. In addition, AdMSC has the capacity to differentiate into Schwann cell [22], pancreatic beta cells [23], and hepatocytes [24]. AdMSCs have important healing properties in injuries and can be an attractive source of stem cells for tissue engineering as well as regenerative therapy for many medical and surgical applications.

A comparison of the yield and differentiation capacity of cells isolated from each tissue was performed to determine whether adipose tissue is a source of MSC bone marrow. De Ugarte *et al.*, [25] found no significant difference in the number of cultured adherent cells between those of stromal cells obtained from bone marrow and those from adipose tissue. However, the average mass of adipose tissue (17 g)

obtained from each patient was higher than the average mass of bone marrow (7 g). Cultured isolated cells in various differentiation media showed no difference in the number of cells developing lipid droplets (adipogenic cells), or osteogenic cell alkaline phosphatase activity. However, in terms of differentiation towards cartilage, adipose-derived cells were positively stained for chondrogenesis whereas bone marrow-derived cells were not.

Using a similar method, several other studies have compared the ability of bone marrow and adipose cells to differentiate along this lineage and demonstrated that cells from both tissues have the same capacity to become adipose, bone, and cartilage tissue [26], [27], [28]. Overall, this is the result of different culture conditions and/or isolation of different parts of MSC. Carbone *et al.* demonstrated that AdMSC cultured in conditioned medium from chondrocyte and osteocytes were capable of producing glycosaminoglycans and mineralized matrix, respectively [29]. These results suggest that AdMSC requires growth factor supplementation from the tissue environment in order to properly differentiate according to the mesodermic lineage. This evidence suggests that compared to bone marrow, a large number of MSCs capable of differentiating into multiple lineages can also be obtained from adipose tissue (Figure 2).

AdMSCs Precondition Concept

Therapy using MSCs is a therapy that provides hope for organ and tissue regeneration. The




Figure 2: Adipose-derived mesenchymal stem cells: Origin, immunophenotype, morphology, and differentiation potential. Lipoaspirate can be easily obtained and processed to obtain a heterogeneous cell population including stromal vascular fraction. Characterization of the cell was by mesenchymal stem cell markers (CD73, CD90, CD105) at the expense of hematopoietic stem cell markers (CD45) and their spindle-shaped morphology with their ability to differentiate into the adipogenic, osteogenic, and chondrogenic lineages. Histological staining was able to analyze the differentiation potential, Oil red O for adipogenic differentiation, Alizarin red for osteogenic differentiation, and Alcian blue for chondrogenic differentiation [30].

conditioned medium or MSCs-EVs exert a protective effect on peripheral nerve tissue and have been shown to promote regeneration in injured tissue. However, transplanting MSCs or administering MSCs-CM or MSCs-EV requires the MSC to have the maximum regenerative capacity. Thus, it is necessary to develop a new strategy to increase the regenerative efficiency of MSCs (Figure 3). *In vitro* preconditioning strategies can improve the survival, engraftment, and paracrine properties of MSCs so as to optimize their reparative and regenerative capacities [4].




Figure 3: Main causes of transplanted cells death. The major factors limiting cell survival during transplantation and strategies to improve cell survival on cell-based therapy. ECM: Extracellular matrix, ROS: Reactive oxygen species [31]

Various *in vitro* preconditioning strategies have been implemented to increase the regenerative capacity of MSCs. Among them is modulation of culture atmosphere (hypoxia or anoxia), three-dimensional (3D) culture, addition of trophic factors (in the form of growth factors, cytokines, or hormones), lipopolysaccharides, and pharmacological agents. Factors secreted by MSCs in response to many different preconditioning and exert anti-apoptosis, pro-angiogenic, and trophic or immunosuppressive immunomodulatory effects (Figure 4).




Figure 4: *In vitro* preconditioning. Several mechanisms are involved in enhancing the therapeutic potency of mesenchymal stem cells under prerequisite condition strategies [4]

The secretome (paracrine profile) of the pre-treated MSC varies according to the preconditioning method. Different preconditioning methods can activate or suppress molecular signals and different signal transduction cascades. The cellular response is very complex and the pre-treatment procedure affects not only one specific molecule or protein but also a number of factors [32].

Hypoxic Preconditions and Hypoxia Inducible Factor (HIF)-1 α

Several strategies were carried out to increase the therapeutic potential of AdMSCs by increasing the survival, grafting, and function of the implanted MSCs by conditioning the cells prior to implantation. Implantation of MSCs into nutrient-deprived and oxygen-deprived environments, such as in recipient tissue, results in extremely low cell survival [32]. Among these strategies, hypoxia preconditioning represents a comprehensive increase in regenerative gene expression in stem cell therapy. Previous studies have shown the successful application of this strategy in several diseases. Furthermore, there are significant advantages in stem cell therapy in the precondition of hypoxia [33].

Hypoxia preconditioning ensures the cells to adapt to the ischemic environment [34]. Preconditioning MSCs can be done by culturing them in a hypoxic environment, which mimics the natural tissue oxygen environment (1–7%) unlike the standard culture conditions (21%). This increases the survival of these cells via HIF-1 α and through Akt-dependent mechanisms. In addition, the hypoxia preconditioning stimulates the secretion of pro-angiogenic growth factor, increases the expression of chemokines stromal cell-derived factor-1 (SDF-1) and its receptor chemokine receptor type 4 (CXCR) - CXCR7 and increases engraftment of MSCs *in vivo* [5], [6], [35]. Hypoxia preconditioning has also been shown to enhance the paracrine effect of MSCs and has been applied to several diseases, such as myocardial infarction and diabetes-related erectile dysfunction [36].

The *in vivo* oxygen concentration is about 0.4%, significantly lower than the oxygen concentration conventionally used *in vitro*, 21% [37]. Most stem cells die within the first 24 h *in vivo*, mainly due to stem cell apoptosis induced by hypoxia [3]. Research by injecting bone marrow stem cells into mice with ventricular myocardial infarction found that 99% of bone marrow stem cells died 4 days later. These results indicate that these stem cells are highly susceptible to ischemia and hypoxia [38]. Based on these results, many studies have used hypoxic state methods to improve stem cell viability. Greijer and van der Wall showed that the severity of hypoxia affects the rate of cell apoptosis versus cell survival during hypoxia. The 0.5% O₂ condition was shown to initiate apoptosis in some cells. In order to prevent the accumulation of hypoxia-induced genetic mutations, there is a balance between pro-apoptotic and anti-apoptotic factors. HIF-1 α as shown in Figure 5, plays an important role in maintaining that balance [39].

Sun *et al.* simulated a hypoxic environment to induce hypoxic precondition in bone marrow stem cells and found that after hypoxia preconditioning, the bone marrow stem cells overexpressed HIF-1 α , leading to decreased apoptosis and minimizing loss




Figure 5: The three-dimensional structure of hypoxia-inducible factor (HIF)-1 α and HIF-1 β [40]

of mitochondrial membrane potential. These results suggest that hypoxia preconditioning has a protective effect on the survival of BM-MSCs *in vivo* under ischemic and hypoxic conditions. Once MSCs are exposed to the precondition of hypoxia (Figure 6), the hypoxia-induced apoptosis pathway is rapidly activated thereby inhibiting feedback from this pathway and then reducing apoptosis in subsequent ischemic and hypoxic conditions [41].




Figure 6: Hypoxia-induced signaling pathways with HIF-1 α Involvement. Schematic depiction of signaling pathways induced by hypoxia leading to apoptosis suggesting the involvement of HIF-1 α . Solid lines represent direct interactions, dotted lines represent indirect interactions. Apaf-1: Apoptotic protease activating factor-1, BNIP3: BCL-2/adenovirus E1B 19 kDa interacting protein 3, HIF-1: Hypoxia-inducible factor 1, IAP-2: Inhibitor of apoptosis protein 2, JNK: c-Jun NH2 terminal kinase, SAPK: Stress activated protein kinase [39]

HIF-1 α is a subunit of a heterodimeric transcription factor HIF-1 α encoded by the HIF-1 α

gene. It is the basic domain of the PAS helix-loop-helix-containing proteins and is considered to be the primary transcription regulator of cellular responses and cellular development responses to hypoxia. Dysregulation and overexpression of HIF-1 α caused by hypoxia or genetic alteration are involved in cancer biology, a number of pathophysiology in the fields of vascularity and angiogenesis, metabolism energy, cell survival, and tumor invasion [42].

HIF-1 α protein plays a major role in regulating the expression of many genes involved in angiogenesis and erythropoiesis, metabolic adaptation to hypoxia, epithelial-mesenchymal transition, extracellular matrix degradation, and chemotaxis through CXCR4 and the CXCL12/SDF-1 axis [43].

Exposure of MSCs to hypoxic conditions may enhance their immunomodulatory and regenerative properties by over-expression of cytoprotective genes and secretory factors. HIF-1 α has a crucial role in upregulating these genes and factors [44]. HIF-1 α can promote MSC migration to ischemic and hypoxic sites by regulating the expression levels of molecules such as stromal cell-derived factor 1 (SDF1) in the microenvironment. Stem cells express the SDF1 receptor C-XC CXCR4 and CXCR7, and serve a key role in tissue repair [45].

Hypoxia activates the hypoxia signaling pathway, which is largely regulated by HIF stabilization (Figure 7). Under normoxic conditions, the proline residues from the HIF-1 α subunit is hydroxylated by oxygen-dependent prolyl-4-hydroxylase (PHD). The Von Hippel-Lindau protein (pVHL), a ubiquitin E3 ligase, binds to the hydroxylated HIF-1 α and acts as a substrate recognition component of the ubiquitin E3 ligase complex, leading to proteosomal degradation of the HIF protein. Asparagine residue from the HIF-1 α subunit is also hydroxylated by




Figure 7: Regulation of hypoxia-inducible factor (HIF) during normoxic and hypoxic states. In the oxygenated state, HIF undergoes hydroxylation of the proline residue by PHDs and is polyubiquitinated by pVHL. This results in the degradation of the HIF by the 26S proteasome system. In a hypoxic state, HIF is stabilized and translocated into the nucleus, where HIF1 α binds to the HIF1 β dimerization partner and increases transcription of the HIF target gene [46]

factor inhibiting HIF (FIH), which inhibits the binding of HIF to the p300/CREB binding protein co-activator [46].

In a hypoxic state, PHD and FIH activity are suppressed, and the HIF- α subunit translocates into the nucleus to bind to HIF-1 β . The heterodimeric complex of transcription factor HIF-1 α : HIF-1 β then looks for hypoxia-responsive elements of its target genes, which results in transcriptional upregulation (Figure 8). There are other HIF-independent signaling pathways that are activated under hypoxia, such as the factor- κ B (NF- κ B) pathway. Early studies reported that I κ B α is phosphorylated during hypoxia and this results in degradation of I κ B α and activation of NF- κ B1. Other studies have shown that I κ B kinase activity is increased via calcium/calmodulin-dependent kinase 2 (calcium/calmodulin-dependent kinase 2) during hypoxia and required kinase 1 activated by TGF- β [47].

(Nobel Prize in Physiology or Medicine 2019: How Cells Sense and Adapt to Oxygen Availability).

Hypoxia can cause tissue inflammation and HIF stabilization can reduce tissue inflammation and promote its repair. HIF elicits a variety of adaptive

responses, particularly focusing on enhancing the transcription cascade which is important for tissue protection and adaptation. HIF-1 α (HIF1A) is known to be associated with the upregulation of glycolytic gene such as phosphoglycerate kinase (CKD) and lactate dehydrogenase A, both of which function metabolically to adapt tissues to oxygen uptake and anaerobic ATP synthesis. HIF-2 α (HIF2A) induces erythropoietin and VEGF, which are important for increasing oxygen supply to hypoxic areas [48].

The secretion of several cytokines associated with neuroprotective and vascular protection was significantly increased from MSCs after hypoxic conditions, namely VEGF, bFGF, BDNF, and GDNF [30]. Hypoxia was reported to increase the angiogenic and mitogenic paracrine activity of AdMSCs. Hypoxia (0.5–2% O₂) significantly increases VEGF, HGF, bFGF, IL-6, angiogenin, NGF, IGFBP-1, IGFBP-2, and the PDGF-b receptors. In contrast, EGF and anti-angiogenic factors are inhibited by hypoxic condition. The increased paracrine factors improve survival, homing, and angiogenesis, which increases the regenerative potential of AdMSCs after transplantation [9], [49].




Figure 8: Hypoxia-inducible factor (HIF)-1 α in Normoxia and Hypoxia State. Under normoxic conditions, HIF-1 α is hydroxylated at two proline residues. The protein is then combined with VHL and then tagged with ubiquitin. After the ubiquitination process, it is transferred to the proteasome which is then undergone degradation. Under hypoxic conditions, no oxygen is available for the hydroxylation of proline. In this situation HIF-1 α translocates to the cell nucleus (nucleus) and combines with HIF-1 β , also called aryl hydrocarbon receptor nuclear translocator. This complex then binds to a region of DNA called the hypoxia-responsive element, which causes the target gene to be transcribed. This gene is involved in the regulation of many processes, including erythropoiesis, glycolysis, and angiogenesis

Flowcytometric analysis showed that the stem cell phenotype of adipose derivatives was not significantly affected by hypoxic conditions for 24 h. Hypoxic conditions can cause MSCs to remain undifferentiated for a certain period of time [50]. Several research reports indicate the existence of hypoxia-induced differentiation of MSCs, however, to grow MSCs in a hypoxic state and in differentiation condition it takes a longer time (can be more than 2 weeks). After transplantation, MSCs in the precondition of hypoxic state significantly reduced vascular and nerve damage compared with MSCs in the normoxic state [33].

In addition to the paracrine effect, AdMSCs have been shown to have the potential to differentiate into a wider range of cell types including endothelial cells, smooth muscle cells, Schwann cells, and neuron cells [51]. Engraftment and pluripotency of transplanted stem cells can contribute to the regeneration of blood vessels and nerves. Hypoxia preconditioning can increase the involvement and viability of MSCs in the host tissue (Figure 9). At 4 weeks after transplantation, the *in situ* differentiation of the transplanted cells was also analyzed by staining the tissue sections with vascular markers (CD31 and α -SMA). It can be concluded that hypoxic pretreatment increases the binding of AdMSCs to the tissue [36], [52].




Figure 9: Functional change of AdMSCs in hypoxic state. Hypoxia induces the proliferation, migration, and secretion of growth factor AdMSCs *in vitro*. Hypoxia preconditioning increases the regenerative potential of AdMSCs *in vivo*. In addition, hypoxia regulates the differentiation of AdMSCs in an oxygen concentration-dependent manner [49]

Hypoxic conditions of 1–2% O_2 increase the survival or proliferation of AdMSCs. The hypoxic state of 2% O_2 increases the phosphorylation of mitogenic signaling molecules, such as Akt and ERK1/2, with the formation of Nox4-mediated reactive oxygen species (ROS); and 2% O_2 is recommended to increase survival, the proliferation rate of AdMSCs, and migration of AdMSCs. Weijers *et al.* (2011) stated that 1% O_2 increases the proliferation of AdMSCs, while 5% O_2 was controversial in terms of increasing or decreasing the proliferation of AdMSCs [53], [54], [55], [56]. Increased mobilization and adhesion to damaged tissue are important factors that have a positive effect on the

therapeutic potential of AdMSCs. As in the case of proliferative AdMSCs, the formation of Nox4-mediated ROS is important in the migration of hypoxia-promoted AdMSCs [49], [57]. The table below summarizes cellular response of AdMSCs due to the hypoxic state *in vitro*.

Table 1: Cellular Response of AdMSCs In Vitro Hypoxic State [49]

O_2 level	Altered functions	Reference
0.1%	Increased cell death	(Folmar <i>et al.</i> , 2006)
0.5%	Increased ROS level and enhanced paracrine effect	(De Barros <i>et al.</i> , 2013)
1%	Increased growth factor secretion	(Behman <i>et al.</i> , 2004), (Lu <i>et al.</i> , 2013), (Stamasson <i>et al.</i> , 2011)
	Increased proliferation, survival, and reduced aging	(Weijers <i>et al.</i> , 2011)
	Differentiation arrest and maintain undifferentiation state	(Lin <i>et al.</i> , 2006)
	Increased proliferation, colony formation	(Pigard <i>et al.</i> , 2009)
	Upregulated leptin expression	(Efimenko <i>et al.</i> , 2011)
	Increased expression of pro-angiogenic factors and decreased expression of anti-angiogenic factors	(Lee <i>et al.</i> , 2009), (Amos <i>et al.</i> , 2008), (Przytyk <i>et al.</i> , 2013), (Chadokov <i>et al.</i> , 2009)
2%	Increased growth factor secretion	(Lee and Kemp, 2006)
	Increased myogenic potential	(Ozu <i>et al.</i> , 2007)
	Increase proliferation and chondrogenesis	(Mallat <i>et al.</i> , 2006)
	Increased proliferation and decreased osteogenic differentiation	(Kim <i>et al.</i> , 2012), (Kim <i>et al.</i> , 2011)
	Increased proliferation and migration	(Oliveira <i>et al.</i> , 2012)
	Decreased ATP content and downregulation of DNA repair gene	(Vokoram <i>et al.</i> , 2012)
	Increased adipogenic and osteogenic potential	(Wan <i>et al.</i> , 2007), (Surgens <i>et al.</i> , 2012), (Buckley <i>et al.</i> , 2010)
	Increased chondrogenesis	(Merceron <i>et al.</i> , 2010)
5%	Increased chondrogenesis, but decreased osteogenic differentiation	(Stalbos <i>et al.</i> , 2012)
	Increased survival and enhanced paracrine effect	(Ranea <i>et al.</i> , 2012)
	Decreased proliferation and keep undifferentiated state	(Yang <i>et al.</i> , 2012b)
	Slightly increased proliferation	(Yang <i>et al.</i> , 2012b)

Several studies have shown that hypoxic or anoxic preconditions substantially increase the regenerative potential of MSCs (Figure 10). Exposure to reduced partial oxygen pressure to MSCs induces expression of genes involved in migration and homing (e.g. CXCR4 and SDF-1), regulated mainly by the activity of HIF-1 α [58]. In addition, MSCs culture in a hypoxic environment resembles the presence of actual MSCs *in situ* than in the standard 'artificial' (21% O_2) culture conditions. Cellular response to hypoxia *in vitro* varies at different oxygen concentrations; 0.1–5% O_2 [4], 1–7% O_2 [5], [6], 1% O_2 [33], [59], 0.5–2% O_2 [49].




Figure 10: Signaling pathways and molecular changes in a hypoxic state. Hypoxia induces ROS formation primarily by Nox4. Acute increase in cellular ROS levels induces PDGFR- β activation and results in phosphorylation of Akt and ERK1/2. Activation of this signaling pathway induces expression of miR-210 via phosphorylation of NF- κ B and Elk1; elevated miR-210 levels downregulate PTPN2 and ISCU2, thereby increasing the proliferation and migration of AdMSCs. On the other hand, inhibition of proline hydroxylase under hypoxic states stabilizes HIF-1 α , which modulates angiogenic growth factor transcription after binding to hypoxia-responsive elements in the nucleus [49]

Hypoxia preconditioning *in vitro* has been shown to stimulate the secretion of growth factors, cytokines and other proteins, and the release of extracellular vesicles (exosomes and microvesicles) from MSCs [60]. Extracellular vesicles also carry various biomolecules such as growth factors, receptors, enzymes, transcription factors, signaling and immunomodulatory molecules, DNA, RNA transcripts, and non-coding RNA transcript including retrotransposons, vault RNA, long non-coding RNA, and long non-coding microRNA [61], [62].

Hypoxic preconditions have also been shown to increase survival and cell proliferation [63], [64], [65]. Pre-incubation of hypoxic MSCs induces metabolic changes resulting in higher *in vivo* cell survival after transplantation [63]. This suggests that culture in a hypoxic state increases the angiogenic potential of MSCs and improves their survival in both *in vitro* and *in vivo* studies [63], [64]. The signaling pathway involved translocation of HIF-1 α to the nucleus by activation of gene expression (e.g. VEGF), as well as the formation of ROS and phosphorylation of Akt - MAPK ERK1/2 (Figure 11) [66].




Figure 11: Axon-Schwann cell interactions and some of the molecules involved [66]

Conclusion

Transplantation of MSCs or administration of MSCs-CM or MSCs-EV requires MSC to have maximum regenerative capacity, so it is necessary to

develop new strategies to increase the regenerative efficiency of MSCs. *In vitro* preconditioning strategies can improve the survival, engraftment, and paracrine properties of MSCs thus optimize their reparative and regenerative capacities. Among these strategies, hypoxia preconditioning represents a comprehensive increase in regenerative gene expression in stem cell therapy. There are significant advantages in stem cell therapy with hypoxia preconditioning.

References

- Zhang R, Rosen JM. The role of undifferentiated adipose-derived stem cells in peripheral nerve repair. *Neural Regen Res.* 2018;13(5):757-63. <https://doi.org/10.4103/1673-5374.232457> PMID:29862994
- Mazini L, Rochette L, Amine M, Malka G. Regenerative capacity of adipose derived stem cells (ADSCs), comparison with mesenchymal stem cells (MSCs). *Int J Mol Sci.* 2019;20(10):2523. <https://doi.org/10.3390/ijms20102523> PMID:31121953
- Toma C, Pittenger MF, Cahill KS, Byrne BJ, Kessler PD. Human mesenchymal stem cells differentiate to a cardiomyocyte phenotype in the adult murine heart. *Circulation.* 2002;105(1):93-8. <https://doi.org/10.1161/hc0102.101442> PMID:11772882
- Schäfer R, Spohn G, Baer PC. Mesenchymal stem/stromal cells in regenerative medicine: Can preconditioning strategies improve therapeutic efficacy? *Transfus Med Hemother.* 2016;43(4):256-67. <https://doi.org/10.1159/000447458> PMID:27721701
- Hu X, Yu SP, Fraser JL, Lu Z, Ogle ME, Wang JA, *et al.* Transplantation of hypoxia-preconditioned mesenchymal stem cells improves infarcted heart function via enhanced survival of implanted cells and angiogenesis. *J Thorac Cardiovasc Surg.* 2008;135(4):799-808. <https://doi.org/10.1016/j.jtcvs.2007.07.071> PMID:18374759
- Hung S, Pochampally RR, Hsu S, Sanchez C, Chen SC, Spees J, *et al.* Short-term exposure of multipotent stromal cells to low oxygen increases their expression of CX3CR1 and CXCR4 and their engraftment *in vivo*. *PLoS One.* 2007;5:e416. <https://doi.org/10.1371/journal.pone.0000416> PMID:17476338
- Horwitz EM, Le Blanc K, Dominici M, Mueller I, Slaper-Cortenbach I, Marini FC, *et al.* Clarification of the nomenclature for MSC: The international society for cellular therapy position statement. *Cytotherapy.* 2005;7(5):393-5. <https://doi.org/10.1080/14653240500319234> PMID:16236628
- Jacobs SA, Pinxteren J, Roobrouck VD, Luyckx A, van't Hof W, Deans R, *et al.* Human multipotent adult progenitor cells are nonimmunogenic and exert potent immunomodulatory effects on alloreactive T-cell responses. *Cell Transplant.* 2013;22(10):1915-28. <https://doi.org/10.3727/096368912x657369> PMID:23031260
- Tsuji W, Rubin JP, Marra KG. Adipose-derived stem cells: Implications in tissue regeneration. *World J Stem Cells.* 2014;6(3):312-21. <https://doi.org/10.4252/wjsc.v6.i3.312> PMID:25126381

10. Meng F, Zhou D, Li W. Adipose-derived stem cells as a potential weapon for diabetic foot ulcers. *Int J Clin Exp Med*. 2017;10(12):15967-73.
11. Lee HC, An SG, Lee HW, Park JS, Cha KS, Hong TJ, et al. Safety and effect of adipose tissue-derived stem cell implantation in patients with critical limb ischemia: A pilot study. *Circ J*. 2012;76:1750-60. <https://doi.org/10.1253/circj.cj-11-1135> PMID:22498564
12. Porada C, Zanjani E, Almeida-Porada G. Adult mesenchymal stem cells: A pluripotent population with multiple applications. *Curr Stem Cell Res Ther*. 2012;1(3):365-9. <https://doi.org/10.2174/157488806778226821> PMID:18220880
13. Zuk P. Adipose-derived stem cells in tissue regeneration: A review. *ISRN Stem Cells*. 2013;2013(1):1-35. <https://doi.org/10.1155/2013/713959>
14. Casadei A, Epis R, Ferroni L, Tocco I, Gardin C, Bressan E, et al. Adipose tissue regeneration: A state of the art. *J Biomed Biotechnol*. 2012;2012:462543. <https://doi.org/10.1155/2012/462543> PMID:23193362
15. Fairbairn NG, Meppelink AM, Ng-Glazier J, Randolph MA, Winograd JM. Augmenting peripheral nerve regeneration using stem cells: A review of current opinion. *World J Stem Cells*. 2015;7(1):11-26. <https://doi.org/10.4252/wjsc.v7.i1.11> PMID:25621102
16. Zuk PA, Zhu M, Mizuno H, Huang J, Futrell JW, Katz AJ, et al. Multilineage cells from human adipose tissue: Implications for cell-based therapies. *Tissue Eng*. 2001;7(2):211-28. <https://doi.org/10.1089/107632701300062859> PMID:11304456
17. Zuk PA, Zhu M, Ashjian P, De Ugarte DA, Huang JI, Mizuno H, et al. Human adipose tissue is a source of multipotent stem cells. *Mol Biol Cell*. 2002;13(12):4279-95. <https://doi.org/10.1091/mbc.e02-02-0105> PMID:12475952
18. Gimble J, Guilak F. Adipose-derived stem cells: Isolation, characterization, and differentiation potential. *Cytotherapy*. 2003;5(5):362-9. <https://doi.org/10.1080/14653240310003026> PMID:14578098
19. Kim YJ, Kim HK, Cho HK, Bae YC, Suh KT, Jung JS. Direct comparison of human mesenchymal stem cells derived from adipose tissues and bone marrow in mediating neovascularization in response to vascular ischemia. *Cell Physiol Biochem*. 2007;20(6):867-76. <https://doi.org/10.1159/000110447> PMID:17982269
20. Kingham PJ, Kolar MK, Novikova LN, Novikov LN, Wiberg M. Stimulating the neurotrophic and angiogenic properties of human adipose-derived stem cells enhances nerve repair. *Stem Cells Dev*. 2013;23(7):741-54. <https://doi.org/10.1089/scd.2013.0396> PMID:24124760
21. Kapur SK, Katz AJ. Review of the adipose derived stem cell secretome. *Biochimie*. 2013;95(12):2222-8. <https://doi.org/10.1016/j.biochi.2013.06.001> PMID:23770442
22. Kingham PJ, Kalbermatten DF, Mahay D, Armstrong SJ, Wiberg M, Terenghi G. Adipose-derived stem cells differentiate into a Schwann cell phenotype and promote neurite outgrowth *in vitro*. *Exp Neurol*. 2007;207(2):267-74. <https://doi.org/10.1016/j.expneurol.2007.06.029> PMID:17761164
23. Chandra V, Swetha G, Muthyala S, Jaiswal AK, Bellare JR, Nair PD, et al. Islet-like cell aggregates generated from human adipose tissue derived stem cells ameliorate experimental diabetes in mice. *PLoS One*. 2011;6(6):e20615. <https://doi.org/10.1371/journal.pone.0020615> PMID:21687731
24. Banas A, Teratani T, Yamamoto Y, Tokuhara M, Takeshita F. Rapid hepatic fate specification of adipose-derived stem cells and their therapeutic potential for liver failure. *J Gastroenterol Hepatol*. 2009;24(1):70-7. <https://doi.org/10.1111/j.1440-1746.2008.05496.x> PMID:18624899
25. De Ugarte DA, Morizono K, Elbarbary A, Alfonso Z, Zuk PA, Zhu M, et al. Comparison of multi-lineage cells from human adipose tissue and bone marrow. *Cells Tissues Organs*. 2003;174(3):101-9. <https://doi.org/10.1159/000071150> PMID:12835573
26. Kern S, Eichler H, Stoeve J, Klüter H, Bieback K. Comparative analysis of mesenchymal stem cells from bone marrow, umbilical cord blood, or adipose tissue. *Stem Cells*. 2006;24(5):1294-301. <https://doi.org/10.1634/stemcells.2005-0342> PMID:16410387
27. Wagner W, Wein F, Seckinger A, Frankhauser M, Wirkner U, Krause U, et al. Comparative characteristics of mesenchymal stem cells from human bone marrow, adipose tissue, and umbilical cord blood. *Exp Hematol*. 2005;33(11):1402-16. <https://doi.org/10.1016/j.exphem.2005.07.003> PMID:16263424
28. Heo JS, Choi Y, Kim HS, Kim HO. Comparison of molecular profiles of human mesenchymal stem cells derived from bone marrow, umbilical cord blood, placenta and adipose tissue. *Int J Mol Med*. 2016;37(1):115-25. <https://doi.org/10.3892/ijmm.2015.2413> PMID:26719857
29. Carbone A, Rucci M, Luigi A, Portincasa A, Conese M. Adipose-derived stem cells and platelet-rich plasma: Inputs for regenerative medicine. *Med Res Arch*. 2017;5(10):1-12.
30. Oberbauer E, Steffenhagen C, Wurzer C, Gabriel C, Redl H, Wolbank S. Enzymatic and non-enzymatic isolation systems for adipose tissue-derived cells: Current state of the art. *Cell Regen*. 2015;4:7. <https://doi.org/10.1186/s13619-015-0020-0> PMID:26435835
31. Baldari S, di Rocco G, Piccoli M, Muraca M, Toietta G. Challenges and strategies for improving the regenerative effects of mesenchymal stromal cell-based therapies. *Int J Mol Sci*. 2017;18(10):2087. <https://doi.org/10.3390/ijms18102087> PMID:28974046
32. Doorn J, Moll G, Le Blanc K, van Blitterswijk C, de Boer J. Therapeutic applications of mesenchymal stromal cells: Paracrine effects and potential improvements. *Tissue Eng Part B Rev*. 2012;18(2):101-15. <https://doi.org/10.1089/ten.teb.2011.0488> PMID:21995703
33. Wang X, Liu C, Li S, Xu Y, Chen P, Liu Y, et al. Hypoxia precondition promotes adipose-derived mesenchymal stem cells based repair of diabetic erectile dysfunction via augmenting angiogenesis and neuroprotection. *PLoS One*. 2015;10(3):e0118951. <https://doi.org/10.1371/journal.pone.0118951> PMID:25790284
34. Sart S, Ma T, Li Y. Preconditioning stem cells for *in vivo* delivery. *Biores Open Access*. 2014;3(4):137-49. PMID:25126478
35. Guo M, Cai C, Zhao G, Qiu X, Zhao H, Ma Q, et al. Hypoxia promotes migration and induces CXCR4 expression via HIF-1 α activation in human osteosarcoma. *PLoS One*. 2014;9(3):e90518. <https://doi.org/10.1371/journal.pone.0090518> PMID:24618817

36. Wei L, Fraser JL, Lu Z, Hu X, Ping S. transplantation of hypoxia preconditioned bone marrow mesenchymal stem cells enhances angiogenesis and neurogenesis after cerebral ischemia in rats. *Neurobiol Dis.* 2012;46(3):635-45. <https://doi.org/10.1016/j.nbd.2012.03.002>
PMid:22426403
37. Lennon DP, Edmison JM, Caplan AI. Cultivation of rat marrow-derived mesenchymal stem cells in reduced oxygen tension: Effects on *in vitro* and *in vivo* osteochondrogenesis. *J Cell Physiol.* 2001;187(3):345-55. <https://doi.org/10.1002/jcp.1081>
PMid:11319758
38. Geng YJ. Molecular mechanisms for cardiovascular stem cell apoptosis and growth in the hearts with atherosclerotic coronary disease and ischemic heart failure. *Ann NY Acad Sci.* 2003;1010:687-97. <https://doi.org/10.1196/annals.1299.126>
PMid:15033813
39. Greijer AE, van der Wall E. The role of hypoxia inducible factor 1 (HIF-1) in hypoxia induced apoptosis. *J Clin Pathol.* 2004;57(10):1009-14. <https://doi.org/10.1136/jcp.2003.015032>
PMid:15452150
40. Martin AR, Ronco C, Demange L, Benhida R. Hypoxia inducible factor down-regulation, cancer and cancer stem cells (CSCs): Ongoing success stories. *Medchemcomm.* 2017;8(1):21-52. <https://doi.org/10.1039/c6md00432f>
PMid:30108689
41. Sun X, Fang B, Zhao X, Zhang G, Ma H. Preconditioning of mesenchymal stem cells by sevoflurane to improve their therapeutic potential. *PLoS One.* 2014;9(3):e90667. <https://doi.org/10.1371/journal.pone.0090667>
PMid:24599264
42. Semenza GL. Targeting HIF-1 for cancer therapy. *Nat Rev Cancer.* 2003;3(10):721-32.
PMid:13130303
43. Romain B, Hachet-Haas M, Rohr S, Brigand C, Galzi JL, Gaub MP, *et al.* Hypoxia differentially regulated CXCR4 and CXCR7 signaling in colon cancer. *Mol Cancer.* 2014;13:58. <https://doi.org/10.1186/1476-4598-13-58>
PMid:24629239
44. Abdolmohammadi K, Mahmoudi T, Nojehdehi S, Tayebi L, Hashemi SM, Noorbakhsh F, *et al.* Effect of hypoxia preconditioned adipose-derived mesenchymal stem cell conditioned medium on cerulein-induced acute pancreatitis in mice. *Adv Pharm Bull.* 2020;10(2):297-306. <https://doi.org/10.34172/apb.2020.036>
PMid:32373500
45. Xue Y, Li Z, Wang Y, Zhu X, Hu R, Xu W. Role of the HIF-1 α /SDF1/CXCR4 signaling axis in accelerated fracture healing after craniocerebral injury. *Mol Med Rep.* 2020;22(16):2767-74. <https://doi.org/10.3892/mmr.2020.11361>
PMid:32945380
46. Lee JW, Ko J. Hypoxia signaling in human diseases and therapeutic targets. *Exp Mol Med.* 2019;51(6):1-13. <https://doi.org/10.1128/mcb.00409-10>
PMid:31221962
47. Culver C, Sundqvist A, Mudie S, Melvin A, Xirodimas D, Rocha S. Mechanism of hypoxia-induced NF-kappa B. *Mol Cell Biol.* 2010;30(20):4901-21.
PMid:20696840
48. Hu C, Wang L, Chodosh LA, Keith B, Simon MC. Differential roles of hypoxia-inducible factor 1 alpha (HIF-1alpha) and HIF-2alpha in hypoxic gene regulation. *Mol Cell Biol.* 2003;23(24):9361-74. <https://doi.org/10.1128/mcb.23.24.9361-9374.2003>
PMid:14645546
49. Kang S, Kim S, Sung J. Cellular and molecular stimulation of adipose-derived stem cells under hypoxia. *Cell Biol Int.* 2014;38(5):553-62. <https://doi.org/10.1002/cbin.10246>
PMid:24446066
50. Dreia K, Sarnowska A, Siedlecka P, Szablowska-Gadomska I, Wielgos M, Jurga M, *et al.* Low oxygen atmosphere facilitates proliferation and maintains undifferentiated state of umbilical cord mesenchymal stem cells in an hypoxia inducible factor-dependent manner. *Cytotherapy.* 2014;16(7):881-92. <https://doi.org/10.1016/j.jcyt.2014.02.009>
PMid:24726658
51. Portalska KJ, Leferink A, Groen N, Fernandes H, Moroni L, van Blitterswijk C, *et al.* Endothelial differentiation of mesenchymal stromal cells. *PLoS One.* 2012;7(10):e46842. <https://doi.org/10.1371/journal.pone.0046842>
PMid:23056481
52. Chang W, Lee CY, Park JH, Park MS, Maeng LS, Yoon CS, *et al.* Survival of hypoxic human mesenchymal stem cells is enhanced by a positive feedback loop involving miR-210 and hypoxia-induced factor 1. *J Vet Sci.* 2013;14(1):69-76. <https://doi.org/10.4142/jvs.2013.14.1.69>
PMid:23388440
53. Kim JH, Park SG, Song S, Kim JK, Sung J. Reactive oxygen species-responsive miR-210 regulates proliferation and migration of adipose-derived stem cells via PTPN2. *Cell Death Dis.* 2013;4(4):e588. <https://doi.org/10.1038/cddis.2013.117>
PMid:23579275
54. Kim JH, Park S, Park SG, Choi J, Xia Y, Sung J. The pivotal role of reactive oxygen species generation in the hypoxia-induced stimulation of adipose-derived stem cells. *Stem Cells Dev.* 2011;20(10):1753-61. <https://doi.org/10.1089/scd.2010.0469>
PMid:21265612
55. Weijers EM, van den Broek LJ, Waaijman T, van Hinsbergh VW, Gibbs S, Koolwijk P. The influence of hypoxia and fibrinogen variants on the expansion and differentiation of adipose tissue-derived mesenchymal stem cells. *Tissue Eng Part A.* 2011;17(21-22):2675-85. <https://doi.org/10.1089/ten.tea.2010.0661>
PMid:21830936
56. Yang J, Zhang H, Zhao L, Chen Y, Liu H, Zhang T. Human adipose tissue-derived stem cells protect impaired cardiomyocytes from hypoxia/reoxygenation injury through hypoxia-induced paracrine mechanism. *Cell Biochem Funct.* 2012;30(6):505-14. <https://doi.org/10.1002/cbf.2829>
PMid:22610511
57. Kim JH, Song S, Park SG, Song SU, Xia Y, Sung J. Primary involvement of NADPH oxidase 4 in hypoxia-induced generation of reactive oxygen species in adipose-derived stem cells. *Stem Cells Dev.* 2012;21(12):2212-21. <https://doi.org/10.1089/scd.2011.0561>
PMid:22181007
58. Ceradini DJ, Kulkarni AR, Callaghan MJ, Tepper OM, Bastidas N, Kleinman ME, *et al.* Progenitor cell trafficking is regulated by hypoxic gradients through HIF-1 induction of SDF-1. *Nat Med.* 2004;10(8):858-64. <https://doi.org/10.1038/nm1075>
PMid:15235597
59. Fotia C, Massa A, Boriani F, Baldini N. Hypoxia enhances proliferation and stemness of human adipose-derived mesenchymal stem cells. *Cytotechnology.* 2015;67(6):1073-84. <https://doi.org/10.1007/s10616-014-9731-2>
PMid:24798810
60. Bruno S, Grange C, Collino F, Deregibus MC, Cantaluppi V, Biancone L, *et al.* Microvesicles derived from mesenchymal stem cells enhance survival in a lethal model of acute kidney injury. *PLoS One.* 2012;7(3):e33115. <https://doi.org/10.1371/journal.pone.0033115>
PMid:22431999

-
61. Chistiakov DA, Chekhonin VP. Extracellular vesicles shed by glioma cells: Pathogenic role and clinical value. *Tumour Biol.* 2014;35(9):8425-38. <https://doi.org/10.1007/s13277-014-2262-9>
PMid:24969563
62. Quesenberry PJ, Goldberg LR, Aliotta JM, Dooner MS, Pereira MG, Wen S, et al. Cellular phenotype and extracellular vesicles: Basic and clinical considerations. *Stem Cells Dev.* 2014;23(13):1429-36. <https://doi.org/10.1089/scd.2013.0594>
PMid:24564699
63. Beegle J, Lakatos K, Kalomoiris S, Stewart H, Isseroff RR, Nolte JA, et al. Hypoxic preconditioning of mesenchymal stromal cells induces metabolic changes, enhances survival, and promotes cell retention *in vivo*. *Stem Cells.* 2015;33:1818-28. <https://doi.org/10.1002/stem.1976>
PMid:25702874
64. Stubbs SL, Hsiao ST, Peshavariya HM, Lim SY, Disting GJ, Dilley RJ. Hypoxic preconditioning enhances survival of human adipose-derived stem cells and conditions endothelial cells *in vitro*. *Stem Cells Dev.* 2012;21(11):1887-96. <https://doi.org/10.1089/scd.2011.0289>
PMid:22165914
65. Zhang W, Liu L, Huo Y, Yang Y, Wang Y. Hypoxia-pretreated human MSCs attenuate acute kidney injury through enhanced angiogenic and antioxidative capacities. *Biomed Res Int.* 2014;2014:462472. <https://doi.org/10.1155/2014/462472>
PMid:25133162
66. Wang L, Sanford MT, Xin Z, Lin G, Lue TF. Role of schwann cells in the regeneration of penile and peripheral nerves. *Asian J Androl.* 2015;17(5):776-82. <https://doi.org/10.4103/1008-682x.154306>
PMid:25999359

Preconditioning of Hypoxic Culture Increases The Therapeutic Potential of Adipose Derived Mesenchymal Stem Cells

ORIGINALITY REPORT

18%

SIMILARITY INDEX

13%

INTERNET SOURCES

12%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1	ijop.net Internet Source	1%
2	theoncologist.alphamedpress.org Internet Source	1%
3	www.journal.ugm.ac.id Internet Source	1%
4	johepal.com Internet Source	1%
5	www.biopreparations.ru Internet Source	1%
6	www.floegel.info Internet Source	1%
7	www.jove.com Internet Source	1%
8	academic-accelerator.com Internet Source	<1%
9	jcp.bmjournals.com Internet Source	<1%

10

Johanna Myllyharju. "Prolyl 4-hydroxylases, key enzymes in the synthesis of collagens and regulation of the response to hypoxia, and their roles as treatment targets", *Annals of Medicine*, 2009

Publication

<1 %

11

Ning Zhang, JinYun Zhu, QunChao Ma, Yun Zhao, YingChao Wang, XinYang Hu, JingHai Chen, Wei Zhu, ZhongChao Han, Hong Yu. "Exosomes derived from human umbilical cord MSCs rejuvenate aged MSCs and enhance their functions for myocardial repair", *Research Square Platform LLC*, 2020

Publication

<1 %

12

jeccr.biomedcentral.com

Internet Source

<1 %

13

discovery.dundee.ac.uk

Internet Source

<1 %

14

repositorio.utad.pt

Internet Source

<1 %

15

Sang eon Park, Hyeong Seop Kim, Soo Jin Kwon, Min-Jeong Kim et al. "Exposure of Mesenchymal Stem Cells to an Alzheimer's Disease Environment Enhances Therapeutic Effects", *Research Square Platform LLC*, 2020

Publication

<1 %

16 So Yoon Ahn, Yun Sil Chang, Won Soon Park. "Stem cells in the treatment of bronchopulmonary dysplasia", Elsevier BV, 2020
Publication <1 %

17 rjptonline.org
Internet Source <1 %

18 www.coursehero.com
Internet Source <1 %

19 advances.umw.edu.pl
Internet Source <1 %

20 docplayer.net
Internet Source <1 %

21 stemcellpowernow.com
Internet Source <1 %

22 www.sciencedaily.com
Internet Source <1 %

23 Dorota Kaniowska, Kerstin Wenk, Phil Rademacher, Ronald Weiß et al. "Extracellular Vesicles of Mesenchymal Stromal Cells Can be Taken up by Microglial Cells and Partially Prevent The Stimulation Induced by β -Amyloid", Research Square Platform LLC, 2021
Publication <1 %

24 Md. Prova Zaman Emon, Rajesh Das, Nuruna Lovely Nishuty, M.M.A. Shalahuddin Qusar, <1 %

Mohiuddin Ahmed Bhuiyan, Md. Rabiul Islam. "Reduced serum BDNF levels are associated with the increased risk for developing MDD: A case-control study with or without antidepressants therapy", Research Square Platform LLC, 2019

Publication

25

fjfsdata01prod.blob.core.windows.net

Internet Source

<1 %

26

Haiying Ma, Shenglu Jiang, Lili Du, Jinfang Liu, Xiaoyan Xu, Xiaomei Lu, Ling Ma, Hua Zhu, Jun Wei, Yanqiu YU. "Conditioned medium from primary cytotrophoblasts, primary placenta-derived mesenchymal stem cells, or placental tissue promoted HUVECs angiogenesis in vitro", Research Square Platform LLC, 2020

Publication

<1 %

27

Zhilai, Zhou, Mo Biling, Qiu Sujun, Dong Chao, Shi Benchao, Huang Shuai, Yao Shun, and Zhang Hui. "Preconditioning in lowered oxygen enhances the therapeutic potential of human umbilical mesenchymal stem cells in a rat model of spinal cord injury", Brain Research, 2016.

Publication

<1 %

28

Jong-Hyuk Sung, Hyo-Sun An, Jin-Hyun Jeong, Soyoung Shin, Seung Yong Song. "Megestrol Acetate Increases the Proliferation, Migration,

<1 %

and Adipogenic Differentiation of Adipose-Derived Stem Cells via Glucocorticoid Receptor", STEM CELLS Translational Medicine, 2015

Publication

29

Joyce Doorn. "Therapeutic Applications of Mesenchymal Stromal Cells: Paracrine Effects and Potential Improvements", Tissue Engineering Part B Reviews, 12/22/2011

Publication

<1 %

30

journal.frontiersin.org

Internet Source

<1 %

31

ki-it.com

Internet Source

<1 %

32

lib.bioinfo.pl

Internet Source

<1 %

33

mdpi-res.com

Internet Source

<1 %

34

stemcellsjournals.onlinelibrary.wiley.com

Internet Source

<1 %

35

www.fbm.msu.ru

Internet Source

<1 %

36

Barbuti, Peter A.. "Stem Cell Therapy for Neurological Disorders: From Bench to Bedside", Stem Cell Therapy for Organ Failure, 2014.

<1 %

37

Dae-Kwan Yoon, Ji-Sung Park, Gyu-Jin Rho, Hyeon-Jeong Lee et al. "The involvement of histone methylation in osteoblastic differentiation of human periosteum-derived cells cultured in vitro under hypoxic conditions", *Cell Biochemistry and Function*, 2017

Publication

<1 %

38

Jeong, Yun-Mi, Young Kwan Sung, Wang-Kyun Kim, Ji Hye Kim, Mi Hee Kwack, Insoo Yoon, Dae-Duk Kim, and Jong-Hyuk Sung. "Ultraviolet B Preconditioning Enhances the Hair Growth-Promoting Effects of Adipose-Derived Stem Cells Via Generation of Reactive Oxygen Species", *Stem Cells and Development*, 2012.

Publication

<1 %

39

Lei Cui. "Expanded Adipose-Derived Stem Cells Suppress Mixed Lymphocyte Reaction by Secretion of Prostaglandin E2", *Tissue Engineering*, 06/2007

Publication

<1 %

40

M Dominici. "Minimal criteria for defining multipotent mesenchymal stromal cells. The International Society for Cellular Therapy position statement", *Cytotherapy*, 8/1/2006

Publication

<1 %

41 RJW Hoogendoorn. "Adipose stem cells for intervertebral disc regeneration: current status and concepts for the future.", *Journal of Cellular and Molecular Medicine*, 2/25/2008
Publication <1 %

42 Samyra El-ftesi. "Aging and Diabetes Impair the Neovascular Potential of Adipose-Derived Stromal Cells", *Plastic and Reconstructive Surgery*, 02/2009
Publication <1 %

43 Tang, Y.L.. "Improved Graft Mesenchymal Stem Cell Survival in Ischemic Heart With a Hypoxia-Regulated Heme Oxygenase-1 Vector", *Journal of the American College of Cardiology*, 20051004
Publication <1 %

44 Werle, Stefanie Bressan, Pedro Chagastelles, Patricia Pranke, and Luciano Casagrande. "The effects of hypoxia on in vitro culture of dental-derived stem cells", *Archives of Oral Biology*, 2016.
Publication <1 %

45 [Www.ijbs.com](http://www.ijbs.com)
Internet Source <1 %

46 Yan Wang, Ying Xie, David Oupický. "Potential of CXCR4/CXCL12 Chemokine Axis in Cancer <1 %

Drug Delivery", Current Pharmacology Reports, 2016

Publication

47

coek.info

Internet Source

<1 %

48

tampub.uta.fi

Internet Source

<1 %

49

www-bioeng.ucsd.edu

Internet Source

<1 %

50

Patrick Wahl, Klara Brixius, Wilhelm Bloch. "Exercise - induced stem cell activation and its implication for cardiovascular and skeletal muscle regeneration", Minimally Invasive Therapy & Allied Technologies, 2009

Publication

<1 %

51

Sarah T. Hsiao, Rodney J. Dilley, Gregory J. Dusting, Shiang Y. Lim. "Ischemic preconditioning for cell-based therapy and tissue engineering", Pharmacology & Therapeutics, 2014

Publication

<1 %

52

Sarah Walsh. "Practical considerations concerning the use of stem cells for peripheral nerve repair", Neurosurgical FOCUS, 02/2009

Publication

<1 %

53	Shih-Chieh Hung. "Short-Term Exposure of Multipotent Stromal Cells to Low Oxygen Increases Their Expression of CX3CR1 and CXCR4 and Their Engraftment In Vivo", PLoS ONE, 05/02/2007 Publication	<1 %
54	cancerres.aacrjournals.org Internet Source	<1 %
55	eprints.qut.edu.au Internet Source	<1 %
56	iris.unige.it Internet Source	<1 %
57	pdffox.com Internet Source	<1 %
58	pure.uva.nl Internet Source	<1 %
59	www.aging-us.com Internet Source	<1 %
60	www.hal.inserm.fr Internet Source	<1 %
61	www.Incrnablog.com Internet Source	<1 %
62	A J Giaccia. "The role of hypoxia-inducible factors in tumorigenesis", Cell Death and Differentiation, 04/2008	<1 %

63

Anastasia Efimenko. "Angiogenic properties of aged adipose derived mesenchymal stem cells after hypoxic conditioning", Journal of Translational Medicine, 2011

Publication

<1 %

64

Maša Čater, Gregor Majdič. "Chapter 656 In Vitro Culturing of Adult Stem Cells: The Importance of Serum and Atmospheric Oxygen", Springer Science and Business Media LLC, 2021

Publication

<1 %

65

Michelle E. McCarthy, Theodore A. Brown, Joanna Bukowska, Bruce A. Bunnell, Trivia Frazier, Xiying Wu, Jeffrey M. Gimble. "Therapeutic Applications for Adipose-Derived Stem Cells in Wound Healing and Tissue Engineering", Current Stem Cell Reports, 2018

Publication

<1 %

66

Caterina Fotia, Annamaria Massa, Filippo Boriani, Nicola Baldini, Donatella Granchi. "Hypoxia enhances proliferation and stemness of human adipose-derived mesenchymal stem cells", Cytotechnology, 2014

Publication

<1 %

67

Deborah Schmitz, Jan W. Robering, Volker Weisbach, Andreas Arkudas et al. "Specific

<1 %

features of ex - obese patients significantly influence the functional cell properties of adipose - derived stromal cells", Journal of Cellular and Molecular Medicine, 2022

Publication

68

Kingham, P.J.. "Adipose-derived stem cells differentiate into a Schwann cell phenotype and promote neurite outgrowth in vitro", Experimental Neurology, 200710

Publication

<1 %

69

Ritbano Ahmed Abdo, Hassen Mosa Halil, Biruk Assefa Kebede, Abebe Alemu Anshebo, Negeso Gebeyehu Gejo. "Birth asphyxia and its contributing factors among the neonates delivered at Negest Eleni Mohammed memorial teaching hospital, Southern Ethiopia: A cross-sectional study", Research Square Platform LLC, 2019

Publication

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

Preconditioning of Hypoxic Culture Increases The Therapeutic Potential of Adipose Derived Mesenchymal Stem Cells

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10

PAGE 11
