

Volume 11 // Number 03 // March 2020

# Indian Journal of Public Health Research & Development

An International Journal

Website:

www.ijphrd.com

### Indian Journal of Public Health Research & Development

#### **EXECUTIVE EDITOR**

#### Prof. Vidya Surwade

Deptt. of Community Medicine, Dr Baba Saheb Ambedkar, Medical College & Hospital, Rohini, Delhi

#### INTERNATIONAL EDITORIAL ADVISORY BOARD

- Dr. Abdul Rashid Khan B. Md Jagar Din, (Associate Professor)
   Department of Public Health Medicine, Penang Medical College, Penang, Malaysia
- 2. Dr. V Kumar (Consulting Physician)
  Mount View Hospital, Las Vegas, USA
- 3. Basheer A. Al-Sum,

Botany and Microbiology Deptt, College of Science, King Saud University, Rivadh, Saudi Arabia

- Dr. Ch Vijay Kumar (Associate Professor)
   Public Health and Community Medicine, University of Buraimi, Oman
- Dr. VMC Ramaswamy (Senior Lecturer)
   Department of Pathology, International Medical University, Bukit Jalil, Kuala Lumpur
- Kartavya J. Vyas (Clinical Researcher)
   Department of Deployment Health Research,
   Naval Health Research Center, San Diego, CA (USA)
- Prof. PK Pokharel (Community Medicine)
   BP Koirala Institute of Health Sciences, Nepal

#### NATIONAL SCIENTIFIC COMMITTEE

- Dr. Anju D Ade (Professor)
   Community Medicine Department, SVIMS, Sri Padamavati Medical College, Tirupati,
   Andhra Pradech
- Dr. E. Venkata Rao (Associate Professor) Community Medicine, Institute of Medical Sciences & SUM Hospital, Bhubaneswar, Orissa.
- Dr. Amit K. Singh (Associate Professor) Community Medicine, VCSG Govt. Medical College, Srinagar – Garhwal, Uttarakhand
- Dr. R G Viveki (Associate Professor) Community Medicine, Belgaum Institute of Medical Sciences, Belgaum, Karnataka
- Dr. Santosh Kumar Mulage (Assistant Professor)
   Anatomy, Raichur Institute of Medical Sciences Raichur(RIMS), Karnataka
- Dr. Gouri Ku. Padhy (Associate Professor) Community and Family Medicine, All India Institute of Medical Sciences, Raipur
- Dr. Ritu Goyal (Associate Professor)
   Anaesthesia, Sarswathi Institute of Medical Sciences, Panchsheel Nagar
- 8. **Dr. Anand Kalaskar** (Associate Professor)
  Microbiology, Prathima Institute of Medical Sciences, AP
- Dr. Md. Amirul Hassan (Associate Professor)
   Community Medicine, Government Medical College, Ambedkar Nagar, UP
- 10. Dr. N. Girish (Associate Professor) Microbiology, VIMS&RC, Bangalore
- 11. Dr. BR Hungund (Associate Professor) Pathology, JNMC, Belgaum.
- 12. Dr Sartaj Ahmad, PhD Medical Sociology, Associate Professor, Swami Vivekananda Subharti University Meerut UP India
- Dr Sumeeta Soni (Associate Professor)
   Microbiology Department, B.J. Medical College, Ahmedabad, Gujarat, India

#### NATIONAL EDITORIAL ADVISORY BOARD

- Prof. Sushanta Kumar Mishra (Community Medicine) GSL Medical College – Rajahmundry, Karnataka
- Prof. D.K. Srivastava (Medical Biochemistry)
   Jamia Hamdard Medical College, New Delhi
- Prof. M Sriharibabu (General Medicine) GSL Medical College, Rajahmundry, Andhra Pradesh
- Prof. Pankaj Datta (Principal & Prosthodentist) Indraprastha Dental College, Ghaziabad

#### NATIONAL EDITORIAL ADVISORY BOARD

- Prof. Samarendra Mahapatro (Pediatrician)
   Hi-Tech Medical College, Bhubaneswar, Orissa
- Dr. Abhiruchi Galhotra (Additional Professor) Community and Family Medicine, All India Institute of Medical Sciences, Raipur
- Prof. Deepti Pruthvi (Pathologist) SS Institute of Medical Sciences & Research Center. Davangere. Karnataka
- 8. Prof. G S Meena (Director Professor)
  Maulana Azad Medical College, New Delhi
- Prof. Pradeep Khanna (Community Medicine)
   Post Graduate Institute of Medical Sciences, Rohtak, Haryana
- Dr. Sunil Mehra (Paediatrician & Executive Director)
   MAMTA Health Institute of Mother & Child. New Delhi
- Dr Shailendra Handu, Associate Professor, Phrma, DM (Pharma, PGI Chandigarh)
- Dr. A.C. Dhariwal: Directorate of National Vector Borne Disease Control Programme, Dte. DGHS, Ministry of Health Services, Govt. of India, Delhi

**Print-ISSN**: 0976-0245-Electronic-ISSN: 0976-5506, Frequency: Quarterly (Four issues per volume)

Indian Journal of Public Health Research & Development is a double blind peer reviewed international journal. It deals with all aspects of Public Health including Community Medicine, Public Health, Epidemiology, Occupational Health, Environmental Hazards, Clinical Research, and Public Health Laws and covers all medical specialties concerned with research and development for the masses. The journal strongly encourages reports of research carried out within Indian continent and South East Asia.

The journal has been assigned International Standards Serial Number (ISSN) and is indexed with Index Copernicus (Poland). It is also brought to notice that the journal is being covered by many international databases. The journal is covered by EBSCO (USA), Embase, EMCare. The journal is now part of DST, CSIR, and UGC consortia.

#### Website: www.ijphrd.com

©All right reserved. The views and opinions expressed are of the authors and not of the Indian Journal of Public Health Research & Development. The journal does not guarantee directly or indirectly the quality or efficacy of any product or service featured in the advertisement in the journal, which are purely commercial.

#### Editor

Dr. R.K. Sharma

Institute of Medico-legal Publications Logix Office Tower, Unit No. 1704, Logix City Centre Mall, Sector- 32, Noida - 201 301 (Uttar Pradesh)

#### Printed, published and owned by

Dr. R.K. Sharma

Institute of Medico-legal Publications Logix Office Tower, Unit No. 1704, Logix City Centre Mall, Sector- 32, Noida - 201 301 (Uttar Pradesh)

#### Published at

#### Institute of Medico-legal Publications

Logix Office Tower, Unit No. 1704, Logix City Centre Mall, Sector- 32, Noida - 201 301 (Uttar Pradesh)



# Indian Journal of Public Health Research & Development

www.ijphrd.com

	Contents	
Vol	ume 11, March 03 March 2	2020
1.	Development and Validation of a Questionnaire on Eating Behaviour for School Children and its Correlation with Nutritional Status	1
2.	Occupational Varicella outbreak at a Tertiary Care Hospital: An Insight	7
3.	An Exploratory Study To Identify Factors Affecting Non Compliance To Dots Therapy Among Tb Patients At Selected Dots Centre Vadodara	13
4.	Assessment of Cognitive Impairment among Elderly in the Selected Rural Community, Kancheepuram District, Tamil Nadu	19
5.	Safety and Tolerability of Two Different Formulations of Mycophenolate (Mycophenolate Mofetil and Mycophenolate Sodium) among Patient with Connective Tissue Disease Associated Interstitial Lung Disease (CTD-ILD) in a Tertiary Care Hospital	22
6.	Assessment of Awareness of Parents on Importance of Dental Care in Pediatric Patients in Ethnic Tamil Population	27
7.	Study of Total Time Taken for OPD Billing Process in a Multi-Specialty Hospital	32
8.	Sleep Quality and Glycemic Control among Patients with Type II Diabetes Mellitus	41
9.	A Comprehensive Break Even Analysis of MRI and CT Unit of a Tertiary Care Hospital in Sikkim	47
10.	A Descriptive Survey on the Consumption of Sweetened Beverages and Contributing Factors among Adolescents in Selected PU Colleges at Mangaluru	53

381.	To Evaluate the Correlation of Stature with Hand & Foot Width in Subjects	.2074
382.	Health Risk Assessment on the Glyphosate Exposure of Knapsack Sprayers	.2077
383.	To Assess Correlation of Stature with Hand & Foot Length	.2083
384.	Teenage Girl Dating Experience: A Qualitative Study	.2086
385.	Diet Compliance Against Blood Pressure Control in Hypertensive Patients in Tegalgundil public Health Center, North Bogor District: A Longitudinal Study	.2091
386.	A Case Study of the Impact of Overweight on Body Image and Self-Esteem in a Population of Moroccan Adolescents and Adults	.2097
387.	Analysis Factors of Nursing Performance at the Mother and Child Hospital in East Java Lembah Andriani Sulistyaning Dyah, Margono Setiawan, Dodi W. Irawanto, Fatchur Rohman, Nursalam Nursalam	.2103
388.	Effectivity of Teacher Motivation on Dental and Oral Hygiene of Elementary School Students	.2109
389.	Study Some of the Factors Affecting the Incidence of Diabetes in the Employed Segment in Basra City Maryam J. Almusawi, Murtadha Gh. Jasim	.2115
390.	Polyunsaturated Fatty Acid Intake and Symptom Severity of Patients with Schizophrenia in Ernaldi Bahar Hospital, South Sumatra, Indonesia	.2119
391.	A Study of Cyber Bullying Behavior in Middle-Schools in Rabat- Morocco: Prevalence and Risk Factors	.2125
392.	The Correlation of Dental Caries with Parents Education Science Level at Ar-Rahmah Islamic Elementary School Tamalanrea Makassar, South Sulawesi	.2131
393.	Prevalence of Diabetes Mellitus among Stroke patients in King Fahad Specialist Hospital, Buraidah, 2018	.2137
394.	Risk Factors and the Pattern of Injuries of Road Traffic Accidents in Holy City of Karbala/Iraq	.2143

## XXX

395.	Alpha-Fetoprotein for Prediction of Placenta Accreta in Women with Complete Placenta Previa Centeralis: A Prospective Study	2148
	Mohammed Hany Mosbeh, Mohammed Abdallah Mohammed, Mo'men Mohamed Hassan, Ahmed Rabie Abd El-Raheim, Heba Reda Mohammed	
396.	Health Promoting School in Surabaya, Indonesia: The Six Elements Implementation	2154
397.	Evaluation Context and Input of National Health Insurance in Ternate City	2159
398.	The Effectiveness Comparison of Type of Treatments in Decreasing of Total Dissolved Solid (TDS) and Total Suspended Solid (TSS) in Household Wastewater	2164
399.	Environmental Health Risk Analysis of Carbon Monoxide Exposure among High Activity Communities Along "X" Street, Yogyakarta	2170
400.	The Effect of Work Satisfaction on the Quality of Health Services (Literature Review)	2176
401.	Respiratory Symptoms of Housewives Exposed to $SO_2$ From Steel Industry in West Cikarang, Indones Nada Amirah, Agustin Kusumayati, Suyud	ia2181
402.	Efficacy of Endoscopic Transantral Versus Transorbital Surgical Approaches in the Repair of Orbital Blow-Out Fractures (Randomized Clinical Trial)	2187
403.	Relationships of Workloads, Working Conditions and Dual Role Conflict with Nursing Stress	2193
404.	Intention of Diabetic Foot Ulcer Prevention Model Based on Social Support and Personal Agency Perspectives	2198
405.	The Development of Diabetic Foot Ulcer Prevention Model Based on Psychosocial Perspectives, Attitude, Intention, Coping Mechanisms	2204
406.	Association of Syphilis and HIV among Indirect Female Sex Worker in Indonesia: Secondary Data Analysis of Integrated Behaviour Biological Survey in 2015	2211
407.	The Knowledge of the Use of the Contraceptive Method among Married Men with Fertility Age 15-54 Years (Analysis of IDHS 2017 Data)	2216
408.	Preliminary Study: Reliability and Validity of CFM-1 Form as Physical Literacy Assessment Instrument  Nur Ikhwan Mohamad, Muhammad Hannan Sazali, Ali Md Nadzalan, Asmadi Ishak, Hishamuddin	2220

# Intention of Diabetic Foot Ulcer Prevention Model Based on Social Support and Personal Agency Perspectives

NasrunPakaya<sup>1,2</sup>, Kusnanto<sup>3</sup>, Hari Basuki Notobroto<sup>4</sup>, Rika Subarniati Triyoga<sup>5</sup>

<sup>1</sup>Student of Doctoral Program of Public Health, Faculty of Public Health, Universitas Airlangga, Mulyorejo Street, Surabaya, Indonesia, <sup>2</sup>Lecturer of Faculty Of Sport and Health, State University of Gorontalo, Indonesia, <sup>3</sup>Lecturer of Faculty of NursingUniversitas Airlangga, Mulyorejo Street, Surabaya, Indonesia, <sup>4</sup>Lecturer of Faculty of Public Health, Universitas Airlangga, Mulyorejo Street, Surabaya, Indonesia, <sup>5</sup>Lecturer of Faculty of Public Health, Universitas Airlangga, Mulyorejo Street, Surabaya, Indonesia

#### **Abstract**

Context: A diabetic foot ulcer is one of the complications occurred on diabetes mellitus patients. The incident rate of diabetic foot ulcer improves each year significantly. The research was to arrange the intention of diabetic foot ulcer prevention model based on social support and personal agency perspectives. This study was included in a cross-sectional study using a questionnaire and a simple random sampling technique on 10 health centers (puskesmas). The research instrument for social support consisted of family's support and friend's support, personal agency consisted of perceived control and self-efficacy, and intention consisted of diet intention, consuming medicine intention, physical activity intention, and foot/blood sugar controlling intention. The data analysis applied SEM-PLS software. The diabetes mellitus patients who did not have ulcer were 329. There was correlation between social support and personal agency of diabetic foot ulcer prevention by having Coefficient value for 0.68, and T value for 16.27, there was correlation between personal agency and intention to prevent the diabetic foot ulcer by having coefficient value for 0,57 and T value for 2.96, and there was correlation between social support and intention to prevent the diabetic foot ulcer by having coefficient value for -0,27 and T value for 2,08. The social support contributed highly to the intention through a personal agency, and the social support contributed directly to intention in preventing the diabetic foot ulcer. It is suggested that diabetes patients should get supports from family and friends to improve the perceived control and self-efficacy hence the intention of diabetic foot ulcer prevention can be improved.

Keywords: Social Support, Personal Agency, Intention, Diabetes

#### Introduction

A diabetic foot ulcer is one of the dead causes in the world, and it can attack whoever the individual is. Diabetes triggers morbidity such as blindness, kidney failure, and non-traumatic amputations<sup>(1)</sup>WHO predicts that the increase of Diabetes Mellitus sufferers in Indonesia reached 8,4 million in 2000 and will be about 21,3 million in 2030<sup>(2)</sup>. In 1990, Indonesia was in the 16<sup>th</sup> place for diabetes, while it was ranked 6<sup>th</sup> in 2010 and changed to the 5<sup>th</sup> place in 2015.

Amputation is done every day for diabetes mellitus patients in the world <sup>(3)</sup> Data from <sup>(4)</sup>. Riskesdas revealed that Indonesia was ranked 10<sup>th</sup> for the world's highest

foot amputation number. Besides treatment and healthy lifestyle, the patients' behavior is one of the determining factors of the success in preventing the diabetic foot ulcer so that it can decrease the amputation incident rate. The effect of amputation that is occurred in patients with diabetic foot ulcer can cause longer treatment periods, the higher treatment costs, and the more decrease the patients' life qualities. The effect of a diabetic foot ulcer is strongly perceived by the patients, thus, the roles and supports of family are really helpful. The support can be instrumental such as the provision of facilities that can support the patients' activities and the companion during treatment periods in health center, and also material and transportation to the treatment place. Good support from

the family improves the intention of diabetic foot ulcer prevention. This study aims at arranging the intention model of the diabetic foot ulcer prevention behavior based on the perspective of social support and personal agency in city of Gorontalo.

#### **Material and Method**

This was a cross-sectional study that had been conducted from December 1<sup>st</sup>, 2018 to May 31<sup>st</sup>, 2019 on respondents suffering from the diabetes mellitus. The samples were 329 respondents out of 1516 population. They were diabetes mellitus sufferers who did not have foot ulcer aged 18 years and over, and had been selected by simple random sampling technique. The variables consisted of social support (X1) which was everything around the individuals that influenced the behaviors of them in preventing the diabetic foot ulcer. The social support (X1) itself comprised of family's support (X1.1), and friend's support (X1.2). The question items included assessment support, instrumental support, informational support, and emotional support.

Other than social support (X1), the personal agency was another independent variable (X2). It was the individual's self-ability to prevent diabetic foot ulcer consisting of a perceived variable (X2.1) and self-efficacy (X2.2). The last was the dependent variable namely intention (Y), the strong desire of the individuals themselves to prevent the diabetic foot ulcer involving the dieting intention (Y.1), physical activities intention (Y.2), consuming medicines intention (Y.3), and foot and blood sugar controlling intention (Y.4). The questionnaire had been ethically tested at Airlangga University of Surabaya, and it had owned its validity and reliability tests. Data analysis was completed by SEM PLS (Partial Keast Square) software.

**Findings:** The research result at Table 1 shows that the diabetes patients for elderly category aged 52 - 65 years are 214 (64,3%), female category consisted of 240 respondents (72,9%), respondents who are Senior High School graduates achieve 223 (67,8%), and those who do not have job (housewives and retired employees) are 215 (65,3%).

Characteristics	Classification	Frequency	Percentage	Mean ± SD Min - Max
	Late Adult (35–45 years)	30	9,0	
<b>A</b>	Early Elderly (46 – 55 years)	62	18,6	
Age	Late Elderly (56 – 65 years)	214	64,3	
	Elderly> 65 years	23	6,9	Mean: 57.29
G.	Male	89	27,1	SD: 8.88
Sex	Female	240	72,9	Min :35
	Unemployment	215	65,3	Max : 84
T.1.	Farmer	61	18,6	
Job	Private Employee	30	9,1	1
	Civil Servant	23	7,0	1

Table 1: Respondents' Charateristics, 2019

Table 2 reveals that respondents who have family's support in less category are 71 respondents (21,3%) and those who receive support from friends in less category are 95 (28,4%), and there are 174 respondents (52,9%) who receive social support in a sufficient category.

Table 2: Social Support, Personal Agency and Intention to Prevent Diabetic Foot Ulcer Variable

Variable	Indicator	Category	Total	Percentage	Mean ± SD Min - Max
	Family's Support	Less	71	21,3	44,65 ± 13,32 20 - 60
		Sufficient	116	34,7	
Social summent		Good	141	42,2	
Social support	Friend's Support	Less	95	28,4	
		Sufficient	107	32	44,61 ± 14,06 25 - 64
		Good	127	38	

Variable	Indicator	Category	Total	Percentage	Mean ± SD Min - Max	
		Less	48	14,6		
Overall Social S	upport Score	Sufficient	174	52,9	89,93 ± 17,68 37 - 128	
		Good	107	32,5		
	Perceived control	Less	157	47,7	$18,89 \pm 6,047 - 34$	
Personal	Perceived control	Good	172	52,3	18,89 ± 0,04 / - 34	
agency	C-16 -65	Less	175	53,2	12.44 + 1.90.9 16	
	Self efficacy	Good	154	46,8	$-13,44 \pm 1,80 \ 8 - 16$	
Orangli Banganal	0 115		168	51,3	22 20 + 6 00 10 46	
Overall Personal	Agency score	Good	161	48,9	$32,38 \pm 6,08 \ 19 - 46$	
	D: (:	Less	119	36,2	10.04 + 1.22 (12	
	Dieting	Good	210	63,8	$10,04 \pm 1,33 \ 6 - 12$	
		Less	121	36,8	(72 + 0.02 4 - 0	
T., 4 4		Good	208	63,2	$6,73 \pm 0,92 \ 4 - 8$	
Intention		Less	119	36,2	(72 + 0.97 4 9	
	Consuming medicines	Good	210	63,8	$6,72 \pm 0,87 \ 4 - 8$	
	Foot and blood sugar controlling	Less	119	36,2	(7(+0.00.2-0	
		Good	210	63,8	$6,76 \pm 0,90 \ 3 - 8$	
0 111			160	48,6	20.27 + 2.04.21 - 24	
Overall Intention Score		Good	169	51,4	$30,27 \pm 2,84\ 21 - 36$	


Table 2 shows that almost all perceived control in a high category are 172 respondents (52,3%), respondents who have a low category of self-efficacy are 175 (53,2%), and the total of personal agency is in less category for 168 respondents (51,1%). Then, 210 respondents (63,8%) are in high category of dieting

intention, 208 respondents (63,2%) are in high category of physical activities intention, 210 respondents (63,8%) are in high category of consuming medicine intention, and 210 respondents (63,8%) are in high category of foot/blood sugar controlling intention.

Table 3: Cross Loadings with Convergent Validity and Reliability Result

Construct and Indicator		Loading (λ)	T-statistics	Chronbach's Alpha	Information
Social aumnout	X1.1	0.968	2.42	0.06	Valid & Reliable
Social support	X1.2	0.971	54,50	0,96	Valid & Reliable
Dorganal aganay	X2.1	0.954	17,97	0,85	Valid & Reliable
Personal agency	X2.2	0.762	41,11		
	Y.1	0.989	62,32		
Intention	Y.2 0.987 5,38	0.07	Valid & Reliable		
Intention	Y.3	0.989	62,32	- 0,97 N	vand & Renable
	Y.4	0.83	27,28		

Table 3 reveals that there is no loading factor for less than 0,5 and T-statistical value is less than 1,96, hence all variables are considered significant and all latent constructs are reliable and marked with the Chronbach's Alpha score which is higher than 0,6.



X1 = Social Support, X2 = Personal Agency, Y = Intention Picture 1; Statistical T Value of Structural Model

#### **Discussion**

Picture 1 reveals that there is direct correlation between social support with personal agency of diabetic foot ulcer prevention for 0,68 unit with T-statistical value for 16,27 ( $T_{count}$  is higher than  $T_{critic}(1,96)$ , there was a direct correlation between personal agency and intention of diabetic foot ulcer prevention for 0,57 unit with T-statistical value for 2,96, and there was direct correlation between social support and intention of diabetic foot ulcer for -0,27 unit with T-statistical value for 2,08.

The social support comprises assessment support, instrumental support, informational support, and emotional support. The research result at picture 1 reveals that there is correlation between social support with the personal agency of diabetic foot ulcer prevention. It is found out that the informational and instrumental supports are required by the patients. The supports can be in the form of discussion among the family members about diabetic foot ulcer prevention and treatment to give when there is an indication of wound to occur and preparing appropriate meals for the diabetes patients. The positive impact felt by the respondents is the improvement of personal agency or individual ability to observe the symptoms of diabetic foot ulcer, and the patients become more confident to do the diabetic foot ulcer prevention.

Family support is required in this phase to assist the patient in preventing potential injury. Diabetic foot ulcer symptoms consisting of less to nothing sensation, dry skin, paralysis in foot area, and callus can be found<sup>(5)</sup>.

Diabetic foot ulcer is one of the sensory nerve defects which can cause the decrease of pain sensation at half to all part of foot area<sup>(6)</sup>. Normally, people who get injured require 2 to 5 days for the inflammation phase till the wound healing process<sup>(7)</sup>.

One of the family support categories experienced less by the respondents is assessment. The support is in the form of dieting support. The family, basically, suggests the patient do a diet, yet there is not any limitation for foods supply, for example, the food containing many calories. Therefore, the respondents are not maximal in running their diet. The research conducted by May beery S.L and Lindsay S., exposes that the behavior of the people around the patients who support the diabetes patient treatment program will increase the obedience of the patients in taking treatment<sup>(8)</sup>.

The research result at picture 1 shows that there is a correlation between personal agency and intention of diabetic foot ulcer prevention. The questionnaire result shows that the respondents are difficult to do physical activities 3 times a week based on the programs of the health center. It is because the respondents are not capable of doing that especially for those who have activities as the housewives. Respondents think that their tasks as housewives are more important than doing exercise. Respondents believe that doing activities as housewives can fulfill the need for physical activities for diabetes sufferers.

The lack of personal Agency, according to the health workers, is initiate by patients having many activities at home. The respondents' ages are in late elderly category for 56 - 65 years. Their household activities should be adjusted with their ability, hence, the patients should focus only on diabetic foot ulcer prevention and other complications. The nurses, in this case, are having crucial roles in improving the personal agency. According to Hsieh Y. L., et al. the health officers are responsible in improving the patients' intention to follow the diabetes complication treatment because a high personal agency will improve the intention of diabetic foot ulcer prevention<sup>(9)</sup>.

Hence, the health officers and the family can do an orientation to patients about how to prevent any injury when doing activities and how to do foot treatment. If it is well-oriented, it will improve the diabetes patients' intention, hence they will obediently conduct the treatment program. It is strengthened by Pakaya which states that orienting the patients to rules and treatments will improve the intention and obedience of patients in following the treatment<sup>(10)</sup>. It is also supported by Pinidiyapathirage J., et al., that personal agency in self-efficacy is one of the important predictors to improve diabetes patients' intention in doing physical activities<sup>(11)</sup>

Table 2 shows that the family's support which is in a low category is 21,3%. The lack of family's support, in terms of instrumental support, strongly influences the behavior of diabetic foot ulcer prevention. The instrumental support is performed by helping the patient to do physical activities; the family can accompany the patient to visit the health center. By doing that, the patient will be more enthusiastic to follow the treatment. The study conducted by Lengerke V., K. en L., states that supports from all parties, including family, will improve the intention of patients to prevent the diabetic foot ulcer<sup>(12)</sup>.

The questionnaire result shows that social support from friends strongly helps the individual to visit the health center. The result is supported by Shuhaida N, M, H., et al., that social support has a significant correlation with blood sugar controlling behavior<sup>(13)</sup>. Social support is a heterogenic concept in which it can help o improve the mental health in terms of intention and physical health in preventing he diabetic foot ulcer<sup>(14)</sup>. The intention is also influenced by attitude, perceived norm, and personal agency<sup>(15)</sup>.

According to Ajzen and Fisben, to do an intention, there should be mutual cooperation with ones who support the intention itself, because intention is

determined by the environment or situational<sup>(16)</sup>. Faries D. M.,has stated that there is often a gap between intention of an individual with the expected result in which the respondents who have intention are difficult to realize it in behavior<sup>(17)</sup>.

#### Conclusion

The developing model of intention to prevent the diabetic foot ulcer is influenced by supports from family and friends in order to improve the intention through personal agency variable. It is suggested that there should be further research related to knowledge and motivation in improving the intention to prevent diabetic foot ulcer.

**Conflict of Interest:** None

**Source of Funding:** None

Ethic and Consent: The ethics commissions of health research at Faculty of Nursing, Universitas Airlangga (Number 1173-KEPK) approves to the protocol by giving informed consent of research and consent form to be respondents by guaranteeing the confidentiality of research.

#### References

- Chand, G., Mishra, A. K., Kumar, S., & Agarwal A. Diabetic foot. Clinical Queries: Nephrology, 2012;1(2), 144–150.
- Perkeni. Konsensus Pengelolaan Dan Pencegahan Diabetes Mellitus tipe 2 di Indonesia. In 2015.
- 3. Huang, Y. Y., Lin, K. Der, Jiang, Y. Der, Chang, C. H., Chung, C. H., Chuang, L. M., Shin SJ. Diabetes-related kidney, eye, and foot disease in Taiwan: An analysis of the nationwide data for 2000-2009., J Formos Med Assoc. 2012;111(11), 637–644.2012.09.006.
- Riskesdas. Riset Kesehatan Dasar (Basic Health Research) 2013;1–306
- Susanto H. Kaki Diabetes, Etiologi dan Patofisiologi (Diabetes foot, etiology and pathophysiology), Simphosium & Workshop, Comprehensive Management Of Diabetes and Its Complication. In 2018.
- Wibiosono S. Diagnosis Diabetes Mellitus, Simphosium & workshop, Comprehensive Management Of Diabetes and Its Complication. In 2018.
- 7. Rabbia J. Special Problem and Interventions,

- Impaired integumentary integrity, article by Mosby, Inc., an affiliate of Elsevier Inc. 2012
- 8. Mayberry L. S, O., CY. Family Support, Medication Adherence, And Glycemic Control Among Adults With type 2 diabetes. Diabetes Care. 2012;35(6):1239–45.
- 9. Hsieh YL, Lee FH, Chen CL, Chang MF, Han PH. Factors Influencing Intention to Receive Examination of Diabetes Complications. Asian Nurs Res (Korean Soc Nurs Sci)[Internet]. 2016;10(4):289–94. Available at: http://dx.doi.org/10.1016/j.anr.2016.10.004
- Pakaya N. Hubungan Orientasi Pasien Dengan Kepatuhan Pasien/ Keluarga Dalam Menjalankan Aturan di rumah sakit Unhas (Relationship Between Patient Orientation / Family Obedience In Carrying Out The Rules At The Hasanuddin University Hospital)Makassar. J Health Sport. 2015;Vol. 9 No.:1507–20.
- 11. Pinidiyapathirage J., Jayasuriya R., Cheung N.W., & Schwarzer R. Self-efficacy and planning strategies can improve physical activity levels in women with a recent history of gestational diabetes mellitus, J Psychol Heal. 2018;
- 12. Lengerke V. T, K. B, L., K. Patients' intention to speak up for health care providers' hand hygiene

- in inpatient diabetic foot wound treatment: a cross-sectional survey in diabetes outpatient centres in Lower Saxony, Germany,. J homepage http://www.tandfonline.com/loi/cphm20,. 2016;
- Shuhaida N., H, M., Suhaila S., Y., M., Azidah A., K., Norhayati M., N., Nani D., Juliawati M. depression, anxiety, stress and socio demographics faktors forr poor glycaemic control in patients with tipe II diabetes,. J Taibah Univ Med Sci. 2019;268–76.
- 14. Pesantes M.A. et al. Family Support and Diabetes: Patient's Experiences From a Public Hospital in Peru. jounal Qual Heal Res 2018. 2018; Vol. 28(12:1871 1882.
- Rimer G & . Health behavior and health education, theory, research and practice. four. C. T Orleans; 2008.
- Ajzen and Fisben. Attitudinal And Normative Variables As Predictors Of Specific Behaviors. J Pers Soc Psychol. 1973;27 no 1:41–57.
- 17. Faries MD. Why We Don't "Just Do It": Understanding the Intention-Behavior Gap in Lifestyle Medicine. Am J Lifestyle Med. 2016;10(5):322–9.