



TABLE OF CONTENTS

ORIGINAL RESEARCH

Self-Management of Patients with Diabetes Mellitus: An Integrative Literature Review <i>Chatarina Hatri Istiari</i>	PDF 901-910
Correlation Between History of ISTC Training on Independent Practicing Doctor With Discovery of TB Suspected Children in Surabaya City <i>Madu Puspita Nuansa Jatu, Fariani Syahrul</i>	PDF 911-917
Navigating the Storm: Integrative Review of Attrition Factors Among Undergraduate Nursing Students <i>Carmela Theresa T. De Leon</i>	PDF 918-926
Knowledge, Attitudes and Application of Child Safety by Families in Karunrung Village, Rappocini, Makassar <i>Baharuddin Baharuddin, Nasrullah Nasrullah, Muhasidah Muhasidah, Sri Angriani</i>	PDF 927-929
Preparation and characterization of BSA-loaded Chitosan Microspheres <i>Esti Hendradi, Retno Sari, Rifka Anggraini Anggai</i>	PDF 930-934
The Efforts to Control and Prevent Cervical Cancer through Early Detection Using the VIA Test in the East Java Provincial Health Office <i>Riza Muhammad Zulham, Chatarina Umbul Wahyuni, Muhammad Atoillah Isfandiari, Hasan Huda</i>	PDF 935-942
Both Acute and Chronic Exercise Decrease Total Cholesterol Level in Human Blood <i>Bambang Purwanto, Wahyu N. Pratiwi, Abdurachman Abdurachman</i>	PDF 943-945
Optimation Ca-Alginate Beads as Drug Reservoir in the Patch System for Model Antiinflammatory Drug <i>Esti Hendradi, Isnaeni Isnaeni, Rifaatul Mahmudah</i>	PDF 946-951
Psychological Factors Affecting the Incidence of Essential Hypertension in General Hospital of Makassar <i>Yonathan Ramba, Suharto Suharto, M. Bagus Qamaruddin, Dewi Retno Suminar, Hendrik Hendrik</i>	PDF 952-955
Risk Factors of HIV and AIDS Mortality At Ibnu Sina Hospital <i>Silvanus Roga Misi, Chatarina Umbul Wahyuni, Hari Basuki Notobroto</i>	PDF 956-960
Evaluation of POMP for Filariasis and Factors Associated with Its Problems in East Manggarai Regency in 2017 <i>Fransiskus Salesius Onggang</i>	PDF 961-968

EDITORIAL TEAM

CHIEF EDITOR (VALID FROM JANUARY 1, 2020)

Prof. Dr. Heru Santoso Wahito NUGROHO, Poltekkes Kemenkes Surabaya (Health Polytechnic of Surabaya), Indonesia

EDITORS (VALID FROM JANUARY 1, 2020)

Joel Rey U. ACOB, Department of Nursing, Visayas State University, Philippines

Dr. Sahrir SILLEHU, Department of Public Health, STIKes (College of Health Science) "Maluku Husada", Indonesia

Tanko Titus AUTA, Department of Nursing, Niger State Ministry of Health, Minna, Nigeria

Wiwin MARTININGSIH, Poltekkes Kemenkes Malang (Health Polytechnic of Malang), Indonesia

Somsak THOJAMPA, School of Nursing, Naresuan University, Thailand

Dr. Hadi PRAYITNO, Department of Public Health, Universitas Jember, Indonesia

Dr. Mfuh Anita Y. LUKONG, Department of Nursing Science, Faculty of Allied Health Sciences, College of Health Sciences, Ahmadu Bello University, Nigeria

Victoria KALUSOPA, School of Nursing, University of Zambia, Zambia

Sanglar POLNOK, Faculty of Nursing, Naresuan University, Thailand

Zhanibek YESSIMBEKOV, Food Engineering Department, Shakarim State University, Kazakhstan

ADMINISTRATOR (VALIF FROM JANUARY 1, 2020)

Suparji SUPARJI, Poltekkes Kemenkes Surabaya, Indonesia



UNIVERSITAS AIRLANGGA

FAKULTAS KESEHATAN MASYARAKAT

Kampus C Mulyorejo Surabaya 601115 Telp.031-5920948, 5920949 Fax 031-5924618

Laman : <http://www.fkm.unair.ac.id>; E-mail: info@fkm.unair.ac.id

SURAT KETERANGAN Nomor : 3148/UN3.1.10/KP/2023

Yang bertandatangan di bawah ini :

Nama : Dr. Santi Martini, dr. M.Kes
NIP : 196609271997022001
Pangkat/Golongan : Pembina / Gol. (IV/a)
Jabatan : Dekan

Dengan ini menerangkan bahwa :

Nama : Dr. Hari Basuki Notobroto, dr., M.Kes
NIP : 196506251992031002
Pangkat/Golongan : Pembina (Gol. IV/a)
Jabatan : Lektor Kepala

Telah melaksanakan penelitian dengan judul sebagai berikut :

No.	Judul Karya Ilmiah	Tahun Pelaksanaan
1	Individual and Neighborhood Socioeconomic status in the prediction of liver transplantation among patients with liver disease (C-18)	2019
2	Survival Analysis of Hemodialysis Patients (C-25)	2016
3	Climate Conditions, Larvae Free Number, DHF Incidence in Surabaya Indonesia (C-37)	2013
4	Development of Air Polluter Model for the Carbon Monoxide (CO) Element Based on Mixed Geographically Temporal Weighted Regression (MGTWR) Kriging (C-38)	2014
5	Comparison Between Hotdeck Method and Regression Method in Handling Health Science Missing Data (C-47)	2016
6	Effect of Climate and Sanitation of Rats Population and Index Flea as Indicator Detection Yersinia Pestis in Port Surabaya (C-51)	2016
7	Factors that Affect the Incidence of HIV/AIDS (An Analysis Using Tobit Regression) (C-57)	2016
8	Demographic Transition and Conditions of Health to Elderly People in East Java Province, Indonesia (C-58)	2016
9	Survival Cox Proportion Hazard in Chronic Kidney Disease with Hemodialysis at Haji General Hospital, Surabaya (C-62)	2018
10	Risk Factors Clustering of Malnutrition Case Based on The Utilization of Toddler Health Service at Lamongan Regency in 2016 (C-63)	2018
11	The Role of the National Health Insurance Program in the Use of Health Services in City X (C-74)	2020
12	Multilevel Analysis of Lifestyle and Household Environment for Toddlers With Symptoms of Acute Respiratory Infection (ARI) in Indonesia in 2007, 2012, and 2017 (C-94)	2022
13	Indikator Karakteristik Fisik Rumah Dominan dalam	2016

	Penentuan Status Kemiskinan untuk Program Rehab Rumah tidak Layak Huni di Kabupaten Sidoarjo (C-110)	
14	Perbandingan Tingkat Konsistensi Normalitas Distribusi Metode Kolmogorov-Smirnov, Lilliefors, Shapiro-Wilk, dan Skewness-Kurtosis (C-105)	2014
15	Penerapan Metode Artificial Neural Network dalam Peramalan Jumlah Kunjungan Ibu Hamil (K4) (C-112)	2019
16	Viability Status of Diabetes Melitus Patients with Complications of Hyperglycaemia, Cetoasidosis, and Gangrene (C-119)	2020
17	Prevalensi Rasio Pelayanan Kesehatan Maternal dan Ketersediaan Fasilitas Kesehatan di ERAJKN/KIS di Indonesia (C-124)	2016
18	Faktor Demografi WUS yang Berhubungan Dengan Status Gizi Berdasarkan LILA di Puskesmas Pacarkeling Tahun 2017 (C-126)	2018
19	Comparison of MICE and Regression Imputation for Handling Missing Data (C-130)	2018
20	The Affecting Factors to Grade of Breast Cancer in Dr. Soetomo Hospital of Surabaya (C-133)	2018
21	Risk Factors of HIV and AIDS Mortality At Ibnu Sina Hospital (C-134)	2018
22	Subset Best Method Regression Analysis with Cp Mallows Statistics on Factors Affecting Life Expectancy (C-135)	2022
23	Hubungan Pola Pengasuhan Orang Tua dengan Tindakan Pencegahan Kekerasan Seksual pada Anak (C-136)	2017

Adapun penelitian tersebut layak dilakukan dan menghasilkan output yang sangat baik, meskipun belum ada *Uji Ethical Clearance* karena merupakan penelitian yang menggunakan data sekunder.

Demikian surat keterangan ini kami buat untuk dapat dipergunakan sebagai persyaratan pengusulan Jabatan Fungsional Guru Besar.

Surabaya, 24 April 2023



Dr. Santi Martini, dr. M.Kes
NIP. 196609271997022001

<http://heanoti.com/index.php/hn>

RESEARCH ARTICLE

URL of this article: <http://heanoti.com/index.php/hn/article/view/hn20910>**Risk Factors of HIV and AIDS Mortality At Ibnu Sina Hospital****Silvanus Roga Misi^{1(CA)}, Chatarina Umbul Wahyuni², Hari Basuki Notobroto³**^{1(CA)}Department of Epidemiology, Faculty of Public Health, Airlangga University, Indonesia;
cellomisi27@gmail.com (Corresponding Author)²Department of Epidemiology, Faculty of Public Health, Airlangga University, Indonesia³Department of Biostatistics and Demography, Faculty of Public Health, Airlangga University, Indonesia**ABSTRACT**

Globally, HIV infection (Human Immunodeficiency Virus) and AIDS (Acquired Immune Deficiency Syndrome) were still a problem in the health. East Java Province's case rate is in the top 10, which is 29.70%. The case rate number was still above the national figure of 26.85% until March 2016. Case Rate of East Java Province was in the 10th position but cumulatively the East Java Province is ranked first with 3,292 deaths until March 2016 compared to other provinces. Gresik Regency was include in the top 10 largest number of AIDS cases with 898 AIDS cases per 2017 increased compared to 2016. The aim of the study was to find out the factors related to mortality in HIV and AIDS patients in Ibnu sina gresik district general hospital. The research method was a retrospective cohort with a total of 299 patients. Data analysis used cox non-proportional hazard with a value of $\alpha = 0.05$. Based on the results of the analysis showed that medication adherence was $p < \alpha (0.05)$ so that H_0 was rejected which meant that medication adherence had an effect on the mortality of HIV and AIDS. Increase the motivation of patients who were less adherent to took medication to be compliant in taking ARV drugs so that patient mortality is reduced. Maintain patients who took ART therapy to be obedient in taking therapy so that assessment of medication adherence can be maintained and patient survival was longer.

Keywords: Mortality, HIV, AIDS, Adherence**INTRODUCTION****Background**

The number of HIV cases up to March 2016 was 198,219 people. Based on 5 provinces with the highest number of HIV, it was DKI Jakarta (40,500), East Java (26,052), Papua (21,474), West Java (18,727) and Central Java (13,547). The cumulative number of AIDS until March was 78,292 people. The highest number of AIDS was reported from the province of East Java (14,499), followed by Papua (13,335), DKI Jakarta (8,105), Bali (5,934), and Central Java (5,049)⁽¹⁾.

AIDS mortality rate decreased from 0.94% in the year to 0.02% in March 2016. The case rate of East Java Province was in the top 10 highest at 29.70%. The case rate number was still above the national figure of 26.85% until March 2016. Case Rate of East Java Province was in the 10th position but cumulatively the East Java Province was ranked first with 3,292 deaths until March 2016 compared to other provinces with the highest case rates, because East Java Province had more population than other provinces. AIDS was the final stage of HIV-infected patients. Delay in ART treatment causes many patients deaths⁽²⁾.

Most patients began treatment in stages III and IV. began treatment at stage III and IV has a risk of death 4.5 times greater than who began treatment at stage I and II. According research clinical conditions were related to stage, CD4 count and functional status. Poor clinical conditions gave affect death greatly. Death often occurred in the clinical condition of bed rest with stages III and IV. Most deaths at median CD4 were 18 cells / mm³. This showed that most patients deaths by CD4 cells less than 200 cells / mm³ or end stage (AIDS)⁽³⁾.

Adherence to ARV treatment was due to several factors, one of which was due to lack of motivation in the patient. The aim of ARV treatment was also to prolong the life of sufferers of HIV and AIDS by preventing complications. The success of the treatment program can be seen from the regular adherence to treatment. PMO (supervisor taking medication) was need to increased patient motivation to taking medication. This can decreased mortality due to HIV and AIDS⁽⁴⁾.

Purpose

The aim of the study was to find out the factors related to mortality in HIV and AIDS patients in Ibnu sina gresik district general hospital.

METHODS

The research method was restropective cohort. Subject of research was HIV and AIDS patients who took ARV treatment for the period 2013-2016. Number of populations by 303 patients. Pregnant patients with HIV and AIDS was 4 patients. Patients with HIV and AIDS who was pregnant excluded from this study because pregnant women have different ART guidelines that were likely counfounding of the study. The number of subject by 299 patients who followed ART at Ibn Sina hospital. CD4 measurement was the CD4 count when the patient was first starts ART. It recorded in the medical records. Stage measurement was the clinical stage when the patient first starts ART based on the category set by the hospital. The measurement of functional status was ability of HIV and AIDS patients to carry out physical activity and self-care when starting ARV therapy. It recorded in the medical records. The measurement of opportunistic infections was presence of other infections that accompany HIV infection in HIV and AIDS patients. It recorded in the medical record. Data analysis used cox non propotional hazard.

RESULTS

Mortality per Year

Based on Figure 1 showed that in 2014 mortality per 100 person per year experienced an increase. Furthermore, until 2016 mortality per 100 person per year had decreased.




Figure 1. Mortality per Year

Mortality of HIV and AIDS Cases per Year

Based on Figure 2 showed that AIDS cases experienced an increase in mortality from 2014 to 2015. While HIV cases experienced a decline from 2016.




Figure 2. Mortality of HIV and AIDS Cases per Year

Risk Factors of HIV and AIDS Mortality

Based on the results of the analysis showed that medication adherence was $p < \alpha$ (0.05) so that H_0 was rejected which meant that medication adherence had an effect on the mortality of HIV and AIDS. Drug adherence hazard rate is 0.961 which means that every increase of 1% adherence to taking medication for HIV and AIDS patients will experience a decrease in mortality by 0.961 times.

Table 1. Risk Factors Mortality of HIV and AIDS

No	Variabel	HR	Nilai P	Keterangan
1.	CD4	1.000	0.929	No significant
2.	Stage			
	Stage I		Referens	
	Stage II	1023.167	0.950	No significant
	Stage III	755.297	0.952	No significant
	Stage IV	3462692	0.941	No significant
3.	Functional Status			
	Working		Referens	
	Ambulatory	1.190	0.856	No significant
	Bedridden	0.634	0.657	No significant
4.	Opportunistic infections			
	Absent		Referens	No significant
	Present	0.799	0.822	
7.	Medication adherence	0.964	0.001	Significant

DISCUSSION

This study showed that from several variables that were tested for their effect on mortality there was only a variable adherence to medication that had an influence on mortality. Research carried out namely survival is a prognosis analysis, it could be HIV and AIDS patients whose initial severity is high because of adherence to drinking can reduce the severity. The level of adherence to medication that is less will be related to the progression of HIV disease so that patients do not survive⁽⁴⁾.

This research showed that stadiums had no effect on the mortality of HIV and AIDS patients. Stadium was a measure of the severity of patients with HIV and AIDS. HIV patients with stage IV enter the final stage, increased the risk of mortality than those who present with stage I, but the stage has no effect on HIV and AIDS. The cause of the stage has no influence on HIV and AIDS was likely to be influenced by the number of censored and missing factors so that the information obtained was also lacking. This is in contrast to other study which showed that patients who started therapy in stage III and IV had a greater risk of death up to 4.5 times than those who started therapy at stage I and II⁽³⁾.

Other factors such as CD4 cell count, functional status, opportunistic infections and adherence to therapy were not related to survival in patients with HIV and AIDS at Ibnu Sina Gresik Hospital. This was likely due to incomplete data collection so that more data is missing and sensors. For example, adherence to therapy was lacking, as was lost to follow-up, as it was found that lost to follow-up was included in the censored data in survival analysis. OIs did not affect survival when tested together with other factors. This was because opportunistic infections were not sensitive enough when tested together with other factors³. In contrast to the other research showed that factors that influence HIV survival in CD4 counts. In addition, OIs are not sensitive in influencing HIV survival. The biggest possibility that influences these factors does not affect the survival of HIV and AIDS due to data regarding CD4, functional status, opportunistic infections, and supervisors taking medication, namely the first data of HIV and AIDS patients first taking ARV therapy⁽⁴⁾.

CONCLUSION

The factors that influence HIV AIDS mortality were adherence to taking medication. Increasing the motivation of patients who are less adherent to take medication to be compliant in taking ARV drugs so that patient mortality is reduced. Maintaining patients who take ART therapy to be obedient in taking therapy so that assessment of medication adherence can be maintained and patient survival is longer.

REFERENCES

1. Ministry of Health of Republic of Indonesia. The Report of HIV and AIDS Development Situation in Indonesia January-March 2016. Jakarta: MoH-RI; 2016.
2. Ministry of Health of Republic of Indonesia. HIV / AIDS Situation in Indonesia. Jakarta: MoH-RI; 2016.
3. Utami S. Predictors of Death of HIV / AIDS Patients with Antiretroviral Therapy (ARV) at Regional General Hospital Period 2006-2014, Badung, Bali. Postgraduate Thesis. Denpasar: Udayana University; 2014.
4. Chakravarty J, Tiwary NK, Shashi RP, Shukla S, Tiwari A, Mishra RN, Sundar S. Determinants of Survival in Adult HIV Patients on Antiretroviral Therapy in Eastern Uttar Pradesh: A Prospective Study. 2013.