

Piracy Threat – Important update to keep your details safe and secure. [Click here for further information.](#)

Table of contents

Volume 679

2021

◀ Previous issue Next issue ▶

The 1st International Conference on Biotechnology and Food Sciences 11 September 2020, Surabaya, Indonesia

Accepted papers received: 08 February 2021

Published online: 26 February 2021

Open all abstracts

Preface

OPEN ACCESS 011001

The 1st International Conference on Biotechnology and Food Sciences (INCOBIFS)
Surabaya Indonesia, 11 September 2020

+ Open abstract  View article  PDF

OPEN ACCESS 011002

Conference Photographs

+ Open abstract  View article  PDF

OPEN ACCESS 011003

Organizing Committee

+ Open abstract  View article  PDF

OPEN ACCESS 011004

Peer review declaration

+ Open abstract  View article  PDF

Papers

OPEN ACCESS 012001

Converting husbandry waste into liquid organic fertilizer using probiotic consortiums
(*Lactobacillus* sp., *Rhodopseudomonas* sp., *Actinomyces* sp., *Streptomyces* sp.)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012002

A review: bioactive compounds of macroalgae and their application as functional beverages

S G Widyaswari, Metusalach, Kasmiati and N Amir

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012003

Improvement quality of sugar cane bagasse as fish feed ingredient

L H Suryaningrum and R Samsudin

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012004

The Growth and Yields of Shallot (*Allium Wakegi* Araki) CV. lembah palu Growing under Hydroponic Substrate Systems

R Yusuf, S A Lasmini, M Sandi, A Rahim and I Wahyudi

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012005

Application of glycerol on bioplastic based carrageenan waste cellulose on biodegradability and mechanical properties bioplastic

S N Fauziyah, A S Mubarak and D Y Pujiastuti

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012006

Gill and skin pathology of hybrid grouper (*E. fuscoguttatus* x *E. lanceolatus*) infested *Zeylanicobdella arugamensis* worms in different infestations degree

M Nisa, G Mahasri and L Sulmartiwi

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012007

Growth performance of tambaqui (*Colossoma macropomum*) supplemented with honey prebiotic in stagnant peat ponds

H Silalahi, R Djauhari and S S Monalisa

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012008

The Concentration of polyethylen glycol (PeG) 400 on bioplastic cellulose based carrageenan waste on biodegradability and mechanical properties bioplastic

D S Maulana, A S Mubarak and D Y Pujiastuti

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.


[+ Open abstract](#) [View article](#) [PDF](#)



-
- OPEN ACCESS** 012009
Fish oil extraction as a by-product of Tilapia (*Oreochromis* sp.) fish processing with dry rendering method
S H Suseno, A K Rizkon, A M Jacob, Kamini and D Listiana
[+ Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012010
Optimization of Extraction Time on The Characteristic of Gelatin from Scales of Red Snapper (*Lutjanus* sp.)
I Safi'i, W Tjahjaningsih and E D Masithah
[+ Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012011
Screening acetylcholinesterase inhibitors from marine-derived actinomycetes by simple chromatography
M Kamaruddin, I Marzuki, A Burhan and R Ahmad
[+ Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012012
Assessment of Heavy Metal Lead (Pb) Contents in Canned Crab Products by Atomic Absorption Spectrophotometry (AAS)
M Agustina, Mulyono and W Tjahjaningsih
[+ Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012013
The concentration of sorbitol on bioplastic cellulose based carrageenan waste on biodegradability and mechanical properties bioplastic
M D Arief, A S Mubarak and D Y Pujiastuti
[+ Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012014
Identification photoprotective activity of marine seaweed: *Eucheuma* sp
R D Kasitowati, A Wahyudi, R Asmara, D Aliviyanti, F Iranawati, M A P Panjaitan, D C Pratiwi and S Arsad
[+ Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012015
Analysis of the density and intensity of metallothionein in *Crassostrea cucullata* oyster hulls in the coastal fishing port of Mayangan Probolinggo, East Java using immunohistochemical techniques
W Isoni, N Maulida and M I Mubarogi
[+ Open abstract](#) [View article](#) [PDF](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



-
- OPEN ACCESS** 012016
The effect of different bait on the catch of traps in the waters of the tip of Pangkah Gresik Regency, East Java
W Isoni, N Maulida and M Huda
[+](#) [Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012017
The Identification and Distribution Components of Polycyclic Aromatic Hydrocarbon Contaminants at the Port of Paotere, Makassar, South Sulawesi
I Marzuki, I Pratama, H E Ismail, I. Paserangi, M Kamaruddin, M. Chaerul and R Ahmad
[+](#) [Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012018
Koi (*Cyprinus carpio*) Hatchery techniques: its performance in BBI Boyolali
M G Laksono, Sugianta and M B Santanumurti
[+](#) [Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012019
Ectoparasite infestation and survival rate of pacific white shrimp (*Litopenaeus vannamei*) that immunized with crude protein *Zoothamnium penaei* in intensive ponds
G Mahasri, A T Mukti, M Nisa, G C Prakosa and W H Satyantini
[+](#) [Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012020
Study of crude extract yield and gonad maturity level of sea cucumber *Phylloporus dobsoni* correlation
S Andriyono, E D Masithah and D Winarni
[+](#) [Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012021
Techniques of additional *Kappaphycus alvarezii* on seaweed face mask production
N R Prima and S Andriyono
[+](#) [Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012022
Ice cream properties affected by carrageenan from seaweed deference type drying methods
I Irawan and Fitriyana
[+](#) [Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012023
The effect of washing on the making of surimi and kamaboko tilapia (*Oreochromis sp.*)
see our Privacy and Cookies policy. 

E Saputra, W Tjahjaningsih and A A Abdillah

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012024

The effect of storage on the making of surimi and kamaboko tilapia (*Oreochromis* sp.)

E Saputra, W Tjahjaningsih and A A Abdillah

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012025

Molecular identification and prevalence of endoparasite worms in Silver pompano (*Trachinotus blochii*) in floating net cages of Mari-culture Center, Lampung

L N F Haryanto, S Subekti, H B Ardiyanti, M K Amiin, R E K Akbar, I Achmadi and M A Yudarana

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012026

Microbiology safety of green mussel, *Perna viridis* after treated with boiling and *sous vide*

A S Samsudin and N U Karim

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012027

The utilization of chitosan from Comb-pen shell (*Atrina pectinata*) as an emulsion stabilizer in the production of hand body cream

Y Supratin, L Sulmartiwi and E Saputra

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012028

The potential of chitosan from comb-pen (*Atrina pectinata*) shell waste on the characteristics of hand body cream

F Fauzi, L Sulmartiwi and E Saputra

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012029

Spawning technique of yellowfin tuna (*Thunnus albacares*) infloating nets cage

B Bramantya, Gunawan and L A Sari

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012030

Enlargement technique of humpback grouper (*Cromileptes altivelis*) with floating nets cage

G Y Pamungkas and L A Sari

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more,

[see our Privacy and Cookies policy](#) [View article](#) [PDF](#)



-
- OPEN ACCESS** 012031
Microencapsulated fish oil powder by spray drying using combination of wall materials in Kasetsart University, Bangkok
I Aprilia, D Y Pujiastuti and W Klaypradit
[+](#) Open abstract [View article](#) [PDF](#)
-
- OPEN ACCESS** 012032
Effect of deacetylation conditions on physicochemical properties of chitosan derived from shrimp shell and squid pen
A R Basarah, D Y Pujiastuti and Yaowapha Waiprib
[+](#) Open abstract [View article](#) [PDF](#)
-
- OPEN ACCESS** 012033
Substitution of patin *Pangasius pangasius* flour in making sticks as an alternative of food high protein and source of calcium for autism patients
V Amelia, S Subekti and L Sulmartiwi
[+](#) Open abstract [View article](#) [PDF](#)
-
- OPEN ACCESS** 012034
Antioxidant properties from seaweeds *Kappaphycus alvarezii*, *Euchema spinosum* and *Sargasum* sp. using different solvent
A A Abdillah, M A Alamsjah and N E Nasution Sugijanto
[+](#) Open abstract [View article](#) [PDF](#)
-
- OPEN ACCESS** 012035
Effect of kappa-carrageenan on physicochemical properties of mantou (Chinese steamed bread)
M H C Putra and A A Abdillah
[+](#) Open abstract [View article](#) [PDF](#)
-
- OPEN ACCESS** 012036
The application of betacyanin microcapsules as natural food colorant on beverage model
S R Nurbaya, W D R Putri, E S Murtini and A Khamidah
[+](#) Open abstract [View article](#) [PDF](#)
-
- OPEN ACCESS** 012037
Microbial quality and diversity of *Caesio cunning* and *Scolopsis taenioptera* harvested by using trap and trawl fishing techniques
M Suhaimi and N U Karim
[+](#) Open abstract [View article](#) [PDF](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



-
- OPEN ACCESS** 012038
Characteristics physicochemical of melanin from squid ink (*loligo sp.*) extracted by ethanol
F Abidin, L Sulmartiwi and E Saputra
[+ Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012039
Analysis of fluctuating asymmetry of black strain tilapia *oreochromis niloticus* and red strain tilapia *Oreochromis niloticus* in Kabat Fish Hatchery Center Banyuwangi, East Java
D Kurniawan and A H Fasya
[+ Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012040
Utilization of fermented Seligi leaf flour *Phyllanthus buxifolius* toward the specific growth rate, daily growth rate and survival rate of siam catfish (*Pangasius pangasius*)
Y G Budiman, M Lamid, B S Rahardja and M Amin
[+ Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012041
Improving crude protein and crude fat content of Seligi leaf (*Phyllanthus buxifolius*) flour through probiotic fermentation
A K Nisa, M Lamid, W P Lokapirnasari and M Amin
[+ Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012042
Addition of turmeric in feed on growth and survival rate of Nilasa red tilapia (*Oreochromis sp.*)
R Cahyani, W H Satyantini, D D Nindarwi and Y Cahyoko
[+ Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012043
Characterization of semi-refined kappa-carrageenan from *Kappaphycus alvarezii* with different solvents in Tanjung Sumenep
H M Noor, M A Alamsjah and S Andriyono
[+ Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012044
The substitution effect of bone fish flour milkfish (*Chanos chanos*) physical and chemical characteristics of cookies
I Muzaki, H Suprpto and R Kusdarwati
[+ Open abstract](#) [View article](#) [PDF](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



-
- OPEN ACCESS** 012045
Identification and the prevalence of fungal gouramy (*Osphronemus gouramy*) in modern market Surabaya
M S Andreas, R Kusdarwati and H Suprpto
[+ Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012046
Substitution of commercial feed with maggot meal (*Hermetia illucens*) to the growth rate, feed conversion ratio and feed efficiency catfish (*Pangasius pangasius*)
M Ulumiah, M Lamid and K T Pursetyo
[+ Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012047
Correlation between water quality with prevalence of black tiger shrimp (*Penaeus monodon*) infested by ectoparasite protozoa in traditional ponds of Wonorejo, Surabaya
M A Luthfi, S Subekti and G Mahasri
[+ Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012048
Study of nitrogen (N) and phosphorus (P) in the land of mangrove sediments in ecotourism area Wonorejo Surabaya and coastal area of Jenu Tuban
N Pradipta, M A Alamsjah and E D Masithah
[+ Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012049
Corellation of water quality to the prevalence of ectoparasite in milkfish (*Chanos chanos*) in Sedati District, Sidoarjo
H Irawan, Kismiyati and G Mahasri
[+ Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012050
Effect of soaking time of blood cockle (*Anadara* sp.) shells powder with hydrochloric acid on the characteristics of nano calcium
D S Herlina, L Sulmartiwi and E D Masithah
[+ Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012051
Addition of crude fish oil (CFO) in feed toward fat and energy retention of mud crab (*Scylla serrata*)
S Hadijah, Agustono and W P Lokapirnasari
[+ Open abstract](#) [View article](#) [PDF](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our [Privacy and Cookies policy](#).



-
- OPEN ACCESS** 012052
Effectiveness of giving clove oil as an anaesthetic for survival rate and number of leucocytes in cantang grouper (*Epinephelus* sp.) in the closed transportation
S Nurkomaria, H Suprpto and Sudarno
[+ Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012053
Effect of the addition of crude fish oil (CFO) in feed to the content of EPA and DHA in mud crab (*Scylla serrata*)
T Wijaya, Agustono and M A A Arif
[+ Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012054
The effectiveness combination of maggot (*Hermetia illucens*) flour with commercial feed on growth rate, feed conversion ratio, and feed efficiency of tilapia (*Oreochromis niloticus*)
V Indriawati, B S Rahardja and Prayogo
[+ Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012055
Depuration of heavy metals Pb and Cd content in blood cockles (*Anadara antiquata*) with different filters
W Arifin, B S Rahardja and K T Pursetyo
[+ Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012056
Substitution of commercial feed with fermented banana peel flour (*Musaceaea* sp.) and fish meal to feed consumption level, specific growth rate, feed efficiency, fat retention, and energy retention in siam catfish (*Pangasius hypophthalmus*)
A Aisyah, A S Gustiningrum, Agustono and M A Al-Arif
[+ Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012057
The effect *Moringa oleifera* leaf extract and *Lactobacillus acidophilus* supplementation on crude protein and crude fat retention in Tambaqui, *Colossoma macropomum*
R A Pitri, E Suyanti, S Septayani, Agustono and W P Lokapirnasari
[+ Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012058
Molecular identification and prevalence of ectoparasite worms in barramundi (*Lates calcarifer*) in Lampung Waters
I Achmadi, S Subekti, H B Ardiyanti, M K Amiin, L N F Haryanto, R E K Akbar and M A Yudarana
[+ Open abstract](#) [View article](#) [PDF](#)
-

-
- OPEN ACCESS** 012059
Morphological profile of L2 *Anisakis typica* on Indian Mackerel (*Rastrelliger kanagurta*) from Sedati Fish Auction, Sidoarjo-East Java, Indonesia using Scanning Electron Microscope (SEM)
N Suryani, S Subekti, S Koesdarto and M K Amiin
[+ Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012060
Occurance of Anisakis of mackarel tuna (*Euthynnus affinis*) from Sendangbiru fishing auction place, East Java, Indonesia
R Bobsaid, P D W Sari and S Subekti
[+ Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012061
The study of virus collation with the polymerase chain reaction (PCR) method in export fishery commodities
L Kurniawati and K T Pursetyo
[+ Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012062
Effect of Culture Combination on Growth and Carrageenan Content of *Kappaphycus alvarezii* Green and Red Variety
A A Abdillah and J Triastuti
[+ Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012063
Nitrate and phosphate dynamics of phytoplankton abundance in Kanceng River, Sepuluh, Bangkalan, East Java, Indonesia
D D Nindarwi, S H Samara and M B Santanumurti
[+ Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012064
Assessment of Seasonal Waters Quality Based on Abundance, Diversity, and Domination of Phytoplankton in Bajulmati Reservoir
E W Pertiwi, E D Masithah and Suciyono
[+ Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012065
Effect of sublethal dose organophosphate pesticides on embryogenesis and hatching rate of silver rasbora eggs (*Rasbora argyrotaenia*)
A B Prastika, L Sulmartiwi and L Lutfiyah
[+ Open abstract](#) [View article](#) [PDF](#)
-

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



OPEN ACCESS

012066

The correlation between temperature and intensity of *Zeylanicobdella arugamensis* on cantang grouper (*E. fuscoguttatus* × *E. lanceolatus*) from traditional ponds in the Kampung Kerapu Lamongan East Java Indonesia

D D Afifah, G Mahasri and W H Satyantini

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012067

Utilization of *Nitrosomonas* sp and *Nitrobacter* sp probiotic towards total suspended solid and ammonia level in Nile tilapia culturing using aquaponic system

K H Dwiardani, Prayogo and B S Rahardja

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012068

The administration of *Caulerpa racemosa* extract on total bacteria and survival rates of white shrimp (*Litopenaeus vannamei*) after infected by *Vibrio parahaemolyticus*

A F Pratiwi, W H Satyantini, G Mahasri, L Sulmartiwi, A T Mukti and Sudarno

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012069

Examination of Taura Syndrome Virus (TSV) in white shrimp (*Litopenaeus vannamei*) and tiger prawn (*Penaeus monodon*) with Polymerase Chain Reaction (PCR) method

A N Fadilah and A H Fasya

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012070

Potential utilization of kombucha as a feed supplement in diets on growth performance and feed efficiency of catfish (*Clarias* sp.)

H U Ramadhan, Prayogo, H Kenconoajati, B S Rahardja, M H Azhar and D S Budi

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012071

Application of edible film from chitosan as biodegradable packaging

E Saputra, W Tjahjaningsih and A A Abdillah

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012072

Substitution of commercial feed with fermented banana peel flour (*Musaceaea* sp.) and fish meal to crude protein, energy, crude lipid and organic matter of meat in siamese catfish (*Pangasius hypophthalmus*)

R Z Darmawan, S M Ghaisani, Agustono and M A Al-Arif

[+ Open abstract](#) [View article](#) [PDF](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



OPEN ACCESS

012073

The effect of garlic (*Allium sativum*) and turmeric (*Curcuma longa*) extract addition in commercial feed on feeding rate, feed efficiency and feed conversion ratio of gouramy fish (*Osphronemus gouramy*)

D Afifah, M Arief and M A Al-Arif

[+](#) [Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012074

The Variability in Population Structure of Gastropods in Sedati Waters, Sidoarjo Regency, East Java

S Fadliyah, L A Sari, K T Pursetyo, A Zein, M H Idris and Y Cahyoko

[+](#) [Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012075

The effect of pH and incubation time on rude protease enzymes activity of *Bacillus mycoides* from anchovy isolates (*Stolephorus* sp.)

D Nirmala, P Yudha and D Cahyanto

[+](#) [Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012076

The influence of garlic (*Allium sativum*) and turmeric (*Curcuma longa*) extract as attractant in commercial feed on feeding rate and fat retention in gouramy (*Osphronemus gouramy*)

G Y Pamungkas, M Arief and M A A Arif

[+](#) [Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012077

Beardless barb *Cyclocheilichthys apogon* (Valenciennes, 1842) (*Cypriniformes*, *Cyprinidae*): Distribution extension and first record from South Bali

V Hasan, A Wijayanti, M B Tamam, R A Islamy and M S Widodo

[+](#) [Open abstract](#) [View article](#) [PDF](#)

JOURNAL LINKS

[Journal home](#)

[Information for organizers](#)

[Information for authors](#)

[Contact us](#)

[Reprint services from Curran Associates](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our [Privacy and Cookies policy](#).



This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



PAPER • OPEN ACCESS

The administration of *Caulerpa racemosa* extract on total bacteria and survival rates of white shrimp (*Litopenaeus vannamei*) after infected by *Vibrio parahaemolyticus*

To cite this article: A F Pratiwi *et al* 2021 *IOP Conf. Ser.: Earth Environ. Sci.* **679** 012068

View the [article online](#) for updates and enhancements.



240th ECS Meeting ORLANDO, FL

Orange County Convention Center **Oct 10-14, 2021**

Abstract submission deadline extended: April 23rd

SUBMIT NOW

The administration of *Caulerpa racemosa* extract on total bacteria and survival rates of white shrimp (*Litopenaeus vannamei*) after infected by *Vibrio parahaemolyticus*

A F Pratiwi¹, W H Satyantini^{2*}, G Mahasri², L Sulmartiwi³, A T Mukti², Sudarno²

¹Aquaculture Program, Faculty of Fisheries and Marine, Universitas Airlangga Campus C UNAIR, Jl.Mulyorejo, Surabaya, Indonesia 60115

²Department of Fish Health Management, Faculty of Fisheries and Marine, Universitas Airlangga, Campus C UNAIR, Jl.Mulyorejo, Surabaya, Indonesia 60115

³Departement of Marine, Faculty of Fisheries and Marine, Universitas Airlangga, Campus C UNAIR, Jl.Mulyorejo, Surabaya, Indonesia 60115

*Corresponding author: woro_hs@fpk.unair.ac.id

Abstract. The purpose of this study was to determine the effect of administration *Caulerpa racemosa* extract on total bacteria and survival rate of white shrimp (*Litopenaeus vannamei*) after infected by *Vibrio parahaemolyticus*. This study used a Completely Randomized Design with five treatments and three replications. Shrimp were divided into four groups, one control group without *C. racemosa* extract, another three groups were administrated with *C. racemosa* extract in different doses; 3 µg/g (P1), 6 µg/g (P2) and 9 µg/g (P3) by injection. After 24 hours, the control group was divided into two groups, negative control (K-) was injected by PBS and positive control (K+) also another three groups (P1, P2 and P3) were infected by *V. parahaemolyticus*. Observed parameters were total bacteria and survival rates. On the 1st day, the total bacteria was lower in the treatment with *C. racemosa* extract than K+ and significantly different ($P < 0.05$). On the 7th day, total bacteria increased again, but in P3 (20.46×10^4 CFU/ml) was lower than K+ and significantly different ($P < 0.05$). The highest survival rates was 93.33% (P2) and the lowest was 56.67% (K+). It can be concluded that *C. racemosa* extract can be given to *L.vannamei* at dose 6 µg/g.

1. Introduction

White shrimp (*Litopenaeus vannamei*) is a fishery commodity that is profitable, including fast growth, high stocking density and has a high selling price [1]. World white shrimp production in 2015 was above 3.5 million tons with the largest contributor to China 1.62 million tons, India 416,000 tons and Indonesia 410,000 tons [2]. The increasing market demand causes white shrimp to be cultured intensively.

Disease is a limiting factor in shrimp farming. Diseases can be caused by bacteria, viruses, fungi or parasites. Vibriosis attacks can cause mortality up to 100% in the larval or juvenile stage [3]. Several studies have reported that acute hepatopancreatic necrosis disease or AHPND caused by *V. parahaemolyticus* caused 80% mortality in China in 2011 [4], while AHPND cases in Mexico resulted in 81% mortality [5].

Antibiotics as disease control have been prohibited from being used, this is because they can cause the emergence of antibiotic-resistant pathogens and leave a residue in the environment and on the body of shrimp [6]. One of the efforts to control and prevent disease in shrimp is through increasing the shrimp defense system by using immunostimulants [7]. Immunostimulants are biological in nature and are synthesized from compounds that can induce a non-specific immune system in shrimp.



Content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](https://creativecommons.org/licenses/by/3.0/). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

Caulerpa racemosa has potential as a source of phytochemicals that have immunological activity, namely sulfate polysaccharides. Sulfate polysaccharides have biological activities such as immunology, antiviral, and anti-oxidation [8]. Sulfate polysaccharides can boost the immune system by activating macrophage activity [9]. The purpose of this study was to determine the effect of administering *Caulerpa racemosa* extract to total bacteria and survival rate of white shrimp (*Litopenaeus vannamei*) after infected with *Vibrio parahaemolyticus*.

2. Material and methods

2.1. Materials

There are 20 aquariums used with a size of 40 x 30 x 30 cm³ for this research. White shrimp (*Litopenaeus vannamei*) DOC 30 with a weight of 2- grams and a length of 5-7 cm. Previously, the shrimp were acclimatized for 15 days, then before using the shrimp, they were isolated first whether they were infected with *Vibrio* or not. *Caulerpa racemosa* is obtained from Balai Besar Perikanan Budidaya Air Payau Jepara. While the *Vibrio parahaemolyticus* isolate was obtained from Balai Perikanan Budidaya Air Payau Situbondo.

2.2. Methods

2.2.1 Research procedures

a. Extraction of *C. racemosa* by hot water method

Caulerpa racemosa powder is put into Erlenmeyer mixed with distilled water with a ratio (g: v / 1:20) then put into a beaker glass filled with hot water 85°C. Extraction was carried out for 2 hours at a temperature of 85°C above the hotplate stirrer. Then the extract was separated between the residue and the supernatant using a refrigerator centrifuge at 4 °C at 3500 rpm for 15 minutes. The supernatant is then separated with the residue and then filtered. The filtrate obtained was then carried out by solvent evaporation using a rotary evaporator and then put in an oven at 35°C to get the extract in the form of a paste.

b. Aquarium preparation and water maintenance media

Before use the aquarium is cleaned by brushing and rinsing water until clean. The aquarium is filled with water that has been added with 10 ppm of chlorine as a disinfectant. The aquarium was immersed for 24 hours, then the chlorine water was removed and rinsed with clean water. Furthermore, drying for 24 hours with an inverted aquarium position.

c. Preparation of anticoagulant

The preparation of the anticoagulant is by mixing the following ingredients 30 mM trisodium citrate, 0.34 M sodium chloride, 10 mM EDTA while maintaining an atmosphere of pH 7.5.

d. *Vibrio parahaemolyticus* Culture and Density Measurement

V. parahaemolyticus culture was carried out by taking one ose of isolate and culturing it on TSB media aseptically, then incubating for 24 hours at room temperature (33 ° C). After 24 hours, the concentration of bacteria was measured using a spectrophotometer.

e. White shrimp rearing

The use of feed in the maintenance of white shrimp uses artificial feed in the form of pellets with a minimum protein content of 36%. White shrimp feed is given as much as 3% of the total body weight. The frequency of feeding was three times a day, namely at 09.00 AM, 01.00 PM and 05.00 PM.

f. Injection with *C. racemosa*

White shrimp was injection with *C. racemosa* intramuscularly in the third segment of the abdomen as much as 0.1 ml / head. Meanwhile, for the control treatment, injected with PBS as much as 0.1 ml / shrimp. *C. racemosa* extract was administered at the beginning of maintenance (T0) on shrimp.

g. Challenge test with *V.parahaemolyticus*

The challenge test with *V. parahaemolyticus* was carried out 24 hours after given *C. racemosa* extract by injection (T1), which was injected intramuscularly into the third segment of the abdomen as much as 0.1 ml / shrimp, for control treatment injected with PBS as much as 0.1 ml / shrimp.

h. Collecting the hepatopankreas

Shrimp hepatopankreas was taken for total bacterial observation. The hepatopankreas was taken by sterilizing the body of the shrimp with alcohol and then the shrimp was dissected to extract the hepatopankreas. The hepatopankreas was placed in a sterile microtube containing 0.9 ml of PBS and then homogenized.

2.2.2 Parameter of Research

The main parameters observed were total bacteria and survival rate. The supporting parameters observed were clinical sign and water quality which included temperature, pH, Dissolved Oxygen and salinity.

2.2.3 Data Analysis

Data processing from the research used Analysis of Variance (Anova) to determine the effect of the treatment given. If the results are significantly different ($p < 0.05$), the calculation is continued with the Duncan Test (DMRT) using the SPSS 20 tool [10].

3. Result and discussion

3.1 Result

3.1.1 Total Bacteria

The average of total bacteria in hepatopankreas of *L. vannamei* could be seen in Table 1.

Table 1. The average of total bacteria in hepatopankreas of *L. vannamei*

Treatment	Total bacteria ($\times 10^4$ CFU/ml)		
	T0 \pm SD	T1 \pm SD	T7 \pm SD
K-	24.56 ^{bc} \pm 0.49	3.33 ^d \pm 0.21	14.20 ^e \pm 0.26
K+	24.86 ^{ab} \pm 0.15	26.20 ^a \pm 0.43	23.46 ^c \pm 0.45
P1	24.13 ^c \pm 0.15	17.03 ^c \pm 0.15	24.90 ^a \pm 0.10
P2	24.56 ^{bc} \pm 0.35	21.36 ^b \pm 0.32	24.26 ^b \pm 0.25
P3	25.26 ^a \pm 0.31	21.26 ^b \pm 0.25	20.46 ^d \pm 0.50

Description: (K-) sterile PBS injection; (K +) infection with VP 10^6 CFU / ml per shrimp; (P1) injection of *C. racemosa* 3 μ g / g + VP infection 10^6 CFU / ml per shrimp; (P2) injection of *C. racemosa* 6 μ g / g + VP infection 10^6 CFU / ml per shrimp; (P3) injection of *C. racemosa* 9 μ g / g + VP infection 10^6 CFU / ml per shrimp; (T0) 24 hours after *C. racemosa* extract administration and prior to infection with *V. parahaemolyticus*; (T1) one day after being infected with *V. parahaemolyticus*; (T7) seven days after being infected with *V. parahaemolyticus*. Different letter notations in the same column showed significantly different results ($p < 0.05$).

Total bacteria on T1 or one day after *V. parahaemolyticus* infection, the K + treatment was significantly different ($p < 0.05$) against K-, P1, P2 and P3. The highest total bacteria shown in the positive control (K +) treatment, shrimp without *C. racemosa* extract and being infected with *V. parahaemolyticus* (26.20×10^4 CFU / ml). Whereas at T7 or seven days after being infected with *V. parahaemolyticus*, the total bacteria in treatment P1 was significantly different ($p < 0.05$) to treatment K-, K +, P2 and P3. Total bacteria decreased at T7, namely in the P3 treatment, namely 20.46×10^4 CFU / ml.

3.1.2 Survival Rates

The survival rates of white shrimp can be seen in Table 2. The survival rates of white shrimp during maintenance after administration of *C. racemosa* extract on T1 or one day after being infected with *V. parahaemolyticus* ranged from 83.33 (P1) to 100% (P2) and the results of the analysis of variability (ANOVA) showed significantly different results ($p < 0.05$). Shrimp survival in P2 treatment (100%) was significantly different from treatment P1 (83.33%), but not significantly different ($p >$

0.05) with treatment K-, K + and P3. The survival rates of white shrimp ranged from 56.67(K +) to 93.33% (P2) and showed significantly different results ($p < 0.05$) at T7.


Table 2. Survival rates of white shrimp (*L. vannamei*)

Treatment	Survival rates (%)		
	T0 \pm SD	T1 \pm SD	T7 \pm SD
K-	100 ^a \pm 0	90.00 ^{ab} \pm 0.00	90.00 ^a \pm 0.00
K+	100 ^a \pm 0	86.67 ^{ab} \pm 11.5	56.67 ^c \pm 11.5
P1	100 ^a \pm 0	83.33 ^b \pm 5.77	73.33 ^b \pm 11.5
P2	100 ^a \pm 0	100.00 ^a \pm 0.00	93.33 ^a \pm 5.77
P3	100 ^a \pm 0	90.00 ^{ab} \pm 10.0	86.67 ^{ab} \pm 5.77


Description: (K-) sterile PBS injection; (K +) infection with VP 10⁶ CFU / ml per shrimp; (P1) injection of *C. racemosa* 3 μ g / g + VP infection 10⁶ CFU / ml per shrimp; (P2) injection of *C. racemosa* 6 μ g / g + VP infection 10⁶ CFU / ml per shrimp; (P3) injection of *C. racemosa* 9 μ g / g + VP infection 10⁶ CFU / ml per shrimp; (T0) 24 hours after *C. racemosa* extract administration and prior to infection with *V. parahaemolyticus*; (T1) one day after being infected with *V. parahaemolyticus*; (T7) seven days after being infected with *V. parahaemolyticus*. Different letter notations in the same column showed significantly different results ($p < 0.05$).

3.1.3 Clinical Sign

The clinical sign emerged at the third day after infected by *V. parahaemolyticus*. Based on the results of the study, white shrimp infected with *V. parahaemolyticus* showed clinical sign, namely the hepatopancreas was pale, the intestines were empty, the uropods turned red and the gnats.



a. sick shrimp; b. normal shrimp



the uropod turned red and the gnats

Figure 1. Clinical sign of shrimp infected by *V. parahaemolyticus*

3.1.4 Water Quality

The data of water quality range value can be seen in Table 3.

Table 3. Water quality during the experiment

Parameter	Treatment					References
	K-	K+	P1	P2	P3	
Temperature (°C)	28-29.2	28.2-29.8	28.2-30.1	28.2-30.1	28.1-29	28-33(SNI. 2014)
Salinity (ppt)	15	15	15	15	15	15-32 (Hukomet <i>et al.</i> 2020)
pH	7-8	7-8	7-8	7-8	7-8	7.5-8.5(SNI. 2014)
DO (ppm)	3.95-5.22	4.14-5.42	4.27-6.26	4.05-5.69	3.87-5.33	>3.0(Mohanty <i>et al.</i> 2018)

Water quality as a supporting parameter in this study was measured in all treatments. Water quality supports the life of aquatic organisms. The water quality all treatments are in accordance with the references which is in the normal range. The temperature of 28-30.1 °C, salinity is 15 ppt, pH is 7-8 and Dissolved Oxygen is in the range of 3.87-6.26 ppm.

3.2 Discussion

Total bacteria were isolated from hepatopancreas of shrimp because of changes in clinical symptoms that appear is pale hepatopancreas. Acute hepatopancreatic necrosis disease (AHPND) or known as Early mortality syndrome (EMS) is known to attack this organ. Total bacteria on T1 (one day after

V. parahaemolyticus infection) decreased in all treatments with the administration of *C. racemosa* extract in P1, P2 and P3 treatments, namely 17.03×10^4 CFU / ml, 21.36×10^4 CFU / ml and 21.26×10^4 CFU / ml, respectively. Meanwhile, in the positive control (K +) treatment, shrimp without *C. racemosa* extract and being infected with *V. parahaemolyticus* showed the highest total bacteria (26.20×10^4 CFU / ml). This indicates that the sulfate polysaccharides contained in *C. racemosa* extract are starting to work as an immunostimulant which can reduce the total bacteria in the hepatopancreas. According to [11], the addition of sulfated polysaccharides from *Ulva lactuca* extract can increase the activity of the immune system in shrimp. This is due to an increase in total hemocytes and phagocytosis activity in shrimp after giving the extract. Hemocytes are a form of cellular defense from shrimp. Haemocyte can extinguish the infection caused pathogen by the synthesis and exocytosis of microbicidal protein bioactive molecule. According to [12], hemocytes consist of granular, semigranular and hyalin cells. Hyalin cells play a role in phagocytosis, semigranular cells play a role in melanization, encapsulation and coagulation, while granular plays a role in activating the ProPO system. The results showed that at T7 or 7 days after being infected with *V. parahaemolyticus* there was an increase in the total bacteria back in treatment P1 (24.90×10^4 CFU / ml) and P2 (24.26×10^4 CFU / ml) but the extract was given at a dose of $9 \mu\text{g} / \text{g}$ showed a lower total bacterial yield than K + (Table 1). This condition indicates that the immune system is back to normal conditions so that in this condition it is necessary to re-administer the *C. racemosa* extract (booster). According to [13], that giving immunostimulants through feed once every 3 days is more effective than giving once every 7 and 10 days. This is because the shrimp do not have memory cells and the short-term nature of the shrimp immune response. So it is necessary to provide periodic immunostimulants so that it can increased non-specific immune systems in shrimp.

The highest survival rates at T1 (one day after being infected with *V. parahaemolyticus*) was found in the treatment with the addition of $6 \mu\text{g} / \text{g}$ (P2) of *C. racemosa* extract which reached 100% and the lowest value was in P1 treatment (83.33%). On the 7th day, the survival rates value in P2 treatment (93.33%) was higher than that of K + treatment (56.67%). According to [14], $6\text{-}20 \mu\text{g} / \text{g}$ of *Spirulina platensis* extract injected into shrimp resulted in a higher survival rates compared to the control. This is because the polysaccharide sulfate content in seaweed can stimulate the shrimp immune system. Polysaccharides sulfate in *Gracilaria verrucosa* can stimulate the immune response of shrimp by increasing resistance to disease so that the survival rate of shrimp is high [15].

The results of the observation of clinical sign of shrimp infected with *V. parahaemolyticus* were among other things: the hepatopancreas was pale, the intestine was empty, the uropod turned red and the gnats. *Vibrio parahaemolyticus* is a halophilic bacterium that attacks white shrimp and causes lethargy, irregular swimming movements, empty intestine and hepatopancreas blanching [16] as shown in Figure 1. The results of measurement of water quality parameters are still in the optimum value and tolerance for the life of white shrimp. The temperature during maintenance is $28\text{-}30.1^\circ\text{C}$. this value is suitable for white shrimp culture, which is between $28 - 33^\circ\text{C}$ [17]. Salinity during the maintenance of white shrimp is 15 ppt. This value is still within the tolerance limit for the maintenance of white shrimp, namely salinity $15 - 32$ ppt [18]. The results of the average pH parameter measurement are $7 - 8$ where these results are in the optimum value for white shrimp culture, namely $7.5 - 8.5$ [17]. The average result of Dissolved Oxygen parameters is $3.87\text{-}5.33$ ppm. The optimum Dissolved Oxygen value in white shrimp culture is > 3 ppm [19].

4 Conclusion

The administration of *C. racemosa* extract by injected can affect the total bacteria in white shrimp after being infected with *Vibrio parahaemolyticus*. On the first day after infection, the total bacteria in the treatment with the administration of *C. racemosa* extract was lower than the positive control. The administration of *C. racemosa* extract can affect the survival rate of white shrimp and the highest was achieved in treatment with a dose of $6 \mu\text{g} / \text{g}$.

5 References

[1] Putri FM, Sarjito, Suminto 2013 J of *Aquacult Manag and Tech.* 2 102-112.

- [2] Food and Agriculture Organization 2017 Fishstat Plus Version 2.30. FAO Fisheries Department. Fishery Information. Data and Statistics Unit. <http://www.fao.org./fi/statist/FISOFT/FISHPLUS.asp>. 24 February 2017.
- [3] Kementerian Kelautan dan Perikanan Indonesia 2018 Diakses dari <http://www.kkp.go.id>
- [4] Zorriehzahra MJ and Banaederakhshan 2015 *Advanc Anim and Vet Sci*. 3 64-72.
- [5] Nunan L, Lightner D, Pantoja C, Gomez-Jimenez S 2014 *Aquatic Organ*. 111 81- 86.
- [6] Satyantini W H, Ananta K, dan Rahayu K 2016 *J Akuakul Indo*. 1(2) 120-129.
- [7] Pujiati. Sarjito dan Suminto 2013 *Jurnal Manaj Akuakul dan Tek*. 2(1) 66-74.
- [8] Xie JH. Wang ZJ, Shen MY, Nie SP, Gong B. Li H 2016 *Food Hydrocolloids*.53 7–15.
- [9] Huang L. Shen M. Morris GA. & Xie J 2019 *Tren in Food Scie & Tech*. 92 1-11.
- [10] Kusrieningrum 2008 *Dasar Perancangan Percobaan dan Rancangan Acak Lengkap. Fakultas Kedokteran Hewan (Universitas Airlangga: Surabaya)* p 92.
- [11]Suleman S. Sri A. and Ating Y 2018 *Res J of Life Sci*. 5(3) 156-162.
- [12] Cheng W, Ka YW and Chang CC 2017 *Fish & Shellfish Immun*. 68 92-101.
- [13] Ojerio VT. Corre VL, Toledo NA, Andrino-Felarca KGS, Nievaes LM & Traifalgar RFM 2018 *Aqua inter*. 26(1) 267-278.
- [14] Tayag C, Yong C, Chang C, Chyng H, Jiann C 2010 *J of Fish & Shellfish Immun*. 28 764-773.
- [15] Jasmanindar Y, Sukenda S, Alimuddin A, Junior MZ, and Utomo NBP 2018 *Omni-Akua*. 14(3).
- [16] Tawut R, Nantavadee B, Somluk A, Boonsirm W, Kanokpan W 2017 *J of Fish and Shellfish Immun*. 65 186-197.
- [17] SNI 8037.1 2014 *Udang Vaname (Litopenaeus vannamei. Boone 1931) Bagian 1: Produksi Induk Model Indoor* Badan Standardisasi Nasional (Jakarta) p 7.
- [18] Hukom V, Nielsen R, Asmild M, and Nielsen M 2020 *Ecologic Eco*. 176 106717.
- [19] Mohanty RK, Ambast SK, Panigrahi P, and Mandal KG 2018 *Aquaculture*. 485 210-219.

6 Acknowledgement

The author would like to thank the supervisors and lecturers, who helped this research. Futhermore thanks to Faculty of Fisheries and Marine, Universitas Airlangga, for conducting the seminar.