

BIOCHEMICAL AND CELLULAR ARCHIVES


Volume No. : 19 (2019) **Issue No. : Supp-02 (December)**

Table Of Contents

Article	Page No.
Table of Contents 19 #Supp#02 BCA : V.19 Supp#02 ABSTRACT PDF HTML	A
Alteplase in preventing posterior capsule opacity by plasminogen activator inhibitor-1 and type I collagen expression inhibition in a posterior capsule opacity model with fibrin reaction in vitro : A experimental laboratory study <i>Rina Wulandari, Indri Wahyuni, Arifa Mustika, Windu Purnomo and Nurwasis</i> ABSTRACT PDF HTML	4695
Erythrocyte-superoxide dismutase (SOD1) among elite combat sport athletes running an intensive training program and the association with micronutrient intake <i>Farapti Farapti, Sri Adiningsih and Mahmud Aditya Rifqi</i> ABSTRACT PDF HTML	4703
Fibrin glue (FG) attenuates fibrosis on human tenon's fibroblasts (HTFs) of glaucomatous eyes : Comparison with mitomycin C <i>Yuyun Rindiasuti, Evelyn Komarath, Helen Susilowati, Hendrian D. Soebagyo and Fedik A. Rantam</i> ABSTRACT PDF HTML	4713
Maternal cardiovascular risk in early and late onset of preeclampsia patients five years after labor : A comparative study <i>Ernawati, Hermanto Tri Joewono, Muhammad Ilham Aldika Akbar, Rozi Aditya aryananda, Manggala Pasca Wardhana, Khanisyah Erza Gumilar, Budi Wicaksono, Nareswari Cininta, Noor Assyifa Zuhjayanti, I Gde Rurus Suryawan, EryGumilar Dachlan and Aditiawarman</i> ABSTRACT PDF HTML	4721
Intracamerar infection of limbal mesenchymal stem cells secretome alleviate inflammation with delayed structural recovery on corneal endothelial cells in phacoemulsified rabbit eyes <i>Arief Fidiarto, Evelyn Komarath, Wimbo Sasono, Willy Sandhika and Hari Basuki Notobroto</i> ABSTRACT PDF HTML	4729
The resveratrol increase of hepatocyte growth factor (HGF) and epidermal growth factor (EGF) levels in Wharton's jelly mesenchymal stem cells (WJ-MSCs) secretome : Toward cell free therapy in dry eye disease (DED) <i>Evelyn Komarath, Yuyun Rindiasuti, Yohanes Widodo Wirohadjojo, Fedik A. Rantam, Aristika Dinaryati, Ni Made Inten Lestari and Cita R. S. Prakoewa</i> ABSTRACT PDF HTML	4737
A comparison of HBA1c in pregestational diabetes patients with obesity <i>A. Rizal D.J., T. J. Hermanto, Sri Murtiwi and Ernawati</i> ABSTRACT PDF HTML	4745
The effect of mesenchymal stem cells for the reduction of liver fibrosis through platelet derived growth factor-b regulation in rats <i>T. Sungkar, A. Putra, D. Lindarto and R. J. Sembiring</i> ABSTRACT PDF HTML	4749
Multipotency of rabbit (Oryctolagus cuniculus) skin -derived mesenchymal stem cells (SMSCs) induced to adipogenic progenitors in vitro <i>Deya Karsari, Aristika Dinaryanti, Nora Ertanti, Igo Syaiful Ihsan, Illyun Illa Renza, Fedik Abdul Rantam and Purwati</i> ABSTRACT PDF HTML	4755
The effect of virgin coconut oil (VCO) on the number of macrophages on post tooth extraction of wistar rats <i>Stanley Santosa Kamadjaja, Kharisma Nisa, AndrySentosa and Indra Mulyawan</i> ABSTRACT PDF HTML	4759
Correlation between the absolute eosinophil counts with cystatin C levels in hyperuricemia <i>Meri</i> ABSTRACT PDF HTML	4763
The effect of lime (Citrus aurantifolia Christm. Swingle) peel extract on osteoblast proliferation and formation matrix-trabecular bone in wistar rat tooth extraction sockets <i>Erlisa Saraswati Hasholan Hutabarat, Jeni Septranti Lolo Allo, Thomas Reynaldi Winanto, Indra Mulyawan and David Buntoro Kamadjaja</i> ABSTRACT PDF HTML	4767
In vitro cytotoxicity study of Platelet Rich Fibrin (PRF) membranes in Mesenchymal Stem Cells (SMSCs) derived from the skin of New Zealand white rabbits <i>Igo Syaiful Ihsan, Qorri Aynay, Deya Karsari, Aristika Dinaryanti, Nora Ertanti, Helen Susilowati, Purwati and Fedik Abdul Rantam</i> ABSTRACT PDF HTML	4771
Antifibrosis effect of fibrin glue on TGF- β and α -SMA expression in human tenon fibroblast, as wound healing model after trabeculectomy : An in vitro study <i>Sylva D. Taqryanka, Evelyn Komarath, Yuyun Rindiasuti, Helen Susilowati, Nurita T. Wijayanti and Fedik A Rantam</i> ABSTRACT PDF HTML	4777
Rabbit Skin-derived Mesenchymal Stem Cells (SMSCs) characterization and differentiation into osteogenic lineage <i>Nora Ertanti, Deya Karsari, Aristika Dinaryanti, Igo Syaiful Ihsan, Eryk Hendrianto, Rafizka Putri Iriandani, Fedik Abdul Rantam and Purwati</i> ABSTRACT PDF HTML	4783
Downregulation of the expression of MMP-3 and TIMP-1 by metformin regardless of TNF- α level in an in vitro lens capsule fibrosis model <i>Razzaqy, Reni Prastyani, Yulia Primitasari and Maftuchah Rochmanti</i> ABSTRACT PDF HTML	4791
Isolation and characterization of Skin derived Mesenchymal Stem Cell (SMSCs) from New Zealand rabbit, Oryctolagus cuniculus : A in vitro study <i>Aristika Dinaryanti, Deya Karsari, Nora Ertanti, Igo Syaiful Ihsan, Aida Ariyanti, Fedik Abdul Rantam, Aulani Aulanni and Purwati</i> ABSTRACT PDF HTML	4797
Amnion membrane-derived Mesenchymal Stem Cells (MSCs) migration and homing in a burn-wounded mouse model as tracked using IVIS imaging <i>Fedik A. Rantam, Abraham Ahmad Ali Firdaus, EviAwwalia, Anak Agung Ayu Meidiary, Purwati Purwati, Eryk Hendrianto, Igo Syaiful Ihsan and Helen Susilowati</i> ABSTRACT PDF HTML	4803
Correlation between obesity and type 2 diabetes mellitus using PPAR γ 2 gene varian <i>Evi Kurniawaty, Sumaryati Syukur, Yanwirasti and EtiYerize</i> ABSTRACT PDF HTML	4807
Immunogenicity evaluation of Polymorphonuclear (PMN) cells, IL-2, IL-10 and IgG of Biodegradable Porous Sponge Cartilage Scaffold (BPSCS), Adipose Derived Mesenchymal Stem Cell (ADMSC) and secretome in New Zealand white rabbits with cartilage defect : In <i>Brilliant Citra Wirashada, Dwikora Novembri Utomo, Purwati, Lukas Widhiyanto and Kuku Dwiputra Hernugrahanto</i> ABSTRACT PDF HTML	4811
Fibrin glue as an antifibrotic agent on Human Tenon Fibroblast (HTFs) through extracellular matrix expression <i>Nurita Tri Wijayanti, Evelyn Komarath, Yuyun Rindiasuti, Sylva Dranindi Taqryanka, Helen S. and Fedik A. Rantam</i> ABSTRACT PDF HTML	4819
A comparative study on the therapeutic potential of ocular and non-ocular stem cell secretome on alkali-induced limbal stem cell niche damage <i>Evelyn Komarath, Yuyun Rindiasuti, Yohanes Widodo Wirohadjojo, Delfitri Lutfi, Nurwasis, Fedik A. Rantam, Nora Ertanti and Cita R. S. Prakoewa</i> ABSTRACT PDF HTML	4825
Pre-clinical trial stem cell metabolites derived from placenta for wound healing <i>Purwati, M. Yulianto Listawan, P. Ardiah Iswanda, Fedik A. Rantam, Helen Susilowati, Eryk Hendrianto, H. Aff Nurul, Ernawati, Imam Susilo and R. Diah Puspita</i> ABSTRACT PDF HTML	4833
Synthesis and anti-cancer activity of copper(II) complex with 2, 4, 5-triphenyl-1H-imidazole ligand <i>T. H. Sucipto, F. Martak, H. Setyawati, N. Ertanti and I. K. Murwani</i> ABSTRACT PDF HTML	4839
Protein profile in self-made fibrin glue as promising biomaterial for wound healing modulation after trabeculectomy <i>Evelyn Komarath, Yuyun Rindiasuti, I. Ketut Sudiana, Cita R. S. Prakoewa and Fedik A. Rantam</i> ABSTRACT PDF HTML	4845
Pitstop-2 and dynasore exposure reduced trehalose entry into human dermal fibroblast <i>I. Kusuma, R. S. Hadi, B. Kiranadi and A. Boediono</i> ABSTRACT PDF HTML	4851
The effects of shiny bush (Peperomia pellucida L. Kunth) extract on fibroblast proliferation post-tooth extraction in wistar rats <i>R. Handhito Satriyo, Andreas Pratama Nugraha, Senitza Anisa Salsabilla, Indra Mulyawan, Coen Pramono D. and Muhammad Subhan Amir</i> ABSTRACT PDF HTML	4859
Snakehead fish extract (Channa striata) increase the number of fibroblasts cells post extraction tooth in wistar rats (Rattus norvegicus) <i>Surya Atmajaya, Dion Sandro Satrya, Retno Kathiningsih and Andra Rizqian</i> ABSTRACT PDF HTML	4863
Effects of lemongrass leaf on macrophages after tooth extraction of wistar rat <i>T. Devianna, A. Purnama, Y. Purnomo and A. Rizqian</i> ABSTRACT PDF HTML	4867
Bovine bone xenograft scaffold seeded with human umbilical cord mesenchymal stem cell to reconstruct segmental defect in dog's mandible : A preliminary study <i>David B. Kamadjaja, Nusdianto Triakoso and Purwati</i> ABSTRACT PDF HTML	4871
Grape seed extract increase osteoblast number in the post-extraction socket healing in rats <i>Olivia Jennifer Gunardi, A. Agustina Putri Kintan, R. Soesanto and Ni Putu Mira Sumarta</i> ABSTRACT PDF HTML	4877
A comparison of the effectiveness of packaged coconut water and UHT milk as a storage media for avulsion tooth <i>A. Setyawan, F. R. Fajriyany and N. P. M. Sumarta</i> ABSTRACT PDF HTML	4883
Intracamerar infection of Limbal Mesenchymal Stem Cells Conditioned Media (LMSCs-CM) improve clinical outcome with delayed on Na-K ATPase corneal endothelial pump recovery in phacoemulsified rabbit eyes <i>Eko Widayanto, Nurwasis, Dicky Hermawan, Arief Fidiarto, Paramita Putri, Yuyun Rindiasuti, Willy Sandhika, Igo Syaiful Ihsan, Hendrian D. Soebagdo and Evelyn Komarath</i> ABSTRACT PDF HTML	4889
Diagnostic of clinically virulent tuberculosis spondylitis with polimerase chain reaction (PCR) <i>Yunus Abdul Bari, Bambang Prijambodo, Ni Made Mertaniasih and Dwikora Novembri Utomo</i> ABSTRACT PDF HTML	4899
Metformin exerts antifibrosis effect on human lens epithelial cells through transforming growth factor-b inhibition <i>Dyah Purwita Trianggadewi, Indri Wahyuni, Deya Karsari, Maftuchah Rochmanti and Nurwasis</i> ABSTRACT PDF HTML	4911
Effects of lime (Citrus aurantifolia Christm. Swingle) peel extract on fibroblast proliferation and angiogenesis in rat's tooth extraction sockets <i>I Dewa Gde Agung Nanda Krismaya, Ramadhan Pramudya, Putri Ylvi Intan Sati and David Buntoro Kamadjaja</i> ABSTRACT PDF HTML	4917
Effect of natrium diclofenac in degrading fibrosis on Human Lens Epithelial Cell (HLEC) in congenital cataract eye to prevent posterior capsule opacification <i>Ryski Meirna, Diany Yogiartoro, Ratna Doemilah and Deya Karsari</i> ABSTRACT PDF HTML	4921

- Current Issue
- Subscription
- Archives **Open Access**
- Editorial Board
- Author Index(Current Vol.)
- Author Guidelines
- View Sample Article

Register For FREE 30 days trail



Published in ENGLISH

Useful Links

OUR SERVICES	SEARCH	PUBLISHER SERVICES	CUSTOMER SERVICES	SUPPORT SERVICES
Home	Quick Search	Publisher Search	User Panel	Contact Us
About Us	Advanced Search	Article Search	Free e-Journals	Post Article
Disclaimer	Title Search	Author Search		Add Your Journal
Privacy Policy	Alphabetical Search	Help		Feedback

BIOCHEMICAL AND CELLULAR ARCHIVES

Executive Council / Editorial Board

EDITOR-IN-CHIEF

Name	DR. P. R. YADAV
Dept./Organisation	Head - Department of Zoology
Address	D A V College Muzaffarnagar - 251 001, India
Email	yadavpry@rediffmail.com, submissionbca@gmail.com

MANAGING EDITOR

Name	MS. UMA YADAV
Address	606/8 South Civil Lines, Muzaffarnagar-251001, India
Email	jexpzool@gmail.com

EXECUTIVE EDITOR

Name	DR. AYAD ALKAIM
Dept./Organisation	Department of Chemistry
Address	Babylon University, Babylon, Iraq
Email	alkaimayad@gmail.com

CO-EDITOR

Name	DR. R. A. BALIKAI
Dept./Organisation	Dept. Ag. Entomology
Address	University of Ag. Sciences, Dharwad - 580 005, India.
Email	rabalikai@gmail.com

EDITOR

Name	DINESH KUMAR
Dept./Organisation	Dept Zoology
Address	Banaras Hindu University, Varanasi - 221 005, India
Email	dines1953@gmail.com

Name	MAYTHAM T. QASIM
Dept./Organisation	Dept Pathological Analysis
Address	College of Science, University of Thi-Qar, Iraq
Email	mtqr86@gmail.com

Name	MONOWAR A KHALID
Dept./Organisation	Dept Environ. Sci.
Address	Integral University, Lucknow - 226 026, India
Email	makhlid@iul.ac.in

Name	SURENDRA YADAV
Dept./Organisation	Dept Botany
Address	M. D. University, Rohtak - 124 001, India
Email	ssyadavindia@gmail.com

Name	ANDANG MIATMOKO
Address	Airlangga University, Campus C UNAIR, Mulyorejo-Surabaya, 60115, Indonesia
Email	andang-m@ff.unair.id

Name	ALEXANDER P. NUGRAHA
Dept./Organisation	Faculty of Dental Medicine
Address	Universitas Airlangga, Surabaya, Indonesia
Email	alexander.sandro11@gmail.com

Name	BHUPENDRA KUMAR
Dept./Organisation	Dept Zoology
Address	Banaras Hindu University, Varanasi - 221 005, India
Email	bhupendrakumar@bhu.ac.in

Name	RAMINDERJIT KAUR
Dept./Organisation	Dept. Cardiovascular & Metabolic Sci.
Address	Cleveland Clinic Main Campus, Ohio, USA
Email	kaurr8@ccf.org

Name	G. TRIPATHI
Dept./Organisation	Dept Zoology
Address	J N V University, Jodhpur-342 001, India
Email	drgst@rediffmail.com

Name	G. ARCHUNAN
Dept./Organisation	Dept Animal Science
Address	Bharathidasan University, Tiruchirappalli-620024, India
Email	garchu56@rediffmail.com

Name	RAED H. OGAILI
Dept./Organisation	Dept Maxillofacial Surgery
Address	College of Dentistry, Kerbala University, Iraq
Email	raedogaili@gmail.com

Name	VELAZHAHAN RETHINASAMY
Dept./Organisation	Dept Crop Sci.
Address	Sultan Qaboos University, Al-Khod, Muscat 123, Sultanate of Oman.

Name	MD. ABDULLAH-AL-MAMUN
Dept./Organisation	Dept. Fish Health Manag.
Address	Sylhet Ag. University, Sylhet - 3100, Bangladesh
Email	mamunff@gmail.com

Name	HASAN S. A. JAWAD
Dept./Organisation	Dept Animal Production
Address	Faculty of Ag. Eng. Sciences, Univ. Baghdad, Iraq
Email	dr.hassan198366@yahoo.com

Name	NIHAD KHALAWA TEKTOOK
Address	College of Medical and Health Tech., Middle Technical University, Iraq
Email	drnihadkhalawe@gmail.com

Name	P. PADMANABHAN
Address	Cognitive Neuroimaging Centre, Nanyang Technological Univ., Singapore -636921
Email	ppadmanabhan@ntu.edu.sg

Name	KRISHNA K. YADAV
Dept./Organisation	Faculty of Sci. & Technol.
Address	Madhyanchal Professional Univ. Bhopal - 462044 India
Email	envirokrishna@gmail.com

Name	KARRAR J HAMZAH
Dept./Organisation	Dept Vet. Internal & Prevent. Medicine
Address	AL-Qasim Green Univ. Babylon, Iraq.
Email	dr.karraraljanabi41@gmail.com


Name	SIVARAMAKRISHNA KOGANTI
Address	Carver College of Medicine, University of Iowa, CBRB, Iowa City, IA , USA.

TECHNICAL EDITOR

Name	PRASHANT KUMAR
Address	Poland
Email	prashantkbio@gmail.com

Name	ABHINAV PRAKASH
Address	India
Email	abhinav.prakash0844@gmail.com

- ▶ [Current Issue](#)
- ▶ [Subscription](#)
- ▶ [Archives](#)
- ▶ [Editorial Board](#)
- ▶ [Author Index\(Current Vol.\)](#)
- ▶ [Author Guidelines](#)
- ▶ [View Sample Article](#)
- ▶ [Register For FREE](#)
- ▶ [30 days trail](#)



Published in ENGLISH

Useful Links

OUR SERVICES
[Home](#)
[About Us](#)
[Disclaimer](#)
[Privacy Policy](#)

SEARCH
[Quick Search](#)
[Advanced Search](#)
[Title Search](#)
[Alphabetical Search](#)

[Publisher Search](#)
[Article Search](#)
[Author Search](#)
[Help](#)

CUSTOMER SERVICES
[User Panel](#)
[Free e-Journals](#)

SUPPORT SERVICES
[Contact Us](#)
[Post Article](#)
[Add Your Journal](#)
[Feedback](#)

INTRACAMERAL INJECTION OF LIMBAL MESENCHYMAL STEM CELLS SECRETOME ALLEVIATE INFLAMMATION WITH DELAYED STRUCTURAL RECOVERY ON CORNEAL ENDOTHELIAL CELLS IN PHACOEMULSIFIED RABBIT EYES

Arief Fidiyanto^{1*}, Evelyn Komaratih¹, Wimbo Sasono¹, Willy Sandhika² and Hari Basuki Notobroto³

¹Department of Ophthalmology, Faculty of Medicine, Airlangga University/Dr. Soetomo General Hospital, Surabaya, Indonesia.

²Department of Anatomic Pathology, Faculty of Medicine, Airlangga University/Dr. Soetomo General Hospital, Surabaya, Indonesia.

³Department of Biostatistic, Faculty of Public Health, Airlangga University, Surabaya, Indonesia.

*e-mail : arief.fidiyanto@gmail.com

(Received 11 August 2019, Revised 18 October 2019, Accepted 25 October 2019)

ABSTRACT : The objective of this research is to investigate the effects of intracameral injection of limbal mesenchymal stem cells (LMSCs) secretome on corneal endothelial cells density and caspase-3 expression after exposure to phacoemulsification ultrasound energy in rabbits. Twenty-four New Zealand white rabbits (n = 24 eyes) were randomly divided into 2 groups and exposed to phaco ultrasound. The control group was treated with intracameral injection of balanced salt solution (BSS), while intracameral injection of LMSCs secretome was utilized on the treatment group. Thereafter, the endothelial cells density was evaluated with specular microscopy before ultrasound exposure on the 3rd day with the cornea used to evaluate the expression of caspase-3 by immunohistochemistry staining. There were no signs of inflammation in both groups. However, exposure of ultrasound energy significantly reduced the corneal endothelial cell density in the control and treatment groups (-93.69 ± 139.04 ; $p = 0.04$ and -119.31 ± 85.41 , $p = 0.001$, respectively). Furthermore, the corneal endothelial cells density ($p = 0.592$) and caspase 3 expression after treatment for 3 days were indifferent in treatment group compared to control with $p = 0.625$; 2.17 ± 0.777 and 1.17 ± 0.322 , respectively. Intracameral injection of LMSCs secretome tends to alleviate ocular inflammation, improved the profile of endothelial cells density and corneal caspase-3 expression in rabbit eyes after exposure to ultrasound energy-phacoemulsification.

Key words : Limbal mesenchymal stem cells secretome, endothelial cells density, phacoemulsification, caspase-3.

INTRODUCTION

Most cataract extraction are currently performed with phacoemulsification techniques which uses ultrasound energy to fragment and emulsify the lens. With advances in the development of surgical devices and techniques, the safety and efficacy of phacoemulsification has rapidly increased. However, its use has been expanded to include cataracts with hard nucleus, thereby, leading to the need for greater ultrasound energy. This results in greater mechanical and thermal risk of corneal endothelial damage which ultimately triggers cell death through the apoptotic pathway. Some studies report that the endothelial cells loss of between 8.0% and 16.7% has a side effect of phacoemulsification. Damage and loss of corneal endothelial cells at some point tends to disrupt the function of the pump and tight junction which causes corneal edema and irreversible damage (Geffen *et al*, 2008; Bonnano, 2012; Takahashi, 2016; Alarfaj, 2017).

Corneal transplantation is currently the only effective treatment for corneal endotheliopathy due to its low regeneration ability. A national bullous keratopathy survey in Japan reported that cataract surgery was the most common cause of penetrating keratoplasty (PK) by 24.2%. Similarly, various other studies showed cataract surgery was ranked second after Fuchs endothelium dystrophy, thereby, making it the biggest cause of secondary endothelial damage. Keratoplasty has several disadvantages including limited corneal donors and the high level of postoperative endothelial cell damage at 49%. (Zavala *et al*, 2013; Zavala *et al*, 2017).

There are various *in vitro* studies aimed at triggering endothelial cell proliferation which showed a promising result. One of such methods is the secretion factors from stem cells also known as conditioned media, secretome, microvesicles or exosome which are expected to provide new therapeutic alternatives for endothelial damage. Furthermore, the stem cell conditioned medium which

have been studied have same benefits. Various studies regarding the secretion factors from stem cells showed that it has the ability to repair various damaged tissue or organ. (Zavala *et al*, 2013; Pawitan, 2014).

The mesenchymal stem cells secrete bioactive factors in conditioned medium contains, growth nutrients, as well as proinflammatory and anti-inflammatory cytokines popullary known as secretome. One of the secreted growth factors is fibroblast growth factor-2 (FGF-2) with various effects on corneal endothelium, include triggering cell proliferation, endothelial migration and morphological changes to mesenchymal or endothelium transition (EnMT). In addition, FGF-2 also inhibites apoptosis through the Phosphoinositide-3-kinase (PI3-K) pathway. Corneal limbal is source of mesenchymal stem cell, LMSCs secretome in the corneal endothelium capabel of improving cell density, while preventingthe apoptosis of damaged corneal endothelial cells (Bonnano, 2012; Pawitan, 2014; Vizoso *et al*, 2017).

METHODS

This is an experimental study using animals with the pre and post test control group design used to investigate the effect of LMSCs secretome on corneal endothelium cells density. Similarly, the post test is designed to evaluate the expression of caspase-3 in corneal endothelium after exposure to ultrasound energy and phacoemulsification. This study was conducted at the Center For Stem Cell Research and Development, Airlangga University in May 2019, with the animal protocol approved by the institutional review board and ethics committees of the Veterinary Faculty, Airlangga University. The inclusion criteria were New Zealand male white rabbits aged 12-18 months weighing 2.5 - 3.5 kg, while theexclusion criteria include animals declared by veterinarians to posses unhealthy eye conditions with transmittable diseases. Furthermore, the drop out criteria consists of unhealthy, dead, corneal perforation, vitreous prolapse, infected, and bleeding rabbits during and after surgery. Twenty-four New Zealand white rabbits (24 eyes) were randomly divided into 2 groups. The control group was intracamerally injected with 0.2ml BSS, while 0.2ml LMSCs secretome was injected in the treatment group.

Surgical procedures

Twenty four rabbits met the inclusion criteria (n=24 eyes), with anterior segment examination carried out using a handheld slit lamp. The rabbits were divided into two groups of 12 with anesthesia corneal endothelial injury model performed using the modified technique from Nemet *et al* (2007) with a NIDEK CV 9000R phacoemulsification machine. The disinfection of eyelid

werecarried out with 10% povidone-iodine. The field of operation was narrowed with sterile ducts and eye speculum mounted. The eyeball was disinfected with Povidone Iodine 5% and irrigated with BSS.

Endothelial injury model using phacoemulsification ultrasound exposure controlled variable parameters as follows: aspiration flow rate 25 ml/s, 70% power and 90cm bottle height using a single operator. The incision used were according to the 2.2 mm phacotip, with its position beveled up in the center of the anterior chamber without touching the endothelium and anterior lens capsule. The power was turned on and alternated every 10 seconds for 2.5 minutes (total time 5 minutes). The corneal incision wound was closed with stromal hydration and its impermeability checked. In the control group, intracameral injection of 0.2 ml BSS was administered, and 0.2ml LMSCs secretome produced and stored at -80°C by Komaratih *et al* (2019) used in the treatment group.

Clinical examinations were performed preoperatively and postoperatively on days 1, 2 and 3. The examination consisted of slit-lamp to check corneal clarity and intraocular inflammation. Endothelial cell density were examined using specular microscopy (CEM-530, Nidek Co., Ltd) on anesthetized rabbit at preoperative and postoperative day 3. Furthermore, all rabbits received topical tetracaine hydrochloride and levofloxacin every 3 hours (Neo Levo; Rohto Lab, Tokyo) on their right eyes during follow-up.

Also, each group was sacrificed with decapitation on the third postoperative 3. The rabbit eyes were enucleated, with blades and scissors used to remove their corneas which was further fixed in a 10% buffered formaldehyde (pH 7.0) for histology examination using caspase-3 antibody (bs-0087R, BIOSS, USA).

Histopathology

The rabbit corneal specimens were embedded in paraffin, sectioned at 4 ìm and stained with caspase-3 antibody (bs-0087R, BIOSS, USA) using standard techniques from the pathology anatomy laboratory of Dr. Soetomo General Hospital. The number of positively stained endothelial cells was assessed using a 40× objective and counted on 3 adjacent sections.

Statistical analysis

Statistical analysis was performed using the statistical package IBM SPSS Statistics for Windows (Version 19.0; IBM Corp., Armonk, NY, USA), while comparisons of the mean baseline values of the ECD between the two groups were performed using the independent t-test.

There was a significant difference in the preoperative and postoperative ECD between the two groups, with its density change assessed at postoperative time point using paired t-test. Furthermore, the positively stained endothelial cell counts from the histology sections were also compared using mann whitneytest with the data presented as mean \pm SD. The significance was set at $p < 0.05$ for this study.

RESULTS

Before surgery, axial length was measured with ultrasound biomicroscopy. Table 1 shows the average length of rabbit's axial in the control group was 15.41 mm and 15.83 mm in the treatment group. The data were normally distributed in the two groups (p control = 0.391 and p treatment = 0.851, $p > 0.05$) without differences ($p = 0.077$, $p > 0.05$) (Table 1).

Furthermore, there were no complications during the phacoemulsification procedure and incision wound impermeable with the anterior chamber was properly formulated after the procedure.

Endothelial Cell Density (ECD) Post Phacoemulsification Exposure

There were no differences in the evaluation time between the control and treatment groups (p day 0 = 0.11 and p day 3 = 0.072, $p > 0.05$). However, in the control group, there was a cell density of 2730.47 ± 139.72 on the first day, which was significantly different from day 3 by 2636.78 ± 209.81 ($p = 0.04$, $p > 0.05$). In the treatment group, the mean density of day 0 at 2582.86 ± 273.05 also showed a significant difference on day-3 with a density of 2463.56 ± 237.81 ($p = 0.001$, $p < 0.05$) (Table 2).

Each control and treatment group showed a decrease in endothelial cell density on day 3. Furthermore, the change in density in the control group was -93.69 ± 139.04 and the treatment group was 11119.31 ± 85.41 , with no significant differences between these two groups ($p = 0.592$, $p > 0.05$).

Expression of caspase-3 in corneal endothelial cells

The examination of caspase-3 expression was carried out by immunohistochemical staining using antibodies. It was calculated based on the number of cells from 3 visual fields under light microscope with 400 times magnification which appears brown in cytoplasm (Fig. 2).

Table 3 shows the mean expression of caspase-3 endothelial cells in each group with no statistical difference between the 2 groups. In the control group its expression was 1.17 ± 0.322 , while in the treatment group it was 2.17 ± 0.777 ($p = 0.625$, $p > 0.05$).

Table 1 : Axial Length Distribution.

Group	n	Axial Length				P
		\bar{x}	SD	Min	Max	
Control	12	15,41	0,60	14,61	16,79	0,077
Treatment	12	15,83	0,52	14,97	16,58	

DISCUSSION

This is the first study to evaluate the effect of intracameral injection of LMSC secretome to the corneal endothelial damage model. It was induced by phacoemulsification energy with endothelial cell loss and apoptosis assessed by density and caspase-3 expression, respectively. In addition, various methods have been used to induce corneal endothelial cell injury, which includes chemical, cryoinjury and mechanical injuries. The main advantage of the injury model of this study its similarity to what patients experience clinically. This phacoemulsification model also has the potential to be modified depending on the desired level. Rabbit's eye possess similar anatomy to the human eye, its corneal endothelium has a fairly high proliferation ability which diminishes with age. In addition, a fairly low proliferation capacity is shown in those over 12 months. Valdez-Garcia *et al* (2015) showed the highest mitotic corneal endothelial cell activity of rabbits at 3 months with 36 hours post-trauma. 12 months old rabbits, showed decreased mitotic activity and 18 months showed no mitotic activity in 72 hours post-trauma event. However, the difficulty faced is that the size of the eyeball is relatively smaller and the lens consumes more space in the eyeball which makes the anterior chamber narrower. (Davis, 1929; Gozkir *et al*, 1997).

Preoperative examination include anterior segment, axial length and endothelial cell density. In the anterior segment examination using a portable slit lamp, all rabbits were found in good condition. Axial length was measured to determine the bottle height for phacoemulsification machine. The mean in the control group was 15.41 mm, and 15.83 in the treatment group. The data were normally distributed (p control = 0.391 and p treatment = 0.851, $p > 0.05$) with no differences between the two groups ($p = 0.077$, $p > 0.05$). This is also consistent with the results of research by Bozkir *et al* (1997) where the average length of the rabbit eyeball was 15.12 ± 0.51 mm. Furthermore, the bottle height for phacoemulsification was set the same for each sample as high as 90 cm due to the homogenous data obtained.

Since the introduction of phaco-emulsification, significant improvement has been made to minimize corneal endothelial cell loss. Their reversible damages of the corneal endothelial are the available option used to







Post Operative (Day-)	Control Group	Treatment Group
1		
2		
3		

Fig. 1 : Anterior segment evaluation on post operative day-1, 2 and 3.




Fig. 2 : Caspase-3 antibody IHC examination, seen with a light microscope 400 times magnification in the control group (A and B) and the treatment group (C and D).

treat the transplantation dysfunction. Possible alternative therapies are conducted using strategies to save endothelial cells which are partially damaged from apoptosis and spurs regeneration. Endothelial cells do not proliferate, though its has the ability to conduct this process with a retained G1 cycle phase due to contact inhibition, TGF- α in aquos humor lacks effective stimulation from growth factors.

Corneal mesenchymal stem cells have been shown to have a positive effect in triggering proliferation with the ability to inhibit corneal endothelial cell loss in invitro and exvivo studies. According to previous studies, the therapeutic effect of mesenchymal stromal cells is mediated through secreted factors. Mesenchymal stem cell conditioned media containing secretion factors from

Table 2 : ECD distribution.

Group	ECD		p	Density change
	day 0	day 3		
Control	2730.47±139.72 (2464,67 – 2936,33)	2636,78±209,81 (2318,67 – 3053,33)	0,040	-93,69 ± 139,04 (-309,33 – 122,33)
Treatment	2582,86±273,05 (2076,33 – 3034,00)	2463,56 ± 237,81 (2010,00 – 2858,67)	0,001	-119,31 ± 85,41 (-276,00 – 27,00)
P	0,110	0,072		0,592

Table 3 : caspase-3 expression.

Group	n	Caspase-3 expression			p
		SD	Min	Max	
Control	12	1,17	0,322	0	0,625
Treatment	12	2,17	0,777	0	

stem cells are also expected to have the same effect in terms of triggering proliferation and preventing death. In this study the corneal endothelium cell densities throughout the sample averaged 2656.67 cells/mm², with a range between 2076.3 to 3034 cells/mm². Sailstad and Peiffer (1981) reported an average rabbit endothelium density of 2998 cells/mm², while Morita (1995) reported its densities were between 2180 and 3460 cells per mm². In the study by Brambatti, it varied between 2336 and 2771.50 cells per mm², while this study showed that the exposure of phacoemulsification ultrasound significantly decreases corneal endothelium cell density. A study by Nemet *et al* (2007) reported a decrease in endothelial cell density up to 551 cells/mm², however, in this study the average cell decline was 93.69 cells/mm² and 119.31 cells/mm² in the control and treatment groups. This difference is mainly due to the varying types of machines used, which showed the safety property of machines against corneal endothelium. Although, it is statistically significant in each group, it seems the damage has not been severe enough to evaluate the therapeutic effect of intracameral LMSCs secretome.

Another factor which influences this study is the measurements, which were limited in the central area, while a research conducted by Kim *et al* (2015) observed changes in corneal endothelial parameters measured in central, superior, temporal, inferior and nasal areas. This showed that corneal endothelial wound healing is mainly observed in the temporal area ie in the primary incision with insignificant changes in the central area do. Usta *et al* (2017) investigated the effect of autologous serum in the anterior chamber of the rabbit’s eye after damaging the phacoemulsification ultrasound energy, which showed no significant difference between the treatment and control groups in terms of endothelium cell density on

daily measurements from day 0, 1 and 7. The difficulty in measuring specular microscopy on rabbit eyeballs was one of the causes of corneal measure in the same area being examined due to the absence of visual fixation and a headrest design not suitable for rabbits. Furthermore, this has the potential to cause a shift in the area examined at different times thereby reducing the quality (Sailstad and Peiffer, 1981; Morita, 1995; Albuquerque *et al*, 2015; Coyo *et al*, 2016).

The effect of LMSCs secretome on endothelial cell density and caspase-3 expression has not been examined using in vivo. Although there are positive results associated with the application of LMSCs secretome, there were no significant differences between the groups receiving LMSCs secretome and BSS as a control in terms of \bar{x} changes in endothelial cell density and caspase-3 expression. In this study, LMSCs secretome proved to have safety property and intracamerally utilized to determine any significant inflammation sign during observation. Nakahara *et al* (2013) and Sun *et al* (2017) showed that giving MSCs secretome provides positive results on invitro corneal endothelial cell proliferation. Corneal endothelial cell cultures originating from human donors provided additional bone marrow MSCs secretome which showed positive effects on corneal endothelial culture in terms of maintaining hexagonal cell morphology, maintaining functional proteins through ZO-1 and Na/K-ATPase expression by increasing cell proliferation through Ki-67 expression by 2 times compared to the control Group.

According to a study by Lu *et al* (2010), human corneal endothelial cells in vitro proliferate on day 2, formed in polygonal groups administered with MSCs secretome and is subcultured to passage 6 while maintaining its shape. While in the culture group without media, proliferation occurred on the 3rd day and in the subculture it reached the 2nd passage. Endothelial cells in both groups showed a function phenotype by expressing ZO-1 and Na-K-ATPase. Furthermore, Sun *et al* (2017) made a model of corneal endothelium dysfunction using rabbit eyes scratched with endothelium

and administered with MSCs secretome therapy. On the 7th day, it was cleared and its thickness rapidly decreased. Cell density was 2898 cells/mm² in the normal group, 2008 cells/mm² in the conditioned media group in day-3, and 2654 cells/mm² in day 7. In contrast, the control group in which the endothelium was damaged had severe edema with the time factor examination noted. Although in the study of Lu *et al* (2010) the endothelial cells were able to show proliferation on day 2, it wasn't enough to provide significant results on endothelium cell density on day 3. Therefore, an examination with longer time period is needed to ensure endothelial cell proliferation affects cell density.

MSC and MSCs secretome are known to have positive effects in decreasing cell apoptosis and increasing its survival rate. Dasari *et al* (2007) examined its effect on the regulation of apoptosis in a mouse model of spinal cord injury and analyzed that the administration reduced apoptosis through caspase-3 pathway regulation. Rouhbakhshzaeri *et al* (2018) investigated its effect on endothelial cell loss after phacoemulsification in the ex vivo pig eye showed that its incubation with MSC-CM significantly reduced the endothelial cell loss rate by an average of 1.29% ± 0.91% compared to the control group with 5.33% ± 3.24% (p < 0.05). Studies which stimulate corneal endothelial cells to proliferate are generally in vitro and ex vivo with the aqueous humor factor ruled out. Conversely, under normal conditions in aqueous humor, the TGF- β is insufficient to hold endothelial cells out of the G1 phase. Liu *et al* (2018) examined changes in aqueous humor after phacoemulsification and obtained a significant increase in malondialdehyde, prostaglandin E2 and interleukin-1. Other researchers found that the TGF- β levels increased significantly after the phacoemulsification action. (Zhang *et al*, 2016). Liu (2002) showed a circadian rhythm effect on differences in TGF- β 2 and FGF-2 levels in rabbit humor aqueous. The total levels of TGF- β 2, active TGF- β 2 and bFGF in aqueous humor changed over a 24-hour period. TGF- β 2 and bFGF levels increase between 4am and 8am. Active TGF- β 2 levels increase between 8 pm and midnight. The same pattern of change was found when the aqueous humor sample was taken in constant dark conditions. In vitreous humor, total TGF- β 2 is active, while bFGF remains unchanged for 24 hours. The influence of aqueous humor and circadian rhythm affects the FGF contained in LMSCs secretome, which was analyzed at the end of the procedure.

Intraocular surgery and anterior segment inflammation was used to reduce the risk associated with endothelial cell density, which was closely related to changes in the levels of cytokines in aqueous humor. A study

by Yaguchi *et al* (2017), which analyzed the relationship between endothelial cell density and increased aqueous humor cytokines showed that endothelial cell was negatively correlated with increased levels of aqueous humor cytokines especially cytokines IL-1 α , IL-4, IL-13, MIP-1 α , TNF- α and E-selectin. Conversely, the administration of conditioned media which is cytokines rich changes the levels of aqueous humor cytokines and affects corneal endothelial cells. The effect depends on the cytokines contents using positive growth factors for corneal endothelium such as FGF-2, which triggers endothelial cells to exit the G1 phase and proliferate. It was observed from in vitro studies that the FGF-2 signaling pathway in promoting proliferation is conducted through the phosphorylation of p27 on Ser10, Thr187, PI3-K and ERK1/2 pathways. The PI3K pathway, leads to activation of Akt, which plays an important role in cell survival, inhibits proapoptotic cytokines such as Bcl-2, Bad, Bax, caspase-9 and activation of caspase-3 as an effector of apoptosis. (Jeong *et al*, 2007; Lee *et al*, 2007; Lee *et al*, 2011; Nakahara *et al*, 2013; Zhang *et al*, 2015). The secretome utilized in this research are expected to possess high FGF-2 content according to research from Bhang *et al* (2014), which showed that using the three-dimensional (3D) spheroid culture method in human adipose stem cells produced conditioned media with VEGF concentrations, FGF-2, as well as HGF and chemokines which were 23 to 27 times higher than the conditioned media produced with conventional monolayer cultures. However, this study failed to take measurements of FGF and other cytokine levels, therefore, further research is needed to ensure the levels are contained and capable of affecting corneal endothelial cells.

Its limitation is associated with, the single observation periods on the 3rd day, while the administration of LMSCs secretome was at the end of the procedure. Specular microscope measurements were carried out in the central area in order not to describe the state of corneal endothelial cell density in other areas. Furthermore, the cytokine level of LMSCs secretome and aqueous humor after the procedure were not assessed.

REFERENCES

- Alarfaj K, Al Osaily A and Magliyah M (2017) The Effect of Phacoemulsification Surgery on Corneal Endothelium. *Journal of Cells* 2, 1-5.
- Bhang S, Lee S, Shin J, Lee T, Jang H and dan Kim B (2014) Efficacious dan Clinically Relevant Conditioned Medium of Human Adipose-derived Stem Cells for Therapeutic Angiogenesis. *Molecular Therapy* 22(4), 862-872.
- Block G, Ohkouchi S, Fung F, Frenkel J, Gregory C, Pochampally R, DiMattia G, Sullivan D and dan Prockop D (2009) Multipotent stromal cells are activated to reduce apoptosis in part by

- upregulation dan secretion of stanniocalcin-1. *Stem Cells* **27**(3), 670-681.
- Bozkir G, Bozkir M, Dogan H, Aycan K and Güler B (1997) Measurements of axial length and radius of corneal curvature in the rabbit eye. *Acta Med Okayama* **51**(1), 9-11.
- Bourne W (2003) Biology of the corneal endothelium in health and disease. *Eye* **17**(8), 912-918.
- Branch M, Hashmani K, Dhillon P, Jones D, Dua H and dan Hopkinson A (2012) Mesenchymal Stem Cells in the Human Corneal Limbal Stroma. *Investigative Ophthalmology & Visual Science* **53**(9), 5109.
- Bulatovic I, Ibarra C, Österholm C, Wang H, Beltrán-Rodríguez A, Varas-Godoy M, Månsson-Broberg A, Uhlén P, Simon A and dan Grinnemo K (2015) Sublethal Caspase Activation Promotes Generation of Cardiomyocytes from Embryonic Stem Cells. *PLOS ONE* **10**(3), e0120176.
- Chen S, Hayashida Y, Chen M, Xie H and Tseng S (2011) A New Isolation Method of Human Limbal Progenitor Cells by Maintaining Close Association with Their Niche Cells. *Tissue Engineering Part C: Methods* **17**(5), 537-548.
- Dasari V, Spomar D, Cady C, Gujrati M, Rao J and dan Dinh D (2007) Mesenchymal Stem Cells from Rat Bone Marrow Downregulate Caspase-3-mediated Apoptotic Pathway After Spinal Cord Injury in Rats. *Neurochemical Research* **32**(12), 2080-2093.
- Ferreira JR, Teixeira G Q, Santos S G, Barbosa MA, Almeida P G and Gonçalves R M (2018) Mesenchymal Stromal Cell Secretome: Influencing Therapeutic Potential by Cellular Pre-conditioning. *Front. Immunol.* **9**, 2837. doi: 10.3389/fimmu.2018.02837
- Geffen N, Topaz M, Kredy-Farhan L, Barequet I, Farzam N, Assia E and dan Savion N (2008) Phacoemulsification-induced injury in corneal endothelial cells mediated by apoptosis: In vitro model. *Journal of Cataract & Refractive Surgery* **34**(12), 2146-2152.
- Gottardi C, Arpin M, Fanning A and Louvard D (1996) The junction-associated protein, zonula occludens-1, localizes to the nucleus before the maturation and during the remodeling of cell-cell contacts. *Proceedings of the National Academy of Sciences* **93**(20), 10779-10784.
- Guo H, Ren H, Liang S, Ji Y, Jiang H, Zhang P and dan Du M (2018) Phosphatidylinositol 3-Kinase/Akt signal pathway resists the apoptosis and inflammation in human extravillous trophoblasts induced by *Porphyromonas gingivalis*. *Molecular Immunology* **104**, 100-107.
- Gwon A (2008) The Rabbit in Cataract/IOL Surgery. *Animal Models in Eye Research*, 1st ed. San Diego: Elsevier, pp.184-204.
- Holan V, Trosan P, Ceika C, Javarkova E, Zaricova A, Hermankova B, Chudickova M and Cekova J (2015) A comparative study of the therapeutic potential of mesenchymal stem cells and limbal epithelial stem cells for ocular surface reconstruction. *Stem Cells Translational Medicine* **4**, 1-12.
- Hoppenreijns V P, Pels E and Vrensen G F (1994) Basic Fibroblast growth factor stimulates corneal endothelium cell growth and endothelium wound healing of human corneas. *Invest Ophthalmol Vis Sci.* **35**, 931-944.
- Jirsova K (2018) *Light and Spekular Microscopy of the Cornea*. Springer.
- Joyce N (2003) Proliferative capacity of the corneal endothelium. *Progress in Retinal and Eye Research* **22**(3), 359-389.
- Joyce N, Zhu C and Harris D (2009) Relationship among Oxidative Stress, DNA Damage, and Proliferative Capacity in Human Corneal Endothelium. *Investigative Ophthalmology & Visual Science* **50**(5), 2116.
- Kim T Y, Kim W I, Smith R E and Kay E P (2001) Role of p27Kip1 in cAMP- and TGF- β 2-Mediated Antiproliferation in Rabbit Corneal Endothelial Cells. *Invest Ophthalmol Vis Sci.* **42**, 3142-3149.
- Kim D, Wee W and Hyon J (2015) The pattern of early corneal endothelial cell recovery following cataract surgery: cellular migration or enlargement?. *Graefe's Archive for Clinical and Experimental Ophthalmology* **253**(12), 2211-2216.
- Komarathih E, Rindiastuti Y, Susilowati H, Wijayanti N T, Dranindi S, Primitasari Y, Nurwasis, Artini W, Rantam F A and Prakoeswa C (2019) Antifibrotic Effects of Limbal Mesenchymal Stem Cells-Conditioned Media (LMSCs-CM) on Human Tenon's Fibroblasts (HTFs) in Glaucomatous Eyes: Comparison with Mitomycin C. *Sch Acad J Biosci.* **7**(1), 4-12.
- Komarathih E, Suhendro G, Eddyanto, Purwati, Prakoeswa C R S, Rindiastuti Y, Hendrianto E, Susilowati H and Rantam F A (2019) Fibrin Glue Maintain Limbal Mesenchymal Stem Cells Survival: A Novel Cell Based Therapy Strategy for Modulating Wound Healing After Trabeculectomy. *Saudi J Biomed Res.* **4**(1), 11-17.
- Lee J G and Kay E P (2006) FGF-2-induced wound healing in corneal endothelium cells requires Cdc42 activation and Rho inactivation through the phosphatidylinositol 3-kinase pathway. *Invest Ophthalmol Vis Sci.* **47**, 1376-1386.
- Lee J G and Kay E P (2006) Cross-talk among Rho GTPases acting downstream of PI 3-kinase induces mesenchymal transformation of corneal endothelium cells mediated by FGF-2. *Invest Ophthalmol Vis Sci.* **47**, 2358-68.
- Lee J G and Kay E P (2007) Two populations of p27 use differential kinetics to phosphorylate Ser-10 and Thr-187 via phosphatidylinositol 3-Kinase in response to fibroblast growth factor-2 stimulation. *J Biol Chem.* **282**, 6444-6454.
- Lee J G, Ko M K and Kay E P (2012) Endothelium mesenchymal transformation mediated by IL-1 beta-induced FGF-2 in corneal endothelium cells. *Exp Eye Res.* **95**, 35-39.
- Li G, Zhou Y, Xie H, Chen S and Tseng S (2012) Mesenchymal stem cells derived from human limbal niche cells. *Investigative ophthalmology & visual Science* **55**(9), 5686-5697.
- Liu J (2002) Circadian variations of transforming growth factor- β 2 and basic fibroblast growth factor in the rabbit aqueous humor. *Current Eye Research* **24**(1), 75-80.
- Liu Y, Setiawan M, Ang M, Yam G and dan Mehta J (2019) Changes in aqueous oxidative stress, prostaglandins and cytokines: Comparisons of low-energy femtosecond laser-assisted cataract surgery versus conventional phacoemulsification. *Journal of Cataract & Refractive Surgery* **45**(2), 196-203.
- Lu J, Lu Z, Reinach P, Zhang J, Dai W, Lu L and Xu M (2006) TGF- β 2 inhibits AKT activation and FGF-2-induced corneal endothelial cell proliferation. *Experimental Cell Research* **312**(18), 3631-3640.
- Lu X, Chen D, Liu Z, Li C, Liu Y, Zhou J, Wan P, Mou Y and Wang Z (2010) Enhanced Survival *in vitro* of Human Corneal Endothelial Cells Using Mouse Embryonic Stem Cell Conditioned Medium. *Molecular Vision* **16**, 611-622.
- Matsubara M and Tanishima T (1983) Wound-healing of corneal

- endothelium in monkey: an autoradiographic study. *Jpn J Ophthalmol.* **27**(3), 444-450.
- Nakahara M, Okumura N, Kay E, Hagiya M, Imagawa K, Hosoda Y, Kinoshita S and Koizumi N (2013) Corneal Endothelial Expansion Promoted by Human Bone Marrow Mesenchymal Stem Cell-Derived Conditioned Medium. *PLoS ONE* **8**(7), e69009.
- Nemet A, Assia E, Meyerstein D, Meyerstein N, Gedanken A and Topaz M (2007) Protective effect of free-radical scavengers on corneal endothelial damage in phacoemulsification. *Journal of Cataract & Refractive Surgery* **33**(2), 310-315.
- Pawitan J A (2014) Prospect of stem cell conditioned medium in regenerative medicine. *Biomed Research International* **965849**, 1-14.
- Polisetty N, Fatima A, Madhira S L, Sangwan V S and Vemuganti G K (2008) Mesenchymal cells from limbal stroma of human eye. *Molecular Vision* **14**, 431-442.
- Redza-Dutordoir M and dan Averill-Bates D (2016) Activation of apoptosis signalling pathways by reactive oxygen species. *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research* **1863**(12), 2977-2992.
- Rouhbakhshzaeri M, Rabiee B, Azar N, Ghahari E, Putra I, Eslani M and Djalilian AR (2018) New ex vivo model of corneal endothelial phacoemulsification injury and rescue therapy with mesenchymal stromal cell secretome. *J Cataract Refract Surg.* **1**, 1-6.
- Roy O, Beaulieu Leclerc V, Bourget J, Theriault M and Proulx S (2015) Understanding the Process of Corneal Endothelial Morphological Change *In Vitro*. *Investigative Ophthalmology & Visual Science* **56**(2), 1228-1237.
- Sailstad D and dan Peiffer R (1981) Specular microscopic observations of the corneal endothelium in the normal rabbit. *Laboratory Animals* **15**(4), 393-395.
- Sarvar D P, Shamsasenan K and Akbarzadehlaleh P (2016) Mesenchymal stem cell-derived exosomes: New opportunity in cell-free therapy. *Adv Pharm Bull.* **6**(3), 293-299.
- Schulze S, Bertelmann T, Manojlovic I, Bodanowitz S, Irle S and Sekundo W (2015) Changes in corneal endothelium cell characteristics after cataract surgery with and without use of viscoelastic substances during intraocular lens implantation. *Clinical Ophthalmology* p.2073.
- Sun P, Shen L, Zhang C, Du L and dan Wu X (2017) Promoting the expansion and function of human corneal endothelial cells with an orbital adipose-derived stem cell-conditioned medium. *Stem Cell Research & Therapy* **8**(1).
- Studzinski (1999) *Apoptosis A practical approach*. Oxford University Press, New York
- Sundquist T, Moravec R, Niles A, O'Brien M and dan Riss T (2006) Timing Your Apoptosis Assay. *Cell Notes* **16**, 18-21.
- Takahashi H (2016) Corneal Endothelium and Phacoemulsification. *Cornea* **35**, S3-S7.
- Usta G, Ogurel R, Onaran Z, Pekcan Z, Ogurel T, Karsli B, Klinigi G H and Hastanesi G D (2017) Autologous serum effect on corneal endothelial damage in the phacoemulsification rabbit model. *Journal of Clinical and Analytical Medicine* **8**(Supplement 4), 330-335.
- Valdez-Garcia J, Lozano-Ramirez J and Zavala J (2015) Adult white New Zealand rabbit as suitable model for corneal endothelial engineering. *BMC Research Notes* **8**(1), 28.
- Van den Bogerd B, Dhuhghaill S, Koppen C, Tassignon M and Zakaria N (2018) A review of the evidence for in vivo corneal endothelium regeneration. *Survey of Ophthalmology* **63**(2), 149-165.
- Vizioso F J, Eiro N, Schneider J and Piere R (2017) Mesenchymal stem cells secretome: toward cell-free therapeutic strategies in regenerative medicine. *Molecular Sciences* **18**(1852), 1-24.
- Zavala J, Hernández-Camarena J C, Loya-García D, Tamez A and Valdez-García J E (2017) Wound Healing in the Corneal Endothelium. *Researchgate* **1**, 1-38.
- Zhang K, Zhu X, Chen M, Sun X, Yang J, Zhou P and dan Lu Y (2016) Elevated Transforming Growth Factor- β 2 in the Aqueous Humor: A Possible Explanation for High Rate of Capsular Contraction Syndrome in High Myopia. *Journal of Ophthalmology* pp.1-6.
- Zhang W, Chen J, Fu Y and Fan X (2015) The signaling pathway involved in the proliferation of corneal endothelium cells. *Journal of Receptor and Signal Transduction* pp.1-7.
- Zhang X, Hu M, Pan K, Li H and Liu R (2016) 3D Spheroid culture enhances the expression of antifibrotic factors in human adipose-derived MSC and improves their therapeutic effects on hepatic fibrosis. *Stem Cells International* **462607**, 1-9.
- Zhang Y, Sun H, Liu Y, Chen S, Cai S, Zhu Y and Guo P (2016) The limbal epithelial progenitors in the limbal niche environment. *Int. J. Med. Sci.* **13**, 835-841.