

2018 Mitochondrial Genome Scolopsis

by Sapto Andriyono

Submission date: 20-Jan-2023 03:48PM (UTC+0800)

Submission ID: 1995904206

File name: ciliata_and_its_phylogenetic_relationship_in_genus_Scolopsis.pdf (916.08K)

Word count: 1286

Character count: 7289



Complete mitochondrial genome of saw-jawed monacle bream *Scolopsis ciliata* and its phylogenetic relationship in genus *Scolopsis*

Sapto Andriyono, Sinar Pagi Sektiana, Md. Jobaidul Alam & Hyun-Woo Kim

To cite this article: Sapto Andriyono, Sinar Pagi Sektiana, Md. Jobaidul Alam & Hyun-Woo Kim (2019) Complete mitochondrial genome of saw-jawed monacle bream *Scolopsis ciliata* and its phylogenetic relationship in genus *Scolopsis*, *Mitochondrial DNA Part B*, 4:1, 393-394, DOI: [10.1080/23802359.2018.1547149](https://doi.org/10.1080/23802359.2018.1547149)

To link to this article: <https://doi.org/10.1080/23802359.2018.1547149>



© 2019 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



Published online: 11 Jan 2019.



[Submit your article to this journal](#)



Article views: 503



[View related articles](#)



[View Crossmark data](#)



[Citing articles: 1 View citing articles](#)

Complete mitochondrial genome of saw-jawed monacle bream *Scolopsis ciliata* and its phylogenetic relationship in genus *Scolopsis*

Sapto Andriyono^{a,b} , Sinar Pagi Sektiana^a, Md. Jobaidul Alam^a and Hyun-Woo Kim^{a,c} 

^aInterdisciplinary Program of Biomedical, Mechanical, and Electrical Engineering, Pukyong National University, Busan, Republic of Korea; ^bDepartment of Marine, Faculty of Fisheries and Marine, Universitas Airlangga C Campus Jl, Surabaya, Indonesia; ^cDepartment of Marine Biology, Pukyong National University, Busan, Republic of Korea

ABSTRACT

The complete mitochondrial genome of the saw-jawed monacle bream, *Scolopsis ciliata* was determined by the next generation sequencing (NGS) technique. The circular mitogenome of *S. ciliata* (16,733 bp) encoded the canonical 37 genes including 13 proteins, 22 *tRNA* genes, 2 rRNAs (12S rRNA and 16S rRNA), and two non-coding regions; the origin of light strand replication (OL) and the putative control region (D-Loop). The gene arrangement of the *S. ciliata* mitogenome was identical to its relative, *Scolopsis vosmeri*. Phylogenetic analysis based on the full mitochondrial genome sequences showed that *S. ciliata* is most closely related to *S. vosmeri* with 83% nucleotide sequence identity. The mitogenome information of *S. ciliata* would be the useful information to understand the evolutionary relationship of fishes in genus *Scolopsis*.

ARTICLE HISTORY

Received 22 October 2018
Accepted 2 November 2018

KEYWORDS

Next generation sequencing; *Scolopsis ciliata*; mitogenome; monacle bream

Approximately 17 species are currently known in the genus *Scolopsis* and most of which inhabit widely in Indo-West Pacific region (Russell 1990). The *Scolopsis ciliata* is one of economically important marine fishes in Indonesia, which found on the sandy fringe of coastal and lagoon reefs (Kuitert and Tonozuka 2001; Allen and Erdmann 2012). Currently, phylogenetic relationship of fishes in genus *Scolopsis* is still not clearly established (Chen and Zheng 1987; Russell 1990) and only one mitochondrial genome sequence in the genus, *Scolopsis vosmeri* is currently reported (Wu et al. 2017). In this study, we report a mitogenome sequence of *S. ciliata*, which would be the useful information to understand the evolutionary relationship of fishes in genus *Scolopsis*.

The next-generation sequencing (NGS) platform was adopted to read the complete mitochondrial genome sequence of *S. ciliata*. The specimen was collected from the coastal water in Malang, East Java, Indonesia (8°26'05,65"S 112°40'55,31"E) and deposited at the Ichthyology Laboratory, Universitas Airlangga, Indonesia. Species of the specimen was identified by both its morphological characteristics and DNA sequence identity in COI region to the database (GenBank Accession number: KY362945). Purified mitochondrial DNA by the commercial kit (Abcam, Cambridge, UK) was further fragmented into the size for library preparation (~350 bp) by Covaris M220 Focused-Ultrasonicator (Covaris Inc., Woburn, MA). A library was constructed by TruSeq[®] RNA library preparation kit V2 (Illumina, San Diego, CA) and its quality and the quantity was analysed

by 2100 Bioanalyzer (Agilent Technologies, Santa Clara, CA). High-throughput sequencing was performed by Illumina MiSeq system (2 × 300 bp).

The complete mitochondrial genome of *S. ciliata* was circular and 16,733 bp in length (GenBank Number: MH995531). It contained the typical 37 genes coding for two ribosomal RNAs (12S rRNA and 16S rRNA), 22 tRNAs, 13 polypeptides, and two non-coding regions; the origin of light strand replication (OL) and the putative control region (D-Loop). OL was located between tRNA-Asn and tRNA-Cys at WANCY cluster, while D-Loop (986 bp) was identified between tRNA-Pro and tRNA-Phe. Besides COX1(GTG) and ATP6 (ATC), 11 protein-coding genes begin with typical ATG. Incomplete stop codons were shown in *ND2*, *COX2*, *COX3*, *ND3*, *ND4*, and *CYTb* genes. The length of tRNAs ranged between 69 and 76 bp, which were located as three conserved tRNA cluster (IQM, WANCY, and HSL)(Satoh et al. 2016). Except for tRNA-Ser, 21 tRNAs were predicted to form the typical clover-leaf secondary structures according to ARWEN (Laslett and Canbäck 2008) which has the typical clover-leaf secondary structures except for tRNA-Ser. Based on the phylogenetic analysis of the mitogenomes by MEGA7 version 7 software (Kumar et al. 2016), *S. ciliata* was most closely related to *S. vosmeri* (GenBank No. NC029489) with 83% identity forming a genus *Scolopsis* cluster, which is distinct from fishes in genus *Nemipterus* (Figure 1). The mitogenome information of *S. ciliata* would help us understand the evolutionary relationship in genus *Scolopsis*.

CONTACT Hyun-Woo Kim  kimhw@pknu.ac.kr  Department of Marine Biology, Interdisciplinary Program of Biomedical, Mechanical and Electrical Engineering, Pukyong National University, Busan 48513, Republic of Korea

© 2019 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.




Figure 1. Phylogenetic tree of *Lutjanus vitta* within Lutjanidae. Phylogenetic tree of *Scolopsis ciliata* complete genome was constructed by MEGA7 software with Minimum Evolution (ME) algorithm with 1000 bootstrap replications. GenBank Accession numbers were shown followed by each scientific name.

Disclosure statement

The authors report that they have no conflicts of interest. The authors alone are responsible for the content and writing of the article.

Funding

This work was supported by an educational grant from the Indonesian Endowment Fund for Education (LPDP) scholarship under Beasiswa Unggulan Dosen Indonesia-Luar Negeri (BUDI-LN) batch I 2016 [Number: PRJ-3635/LPDP.3/2016] and a part of the project titled "Long-term change of structure and function in marine ecosystems of Korea", funded by the Ministry of Oceans and Fisheries, Korea.

ORCID

Sapto Andriyono  <https://orcid.org/0000-0002-2566-1636>
Hyun-Woo Kim  <http://orcid.org/0000-0003-1357-5893>

References

- Allen GR, Erdmann MV. 2012. Reef fishes of the East Indies. Perth, Australia: Tropical Reef Research.
- Chen Q, Zheng B. 1987. Systematic synopsis of Chinese fishes. Beijing: Science Press; p. 1–2.
- Kuiter R, Tonozuka T. 2001. Indonesian reef fishes (part 1 dan 2). Melbourne, Australia: Zoonetic.
- Kumar S, Stecher G, Tamura K. 2016. MEGA7: molecular evolutionary genetics analysis version 7.0 for bigger datasets. *Mol Biol Evol.* 33:1870–1874.
- Laslett D, Canbäck B. 2008. ARWEN: a program to detect tRNA genes in metazoan mitochondrial nucleotide sequences. *Bioinformatics.* 24:172–175.
- Russell BC. 1990. Nemipterid fishes of the world. Vol. 12. Rome, Italy: FAO Fisheries Synopsis; p. 1.
- Satoh TP, Miya M, Mabuchi K, Nishida M. 2016. Structure and variation of the mitochondrial genome of fishes. *BMC Genom.* 17:719–738.
- Wu R, Liu J, Niu S, Zheng J. 2017. The complete mitochondrial genome of *Scolopsis vosmeri* and phylogenetic relationship of genus *Scolopsis*. *Mitochondrial DNA A DNA Mapp Seq Anal.* 28:81–82.

2018 Mitochondrial Genome Scolopsis

ORIGINALITY REPORT

2%

SIMILARITY INDEX

0%

INTERNET SOURCES

2%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1

Kinga Székvári, Zoltán Szabolcsi, Barbara Kutasy, Géza Hegedűs, Eszter Virág. "New complete mitogenome datasets and their characterization of the European catfish (*Silurus glanis*)", Data in Brief, 2021

Publication

2%

Exclude quotes On

Exclude matches Off

Exclude bibliography On

2018 Mitochondrial Genome Scolopsis

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3
