

TROPICAL JOURNAL OF
**NATURAL
PRODUCT
RESEARCH**

TJNPR

www.tjnpr.org VOL. 11(6) 1, JULY 2017



**Tropical Journal of
NATURAL PRODUCT RESEARCH**

This journal is indexed in Open J-Gate, Crossref, Creative Commons, Index Copernicus, Google Scholar.

All published papers are DOI assigned.

Official Journal of Natural Product Research Group
University of Benin

Editorial Team

Editorial Board

Editor-in-Chief

Professor Abiodun Falodun

Professor of Pharmaceutical Chemistry
Natural product Research Group
University of Benin, Nigeria.

Email: editor.tjnpr@gmail.com; editor.tjnpr@uniben.edu
(<mailto:editor.tjnpr@gmail.com>); (editor.tjnpr@uniben.edu)
Phone: +2348073184488

Associate Editors:


- **Professor Dr. Dr. Peter Langer**, Institute of Organic Chemistry, University of Rostock, (Germany)
- **Professor Frederick O. Ekhaize**, Microbiology, University of Benin, Nigeria.
- **Professor Martins Emeje**: Professor of Drug Delivery/Nanomedicine in the Department of Pharmaceutical Technology and Raw Materials Development (PT&RMD) at the National Institute of Pharmaceutical Research and Development.

Editorial Assistant:

- **Erharuyi Osayemwenre**, Faculty of Pharmacy, University of Benin, Nigeria.

Board Members

- **Professor Ikhlas A. Khan**, National Center for Natural Product Research, Mississippi (USA)
- **Professor Nosa Egiebor**, College of Environmental Science & Forestry, State University of New York
- **Professor Samuel Qiu**, South China Botanical Gardens, Chinese Academy of Sciences (China)
- **Professor Xavier Barril**, De Fisicoquimica-Facultat De Farmacia Universitat de Barcelona, (Spain)
- **Professor Abiodun Ogundiani**, Pharmaceutical Chemistry, OAU, Ile-Ife, (Nigeria)
- **Professor Thomas Kodadek**, The Scripps Research Institute, Scripps Florida (USA)
- **Professor Anthony B Ebeigbe**, Physiology, College of Medical Sciences, University of Benin, (Nigeria)
- **Professor Eric Kl Omogbai**, Pharmacology and Toxicology, University of Benin, Nigeria
- **Professor Dr. Udo Kragl**, Institute of Organic Chemistry, University of Rostock, (Germany)
- **Professor Cyril O. Usifoh**, Faculty of Pharmacy, University of Benin, Nigeria.
- **Professor Azuka C Opara**, Clinical Pharmacy & Pharmacy Practice, University of Benin, (Nigeria).
- **Professor Ikhide G. Imumorin**, Biological Sciences, Georgia Institute of Technology, Atlanta (USA)
- **Professor Mark T. Hamann**, Medical University College, South Carolina (USA)
- **Professor Barbara Nebe**, University of Rostock, (Germany)
- **Professor Anthony I. Okoh**, University of Fort Hare, Alice (South Africa)
- **Professor Dr. M. Iqbal Choudhary**, HEJ, University of Karachi, (Pakistan)
- **Professor Omoanghe S. Isikhuemhen**, North Carolina A&T State University, (USA)
- **Professor Ezekiel Green**, University of Johannesburg, (South Africa)
- **Professor John Igoli**, Strathclyde Institute of Pharmacy and Biomedical Sciences, UK
- **Professor Peter Akah**, Pharmacology & Toxicology, University of Nigeria, Nigeria
- **Professor H.A.B. Coker**, Faculty of Pharmacy, University of Lagos (Nigeria)
- **Dr Kingsly Agho**, School of Science and Health, Western Sydney University (Australia)
- **E. Igbiosa**, Microbiology, University of Benin, Nigeria
- **Pius Fasinu**, School of Pharmacy, Campbell University (USA)
- **Professor Larry A Walker**, National Center for Natural Products Research, Mississippi, USA
- **Dr. Alireza Heidari**, Faculty of Chemistry, California South University (CSU), Irvine, California, USA
- **Professor Iyere O Onoagbe**, Faculty of Life Sciences, University of Benin, (Nigeria)
- **Professor Broderick Eribo**, Department of Biology, Howard University, Washington DC, (USA)
- **Professor Simon Gibbons**, School of Pharmacy, University College London, UK
- **Professor Masashi Mizuno**, Laboratory of Food & Nutritional Chemistry, Kobe University, Japan
- **Professor FBC Okoye**, Professor of Medicinal and Pharmaceutical Chemistry, Nnamdi Azikiwe University, Awka



(<https://tjnpr.org/index.php/home/about/submissions>)



More
Graphical
Abstracts

(<https://tjnpr.dynetng.org/index.php/home/graphicalabstra>)

Indexing & Abstracting



(<https://www.scopus.com/sourceid/21100933230>)



(<https://www.scopus.com/sourceid/21100933230>)

Keywords



Home (<https://tjnpr.org/index.php/home/index>)
/ Archives (<https://tjnpr.org/index.php/home/issue/archive>)
/ Vol. 6 No. 6 (2022): Tropical Journal of Natural Product Research

Published: 2022-07-06

Articles

Prospects for the Use of Essential Oils as Repellants and/or Insecticides (<https://tjnpr.org/index.php/home/article/view/4>)

Kirill Tkachenko, Elizaveta Varfolomeeva
831-835

PDF (<https://tjnpr.org/index.php/home/article/view/4/3>)

DOI (<https://tjnpr.org/index.php/home/article/view/4/4>)

Overview of the Potential Role of Trace Elements in COVID-19 doi.org/10.26538/tjnpr/v6i6.2

(<https://tjnpr.org/index.php/home/article/view/6>)

Sara T. Ismail, Eman A. Sulaiman
836-841

PDF (<https://tjnpr.org/index.php/home/article/view/6/5>)

DOI (<https://tjnpr.org/index.php/home/article/view/6/6>)

Croton gratissimus Burch. (Lavender croton): A Review of the Traditional Uses, Phytochemistry, Nutritional Constituents and Pharmacological Activities doi.org/10.26538/tjnpr/v6i6.3

(<https://tjnpr.org/index.php/home/article/view/9>)

Joseph O. Erhabor, Motlalepula G. Matsabisa, Omolola R. Oyenih, Ochuko L. Erukainure
842-855

PDF (<https://tjnpr.org/index.php/home/article/view/9/7>)

DOI (<https://tjnpr.org/index.php/home/article/view/9/9>)

Isolation and Characterization of Sugarcane (*Saccharum officinarum* L.) Bagasse Cellulose Hydrolyzed with Acid Variation (<https://tjnpr.org/index.php/home/article/view/10>)

Begum Fauziyah, Mohammad Yuwono, Isnaeni, Nadhifaton Nahdhia, Fatimatus Sholihah
856-862

PDF (<https://tjnpr.org/index.php/home/article/view/10/8>)

DOI (<https://tjnpr.org/index.php/home/article/view/10/10>)

Phytochemical Screening and Biological Activities of a Remedy from A-thi-sa-ra-wak Scripture as a Folkloric Diabetic Medicine doi.org/10.26538/tjnpr/v6i6.5

(<https://tjnpr.org/index.php/home/article/view/14>)

Piyapong Yupparach, Adisak Sumalee, Ampa Konsue
863-867

PDF (<https://tjnpr.org/index.php/home/article/view/14/11>)

DOI (<https://tjnpr.org/index.php/home/article/view/14/12>)

Phytochemical Screening and Antioxidant Activity of Longevity Remedy from National Thai Traditional Medicine Scripture (Formulary Special Edition) doi.org/10.26538/tjnpr/v6i6.6

(<https://tjnpr.org/index.php/home/article/view/15>)

Ampa Konsue, Ladachart Taeponsorat
868-871



(<https://tjnpr.org/index.php/home/about/submissions>)



More
Graphical
Abstracts

(<https://tjnpr.dyntng.org/index.php/home/graphicalat>)

Indexing & Abstracting



(<https://www.scopus.com/sourceid/21100933230>)



0.3 cases2022
100 percent
powered by Scopus

(<https://www.scopus.com/sourceid/21100933230>)

Keywords



PDF (<https://tjnpr.org/index.php/home/article/view/15/13>)

DOI (<https://tjnpr.org/index.php/home/article/view/15/14>)

The Effect of Red Dragon Fruit (*Hylocereus polyrhizus*) Peel Ethanol Extract on Oxidative Stress in Sprague Dawley Rats (*Rattus norvegicus*)

doi.org/10.26538/tjnpr/v6i6.7

(<https://tjnpr.org/index.php/home/article/view/16>)

Seno Santoso, Santoso Jaeri, Amallia Nuggetsiana Setyawati, Nani Maharani
872-874

PDF (<https://tjnpr.org/index.php/home/article/view/16/15>)

The Potential of Lampung Robusta Green Coffee (*Coffea Canephora*) Extract Toward T Cell Activation in ISA-brown Laying Chickens
(<https://tjnpr.org/index.php/home/article/view/17>)

Djalal Rosyidi, Lilik E. Radiati, Dahliatul Qosimah, Indah A. Amri, Dodik Prasetyo, Agri K. Anisa
875-878

PDF (<https://tjnpr.org/index.php/home/article/view/17/16>)

DOI (<https://tjnpr.org/index.php/home/article/view/17/17>)

Synthesis of Nanoytria Nanoparticles Using *Moringa oleifera* Seed as a Biological Factory and the Biocontrol Impact of the Nanoparticles on Houseflies (*Musca domestica*)

doi.org/10.26538/tjnpr/v6i6.9

(<https://tjnpr.org/index.php/home/article/view/18>)

Rasha S. Nuaman, Raghad J. Fayyad, Rasha S. Hameed, Maan A.A. Shafeeq
879-883

PDF (<https://tjnpr.org/index.php/home/article/view/18/18>)

DOI (<https://tjnpr.org/index.php/home/article/view/18/19>)

Anti-Hyperglycemic and Anti-Inflammatory Activities of *Moringa oleifera* Lam Leaves Extract

doi.org/10.26538/tjnpr/v6i6.10

(<https://tjnpr.org/index.php/home/article/view/1496>)

Jessica Novia, Denny Juvi, Riskianto Riskianto, Marcellia Sugata, Dela Rosa
884-888

PDF (<https://tjnpr.org/index.php/home/article/view/1496/2333>)

DOI (<https://tjnpr.org/index.php/home/article/view/1496/2334>)

Variations in Phytochemicals and Antioxidant Activity of *Clitoria ternatea* Flowers and Leaves

doi.org/10.26538/tjnpr/v6i6.11

(<https://tjnpr.org/index.php/home/article/view/1571>)

Danh Vu, Dien To
889-893

PDF (<https://tjnpr.org/index.php/home/article/view/1571/2401>)

DOI (<https://tjnpr.org/index.php/home/article/view/1571/2402>)

Taxonomic and Biodiversity Study of Sandfly Species in the Fez Region, Central Morocco

doi.org/10.26538/tjnpr/v6i6.12

(<https://tjnpr.org/index.php/home/article/view/20>)

Najoua Darkaoui, Abdellatif J. Idrissi, Fatima Z. Talbi, Youness El Fattouhi, Abdelkarim Taam, Nordine Nouayti, Moussa Benboubker, Abdelhakim E.O. Lalami
894-899

PDF (<https://tjnpr.org/index.php/home/article/view/20/20>)

DOI (<https://tjnpr.org/index.php/home/article/view/20/21>)

Classification of Dysmenorrhea among Students at Ubon Ratchathani Rajabhat University, Thailand According to the Māhaachortārat Scripture

doi.org/10.26538/tjnpr/v6i6.13

(<https://tjnpr.org/index.php/home/article/view/21>)

Phanida Kamuttachat, Pitchanan Thiantongin
900-905



PDF (<https://tjnpr.org/index.php/home/article/view/21/22>)

DOI (<https://tjnpr.org/index.php/home/article/view/21/23>)

Antibacterial Activity and Mechanism of Action of Crude Aqueous Extracts from Agricultural Waste against Foodborne Pathogenic Bacteria

doi.org/10.26538/tjnpr/v6i6.14

(<https://tjnpr.org/index.php/home/article/view/23>)

Nawal E. Al-Hazmi
906-909

PDF (<https://tjnpr.org/index.php/home/article/view/23/24>)

DOI (<https://tjnpr.org/index.php/home/article/view/23/25>)

Probiotics from Thai Fermented Foods Reduced Anxiety and Enhanced Neuroplasticity in a Wistar Rat Model

doi.org/10.26538/tjnpr/v6i6.15

(<https://tjnpr.org/index.php/home/article/view/24>)

Vijitra L.-In, Worachot Saengha, Thipphiya Karirat, Ampa Konsue, Eakapol Wangkahart, Teeraporn Katisart
910-914

PDF (<https://tjnpr.org/index.php/home/article/view/24/26>)

DOI (<https://tjnpr.org/index.php/home/article/view/24/27>)

Analysis of Potential Poly (ADP-Ribose) Polymerase 2 (PARP2) Inhibitor in Nyale Worm (Eunice sp.) Extract for Ovarian Cancer: An In Silico Approach

doi.org/10.26538/tjnpr/v6i6.16

(<https://tjnpr.org/index.php/home/article/view/25>)

Putu D. Arijita, Rozikin Rozikin, Gede A. Adnyana, Putu B.A. Saputra, Sabrina I. Zoraya
915-920

PDF (<https://tjnpr.org/index.php/home/article/view/25/28>)

DOI (<https://tjnpr.org/index.php/home/article/view/25/29>)

Evaluation of Antioxidant and Cholinesterase Inhibitory Activities of Various Extracts of *Cassia spectabilis*

doi.org/10.26538/tjnpr/v6i6.17

(<https://tjnpr.org/index.php/home/article/view/26>)

Suciati Suciati, Hanifa R. Putri, Wachidatur Rizqiyah, **Chrimawan Ardianto**
921-925

PDF (<https://tjnpr.org/index.php/home/article/view/26/30>)

DOI (<https://tjnpr.org/index.php/home/article/view/26/31>)

Evaluation of In vivo Antiplasmodial Activity of the Methanol Root Bark Extract and Fractions of *Bombax costatum* (Bombacaceae) in *Plasmodium berghei*-Infected Mice

doi.org/10.26538/tjnpr/v6i6.18

(<https://tjnpr.org/index.php/home/article/view/27>)

Bila H. Ali, Ilyas Mohammed, Musa A. Muhammed, Sani Y. Mohammed, Dauda Garba, Olorukooba A. Busola, Imam I. Khadijah, Mailafiya M. Manager
926-930

PDF (<https://tjnpr.org/index.php/home/article/view/27/32>)

DOI (<https://tjnpr.org/index.php/home/article/view/27/33>)

Phytochemistry and Acute Toxicity Study of Aqueous and Methanol Stem Bark Extract of *Terminalia catappa*

doi.org/10.26538/tjnpr/v6i6.19

(<https://tjnpr.org/index.php/home/article/view/29>)

Rosephine Enadeghe, Ekene V. Asoya, Joshua Iluma, Oluwadamilola Kolawole, Andrew E. Omoregie
931-937

PDF (<https://tjnpr.org/index.php/home/article/view/29/34>)

DOI (<https://tjnpr.org/index.php/home/article/view/29/35>)

Dodoneine, its Bicyclic Lactone and a Dihydroxyl-lupeol Palmitate from *Tapinanthus globiferus*

doi.org/10.26538/tjnpr/v6i6.20

(<https://tjnpr.org/index.php/home/article/view/30>)

John V. Anyam, John B. Nvau, Kachollom Thomas, Eman Santali Irvine A. Gray, John. O. Igoli
938-942

PDF (<https://tjnpr.org/index.php/home/article/view/30/36>)

DOI (<https://tjnpr.org/index.php/home/article/view/30/37>)

Effects of Aframomum melegueta K. Schum. Leaf on Monosodium Glutamate and High Fat Diet-Induced Obesity in Wistar Rat
doi.org/10.26538/tjnpr/v6i6.21

(<https://tjnpr.org/index.php/home/article/view/32>)

Adetoun E. Morakinyo, Bolajoko A. Akinpelu, Oluokun O. Oyedapo
943-950

PDF (<https://tjnpr.org/index.php/home/article/view/32/38>)

DOI (<https://tjnpr.org/index.php/home/article/view/32/39>)

Biocidal Evaluation of Ethanol Leaf Extract of Jatropha tanjorensis by Inhibition of Dehydrogenase Activity of Staphylococcus aureus and Candida albicans
doi.org/10.26538/tjnpr/v6i6.22

(<https://tjnpr.org/index.php/home/article/view/33>)

Cosmas O. Ujowundu, Arthur C. Morah, Favour N. Ujowundu, Ignatious O. Onyeocha, Kalu O. Igwe, Emeka S. Asiwe, James O. Kalu, Viola A. Onwuliri
951-956

PDF (<https://tjnpr.org/index.php/home/article/view/33/40>)

DOI (<https://tjnpr.org/index.php/home/article/view/33/41>)

Proanthocyanidin-Rich-Fraction of Vitis vinifera Seed Abrogates Convulsion Indices: Glutamatergic/ NMDA Inhibition, Enhancement of Anti-Neu N, and NRF2 Expression
doi.org/10.26538/tjnpr/v6i6.23

(<https://tjnpr.org/index.php/home/article/view/31>)

Opeyemi S. Osuntokun, Samuel O. Oyedokun, Gbola Olayiwola, Damilare A. Adekomi
957-96

PDF (<https://tjnpr.org/index.php/home/article/view/31/42>)

Phytochemical Analysis, Antibacterial and Antioxidant Activities of Leaf Extracts of Strychnos Innocua Del.
doi.org/10.26538/tjnpr/v6i6.24

(<https://tjnpr.org/index.php/home/article/view/34>)

Rachel G. Ayo, Jonathan I Achika, Olubunmi O. Bolarin-Akinwande, Dollapo Fawole
962-968

PDF (<https://tjnpr.org/index.php/home/article/view/34/44>)

DOI (<https://tjnpr.org/index.php/home/article/view/34/45>)

Gotu Kola Supplement Ameliorates Stress-Induced Liver Injury in Mice
doi.org/10.26538/tjnpr/v6i6.25

(<https://tjnpr.org/index.php/home/article/view/35>)

Anthony T. Eduviere, Mamerhi T. Enaohwo, Prosper E. Awhin, Lily O. Otomewo, Omogor Iwalio
969-973

DOI (<https://tjnpr.org/index.php/home/article/view/35/47>)

Evaluation of the Antidiarrhoeal Activity of the Methanol Pulp Extract and Fractions of Annona muricata Linn (Annonaceae)
doi.org/10.26538/tjnpr/v6i6.26

(<https://tjnpr.org/index.php/home/article/view/36>)

Igweonu A. Chinyere, Peter I. Emmanuel, Onyeali V. Chidinma, Nworu C. Sylvester
997-1003

PDF (<https://tjnpr.org/index.php/home/article/view/36/48>)

DOI (<https://tjnpr.org/index.php/home/article/view/36/49>)

Antibacterial Activity and Molecular Docking Analysis of the Stem Bark Extracts of Persea americana Mill (Lauraceae)
doi.org/10.26538/tjnpr/v6i6.27

(<https://tjnpr.org/index.php/home/article/view/37>)

Adewale O. Olatayo, Ayantade D.V. Ayansina, Samuel O. Dahunsi

PDF (<https://tjnpr.org/index.php/home/article/view/37/50>)

DOI (<https://tjnpr.org/index.php/home/article/view/37/51>)

Protective Effect of Aqueous Extract of Hypoestes rosea Against Indomethacin- Induced Gastric Ulcer in Sprague Dawley Rats
doi.org/10.26538/tjnpr/v6i6.28

(<https://tjnpr.org/index.php/home/article/view/38>)

Agala E. Egbe, Joseph I. Inyang, Okon A. Eyo
991-996

PDF (<https://tjnpr.org/index.php/home/article/view/38/52>)

DOI (<https://tjnpr.org/index.php/home/article/view/38/53>)

Effects of Thesium viride Extract and Fractions on Some Liver Biochemical Parameters in CCl4-Induced Damage in Wistar Rats
doi.org/10.26538/tjnpr/v6i6.30

(<https://tjnpr.org/index.php/home/article/view/41>)

Sani Shehu, Ezzeldin M. Abdurahman, Umar H. Danmalam, Abdullahi Mohammed, Salisu Shehu, Uwaisu Iliyasu
1004-1009

PDF (<https://tjnpr.org/index.php/home/article/view/41/59>)

DOI (<https://tjnpr.org/index.php/home/article/view/41/60>)

Assessment of Post-Thaw In-Vitro Quality of Male Wistar Rat Spermatozoa Preserved in Diluent with Natural Honey as Supplement
doi.org/10.26538/tjnpr/v6i1.21

(<https://tjnpr.org/index.php/home/article/view/207>)

Cyril A. Agbor, Christe E. Fischer, Eric A. Agaba, Williams A. Nnenna
133-137

PDF (<https://tjnpr.org/index.php/home/article/view/207/283>)

DOI (<https://tjnpr.org/index.php/home/article/view/207/284>)

ABOUT TJNPR

About the Journal (<https://tjnpr.dynetng.org/index.php/home/about>)
Aims & Scope (<https://tjnpr.dynetng.org/index.php/home/aimandscope>)
Editor Profile (<https://tjnpr.dynetng.org/index.php/home/editorprofile>)
Editorial Board (<https://tjnpr.dynetng.org/index.php/home/about/editorialTeam>)

RESOURCES

Editorial Policy & Malpractice Statement (<https://tjnpr.dynetng.org/index.php/home/ethicsandmalpractices>)
Guide for Authors (<https://tjnpr.dynetng.org/index.php/home/authorguidelines>)
Guidelines for Reviewers (<https://tjnpr.dynetng.org/index.php/home/reviewersguidelines>)
Open Access Policy (<https://tjnpr.dynetng.org/index.php/home/openaccesspolicy>)

MY ACCOUNT

Authors (<https://tjnpr.dynetng.org/index.php/home/login>)
Editors (<https://tjnpr.dynetng.org/index.php/home/login>)
Reviewers (<https://tjnpr.dynetng.org/index.php/home/login>)
Subscribers (<https://tjnpr.dynetng.org/index.php/home/login>)

ISSUES

Current (<https://tjnpr.dynetng.org/index.php/home/issue/current>)
Archives (<https://tjnpr.dynetng.org/index.php/home/issue/archive>)



**Evaluation of Antioxidant and Cholinesterase Inhibitory Activities of Various Extracts of *Cassia spectabilis***Suciati Suciati^{1,2*}, Hanifa R. Putri³, Wachidatur Rizqiyah², Chrismawan Ardianto⁴, Aty Widawaruyanti^{1,2}¹Department of Pharmaceutical Sciences, Faculty of Pharmacy, Universitas Airlangga, Surabaya 60115, East Java, Indonesia²Center for Natural Product Medicine Research and Development, Institute of Tropical Diseases, Universitas Airlangga, Surabaya, 60115, East Java, Indonesia³Master Program in Pharmaceutical Science, Faculty of Pharmacy, Universitas Airlangga, Surabaya 60115, East Java, Indonesia⁴Department of Pharmacy Practice, Faculty of Pharmacy, Universitas Airlangga, Surabaya 60115, East Java, Indonesia

ARTICLE INFO

ABSTRACT

Article history:

Received 22 May 2022

Revised 13 June 2022

Accepted 23 June 2022

Published online 02 July 2022

Copyright: © 2022 Suciati *et al.* This is an open-access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Alzheimer's disease (AD) is a progressive neurodegenerative disorder that mainly occurs in elderly people. The increasing number of people suffering from AD causes health, social and economic problems. Therapeutic strategies implemented to slow down the progress of AD are by using cholinesterase inhibitors and antioxidants. The current study aimed to investigate the antioxidant and cholinesterase inhibitory properties of the leaves and stems of *Cassia spectabilis* as well as to determine the total phenolic contents in the samples. The leaves and stems of *C. spectabilis* were extracted with 96% ethanol by the maceration method. The cholinesterase inhibitory assay was performed by using the modified Ellman's method against two cholinesterase enzymes, namely acetylcholinesterase (AChE) and butyrylcholinesterase (BChE). The antioxidant properties of the samples were evaluated using 2,2-diphenyl-1-picryl hydrazyl (DPPH) and 2,2'-azinobis-(3-ethylbenzothiazoline-6-sulfonic acid) (ABTS) methods. The total phenolic contents (TPC) were determined by a colorimetric assay using gallic acid as a reference. The results showed that the leaves and stems of *C. spectabilis* exerted significant inhibitory effects against both AChE and BChE enzymes with IC₅₀ values of 24.3 and 9.8 µg/mL for the leaves and 58.6 and 47.8 µg/mL for the stems, respectively. The extracts showed moderate antioxidant activity in both DPPH and ABTS assays with IC₅₀ ranging from 117 – 313 µg/mL. The presence of phenolic content in the samples may contribute to the antioxidant potency of the samples. The presence of non-phenolic compounds, such as alkaloids may be responsible for the cholinesterase inhibitory properties of the extracts.

Keywords: Alzheimer's disease, *Cassia spectabilis*, Cholinesterase inhibitor, Antioxidant.

Introduction

For many years plants have shown a great contribution to the treatment of many diseases. Herbal medicines have been reported to show significant effects in the treatment of Alzheimer's disease, such as the well-known *Ginkgo biloba*, *Bacopa monnieri*, and lately Chinese medicinal plant *Huperzia serrata*.^{1,2} Alzheimer's disease (AD) is a neurodegenerative disorder that commonly affects elderly people. AD produces progressive and irreversible memory and cognitive decline and is associated with progressive behavioral disturbances and restrictions in activities of daily living. One of the common features of this disease is the low level of acetylcholine in the brain. Acetylcholine (ACh) is a neurotransmitter produced in the nerve ending of the presynaptic nerve from choline and acetyl coenzyme A. ACh is hydrolyzed to choline and ethanoic acid by AChE in the postsynaptic nerve.³ Another enzyme that also plays an important role in the pathogenesis of AD is BChE which is co-regulated with AChE in the metabolism of ACh.⁴ Growing evidence showed the relation between oxidative stress and AD.

*Corresponding author. E mail: suciati@ff.unair.ac.id
Tel: +62-315933150

Citation: Suciati S, Putri HR, Rizqiyah W, Ardianto C, Widawaruyanti A. Evaluation of Antioxidant and Cholinesterase Inhibitory Activities of Various Extracts of *Cassia spectabilis*. Trop J Nat Prod Res. 2022; 6(6):921-925. doi.org/10.26538/tjnpr/v6i6.17

Official Journal of Natural Product Research Group, Faculty of Pharmacy, University of Benin, Benin City, Nigeria.

The overproduction of reactive oxygen species (ROS) or lack of antioxidants leads to oxidative stress which eventually damages the brain cells, and leads to the progression of AD to dementia.^{5,6} The presence of a toxic peptide, β-amyloid in the brain of patients with Alzheimer's disease is also contributed by oxidative stress.⁵ Therefore, strategies for the treatment of AD have involved the use of cholinesterase inhibitors such as donepezil, rivastigmine, and galantamine, as well as the use of antioxidants.^{7,8}

Cassia spectabilis (sin. *Senna spectabilis*) (DC.) H.S.Irwin & Barneby family Fabaceae is a flowering plant that can be found in tropical and sub-tropical regions.⁹ The flowers are bright yellow therefore it is usually used as ornamental plants in some regions. *C. spectabilis* is also known for its medicinal purposes in the folk medicines of several countries such as Indonesia, Brazil, and Thailand. The leaves are commonly used as laxatives and purgatives. Other traditional uses are for the treatment of skin diseases, edema, as well as poisoning, and protozoic infection of the gut.¹⁰ A phytochemical study of the plant revealed the presence of alkaloids, terpenoids, flavonoids, anthraquinones, and steroids from various parts of *C. spectabilis* including the leaves, flowers, fruits, seeds, stems, and roots.¹⁰⁻¹² Pharmacological studies of the plant have shown promising antimicrobial¹³⁻¹⁶ anticonvulsant,^{17,18} antinociceptive, and anti-inflammatory activities.¹⁹ In our previous study, we screened the potency of the leaves of several *Cassia* species as cholinesterase inhibitors. It was discovered that *C. spectabilis* showed the best potency among other *Cassia* species tested.²⁰ Leaves and stems are the part of this plant that show in abundance the presence of various metabolites. Therefore, in the current study, the potency of the leaves and the stems of *C. spectabilis* as cholinesterase inhibitors was

compared as well as to determine the antioxidant and quantify the phenolic contents in the extracts.

Materials and Methods

Materials

The reagents used for cholinesterase assays were acetylcholinesterase from electric eel (AChE type VI-S), acetylthiocholine iodide (ATCI), horse-serum butyrylcholinesterase (BChE), butyrylthiocholine iodide (BTCl), 5,5'-dithiobis[2-nitrobenzoic acid] (DTNB), bovine serum albumin (BSA), tris buffer, and galantamine. The chemicals used for antioxidant assays were 2,2-Diphenyl-1-picrylhydrazyl (DPPH), 2,2'-Azino-bis(3-ethylbenzothiazoline-6-sulfonic acid) (ABTS), and potassium persulfate. Folin-Ciocalteu's phenol reagent, sodium carbonate, and gallic acid were used for the determination of total phenolics. All reagents were purchased from Sigma-Aldrich.

Sample collection

The leaves of *Cassia spectabilis* were collected from Purwodadi Botanic Garden, East Java, Indonesia on March 9th, 2019. The voucher specimen (PWD 02) was stored at the Faculty of Pharmacy, Universitas Airlangga. The plant was identified by Purwodadi Botanic Garden, Indonesian Institute of Sciences with identification letter number: 0371/IPH.06/HM/III/2019.

Preparation of extracts

Freshly collected leaves and stems of *Cassia spectabilis* were air-dried at room temperature for approximately seven days and then pulverized. Four hundred grams of the powdered leaves and stems were each extracted with 2 L of 96% ethanol. The samples were soaked in the solvent for 24 hours, followed by vacuum filtration. Then residues were each re-extracted with 1 L of ethanol using the same procedure. This process was repeated twice. All collected filtrates were then concentrated in a rotary evaporator at 40°C to yield crude ethanolic extracts of the leaves (27.5 g) and stems (9.6 g).

Anticholinesterase Assay

The assay was carried out according to the modified Ellman's method.²¹⁻²³ The extracts were dissolved in methanol at a concentration of 10 mg/mL and were then diluted with water to obtain serial concentrations of samples containing not more than 10% of methanol. The final test concentrations in the wells were: 300, 200, 100, 30, 20, 10, 3, 2, 1, 0.3 and 0.1 µg/mL. Twenty-five microliters of sample solutions were added to a 96-well microplate, followed by the addition of 25 µL substrates 1.5 mM ATCI or 1.5 mM BTCl, 125 µL of 3 mM DTNB, and 50 µL Tris buffer, and finally 25 µL of 0.22 U/mL AChE or BChE. The solutions were placed in a microplate reader (Thermo Scientific Multiskan FC) and shaken for 30 s before measurement. The absorbances were measured at 405 nm every 5 s for 2 mins. Experiments were carried out in triplicates. Galantamine was used as a positive control, and 10% methanol was used as a negative control. The percentage of inhibition was then calculated as follows:

$$\% \text{Inhibition} = \frac{(\text{Mean velocity of control} - \text{Mean velocity of sample})}{\text{Mean velocity of control}} \times 100$$

DPPH radical scavenging assay

The DPPH assay was performed according to the modified method of Herald *et al.* (2012) and Lee *et al.*^{24,25} The extracts were dissolved with methanol to make a series of concentrations of 5 – 500 µg/mL. Gallic acid was employed as a standard. The samples (100 µL) were added to 96 microwell plates and mixed with 0.25 mM DPPH reagent (100 µL). The DPPH reagent (100 µL) was mixed with methanol (100 µL) as a control, while methanol (200 µL) was used as a blank. The mixtures were then incubated in the dark at room temperature for 30 mins. The solutions were shaken for 30 s in a microplate reader (Thermo Scientific Multiskan FC). The absorbances were then recorded at 517 nm. The DPPH scavenging effect was calculated using the following formula.

$$\text{DPPH Radical Scavenging activity (\%)} = \frac{(\text{abs control} - \text{abs sample})}{\text{abs control}} \times 100$$

ABTS radical scavenging assay

The ABTS assay was carried out based on Lee *et al.* with some modifications.²⁵ ABTS solution (5 mL, 7 mM) was mixed with potassium persulfate (88 µL, 140 nM), and the mixture was kept in the dark at room temperature for 16 h to produce ABTS radical. Solution of samples at the concentration range 2.5 – 250 µg/mL was prepared in methanol. The samples (100 µL) were then mixed with 100 µL of ABTS in a 96-well microplate followed by incubation for 6 mins in the dark at room temperature. The absorbance was measured at 734 nm in a microplate reader and the plates were shaken for 30 s before reading. Gallic acid was used as standard. Experiments were done in triplicate. The ABTS radical scavenging activity was calculated using the equation as follows.

$$\text{ABTS Radical Scavenging activity (\%)} = \frac{(\text{abs control} - \text{abs sample})}{\text{abs control}} \times 100$$

Determination of total phenolic content (TPC)

The TPC of the extracts was determined according to the method by Herald *et al.* with slight modification.²⁴ Briefly, twenty-five microliters of serial concentrations of gallic acid (25 – 500 µg/mL) or samples (1000 µg/mL) were added to a 96-well microplate, followed by the addition of water (75 µL) and Folin-Ciocalteu's phenol reagent (25 µL). The solutions were incubated for 6 mins at room temperature. Then 100 µL of Na₂CO₃ solution (75 g/L) was added to each well, followed by incubation for 90 mins in the dark at room temperature. The mixtures were shaken for 30 s before measurement of the absorbance at 765 nm in a microplate reader (Thermo Scientific Multiskan FC). The TPC of samples was expressed as milligrams of gallic acid equivalents (GAE) per gram extract.

Statistical analysis

The 50% inhibitory concentration (IC₅₀) values were determined using GraphPad Prism 8.0 software by plotting log concentrations as axis and % inhibition as ordinate for the cholinesterase inhibitory assays, and concentration of extracts as axis and % scavenging DPPH or ABTS for the antioxidant assays. Results were expressed as mean ± standard error of the mean (SEM) of three experiments.

Results and Discussion




Cholinesterase Inhibitory Activities

Despite the multifactorial nature of Alzheimer's disease, the most current therapeutic approach used is based on the cholinergic hypothesis. The use of cholinesterase inhibitors becomes the main therapeutic agent for this disease. Cholinergic neurotransmission is terminated by two cholinesterase enzymes, namely acetylcholinesterase (AChE) and butyrylcholinesterase (BChE), that play an important role in the hydrolysis of acetylcholine (ACh).²⁶ Several plants from the genus *Cassia* have been reported as cholinesterase inhibitors such as *C. obtusifolia* inhibit AChE, BChE, and BACE1^{27,28} and *Cassia alata* leave which showed inhibition against AChE.²⁹ In the present study, the *in vitro* cholinesterase inhibitory activity of the ethanolic extracts of the leaves and stems of *C. spectabilis* was evaluated based on the modified Ellman's method. The results show that all samples tested were able to inhibit both AChE and BChE enzymes in a concentration-dependent manner (Figures 1 and 2). As can be seen in Table 1 and Figures 1 and 2 the ethanolic extracts of the leaves and the stems demonstrated stronger inhibition against BChE compared to AChE. Inhibition of both AChE and BChE enzymes will prevent rapid hydrolysis of acetylcholine so that the amount of ACh will increase in the brain. BChE is also associated with neuritic plaques and the accumulation of fibrillar Aβ plaques. Therefore, inhibition against BChE will serve two roles, increasing the amount of acetylcholine as well as inhibiting fibrillar Aβ deposition.⁴

Table 1: Cholinesterase inhibitory activity of *C. spectabilis* extracts

Samples	IC ₅₀ (µg/mL) ^a	
	AChE	BChE
Leaves extract	24.3 ± 0.4	9.8 ± 0.8
Stems extract	58.6 ± 0.4	47.8 ± 1.2
Galantamine	0.4 ± 0.1	2.2 ± 0.3

^aData presented as mean ± SEM of three experiments, each done in triplicates.

**Figure 1:** Concentration-dependent response of *C. spectabilis* leaves (a) and stems (b) extracts against AChE**Figure 2:** Concentration-dependent response of *C. spectabilis* leaves (a) and stems (b) extracts against BChE, each value is expressed as means ± SEM ($n = 3$)**Figure 3:** Scavenging effect of *C. spectabilis* extracts on DPPH free radicals, each value is expressed as mean ± SEM ($n = 3$)

The leaves extract gave IC₅₀ values of 24.3 and 9.8 µg/mL against AChE and BChE, respectively compared to the stem extracts with IC₅₀ values of 58.6 and 47.8 µg/mL. These results suggested that the leaves extract of *C. spectabilis* has better potency compared to the extract of the stems. Selegato *et al.* (2017) have summarized secondary metabolites reported from various parts of *C. spectabilis*.¹¹ It was found that the leaves contain more alkaloids compared to other parts of this plant, and there are no alkaloids reported from the stem. Numerous classes of compounds have been reported as cholinesterase inhibitors, however, the majority of these are alkaloids.^{2,30} Alkaloids have shown promising anticholinesterase activities, even the current drug uses for AD therapy are alkaloids, galantamine, rivastigmine, and donepezil. The higher cholinesterase inhibitory activities of the leaves of *C. spectabilis* compared to the stems are possibly due to the presence of alkaloids in the extracts. Non-alkaloidal compounds present in the stem may also contribute to the anticholinesterase activity of the extract. This finding is in accordance with our previous study that based on the LC-MS/MS analysis the leaves of *C. spectabilis* contain alkaloid cassine, spectraline, and 3-*O*-acetylspectraline.²⁰ Further study is needed to investigate the chemical composition of the stems of *C. spectabilis* since there is a limited report.

Antioxidant Activity

In this study, the radical-scavenging activities of the ethanolic and aqueous extracts of *C. spectabilis* leaves were evaluated using the DPPH and ABTS assays. In the DPPH assay, the samples with antioxidant compounds provide a hydrogen atom, which can react with the stable radical DPPH to form a yellow-colored non-radical diphenylpicrylhydrazine. The degree of discoloration indicates the radical-scavenging potential of the sample.³¹ The principle of ABTS assay is similar to that of DPPH assay, in which the antioxidant acts as a hydrogen donor to form a non-radical ABTS. The reduction of a dark-bluish color of ABTS radical can be monitored by spectrophotometer.³² The results of the DPPH and ABTS scavenging activity of the extracts are shown in Table 2, Figures 3 and 4. The ethanolic extracts of *C. spectabilis* exhibited concentration-dependent antiradical activities in both DPPH and ABTS assays. The extracts showed lower antioxidant potency compared to the standard antioxidant gallic acid with inhibitory concentration 50% (IC₅₀) values ranging from 233.2-313.8 µg/mL and 117.2-214.0 µg/mL in the DPPH and ABTS assays, respectively.

Total Phenolic Content (TPC)

The total phenolic contents in the leaves and stems extracts were evaluated using a Folin-Ciocalteu reagent with gallic acid as a standard. The calculation was based on the standard curve of equation ($y = 0.0059x - 0.1105$, $R^2 = 0.998$). The results as can be seen in Table 3 showed that the leaves and stems of *C. spectabilis* contain a slightly low amount of phenolics, which may contribute to the moderate antioxidant activities of the extracts.

Phenolic compounds have been reported to play a significant role in the antioxidant activities of plants. The antioxidant potential of the phenolic compound is predominantly due to its redox capability so that it can absorb and neutralize free radicals, decompose peroxide, and quench singlet or triplet oxygen.³³ Studies revealed that the antioxidant capacity of the phenolic compounds depends on the number and arrangement of the hydroxyl groups in this compound. The relation between phenolic contents and the antioxidant activity of several *Cassia* species has been documented.³⁴⁻³⁸ The antioxidant potency, as well as phenolic contents of seven *Cassia* species, have been reported. *C. glauca* was reported to show the strongest antioxidant capacity compared to the other six *Cassia* species, which was related to its high content of phenolic compounds.³⁴ Several phenolic compounds such as anthraquinones and flavonoids have been reported from *C. spectabilis*.¹¹ The results from our study are in accordance with that reported in the previous study. Jothy *et al.* investigated the antioxidant potency of the leaves of *C. spectabilis* which showed moderate antioxidant activity with an IC₅₀ value of 30.178 ± 0.129 mg/mL in the DPPH assay. The antioxidant activity of the stems of *C. spectabilis* has not been reported.³³

Table 2: Antioxidant activity of *C. spectabilis* extracts

Samples	IC ₅₀ (µg/mL) ^a	
	DPPH	ABTS
Leaves extract	313.8 ± 5.9	214.0 ± 1.0
Stems extract	233.2 ± 2.2	117.2 ± 0.4
Gallic acid	2.76 ± 0.02	0.97 ± 0.03

^a Data presented as mean ± SEM of three experiments, each done in triplicates.




Figure 4: Scavenging effect of *C. spectabilis* extracts on DPPH free radicals, each value is expressed as mean ± SEM ($n = 3$)

Table 3: Extract yield and total phenolic content (TPC) of *C. spectabilis* extracts

Samples	Extract Yield (%)	TPC (mg GAE/g extract) ^a
Leaves extract	6.9	29.9 ± 0.4
Stems extract	2.4	25.2 ± 0.2

^aData presented as mean ± SEM of three experiments, each done in triplicate.

Conclusion

The leaves and stems extracts of *Cassia spectabilis* exhibited significant cholinesterase inhibitory activity against AChE and BChE. Both extracts demonstrated moderate antioxidant activity. The presence of non-phenolic compounds, such as alkaloids in the leaves may be responsible for the higher potency of the extract as a cholinesterase inhibitor.

Conflict of Interest

The authors declare no conflict of interest.

Authors' Declaration

The authors hereby declare that the work presented in this article is original and that any liability for claims relating to the content of this article will be borne by them.

Acknowledgements

Authors acknowledge Purwodadi Botanic Garden Indonesian Institute of Sciences for sample collection, Directorate of Research and Community Services, the Ministry of Research and Technology/National Research and Innovation Agency Republic of Indonesia for research grant PDUPT 2021 contract number 401/UN3.1.5/PT/2021.

References

- Akram M and Nawaz A. Effects of medicinal plants on Alzheimer's disease and memory deficits. *Neural Regen Res.* 2017; 12(4):660-670.
- dos Santos TC, Gomes TM, Pinto BAS, Camara AL, Paes AMA. Naturally occurring acetylcholinesterase inhibitors and their potential use for Alzheimer's disease therapy. *Front Pharmacol.* 2018; 9:1192.
- Dev K and Maurya R. Marine-derived antialzheimer's agents of promise. In: Brahmachari G (ed). *Neuroprotective Natural Products, Clinical Aspects and Mode of Action.* Weinheim: Wiley-VCH; 2017; 153-184p.
- Darvesh S. Butyrylcholinesterase as a diagnostic and therapeutic target for Alzheimer's disease. *Curr Alzheimer Res.* 2016; 13(10):1173-1177.
- Cassidy L, Fernandez F, Johnson JB, Naiker M, Owoola AG, Broszczak DA. Oxidative stress in Alzheimer's disease: A review on emergent natural polyphenolic therapeutics. *Compl Ther Med.* 2020; 49:102294.
- Sinyor B, Mineo J, Ochner C. Alzheimer's disease, inflammation, and the role of antioxidants. *J Alzheimers Dis Rep.* 2020; 4(1):175-183.
- Sharma K. Cholinesterase inhibitors as Alzheimer's therapeutics. *Mol Med Rep.* 2019; 20(2):1479-1487.
- Juszczak G, Mikulska J, Kasperek K, Pietrzak D, Mrozek W, Herbet M. Chronic stress and oxidative stress as common factors of the pathogenesis of depression and Alzheimer's disease: The role of antioxidants in prevention and treatment. *Antioxid (Basel).* 2021; 10(9):1439.
- Torey A, Sasidharan S, Yeng C, Latha LY. Standardization of *Cassia spectabilis* with respect to authenticity, assay and chemical constituents analysis. *Mol.* 2010; 15(5):3411-3420.
- Jothy SL, Torey A, Darah I, Choong YS, Saravanan D, Chen Y, Latha LY, Deivanai S, Sasidharan S. *Cassia spectabilis* (DC) Irwin et Barn: A promising traditional herb in health improvement. *Mol.* 2012; 17(9):10292-10305.
- Selegato DM, Monteiro AF, Vieira NC, Cardoso P, Pavani VD, Bolzani VS, Castro-Gamboa I. Update: biological and chemical aspects of *Senna spectabilis*. *J Braz Chem Soc.* 2017; 28(3):415-426.
- Franca MGA, Cavalheiro AJ, Silva MG. A comprehensive LC-DAD-QTOF-MS method for dereplication of bioactive compounds in *Senna* extracts. *Rev Bras Farmacogn.* 2021; 31:32-39.
- Sangetha S, Zuraini Z, Sasidharan S, Suryani S. Antibacterial, antifungal and cytotoxic activities of *Cassia spectabilis*. *Asian J Pharm Clin Res.* 2008; 1:17-20.
- Sangetha S, Zuraini Z, Suryani S, Sasidharan S. In situ TEM and SEM studies on the antimicrobial activity and prevention of *Candida albicans* biofilm by *Cassia spectabilis* extract. *Micron.* 2009; 40(4):439-443.
- Krishnan N, Ramanathan S, Sasidharan S, Murugaiyah V, Mansor SM. Antimicrobial activity evaluation of *Cassia spectabilis* leaf extracts. *Int J Pharmacol.* 2010; 6(4):506-510.
- Torey A and Sasidharan S. Anti-candida albicans biofilm activity by *Cassia spectabilis* standardized methanol extract: An ultrastructural study. *Eur Rev Med Pharmacol Sci.* 2011; 15(8):875-882.
- Bum EN, Nkantchoua GN, Njikam N, Taiwe GS, Ngoupaye GT, Pelanken MM, Nanga, Maidawa F, Rakotonirina A, Rakotonirina SV. Anticonvulsant and sedative activity of leaves of *Senna spectabilis* in mice. *Int J Pharmacol.* 2010; 6(2):123-128.
- de Oliveira Silva F, de Vasconcelos Silva MG, Feng D, de Freitas RM. Evaluation of central nervous system effects of iso-6-cassine isolated from *Senna spectabilis* var. *excelsa* (Schrad) in mice. *Fitoter.* 2011; 82(2):255-259.

19. da Silva KA, Manjavachi MN, Paszcuk AF, Pivatto M, Viegas C Jr, Bolzani VS, Calixto JB. Plant derived alkaloid (–)-cassine induces anti-inflammatory and anti-hyperalgesics effects in both acute and chronic inflammatory and neuropathic pain models. *Neuropharmacol.* 2012; 62(2):967-977.
20. Suciati S, Laili ER, Poerwantoro D, Hapsari AP, Gifanda LZ, Rabgay K, Ekasari W, Ingkaninan K. Evaluation of cholinesterase inhibitory activity of six Indonesian *Cassia* species. *J Res Pharm.* 2020; 24(4):472-478.
21. Ellman GL, Courtney KD, Andres V Jr, Featherstone RM. A new and rapid colorimetric determination of acetylcholinesterase activity. *Biochem Pharmacol.* 1961; 7(2):88-95.
22. Suciati S, Poerwantoro D, Widyawaruyanti A, Ingkaninan K. Acetylcholinesterase inhibitory activity of extract and fractions from the root of *Rauvolfia serpentina* (L.) Bth.ex Kurz. *J Basic Clin Physiol Pharmacol.* 2021; 32(4):313-317.
23. Aristyawan AD, Setyaningtyas VF, Wahyuni TS, Widyawaruyanti A, Ingkaninan K, Suciati S. In vitro acetylcholinesterase inhibitory activities of fractions and iso-agelasine C isolated from the marine sponge *Agelas nakamurai*. *J Res Pharm.* 2022; 26(2):279-286.
24. Herald TJ, Gadgil P, Tilley M. High-throughput micro plate assays for screening flavonoid content and DPPH-scavenging activity in sorghum bran and flour. *J Sci Food Agric.* 2012; 92(11):2326-2331.
25. Lee KJ, Oh YC, Cho WK, Ma JY. Antioxidant and anti-inflammatory activity determination of one hundred kinds of pure chemical compounds using offline and online screening HPLC assay. *Evid-Based Compl Altern Med.* 2015; 2015:165457.
26. Stanciu GD, Luca A, Rusu RN, Bild V, Beschea Chiriac SI, Solcan C, Bild W, Ababei DC. Alzheimer's disease pharmacotherapy in relation to cholinergic system involvement. *Biomol.* 2019; 10(1):40.
27. Jung HA, Ali MY, Jung HJ, Jeong HO, Chung HY, Choi JS. Inhibitory activities of major anthraquinones and other constituents from *Cassia obtusifolia* against β -secretase and cholinesterases. *J Ethnopharmacol.* 2016; 191:152-160.
28. Shrestha S, Seong SH, Paudel P, Jung HA, Choi JS. Structure related inhibition of enzyme systems in cholinesterases and BACE1 *in vitro* by naturally occurring naphthopyrone and its glycosides isolated from *Cassia obtusifolia*. *Mol.* 2017; 23(1):69-86.
29. Feitosa CM, Freitas RM, Luz NN, Bezerra MZ, Trevisan MT. Acetylcholinesterase inhibition by some promising Brazilian medicinal plants. *Braz J Biol.* 2011; 71(3):783-789.
30. Tamfu AN, Kucukaydin S, Yeskalyeva B, Ozturk M, Dinica RM. Non-alkaloid cholinesterase inhibitory compounds from natural sources. *Mol.* 2021; 26(18):5582.
31. Molyneux P. The use of the stable free radical diphenylpicrylhydrazyl (dpph) for estimating antioxidant activity. *Songklanakarin J Sci Technol.* 2004; 26(2):211-219.
32. Re R, Pellegrini N, Proteggente A, Pannala A, Yang M, Rice-Evans C. Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Rad Biol Med.* 1999; 26(9-10):1231-1237.
33. Jothy SL, Aziz A, Chen Y, Sasidharan S. Antioxidant activity and hepatoprotective potential of *Polyalthia longifolia* and *Cassia spectabilis* leaves against paracetamol-induced liver injury. *Evid-Based Compl Altern Med.* 2012; 2012:561284.
34. El-Hashash MM, Abdel-Gawad MM, El-Sayed MM, Sabry WA, Abdel-Hameed el-SS, Abdel-Lateef Eel-S. Antioxidant properties of methanolic extracts of the leaves of seven Egyptian *Cassia* species. *Acta Pharm.* 2010; 60(3):361-367.
35. Irshad Md, Zafaryab Md., Singh M, Rizvi MM. Comparative analysis of the antioxidant activity of *Cassia fistula* extracts. *Int J Med Chem.* 2012; 2012:157125.
36. Ahmed SI, Hayat MQ, Tahir M, Mansoor Q, Ismail M, Keck K, Bates RB. Pharmacologically active flavonoids from the anticancer, antioxidant and antimicrobial extracts of *Cassia angustifolia* Vahl. *BMC Compl Altern Med.* 2016; 16(1):460.
37. Kolar FR, Gogi CL, Khudavand MM, Choudhari MS, Patil SB. Phytochemical and antioxidant properties of some *Cassia* species. *Nat Prod Res.* 2017; 32(11):1324-1328.
38. Mehta JP, Parmar PH, Vadia SH, Patel MK, Tripathi CB. *In-vitro* antioxidant and *in-vivo* anti-inflammatory activities of aerial parts of *Cassia* species. *Arab J Chem.* 2017; 10(supp.2):S1654-S1662.


IDP Adalah Co-Owner IELTS

Tropical Journal of Natural Product Research

COUNTRY

Nigeria

 Universities and research institutions in Nigeria


 Media Ranking in Nigeria

PUBLISHER

Faculty of Pharmacy,
University of Benin

H-INDEX

5



SUBJECT AREA AND CATEGORY

Biochemistry, Genetics and Molecular Biology

Biochemistry
Molecular Medicine

Chemistry

Analytical Chemistry

Medicine

Complementary and Alternative Medicine

Pharmacology, Toxicology and Pharmaceutics

Drug Discovery
Pharmaceutical Science
Pharmacology

PUBLICATION TYPE

Journals

ISSN

26160684, 26160692

COVERAGE

2017-2021

INFORMATION

[Homepage](#)

[How to publish in this journal](#)

[Contact](#)



SCOPE

The Tropical Journal of Natural Product Research is open access, peer-reviewed international journal aimed at making important contributions in the field of Natural Product Research, Pharmaceutical and Natural Sciences. The journal covers all aspects of Pharmaceutical research, chemistry and biochemistry of naturally occurring compounds, ethnomedicine,

welcomed. Although the journal focuses mainly on original research articles, timely, concise and focused reviews on recent progress in active areas of natural Product is also encouraged.


Join the conversation about this journal

Quartiles

FIND SIMILAR JOURNALS

options

1 Pharmacognosy Journal IND 76% similarity	2 Journal of Traditional and Complementary Medicine NLD 76% similarity	3 Pharmacologyonline ITA 75% similarity	4 Pharmacognosy Research IND 73% similarity	5 Oriental Pharmacy and Experimental Medicine USA 72% similarity
--	--	---	---	--




Tropical Journal of Natural Product Research
Analytical Chemistry
Q4 best quartile
SJIR 2021 0.13
powered by scimagojr.com

Show this widget in your own website
Just copy the code below and paste within your html code:

SCImago Graphica

Explore, visually communicate and make sense of data with our **new data visualization tool.**



Metrics based on Scopus® data as of April 2022